Labetalol

****This monograph is under review****

Neonatal

FORM Ampoule containing 100mg in 20ml

INDICATION Hypertension

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 1 month	500 microgram/kg/hr up to a maximum of 4mg/kg/hr	Continuous infusion	IV
1 – 6 months	500 microgram/kg/hr -3 mg/kg/hr	Continuous infusion	IV

PRESCRIPTION OF CONTINUOUS INFUSION

50mg/kg of labetalol in 50ml of infusion fluid

This gives:-

- 500 micrograms/kg/hr at 0.5ml/hr
- 1mg/kg/hr at 1ml/hr

Seek advice for prescription in babies greater than 5kg

RECONSTITUTION Already in solution

DILUTION 10 x wt (kg) is the number of ml of labetalol 100mg/20ml to be diluted up to 50ml total with infusion fluid (equivalent to 50mg/kg in 50ml)

METHOD OF ADMINISTRATION By continuous intravenous infusion, flow rate adjusted according to the baby's response (see prescription section for details).

Seek advice for administration guidance in babies greater than 5kg

COMPATIBILITY

Solution compatibility	Sodium Choride and Glucose infusion, Glucose 5%, Sodium Chloride 0.9%
Solution incompatibility	Sodium bicarbonate 4.2%, furosemide
IV Line compatibility	Dobutamine, dopamine, insulin, linezolid, midazolam, morphine, noradrenaline, Rocuronium, sodium nitroprusside
IV Line incompatibility	Heparin, sodium bicarbonate

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

See Summary of Product Characteristics and most recent edition of BNF for Children

.

FURTHER INFORMATION

PH 3.5 – 4.2

LICENSED STATUS Off label use in children and neonates

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:
Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo pharm group	Checked by	Peter Mulholland
Date prepared	May 2016	Date updated	September 2019
Updated by	Peter Mulholland	Review Date	September 2022
Extension Agreed By	WoS NPG	Review Date	March 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department