Vitamin K - Phytomenadione

FORM

Phytomenadione 2mg/0.2ml– solution for IM or IV injection.
Caution: adult strength injection also available

INDICATION

1. Prevention of haemorrhagic disease of the newborn. See protocol for full details
2. Use of intravenous Vitamin K where clinically indicated

DOSE RANGE

1. Prevention of haemorrhagic disease of the newborn. (Patient Group Direction also available)

AGE	DOSE	FREQUENCY	ROUTE
Preterm (< 36weeks)	0.5 mg (0.05 ml)	Single dose *	IM/(IV)**
Term (≥ 36 weeks)	1 mg (0.1 ml)	Single dose *	IM/(IV)**

* **GIVE SOON AFTER BIRTH. Use Phytomenadione 2 mg/0.2 ml solution for injection**

** **INTRAVENOUS** administration can be considered in very low birth weight infants or where IM route is not possible. Where IV route used, subsequent PO doses should be given at day 7 and day 28 of life

2. Use of intravenous Vitamin K where clinically indicated

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6months	1mg	As required depending on clinical picture and coagulation status	IV over 5 to 10minutes

RECONSTITUTION

Not required

DILUTION

For IM use – no dilution required.

For IV use - dilute with Glucose 5%. The infusion must be freshly prepared prior to use.

Phytomenadione 2mg/0.2ml	0.2ml
Glucose 5%	up to 2ml total volume

Gives 1mg/ml solution.

METHOD OF ADMINISTRATION

The line should be flushed with IV glucose 5% before and after administration.
Slow IV bolus over 3-5 minutes.

Neonatal

COMPATIBILITY

Solution compatibility	Glucose 5%; sodium chloride 0.9%
Solution incompatibility	Lipid
IV Line compatibility	Do not run with any other medicines.
IV Line incompatibility	Do not run with any other medicines.

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Care required over selection of the correct product as administration of the wrong product can lead to anaphylactic shock.
- Parenteral administration to premature babies weighing <2.5kg may increase the risk for the development of kernicterus.
- If contents of ampoule are turbid or separated, the ampoule must be discarded.

LICENSED STATUS

Phytomenadione 2mg/0.2ml is licensed for:

- Prophylaxis of haemorrhagic disease (VKDB) of the newborn in healthy neonates > or = 36 weeks gestation *(IM and oral administration)*
- Prophylaxis of haemorrhagic disease of the newborn <36 weeks or term neonates at special risk *(IM and IV administration)*. For this indication, the dose above is off-label but widely recognised practice. See guideline below.
- Haemorrhagic disease of the newborn *(IV administration)*
- Neonatal hypoprothrombinaemia *(IV administration, off label use)*

APPLICABLE POLICIES

Vitamin K Prophylaxis for Neonates, Wos MCN (695) | NHSGGC

Document Number:	002	Supersedes:	001
Prepared/Checked by:	Fraces Lowrie/Susan Kafka/Andrew Stockton	Approved by	WoS Neonatal Pharmacist Group
Date prepared	September 2026	Review date	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department