Biotin

**** This monograph is under review ****

FORM Vial containing 5mg in 1ml

INDICATION

1) Seizures (under specialist neurology advice)
2) Metabolic disorders (under specialist metabolic advice)

DOSE RANGE

1) Seizures (under specialist neurology advice)

AGE	DOSE	FREQUENCY	ROUTE
0 – 6months	10mg	Once daily (as per neurology advice)	IV

2) Metabolic disorders (under specialist metabolic advice)

AGE	DOSE	FREQUENCY	ROUTE
0 – 6months	5 – 10mg	6 hourly – 24 hourly As per metabolic advice (max 20mg daily)	IV

RECONSTITUTION Already in solution

DILUTION Not required

METHOD OF ADMINISTRATION Slow intravenous bolus over 2 minutes

(note the IM preparation can be given intravenously)

COMPATIBILITY No compatibility information available

LICENSED STATUS Unlicensed preparation

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared/Checked by:	Anisa Patel/Susan Kafka	Approved by	WoS NPG
Date prepared	Oct 2020	Updated date	October 2023
Extension agreed	WoS NPG	Review date	August 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department