


Exercise 1. Express the following UML class diagram in FOL:




Exercise 2. Consider the above UML class diagram and the following (partial) instantiation:

<i>Customer</i>	<i>BCustomers</i>	<i>Services</i>	<i>Provider</i>	<i>provides</i>	<i>contacts/cost</i>																																								
<table><tr><td>c1</td></tr><tr><td>c2</td></tr><tr><td>c3</td></tr><tr><td>c4</td></tr></table>	c1	c2	c3	c4	<table><tr><td>b1</td></tr><tr><td>b2</td></tr><tr><td>b3</td></tr></table>	b1	b2	b3	<table><tr><td>s1</td></tr><tr><td>s2</td></tr><tr><td>s3</td></tr></table>	s1	s2	s3	<table><tr><td>p1</td></tr><tr><td>p2</td></tr></table>	p1	p2	<table><tr><td>p1</td><td>s1</td></tr><tr><td>p1</td><td>s2</td></tr><tr><td>p1</td><td>s3</td></tr><tr><td>p2</td><td>s2</td></tr></table>	p1	s1	p1	s2	p1	s3	p2	s2	<table><tr><td>c1</td><td>s1</td><td>p1</td><td>90.0</td></tr><tr><td>c1</td><td>s2</td><td>p1</td><td>80.0</td></tr><tr><td>c1</td><td>s3</td><td>p1</td><td>50.0</td></tr><tr><td>b2</td><td>s1</td><td>p2</td><td>170.0</td></tr><tr><td>b2</td><td>s2</td><td>p2</td><td>100.0</td></tr></table>	c1	s1	p1	90.0	c1	s2	p1	80.0	c1	s3	p1	50.0	b2	s1	p2	170.0	b2	s2	p2	100.0
c1																																													
c2																																													
c3																																													
c4																																													
b1																																													
b2																																													
b3																																													
s1																																													
s2																																													
s3																																													
p1																																													
p2																																													
p1	s1																																												
p1	s2																																												
p1	s3																																												
p2	s2																																												
c1	s1	p1	90.0																																										
c1	s2	p1	80.0																																										
c1	s3	p1	50.0																																										
b2	s1	p2	170.0																																										
b2	s2	p2	100.0																																										

1. Check whether the above instantiation, once completed, is correct, and explain why it is or it is not.
2. Express in FOL the following queries and evaluate them over the completed instantiation:
 - (a) Check that, for every provider x and service y involved in a contract, provider x does provide service y .
 - (b) Return those customers that have contracts only for services provided by $p1$.
 - (c) Return those customers that have a contract for all services.

Exercise 3. Model check the Mu-Calculus formula $\nu X. \mu Y. ((a \wedge [next]X) \vee (b \wedge \langle next \rangle Y))$ and the CTL formula $AF(EG(a \supset AXEX\neg a))$ (showing its translation in Mu-Calculus) against the following transition system:




Exercise 4. Check whether the Hoare triple below is correct, by using $(x \geq 0 \wedge y \geq 0 \wedge x + y = 23)$ as invariant:

$$\{x = 23 \wedge y = 0\} \text{ while}(x > 0) \text{ do } (x = x - 1; y := y + 1) \{y = 23\}$$


Exercise 5. Check whether the following FOL formula is valid, by using tableaux:

$$(\forall x. P(x) \wedge \forall x. Q(x)) \equiv \forall x. (P(x) \wedge Q(x))$$

Exercise 6 (optional). Model check the LTL formula $\Diamond \Box \neg a$ against the following transition system, by considering that the Büchi automaton for $\neg(\Diamond \Box \neg a)$ is the one below:



Exercise 1. Express the following UML class diagram in FOL:



$C(x) \quad BC(x) \quad P(x) \quad S(x)$

$CON(x, y, z)$

$COST(x, y, z, w)$

$PROV(x, y)$

$\forall x, y, z. CON(x, y, z) \supset C(x) \wedge P(y) \wedge S(z)$

$\forall x, y, y', z. CON(x, y, z) \wedge CON(x, y', z) \supset y = y'$

$\forall x, y, z, w. COST(x, y, z, w) \supset CON(x, y, z) \wedge REAL(w)$

$\forall x, y. PROV(x, y) \supset P(x) \wedge S(y)$

$\forall x. P(x) \supset 1 \leq \# \{y \mid PROV(x, y)\} \leq 10$

$\forall y. S(y) \supset 1 \leq \# \{x \mid PROV(x, y)\}$

$\forall x. BC(x) \supset C(x)$

Exercise 2. Consider the above UML class diagram and the following (partial) instantiation:

Customer	BCustomers	Services	Provider	provides	contacts/cost
c1	b1	s1	p1	p1 s1	c1 s1 p1 90.0
c2	b2	s2	p2	p1 s2	c1 s2 p1 80.0
c3	b3	s3		p1 s3	c1 s3 p1 50.0
c4				p2 s2	b2 s1 p2 170.0
					b2 s2 p2 100.0

1. Check whether the above instantiation, once completed, is correct, and explain why it is or it is not.
2. Express in FOL the following queries and evaluate them over the completed instantiation:
 - (a) Check that, for every provider x and service y involved in a contract, provider x does provide service y .
 - (b) Return those customers that have contracts only for services provided by $p1$.
 - (c) Return those customers that have a contract for all services.

1) $C = \{c_1, c_2, c_3, c_4, b_1, b_2, b_3\}$

$$\forall x, y. \text{PROV}(x, y) \supset P(x) \wedge S(y)$$

p_1, p_2 ARE PROVIDERS

→ CARDINALS OK!

s_1, s_2, s_3 ARE SERVICES

$$\forall x, y, z. \text{CON}(x, y, z) \supset C(x) \wedge P(y) \wedge S(z)$$

c_1, b_2 ARE CUSTOMERS

p_1, p_2 ARE PROVIDERS

→ CARDINALS OK!


s_1, s_2, s_3 ARE SERVICES

2) a. $\forall c, p, s. \text{CON}(c, s, p) \supset \text{PROV}(p, s)$
 $\{\text{FALSE}\}$

b. $C(x) \wedge \exists s, p. \text{CON}(x, s, p) \wedge \forall s, p. (\text{CON}(x, s, p) \supset \text{PROV}(p, s))$
 $\{c_1\}$

c. $\forall s. C(x) \wedge (S(s) \supset \exists p \text{CON}(x, s, p))$
 $\{c_1\}$

Exercise 3. Model check the Mu-Calculus formula $\nu X. \mu Y. ((a \wedge [next]X) \vee (b \wedge \langle next \rangle Y))$ and the CTL formula $AF(EG(a \supset AX EX \neg a))$ (showing its translation in Mu-Calculus) against the following transition system:



a) $\nu X. \mu Y. ((a \wedge [next]X) \vee (b \wedge \langle next \rangle Y))$

$$[X_0] = \{0, 1, 2, 3, 4\}$$

$$[X_1] = [\mu Y. ((a \wedge [next]X_0) \vee (b \wedge \langle next \rangle Y))]$$

$$[Y_0] = \phi$$

$$[Y_1] = ([a] \cap PREA(next, X_0)) \cup ([b] \cap FREE(next, Y_0)) = \\ = (\{1, 2\} \cap \{0, 1, 2, 3, 4\}) \cup (\{3, 4\} \cap \phi) = \{1, 2\}$$

$$[Y_2] = ([a] \cap PREA(next, X_0)) \cup ([b] \cap FREE(next, Y_1)) = \\ = (\{1, 2\} \cap \{0, 1, 2, 3, 4\}) \cup (\{3, 4\} \cap \{0, 1, 2\}) = \{1, 2\}$$

$$[Y_1] = [Y_2] = [X_1] = \{1, 2\}$$

$$[X_2] = [\mu Y. ((a \wedge [next]X_1) \vee (b \wedge \langle next \rangle Y))]$$

$$[Y_0] = \phi$$

$$[Y_1] = ([a] \cap PREA(next, X_1)) \cup ([b] \cap FREE(next, Y_0)) = \\ = (\{1, 2\} \cap \{1\}) \cup (\{3, 4\} \cap \phi) = \{1\}$$

$$[Y_2] = ([a] \cap PREA(next, X_1)) \cup ([b] \cap FREE(next, Y_1)) = \\ = (\{1, 2\} \cap \{1\}) \cup (\{3, 4\} \cap \{0\}) = \{1\}$$

$$[Y_1] = [Y_2] = [X_2] = \{1\}$$

$$[X_3] = [\mu Y. ((a \wedge [next]X_2) \vee (b \wedge \langle next \rangle Y))]$$

$$[Y_0] = \phi$$

$$[Y_1] = ([a] \cap PREA(next, X_2)) \cup ([b] \cap FREE(next, Y_0)) = \\ (\{1, 2\} \cap \phi) \cup (\{3, 4\} \cap \phi) = \phi$$

$$[Y_0] = [Y_1] = [X_3] = \phi$$

$$[X_4] = [\mu Y. ((a \wedge [next]X_3) \vee (b \wedge \langle next \rangle Y))]$$

$$[Y_0] = \phi$$

$$[Y_i] = ([a] \cap \text{PREA}(\text{NEXT}, X_3)) \cup ([b] \cap \text{PREE}(\text{NEXT}, Y_0)) =$$



$$(\{1, 2\} \cap \phi) \cup (\{3, 4\} \cap \phi) = \phi$$

$$[Y_0] = [Y_i] = [X_4] = \phi$$

$$[X_3] = [X_4] = \phi$$

$$S_0 \in [\cup X. \mu Y. ((a \wedge [\text{NEXT}]X) \vee (b \wedge \langle \text{NEXT} \rangle Y))] = \phi? \text{ NO!}$$

b) $\text{AF}(\text{EG}(a \supset \text{AXEX} \neg a))$



$$[\alpha] = [EX \neg a] = [\langle \text{NEXT} \rangle \neg a] = \text{PREE}(\text{NEXT}, \neg a) = \{0, 2, 3, 4\} = [\alpha]$$

$$[\beta] = [AX \alpha] = [[\text{NEXT}] \alpha] = \text{PREA}(\text{NEXT}, \alpha) = \{1, 2, 3, 4\} = [\beta]$$

$$[\gamma] = [a \supset \beta] = [\neg a] \cup [\beta] = \{0, 3, 4\} \cup \{1, 2, 3, 4\} = \{0, 1, 2, 3, 4\} = [\gamma]$$

$$[\delta] = [\text{EG} \gamma] = [\cup Z. \gamma \wedge \langle \text{NEXT} \rangle Z]$$

$$[Z_0] = \{0, 1, 2, 3, 4\}$$

$$[Z_1] = [\gamma] \cap \text{PREE}(\text{NEXT}, Z_0) =$$

$$= \{0, 1, 2, 3, 4\} \cap \{0, 1, 2, 3, 4\} = \{0, 1, 2, 3, 4\}$$

$$[Z_0] = [Z_1] = [\delta] = \{0, 1, 2, 3, 4\}$$

$$[\epsilon] = [\text{AF} \delta] = [\mu Z. \delta \vee [\text{NEXT}] Z]$$

$$[Z_0] = \phi$$

$$[Z_1] = [\delta] \cup \text{PREA}(\text{NEXT}, Z_0) =$$

$$= \{0, 1, 2, 3, 4\} \cup \phi = \{0, 1, 2, 3, 4\}$$

$$[Z_2] = [\delta] \cup \text{PREA}(\text{NEXT}, Z_1) =$$

$$= \{0, 1, 2, 3, 4\} \cup \{0, 1, 2, 3, 4\} = \{0, 1, 2, 3, 4\}$$

$$[Z_1] = [Z_2] = [\epsilon] = \{0, 1, 2, 3, 4\}$$

$$\neg S_0 \models \epsilon? \rightarrow S_0 \in [\epsilon] = \{0, 1, 2, 3, 4\}? \text{ YES!}$$

Exercise 4. Check whether the Hoare triple below is correct, by using $(x \geq 0 \wedge y \geq 0 \wedge x + y = 23)$ as invariant:

$$\{x = 23 \wedge y = 0\} \text{ while}(x > 0) \text{ do } (x = x - 1; y = y + 1) \{y = 23\}$$

1. $P \supset I$

1. $\{x = 23 \wedge y = 0\} \supset (x \geq 0 \wedge y \geq 0 \wedge x + y = 23)$ ✓

2. $\neg g \wedge I \supset Q$

2. $(x \leq 0 \wedge (x \geq 0 \wedge y \geq 0 \wedge x + y = 23)) \supset y = 23$

3. $\{g \wedge I\} \delta \{I\}$

$(x = 0 \wedge y \geq 0 \wedge x + y = 23) \supset y = 23$ ✓

3. $\{x > 0 \wedge (x \geq 0 \wedge y \geq 0 \wedge x + y = 23)\} (x = x - 1; y = y + 1) \{x \geq 0 \wedge y \geq 0 \wedge x + y = 23\}$
 $\{x > 0 \wedge y \geq 0 \wedge x + y = 23\} \supset \text{wp}(x = x - 1; y = y + 1) \{x \geq 0 \wedge y \geq 0 \wedge x + y = 23\} ?$


$$\begin{aligned} &\{x \geq 0 \wedge y \geq -1 \wedge x + y = 22\} [x/x-1] = \{x \geq 1 \wedge y \geq -1 \wedge x + y = 23\} \\ &\quad x = x - 1; \\ &\{x \geq 0 \wedge y \geq 0 \wedge x + y = 23\} [y/y+1] = \{x \geq 0 \wedge y \geq -1 \wedge x + y = 22\} \\ &\quad y = y + 1; \\ &\{x \geq 0 \wedge y \geq 0 \wedge x + y = 23\} \end{aligned}$$

$\{x > 0 \wedge y \geq 0 \wedge x + y = 23\} \supset \{x \geq 1 \wedge y \geq -1 \wedge x + y = 23\} ?$ ✓

$(x \geq 0 \wedge y \geq 0 \wedge x + y = 23)$ IS AN INVARIANT


Exercise 5. Check whether the following FOL formula is valid, by using tableaux:

$$(\forall x.P(x) \wedge \forall x.Q(x)) \equiv \forall x.(P(x) \wedge Q(x))$$



SINCE ALL BRANCHES ARE CLOSED, THE FORMULA IS **VALID**

Exercise 6 (optional). Model check the LTL formula $\Diamond \Box \neg a$ against the following transition system, by considering that the Büchi automaton for $\neg(\Diamond \Box \neg a)$ is the one below:



$$\cup X. \mu Y (F \wedge \langle \text{NEXT} \rangle X \vee \langle \text{NEXT} \rangle Y)$$

$$[X_0] = \{1, 2, 3, 4, 5, 6\}$$

$$[X_i] = [\mu Y (F \wedge \langle \text{NEXT} \rangle X_0 \vee \langle \text{NEXT} \rangle Y)]$$

$$[Y_0] = \emptyset$$

$$[Y_1] = [F] \wedge \text{FREE}(\text{NEXT}, X_0) \cup \text{FREE}(\text{NEXT}, Y_0) =$$

$$= \{4, 6\} \cap \{1, 2, 3, 4, 5, 6\} \cup \emptyset = \{4, 6\}$$

$$[Y_2] = [F] \wedge \text{FREE}(\text{NEXT}, X_0) \cup \text{FREE}(\text{NEXT}, Y_1) =$$

$$= \{4, 6\} \cap \{1, 2, 3, 4, 5, 6\} \cup \{2, 3, 4, 5, 6\} = \{2, 3, 4, 5, 6\}$$

$$[Y_3] = [F] \wedge \text{FREE}(\text{NEXT}, X_0) \cup \text{FREE}(\text{NEXT}, Y_2) =$$

$$= \{4, 6\} \cap \{1, 2, 3, 4, 5, 6\} \cup \{1, 2, 3, 4, 5, 6\} = \{1, 2, 3, 4, 5, 6\}$$

$$[Y_4] = [F] \wedge \text{FREE}(\text{NEXT}, X_0) \cup \text{FREE}(\text{NEXT}, Y_3) =$$

$$= \{4, 6\} \cap \{1, 2, 3, 4, 5, 6\} \cup \{1, 2, 3, 4, 5, 6\} = \{1, 2, 3, 4, 5, 6\}$$

$$[Y_3] = [Y_4] = [X_i] = \{1, 2, 3, 4, 5, 6\}$$

$$[X_0] = [X_i] = \{1, 2, 3, 4, 5, 6\}$$

$$S_i \in [\cup X. \mu Y (F \wedge \langle \text{NEXT} \rangle X \vee \langle \text{NEXT} \rangle Y)] = \{1, 2, 3, 4, 5, 6\} \quad \text{YES!}$$

$$\text{SO } (A\gamma \wedge A\neg\varphi) \neq \emptyset \rightarrow \gamma \neq \varphi$$