

# TD-Probability

Marco Biroli

September 3, 2020



# Chapter 1

## TD 1

### 1.1 A strategic choice.

Let  $X \in \{0, 1\}^3$  (resp.  $Y$ ) be the random variable corresponding to the results of the matches using the first strategy (resp. the second strategy). Then we have that (let  $D = \{(1, 1, 1), (1, 1, 0), (0, 1, 1)\}$ ):

$$P(X \in D) = a^2b + ab(1 - a) + (1 - a)ba = ab(2 - a)$$

Similarly:

$$P(Y \in D) = b^2a + ba(1 - b) + (1 - b)ab = ba(2 - b)$$

Then since  $a > b$  we have that  $P(X \in D) < P(Y \in D)$ , hence the winning strategy is BAB.

### 1.2 Derangements

#### 1.2.1

Let  $E$  be a finite set and  $A, B \subseteq E$ . We denote by  $1_A$  the indicator function of  $A$  and  $\bar{A}$  the complement of  $A$ . Then we have that:

$$1_{\bar{A}} = 1 - 1_A \quad \text{and} \quad 1_{A \cap B} = 1_A \cdot 1_B \quad \text{and} \quad 1_{A \cup B} = 1_A + 1_B - 1_{A \cap B}$$

#### 1.2.2

We will prove this by induction on  $n$ . The base case  $n = 1$  as well as  $n = 2$  are trivially satisfied. Now assume that this is satisfied for  $n$  then we have that (using the induction hypothesis for  $n = 2$ ):

$$\text{card} \left( \bigcup_{i=1}^n A_i \cup A_{n+1} \right) = \text{card} \left( \bigcup_{i=1}^n A_i \right) + \text{card}(A_{n+1}) - \text{card} \left( \left( \bigcup_{i=1}^n A_i \right) \cap A_{n+1} \right)$$

Now we develop the last term into:

$$\left( \bigcup_{i=1}^n A_i \right) \cap A_{n+1} = \bigcup_{i=1}^n (A_i \cap A_{n+1})$$

Now applying the induction hypothesis gives the desired result.

#### 1.2.3

Let  $A_i$  be the set of permutations that fixes point  $i$ . Then from the inclusion-exclusion principle we have:

$$D_n = n! - \text{card} \left( \bigcup_{i=1}^n A_i \right) = n! - \sum_{k=1}^n (-1)^{k-1} \binom{n}{k} (n-k)! = n! \sum_{k=2}^n \frac{(-1)^k}{k!}$$

#### 1.2.4

The probability that no one gets their jacket corresponds to the probability of having a derangement in other words:

$$p_n = \frac{D_n}{n!} = \sum_{k=2}^n \frac{(-1)^k}{k!} \xrightarrow{n \rightarrow \infty} \frac{1}{e}$$

### 1.2.5

We have that:

$$D_{n,l} = \binom{n}{l} D_{n-l} = \binom{n}{l} (n-l)! \sum_{k=2}^{n-l} \frac{(-1)^k}{k!} = \frac{n!}{l!} \sum_{k=2}^{n-l} \frac{(-1)^k}{k!}$$

Hence the probability that exactly  $l$  people leave with their jackets is:

$$p_l = \frac{D_{n,l}}{n!} = \frac{1}{l!} \sum_{k=2}^{n-l} \frac{(-1)^k}{k!}$$

### 1.2.6

The probability that a given person gets back their jacket is  $p_s = \frac{1}{n}$ . The probability that at least one person gets back their jacket is:

$$p_a = 1 - p_n = 1 - \sum_{k=2}^n \frac{(-1)^k}{k!}$$

Notice that  $p_s < p_a$ .

## 1.3 Balls in bins

### 1.3.1

- (a) If all the balls are distinguishable then we have  $\Omega = \llbracket 1, n \rrbracket^r$  is the set of tuples where each element corresponds to where the  $i$ -th ball has been sent to. Then  $\mathcal{F} = \mathcal{P}(\Omega)$  and since each event is sampled uniformly at random we have that:

$$\forall \omega \in \Omega, P(\omega) = \frac{1}{|\Omega|} = \frac{1}{n^r}$$

Then the probability of  $(r_1, \dots, r_n)$  is given by:

$$P[(r_1, \dots, r_n)] = P[\{\omega \in \Omega : \forall i \in \llbracket 1, n \rrbracket \# \{b \in \Omega : b = i\} = r_i\}] = \frac{1}{n^r} \binom{r}{r_1, r_2, \dots, r_n}$$

- (b) Now we have that  $\Omega = \{(r_i \in \mathbb{N} : i \in \llbracket 1, n \rrbracket) : \sum_{i=1}^n r_i = r\}$ . Again we have that  $\mathcal{F} = \mathcal{P}(\Omega)$ . Then we have that:

$$P[(r_1, \dots, r_n)] = \frac{1}{|\Omega|} = \frac{1}{\binom{r+n-1}{n-1}}$$

- (c) Now we have that  $\Omega = \{s \in \{0, 1\}^n : \sum_{i=1}^n s_i = r\}$  corresponding to the tuple indicating if each state is occupied or not. Once again  $\mathcal{F} = \mathcal{P}(\Omega)$ . Now the probability is given by:

$$P[(r_1, \dots, r_n)] = \frac{1}{|\Omega|} = \frac{1}{\binom{n}{r}}$$

### 1.3.2

The probability that at least two have the same birthday is the 1 minus the probability that none of them share a birthday. The probability that none of them share a birthday is given by  $\frac{r!}{n^r} \binom{n}{r}$ . Hence the probability that at least two people share a birthday is given by:  $1 - \frac{r!}{n^r} \binom{n}{r}$ .

### 1.3.3

Days are bins, accidents are distinguishable balls hence the probability is given by:

$$\frac{\binom{r}{n} n^{r-n}}{n^r} = n^{-n} \binom{r}{n}$$

### 1.3.4