

Mathematical and Physical Modeling of Entangled Photon Propagation in Optical Fibers for MATLAB Simulation

Marco Cioci
Politecnico di Milano
Student ID: 10694772
`marco.cioci@mail.polimi.it`

Contents

1 General Framework

1.1 Purpose and Scope

The purpose of this document is to define a coherent working framework for the mathematical modeling and numerical simulation of polarization-entangled photon pairs propagating through optical fibers.

This text is not intended to appear directly in the final thesis manuscript. Instead, it serves as an internal technical guide throughout the development of the project. Its role is to support communication, maintain consistency of notation and assumptions, and provide a stable reference for the progressive validation of both the theoretical model and the MATLAB implementation.

The central objective is to organize the physical assumptions, mathematical formalism, and simulation strategy into a reproducible structure, so that each computational step can be traced back to a precise theoretical element. In this way, the simulator is not developed as an isolated numerical tool, but as a direct operational translation of the underlying quantum model.

Simulation notes. At the MATLAB level, this document should be treated as the reference specification for notation, model hierarchy, and implementation priorities. The simulator should therefore be organized into clearly identifiable modules, each corresponding to a specific theoretical block. This modular structure is essential both for step-by-step validation of intermediate results and for future extensions of the model.

1.2 Theoretical Formalism and Notation

The system is described within the framework of quantum information theory. The relevant mathematical structures include finite-dimensional Hilbert spaces, quantum states, tensor products, unitary operators, density operators, and projective measurements. Elements of classical optics are introduced only when required to justify the physical origin of the effective quantum transformations.

The notation and formal tools are aligned, as far as possible, with the framework of *Quantum Computation and Quantum Information* by Nielsen and Chuang.

In this formalism:

- single subsystems are represented by vectors in $\mathbb{C}^2$,
- composite systems are described by tensor-product spaces,
- evolutions are represented by linear operators (unitary in the ideal case),
- measurements are described through Hermitian operators.

All states and operators are represented as vectors and matrices over $\mathbb{C}$, with dimensions determined by the tensor-product structure of the system.

Simulation notes. The simulator adopts a quantum-information-oriented representation from the outset. Pure states are represented as complex vectors, mixed states as density matrices, and evolutions and measurements as matrix operators.

This abstract structure defines the numerical backbone of the simulator and is independent of the specific physical encoding used.

1.3 Physical Scenario and Experimental Setting

The physical setting considered in this work is the following. A spontaneous parametric down-conversion (SPDC) source generates pairs of polarization-entangled photons, ideally prepared in a Bell state. The two photons are injected into distinct optical paths, denoted A and B , and propagate through independent optical fibers.

Each fiber acts on the polarization degree of freedom of the corresponding photon. In a general description, this action can be represented by a unitary operator acting on a single-qubit Hilbert space. However, motivated by experimental constraints and by the limited degree of control available on the full polarization transformation, the model is initially restricted to a simplified effective description.

After compensation of global polarization rotations, the dominant residual effect can be modeled as a relative phase shift between the horizontal and vertical polarization components. In the qubit representation, this corresponds to a rotation around the computational Z -axis of the Bloch sphere. The objective at this stage is therefore not to reproduce the full electromagnetic propagation in optical fibers, but to construct an effective quantum model that captures the relevant impact of fiber propagation on entanglement and measurable correlations.

Simulation notes. At the simulator level, the physical scenario is reduced to the minimal set of elements required for computation: an initial bipartite quantum state, two labeled channels A and B , and one local transformation acting on each subsystem.

In the first modeling stage, unnecessary optical detail is neglected and each fiber is described exclusively through its effective action on polarization. This leads to a local unitary representation of the form

$$U_A \otimes U_B,$$

where $U_A, U_B \in \mathbb{C}^{2 \times 2}$ are unitary operators.

This abstraction defines the fundamental interface between the physical system and the simulator, and will later allow the introduction of more refined models, including stochastic phase evolution and non-unitary effects.

2 Modeling Strategy

2.1 Progressive Modeling Approach


The modeling approach follows a progressive refinement strategy, in which the system is described through successive levels of approximation of increasing physical complexity.

Each stage introduces additional physical effects while remaining consistent with the previous ones. This ensures analytical transparency and preserves continuity between simplified and refined models.

From a theoretical perspective, this approach isolates the contribution of individual mechanisms. From a computational perspective, it enables a layered simulator structure, where each refinement corresponds to a well-defined extension of the existing codebase.

2.2 Hierarchy of Modeling Levels

The modeling workflow is structured as a sequence of progressively refined descriptions:



More explicitly:

- **Level 1: Minimal phase model**
Pure, maximally entangled state. Fiber action reduced to a relative phase. Analytical dependence of correlations on a single parameter.
- **Level 2: General unitary dynamics**
Arbitrary polarization rotations described by local operators $U_A \otimes U_B$.
- **Level 3: Mixed-state description**
Density operators used to represent statistical mixtures and loss of coherence.

- **Level 4: Noisy quantum channels**

Effective completely positive trace-preserving (CPTP) maps modeling depolarization and decoherence.

- **Level 5: Effective fiber models**

Physically motivated descriptions, such as concatenations of random unitary segments or stochastic polarization evolution.

This hierarchy provides both a conceptual and an implementation roadmap. The initial levels remain analytically tractable and allow direct validation, while the later levels increase physical realism and connect to experimentally relevant effects.

Simulation notes. This hierarchy should be mirrored explicitly in the codebase. Each level should correspond to a distinct layer of functionality: state preparation, local unitary evolution, density-matrix handling, channel application, and advanced propagation models. This structure ensures that each refinement appears as an additional computational step rather than a structural modification of the simulator.

2.3 Baseline Model: Pure-State Description

At the first stage, the bipartite system is described as a pure state in $\mathbb{C}^2 \otimes \mathbb{C}^2$, and fiber propagation is modeled through local unitary transformations. Particular emphasis is placed on relative phase shifts induced by the fibers.

The main objective is to analyze how these phase shifts affect measurable two-photon correlations. The relevant observables are operators of the form

$$\sigma_i \otimes \sigma_j, \quad i, j \in \{x, y, z\},$$

as well as operators corresponding to arbitrary measurement directions on the Bloch sphere.

This baseline model constitutes the simplest regime in which entanglement is preserved and can be probed exclusively through correlation observables. It provides a controlled setting for deriving analytical expressions and validating the numerical implementation.

Simulation notes. At this level, the simulator operates on pure two-qubit states represented as 4×1 complex vectors. The core tasks are: generation of the Bell state, application of local operators U_A and U_B , and evaluation of expectation values.

In this regime, the simulator enables direct comparison between analytical predictions and numerical results for correlation functions, establishing a first level of validation that will be extended in Chapter 4 through quantitative state characterization.

2.4 Extension to Mixed-State Formalism

At the next level, the model incorporates statistical effects and decoherence. In this regime, the pure-state description is no longer sufficient and must be replaced by a density-operator formalism. This extension enables the description of:

- reduced states of individual photons,
- loss of coherence due to unresolved degrees of freedom,
- effective non-unitary dynamics arising from environmental interactions or statistical averaging.

Within this framework, physically relevant quantities such as purity, fidelity, and concurrence can be defined and computed, allowing a quantitative characterization of mixedness and entanglement (see Chapter 4).

An important derived quantity is the reduced density operator of a subsystem, obtained via partial trace. For instance, the reduced state of subsystem A is

$$\rho_A = \text{Tr}_B(\rho_{AB}).$$

For maximally entangled states such as $|\Psi^+\rangle$, the reduced density operators are maximally mixed:

$$\rho_A = \rho_B = \frac{1}{2}I.$$

This property reflects the absence of local information despite strong nonlocal correlations, and provides the conceptual link between entanglement and mixedness that will be explored quantitatively in later sections.

Expectation values in the density-operator formalism are computed as

$$\langle O \rangle = \text{Tr}(\rho O),$$

which generalizes the pure-state expression $\langle \psi | O | \psi \rangle$.

Simulation notes. The simulator must be extended so that all routines operate on density matrices, including 4×4 representations, partial trace operations, and observable evaluation. This generalized framework enables the transition from deterministic evolution to ensemble-based descriptions, and supports the computation of state metrics used for validation and physical interpretation.

The pure-state implementation is preserved as a special case, ensuring consistency across all modeling levels.

2.5 Simulator Design and Architecture

The simulator is designed as a modular implementation of the theoretical model.

The initial modules include:

- Bell-state generation,
- local unitary propagation,
- correlation evaluation.

More advanced modules introduce:

- quantum channels,
- reduced density operators,
- entanglement measures,
- experimentally relevant observables such as Bell parameters and Stokes-like quantities.

The theoretical structure is directly reflected in the computational architecture, ensuring consistency, traceability, and extensibility.

The implementation distinguishes clearly between reusable computational kernels and higher-level experiment scripts, allowing the simulator to evolve into a structured and reusable research tool.

Simulation notes. At the current stage, this modular design is reflected in the directory structure:

- **Core:** state vectors, operators, expectation values, correlation routines;
- **Models:** physical propagation models;
- **Utils:** validation routines and auxiliary tools;
- **Experiments:** scripts implementing complete simulation studies.

This separation enforces a strict distinction between general mathematical operations and physically specific modeling choices.

3 Physical and Mathematical Representation

3.1 Quantum System and Hilbert Space Structure

The physical system considered consists of polarization-entangled photon pairs generated by spontaneous parametric down-conversion (SPDC). The two photons propagate along distinct paths, denoted A and B , forming a bipartite system.

In this work, the qubit is encoded in the polarization degree of freedom:

$$|H\rangle \equiv |0\rangle, \quad |V\rangle \equiv |1\rangle.$$

Accordingly, the computational basis of the bipartite system is

$$\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\},$$

corresponding to

$$\{|H\rangle_A |H\rangle_B, |H\rangle_A |V\rangle_B, |V\rangle_A |H\rangle_B, |V\rangle_A |V\rangle_B\}.$$

The Hilbert space of the total system is therefore

$$\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B \cong \mathbb{C}^2 \otimes \mathbb{C}^2,$$

which has dimension 4.

As a consequence:

- pure states are represented by complex vectors of length 4,
- operators acting on the full system are 4×4 matrices.

This construction specifies the concrete realization of the abstract quantum-information framework introduced previously.

Simulation notes. At MATLAB level, bipartite pure states are represented as 4×1 complex vectors and operators as 4×4 matrices. The basis ordering is fixed as

$$\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\},$$

and must remain invariant, since all tensor products, observables, and density matrices depend on this convention.

This ordering is already enforced in the current codebase and must be preserved in all future extensions.

3.2 Bell-State Initialization

As a reference initial condition, we consider the Bell state

$$|\Psi^+\rangle = \frac{1}{\sqrt{2}} (|H\rangle_A |V\rangle_B + |V\rangle_A |H\rangle_B),$$

which, in computational notation, is

$$|\Psi^+\rangle = \frac{1}{\sqrt{2}} (|01\rangle + |10\rangle).$$

In the ordered computational basis

$$\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\},$$

this state is represented by the column vector

$$|\Psi^+\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix}.$$

This state is maximally entangled and provides the natural input for the baseline version of the simulator. It also serves as the reference state with respect to which later quantities such as fidelity and Bell-inequality violation may be evaluated.

The choice of a Bell state is especially convenient because it provides both a clean theoretical benchmark and a physically relevant entangled resource. It also allows the effect of fiber propagation to be interpreted directly in terms of coherence and two-photon correlation changes.

Simulation notes. The simulator should include a dedicated state-preparation routine returning the Bell-state vector in canonical basis ordering. This reference state is useful for normalization checks, debugging, comparison with analytical formulas, and later evaluation of propagated-state fidelity.

At the current stage, the simulator already includes a Bell-state generator returning the four standard Bell states:

$$|\Psi^\pm\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle), \quad |\Phi^\pm\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle),$$

although the present baseline model uses $|\Psi^+\rangle$ as the default reference input.

3.3 Fiber Propagation as Local Unitary Evolution

Each photon propagates through its own optical fiber. At the level of the polarization qubit, the action of each fiber is described by a linear operator acting locally on the corresponding single-photon Hilbert space. In the most general ideal lossless description, this operator is unitary. Therefore, if U_A and U_B denote the single-qubit operators associated with the two fibers, the evolution of the initial two-photon state is

$$|\psi_{\text{out}}\rangle = (U_A \otimes U_B) |\psi_{\text{in}}\rangle.$$

This is the natural bipartite extension of single-qubit evolution and follows directly from the tensor-product structure of composite quantum systems. It is the appropriate starting point for later refinements, including random unitary perturbations, concatenated fiber sections, and effective channel descriptions.

In this framework, the two fibers act independently on the two polarization subsystems, and the total evolution is obtained by tensor composition of the corresponding local transformations. This description is valid as long as the fibers do not induce any direct interaction between the two photons.

Simulation notes. The fundamental evolution primitive in MATLAB should be the tensor-product action

$$\psi_{\text{out}} = (U_A \otimes U_B) \psi_{\text{in}}.$$

It is convenient to isolate this in a dedicated routine, so that any later modification of the single-fiber model automatically propagates to the full bipartite simulation.

This abstraction is already present in the implementation, where local single-qubit matrices are combined into the global operator $U_A \otimes U_B$ before being applied to the input state.

3.4 Reduced Phase Model

Although a general fiber transformation may involve arbitrary polarization rotations, birefringence, and mode-coupling effects, the first approximation adopted here is simpler and directly motivated by the experimental control scheme. After partial compensation of polarization drifts, the dominant residual effect is modeled as a relative phase shift between the horizontal and vertical components, corresponding to an effective rotation around the Z -axis of the Bloch sphere.

At the single-qubit level, this is represented by the diagonal unitary

$$U(\theta) = \begin{pmatrix} 1 & 0 \\ 0 & e^{i\theta} \end{pmatrix},$$

up to an irrelevant global phase. If the two fibers induce phases θ_A and θ_B , then the total transformation is

$$U_A \otimes U_B = U(\theta_A) \otimes U(\theta_B).$$

Applying this operator to the reference Bell state $|\Psi^+\rangle$ gives

$$|\psi(\theta_A, \theta_B)\rangle = \frac{1}{\sqrt{2}}(e^{i\theta_B}|01\rangle + e^{i\theta_A}|10\rangle).$$

By factoring out a global phase, which has no effect on measurement statistics, this state can be rewritten as

$$|\psi(\theta)\rangle = \frac{1}{\sqrt{2}} (|01\rangle + e^{i\theta}|10\rangle),$$

where

$$\theta = \theta_A - \theta_B.$$

Thus, in the reduced model, the physically relevant fiber action is encoded in a single parameter: the relative phase θ between the two components of the entangled state. This parameter depends on propagation conditions such as birefringence, path length, and environmental fluctuations, and it becomes the main control variable in the first stage of the simulator.

This reduced description is useful because it isolates the dominant effect of interest while preserving the full two-qubit entangled structure.

Simulation notes. This is the first concrete propagation model to implement. The simulator should allow θ_A and θ_B as input parameters, while also recognizing that the observable behavior depends only on $\theta = \theta_A - \theta_B$. In practice, both forms are useful: the two-phase form for physical interpretation, and the reduced one-parameter form for analytical checks and parameter sweeps. At the current implementation stage, the reduced model is realized through a dedicated single-qubit phase-unitary routine and is used operationally in the first complete experiment by fixing one arm to the identity and sweeping the phase in the other arm.

3.5 Interpretation and Role of the Reduced Model

The reduced phase model is particularly useful for three reasons. First, it preserves a pure-state description, keeping the initial implementation analytically transparent. Second, it isolates the effect most directly relevant for the first experimental analysis, namely the deformation of two-photon correlations induced by a relative phase. Third, it provides the natural starting point for later extensions to density operators and noisy channels without changing the basic bipartite structure of the problem.

For this reason, the first stage of the simulator is built around the family of states

$$|\psi(\theta)\rangle = \frac{1}{\sqrt{2}} (|01\rangle + e^{i\theta}|10\rangle),$$

from which one computes expectation values of local and nonlocal observables, correlation functions, and Bell-type quantities as functions of θ .

The reduced model therefore plays a dual role: it is both a physically motivated approximation to fiber propagation and the minimal operational model from which one can derive clean and testable predictions.

Simulation notes. The first validated MATLAB study should therefore consist of a parameter sweep in θ , with numerical evaluation of state vectors, expectation values, and correlation quantities. This provides a minimal but complete simulator: one input family of states, one effective propagation parameter, and a set of measurable outputs that can be directly plotted and compared with theory.

At the current stage, this is implemented through a sweep routine that returns a structured collection of sampled phase values and corresponding correlation outputs, together with a plotting utility for visualizing selected correlations as functions of θ .

3.6 Correlation Observables: Theory and Implementation

At the present stage of development, the observables explicitly implemented in the simulator are the standard aligned two-qubit Pauli correlations:

$$\sigma_x \otimes \sigma_x, \quad \sigma_y \otimes \sigma_y, \quad \sigma_z \otimes \sigma_z.$$

These are constructed from the single-qubit Pauli matrices

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Given a pure two-qubit state $|\psi\rangle$, the expectation values are defined as

$$\langle \sigma_i \otimes \sigma_i \rangle = \langle \psi | (\sigma_i \otimes \sigma_i) | \psi \rangle, \quad i \in \{x, y, z\}.$$

Analytical behavior under the reduced phase model. Within the reduced phase model, the initial Bell state

$$|\Psi^+\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}}$$

evolves into

$$|\psi(\theta)\rangle = \frac{|01\rangle + e^{i\theta}|10\rangle}{\sqrt{2}}.$$

For this family of states, the correlation observables can be computed analytically:

$$\langle \sigma_x \otimes \sigma_x \rangle = \cos \theta, \quad \langle \sigma_y \otimes \sigma_y \rangle = \cos \theta, \quad \langle \sigma_z \otimes \sigma_z \rangle = -1.$$

These results show that the fiber-induced phase θ affects only the transverse correlations in the x - and y -directions, while the longitudinal correlation along z remains invariant. This reflects the fact that the relative phase modifies coherence between computational-basis components without changing their populations.

Simulation notes. At implementation level, the simulator contains routines for:

- generating the Pauli matrices,
- constructing tensor-product operators,
- evaluating expectation values,
- packaging correlation outputs into structured data containers.

Additional helper routines verify normalization of input state vectors and unitarity of operators used in the fiber model.

These observables constitute the primary diagnostic outputs of the simulator and provide the most direct bridge between analytical predictions and numerical results. Their dependence on θ defines the first validation pipeline of the code.

3.7 From Deterministic Evolution to Ensemble Description

The reduced phase model introduced in the previous sections describes a deterministic unitary evolution of a pure entangled state, parametrized by a single phase θ .

In realistic fiber propagation, however, this phase is not constant but fluctuates due to environmental effects such as birefringence variations and thermal instability. As a consequence, each photon pair experiences a different phase shift.

When these fluctuations occur on timescales shorter than the measurement resolution, individual realizations cannot be resolved experimentally. The observed system is therefore not described by a single pure state, but by an ensemble of states corresponding to different phase realizations.

In this regime, the effective description is obtained by averaging over the phase distribution:

$$\rho = \mathbb{E}_\theta [|\psi(\theta)\rangle\langle\psi(\theta)|].$$

This transition does not introduce new dynamics at the level of individual realizations, which remain unitary, but reflects a loss of accessible phase information. The resulting state is mixed, even though each underlying realization is pure.

This mechanism provides the physical origin of mixed-state descriptions in the present model and establishes the connection between fiber-induced phase fluctuations and decoherence.

Expectation values are therefore computed using the density-operator formalism:

$$\langle O \rangle = \text{Tr}(\rho O),$$

which accounts for the statistical nature of the ensemble.

This framework forms the basis for the modeling and analysis of phase-noise effects, which will be investigated quantitatively in Chapter 4.

4 Quantitative Characterization of Bipartite Quantum States

4.1 Reduced Density Operators

In bipartite quantum systems, the state of an individual subsystem is generally described by a density operator obtained through a partial trace over the complementary subsystem. This formalism accounts for correlations that may be present at the global level.

Let the composite system be defined on the Hilbert space

$$\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B,$$

with $\mathcal{H}_A \cong \mathbb{C}^2$ and $\mathcal{H}_B \cong \mathbb{C}^2$. A generic global state is described by a density operator

$$\rho_{AB} \in \mathbb{C}^{4 \times 4}.$$

The reduced density operators are defined as

$$\rho_A = \text{Tr}_B(\rho_{AB}), \quad \rho_B = \text{Tr}_A(\rho_{AB}).$$

These operators reproduce all expectation values of observables acting locally on the subsystem. For any observable O_A acting on subsystem A ,

$$\langle O_A \rangle = \text{Tr}(\rho_{AB}(O_A \otimes I_B)) = \text{Tr}(\rho_A O_A).$$

An explicit construction can be written as

$$\rho_A = \sum_{j=0}^1 (I_A \otimes \langle j|) \rho_{AB} (I_A \otimes |j\rangle),$$

which shows that the partial trace eliminates the degrees of freedom associated with the unobserved subsystem.

For maximally entangled Bell states, such as

$$|\Psi^+\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}},$$

one obtains

$$\rho_A = \rho_B = \frac{I}{2},$$

indicating that each subsystem is locally maximally mixed despite the global state being pure. The same result holds for the phase-evolved state

$$|\psi(\theta)\rangle = \frac{|01\rangle + e^{i\theta}|10\rangle}{\sqrt{2}},$$

where the relative phase modifies global coherence but leaves local states unchanged:

$$\rho_A(\theta) = \rho_B(\theta) = \frac{I}{2}.$$

A detailed derivation of this result, including the explicit matrix representation and term-by-term partial trace, is provided in Appendix A.

By linearity of the partial trace, this property is preserved under convex combinations. Thus, for the ensemble state

$$\rho_{AB}^{(\text{ens})} = \frac{1}{N} \sum_{k=1}^N |\psi(\theta_k)\rangle \langle \psi(\theta_k)|,$$

one finds

$$\rho_A^{(\text{ens})} = \rho_B^{(\text{ens})} = \frac{I}{2}.$$

This leads to an important conceptual point: the same reduced density operator

$$\rho_A = \frac{I}{2}$$

can arise from physically distinct global states.
On one hand, it may describe a classical ensemble

$$\rho^{(\text{class})} = \frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|1\rangle\langle 1|,$$

where the system is in a definite state but the observer lacks knowledge.
On the other hand, it may result from entanglement:

$$\rho_A^{(\text{ent})} = \text{Tr}_B(|\Psi^+\rangle\langle\Psi^+|) = \frac{I}{2}.$$

Since both cases yield identical expectation values for all local observables,

$$\text{Tr}(\rho^{(\text{class})}O_A) = \text{Tr}(\rho_A^{(\text{ent})}O_A),$$

they are indistinguishable by local measurements alone.

The distinction emerges only at the global level, where coherence and correlations must be analyzed through quantities such as purity and bipartite correlation functions (see Appendix A for an explicit comparison).

Finally, the reduced density operator directly determines measurable local quantities. For Pauli observables,

$$\langle\sigma_x\rangle = \text{Tr}(\rho_A\sigma_x), \quad \langle\sigma_y\rangle = \text{Tr}(\rho_A\sigma_y), \quad \langle\sigma_z\rangle = \text{Tr}(\rho_A\sigma_z),$$

and for $\rho_A = I/2$ all expectation values vanish, indicating complete local depolarization.

4.2 Purity and Mixedness

Having introduced reduced density operators in Section 4.1, we now require a quantitative criterion to distinguish whether a quantum state is pure or mixed. This distinction is essential for interpreting the physical effects of fiber propagation, especially when extending the model beyond deterministic unitary evolution.

A quantum state described by a density operator ρ is said to be *pure* if it can be written as

$$\rho = |\psi\rangle\langle\psi|,$$

for some normalized state vector $|\psi\rangle$. Otherwise, the state is said to be *mixed*, meaning that it cannot be represented by a single state vector but only as a statistical ensemble

$$\rho = \sum_k p_k |\psi_k\rangle\langle\psi_k|, \quad p_k \geq 0, \quad \sum_k p_k = 1.$$

A convenient and operational way to distinguish between these two cases is provided by the *purity*, defined as

$$\gamma(\rho) = \text{Tr}(\rho^2).$$

This quantity satisfies the fundamental properties

$$\gamma(\rho) = 1 \quad \Longleftrightarrow \quad \rho \text{ is pure,}$$

$$\gamma(\rho) < 1 \quad \Longleftrightarrow \quad \rho \text{ is mixed.}$$

This follows directly from the spectral decomposition of ρ . If

$$\rho = \sum_i \lambda_i |i\rangle\langle i|,$$

with $\lambda_i \geq 0$ and $\sum_i \lambda_i = 1$, then

$$\gamma(\rho) = \sum_i \lambda_i^2.$$

The purity reaches its maximum value 1 only when a single eigenvalue is equal to 1 and all others vanish, corresponding to a pure state. Otherwise, the sum of squares is strictly smaller than 1.

More generally, for a system with Hilbert space dimension d , the purity is bounded by

$$\frac{1}{d} \leq \gamma(\rho) \leq 1,$$

where the lower bound is attained by the maximally mixed state $\rho = I/d$.

For single-qubit systems, purity admits a particularly transparent interpretation in terms of the Bloch vector representation introduced in Section 4.1. Any qubit density operator can be written as

$$\rho = \frac{1}{2} (I + \vec{r} \cdot \vec{\sigma}),$$

where $\vec{r} \in \mathbb{R}^3$ is the Bloch vector and $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is the vector of Pauli matrices. In this representation, one finds

$$\gamma(\rho) = \frac{1 + \|\vec{r}\|^2}{2}.$$

Therefore,

$$\|\vec{r}\| = 1 \Rightarrow \rho \text{ is pure}, \quad \|\vec{r}\| = 0 \Rightarrow \rho = \frac{I}{2} \text{ is maximally mixed}.$$

This result provides a direct connection between the Bloch/Stokes parameters computed in the simulator and the degree of mixedness of the state.

As already emphasized in Section 4.1, mixedness does not necessarily imply classical noise. A state may be mixed either because it belongs to a classical statistical ensemble or because it arises as the reduced state of a larger entangled system.

In the present context, both mechanisms are relevant but play distinct roles. Mixedness induced by entanglement is intrinsic to the bipartite structure of the system and is already present for pure global states. By contrast, ensemble-induced mixedness arises from averaging over random parameters, such as phase fluctuations in fiber propagation, and leads to a genuine loss of global coherence.

To illustrate the first mechanism, we consider a paradigmatic example provided by the Bell state

$$|\Psi^+\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}}.$$

Its global density operator

$$\rho_{AB} = |\Psi^+\rangle\langle\Psi^+|$$

is pure, and therefore satisfies

$$\gamma(\rho_{AB}) = 1.$$

However, its reduced density operators are

$$\rho_A = \rho_B = \frac{I}{2},$$

which are maximally mixed and satisfy

$$\gamma(\rho_A) = \gamma(\rho_B) = \frac{1}{2}.$$

Thus, a globally pure entangled state may generate locally mixed subsystem states. This distinction between global purity and local mixedness is central to the interpretation of bipartite entangled systems.

In the context of the present model, this distinction plays a crucial role. For the phase-evolved Bell-like state

$$|\psi(\theta)\rangle = \frac{|01\rangle + e^{i\theta}|10\rangle}{\sqrt{2}},$$

the evolution is generated by local unitary transformations, and therefore the global state remains pure for all values of θ :

$$\gamma(\rho(\theta)) = 1.$$

At the same time, as shown in Section 4.1, the reduced states remain

$$\rho_A(\theta) = \rho_B(\theta) = \frac{I}{2},$$

so that their purity is constant and minimal:

$$\gamma(\rho_A(\theta)) = \gamma(\rho_B(\theta)) = \frac{1}{2}.$$

Therefore, the introduction of a relative phase modifies the correlation observables without affecting either global purity or local mixedness.

The notion of purity becomes particularly relevant when extending the model to include more realistic physical effects. While deterministic unitary evolution preserves purity, ensemble averaging over random parameters or the inclusion of frequency-dependent propagation effects, such as polarization mode dispersion, can lead to genuine mixed states.

In particular, for the random-phase ensemble introduced above, the global density operator depends on the ensemble coherence

$$\mu = \langle e^{i\theta} \rangle,$$

and its purity is given by

$$\gamma(\rho_{AB}^{(\text{ens})}) = \frac{1 + |\mu|^2}{2}.$$

Equivalently,

$$\gamma(\rho_{AB}^{(\text{ens})}) = \frac{1 + \langle \cos \theta \rangle^2 + \langle \sin \theta \rangle^2}{2}.$$

This shows that the loss of purity is directly governed by the reduction of phase coherence in the ensemble. When $|\mu| = 1$, the state remains pure, while for $|\mu| < 1$ the ensemble becomes mixed. For this reason, purity will serve as a fundamental diagnostic tool in the numerical simulator, complementing correlation observables and preparing the ground for more advanced quantities such as fidelity and entanglement measures.

4.3 Fidelity with Respect to Bell States

Having introduced reduced density operators and purity as measures of local information and mixedness, we now require a quantitative tool to evaluate how close a given quantum state is to a reference target state. This role is fulfilled by the notion of fidelity.

In general, the fidelity between two quantum states described by density operators ρ and σ is defined as

$$F(\rho, \sigma) = \left(\text{Tr} \sqrt{\sqrt{\rho} \sigma \sqrt{\rho}} \right)^2.$$

This quantity satisfies

$$0 \leq F(\rho, \sigma) \leq 1,$$

with $F = 1$ if and only if $\rho = \sigma$, and $F = 0$ for perfectly distinguishable (orthogonal) states. Fidelity therefore provides a measure of similarity between quantum states.

In the present context, the reference state is a pure Bell state, and the fidelity simplifies considerably. Let

$$\sigma = |\phi\rangle\langle\phi|,$$

with $|\phi\rangle$ a normalized state vector. Then one obtains (see Appendix B for a detailed derivation)

$$F(\rho, |\phi\rangle) = \langle\phi|\rho|\phi\rangle.$$

This expression admits a direct physical interpretation: it corresponds to the probability that the state ρ is found in the state $|\phi\rangle$ upon measurement.

In this work, we focus on the Bell state

$$|\Psi^+\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}},$$

and define the fidelity of a generic state ρ with respect to this target as

$$F_{\Psi^+}(\rho) = \langle \Psi^+ | \rho | \Psi^+ \rangle.$$

We now analyze this quantity for the phase-evolved Bell-like state introduced in Section 3.4:

$$|\psi(\theta)\rangle = \frac{|01\rangle + e^{i\theta}|10\rangle}{\sqrt{2}},$$

with associated density operator

$$\rho(\theta) = |\psi(\theta)\rangle\langle\psi(\theta)|.$$

The fidelity with respect to $|\Psi^+\rangle$ is given by

$$F_{\Psi^+}(\rho(\theta)) = |\langle \Psi^+ | \psi(\theta) \rangle|^2.$$

A direct computation yields

$$\langle \Psi^+ | \psi(\theta) \rangle = \frac{1}{2}(1 + e^{i\theta}),$$

and therefore

$$F_{\Psi^+}(\rho(\theta)) = \frac{1 + \cos \theta}{2}.$$

This result shows that the relative phase θ modifies the overlap with the reference Bell state, even though the global state remains pure for all values of θ . In particular,

$$F_{\Psi^+} = 1 \quad \text{for } \theta = 0, \quad F_{\Psi^+} = 0 \quad \text{for } \theta = \pi.$$

Thus, fidelity captures variations in global coherence that are not reflected in the purity or in the reduced density operators.

For ensemble states arising from random phase fluctuations, described by

$$\rho_{AB}^{(\text{ens})} = \frac{1}{N} \sum_{k=1}^N |\psi(\theta_k)\rangle\langle\psi(\theta_k)|,$$

the fidelity can be computed using the linearity of the expectation value:

$$F_{\Psi^+}(\rho_{AB}^{(\text{ens})}) = \frac{1}{N} \sum_{k=1}^N F_{\Psi^+}(\rho(\theta_k)).$$

Introducing the ensemble coherence parameter

$$\mu = \langle e^{i\theta} \rangle,$$

one finds

$$F_{\Psi^+} = \frac{1}{2} (1 + \text{Re}(\mu)) = \frac{1}{2} (1 + \langle \cos \theta \rangle).$$

This shows that fidelity directly quantifies the loss of coherence induced by phase fluctuations. In particular, for a uniform distribution over $[0, 2\pi]$, one has $\langle \cos \theta \rangle = 0$, leading to

$$F_{\Psi^+} = \frac{1}{2}.$$

In summary, fidelity provides a quantitative measure of how closely a given state approximates the target Bell state. Unlike purity, which characterizes mixedness, fidelity is sensitive to coherent deformations of the state and therefore plays a central role in the validation and interpretation of the numerical simulations.

Simulation notes. Within the simulator architecture, fidelity is introduced as a general-purpose quantitative tool for comparing quantum states in a basis-independent manner. Given a numerically generated state either in pure vector form or as a density operator, fidelity provides a scalar measure of proximity to a chosen reference state, typically a Bell state in the present framework. From an implementation standpoint, this requires supporting both pure-state overlaps and density-matrix evaluations, allowing the same routine to be applied uniformly across different modeling regimes (deterministic evolution, ensemble averaging, or noisy channels). Unlike correlation observables, which probe specific measurement directions, fidelity captures a global property of the state and therefore serves as a compact diagnostic for validating transformations, monitoring deviations from ideal entanglement, and benchmarking different physical models within the simulator pipeline.

4.4 Concurrence as an Entanglement Measure

Reduced density operators, purity, and fidelity provide complementary information about a bipartite quantum state. Reduced states describe locally accessible information, purity quantifies mixedness, and fidelity measures proximity to a chosen reference state. However, none of these quantities alone provides a direct measure of bipartite entanglement. In particular, fidelity with respect to a Bell state may change under local unitary evolution even when the amount of entanglement remains unchanged. For this reason, a genuine entanglement measure is required.

For two-qubit systems, one of the most widely used entanglement measures is the *concurrence*. It is specifically designed to quantify bipartite entanglement and satisfies the fundamental bounds

$$0 \leq C \leq 1,$$

where

$$C = 0$$

corresponds to a separable state, while

$$C = 1$$

corresponds to a maximally entangled two-qubit state.

We first consider the case of a pure two-qubit state

$$|\psi\rangle = a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle,$$

with complex coefficients satisfying

$$|a|^2 + |b|^2 + |c|^2 + |d|^2 = 1.$$

For such a state, the concurrence is given by

$$C(|\psi\rangle) = 2|ad - bc|.$$

This expression immediately distinguishes separable from entangled states. Indeed, if

$$|\psi\rangle = |\phi_A\rangle \otimes |\phi_B\rangle,$$

then the coefficients satisfy

$$ad - bc = 0,$$

and therefore

$$C(|\psi\rangle) = 0.$$

Conversely, for maximally entangled Bell states the concurrence is equal to one.

For example, for the Bell state

$$|\Psi^+\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}},$$

one has

$$a = 0, \quad b = \frac{1}{\sqrt{2}}, \quad c = \frac{1}{\sqrt{2}}, \quad d = 0,$$

and thus

$$C(|\Psi^+\rangle) = 2 \left| 0 - \frac{1}{2} \right| = 1.$$

The same conclusion holds for the phase-evolved Bell-like state

$$|\psi(\theta)\rangle = \frac{|01\rangle + e^{i\theta}|10\rangle}{\sqrt{2}}.$$

In this case,

$$a = 0, \quad b = \frac{1}{\sqrt{2}}, \quad c = \frac{e^{i\theta}}{\sqrt{2}}, \quad d = 0,$$

so that

$$C(|\psi(\theta)\rangle) = 2 \left| -\frac{e^{i\theta}}{2} \right| = 1.$$

This is an important result for the present model. The relative phase θ changes the overlap with the reference Bell state $|\Psi^+\rangle$, and therefore modifies the fidelity introduced in Section 4.3. However, it does *not* change the amount of entanglement. The state remains maximally entangled for all values of θ . This reflects the fact that the phase evolution considered in the reduced model is generated by local unitary transformations, and bipartite entanglement is invariant under local unitaries.

For pure bipartite states, concurrence is also directly connected to the reduced density operators introduced in Section 4.1 and to the purity discussed in Section 4.2. In particular, if

$$\rho_A = \text{Tr}_B(|\psi\rangle\langle\psi|), \quad \rho_B = \text{Tr}_A(|\psi\rangle\langle\psi|),$$

then one may write

$$C(|\psi\rangle) = 2\sqrt{\det(\rho_A)} = 2\sqrt{\det(\rho_B)},$$

or equivalently,

$$C(|\psi\rangle) = \sqrt{2(1 - \gamma(\rho_A))} = \sqrt{2(1 - \gamma(\rho_B))}.$$

This relation shows that, for pure two-qubit states, entanglement is directly reflected in the mixedness of the reduced states. In particular, if a pure bipartite state is separable, then the reduced states are pure and

$$\gamma(\rho_A) = \gamma(\rho_B) = 1,$$

which implies

$$C = 0.$$

By contrast, for a maximally entangled Bell state,

$$\rho_A = \rho_B = \frac{I}{2},$$

and therefore

$$\gamma(\rho_A) = \gamma(\rho_B) = \frac{1}{2},$$

which yields

$$C = 1.$$

Thus, Sections 4.1 and 4.2 naturally prepare the ground for concurrence: reduced mixedness of a subsystem, when the global state is pure, is a direct signature of bipartite entanglement.

We now turn to the more general case of mixed two-qubit states, described by a density operator

$$\rho \in \mathbb{C}^{4 \times 4}.$$

For mixed states, concurrence is defined through the *spin-flipped* matrix

$$\tilde{\rho} = (\sigma_y \otimes \sigma_y) \rho^* (\sigma_y \otimes \sigma_y),$$

where ρ^* denotes the complex conjugate of ρ in the computational basis. One then considers the eigenvalues of the matrix

$$\rho \tilde{\rho}.$$

Let $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ denote the square roots of these eigenvalues, ordered such that

$$\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4 \geq 0.$$

The concurrence of ρ is then defined as

$$C(\rho) = \max(0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4).$$

This formula provides a complete entanglement measure for arbitrary two-qubit states, both pure and mixed. In the pure-state limit, it reduces to the simpler expression

$$C(|\psi\rangle) = 2|ad - bc|.$$

In the present simulator, this mixed-state formulation becomes essential when the global state is no longer described by a single state vector. This occurs, for example, when one introduces ensemble averaging over random phases,

$$\rho_{AB}^{(\text{ens})} = \frac{1}{N} \sum_{k=1}^N |\psi(\theta_k)\rangle \langle \psi(\theta_k)|,$$

or more generally when propagation effects lead to decoherence and partial loss of phase information.

For the random-phase ensemble considered in this work, the density operator depends on the coherence parameter

$$\mu = \langle e^{i\theta} \rangle,$$

introduced in Sections 4.2 and 4.3. In the computational basis, the ensemble state has the form

$$\rho_{AB}^{(\text{ens})} = \frac{1}{2} |01\rangle \langle 01| + \frac{1}{2} |10\rangle \langle 10| + \frac{\mu}{2} |10\rangle \langle 01| + \frac{\mu^*}{2} |01\rangle \langle 10|.$$

For this family of states, the concurrence can be computed analytically and is found to be

$$C(\rho_{AB}^{(\text{ens})}) = |\mu|.$$

This result is particularly significant because it reveals that the same coherence parameter controlling mixedness and Bell-state fidelity also controls entanglement degradation in the reduced random-phase model. Indeed, from the previous sections we already found that

$$\gamma(\rho_{AB}^{(\text{ens})}) = \frac{1 + |\mu|^2}{2}, \quad F_{\Psi^+} = \frac{1 + \text{Re}(\mu)}{2}.$$

4.5 Interplay Between Purity, Fidelity, and Concurrence

The previous sections introduced three quantitative descriptors of bipartite quantum states: purity, fidelity, and concurrence. Each of these quantities captures a different physical aspect of the state. Purity, defined as $\gamma(\rho) = \text{Tr}(\rho^2)$, characterizes mixedness, fidelity measures similarity to a chosen reference state, and concurrence provides a direct quantification of bipartite entanglement.

In general, these quantities are independent. However, within the random-phase ensemble model considered in this work, they become closely related through a single complex parameter, namely the ensemble coherence

$$\mu = \langle e^{i\theta} \rangle = \langle \cos \theta \rangle + i \langle \sin \theta \rangle.$$

This parameter encapsulates the effect of phase fluctuations on the global quantum state.

For the family of ensemble states introduced in Section 4.4, the relevant quantities (purity, concurrence, and Bell-state fidelity) are all functions of the same coherence parameter μ , but depend on it in qualitatively different ways.

Purity depends on the squared magnitude $|\mu|^2$, and therefore quantifies the overall degree of coherence retained after ensemble averaging through

$$\gamma(\rho) = \frac{1 + |\mu|^2}{2}.$$

Concurrence depends linearly on $|\mu|$, and directly measures the amount of bipartite entanglement present in the state. By contrast, the fidelity with respect to the Bell state $|\Psi^+\rangle$ depends only on the real part $\text{Re}(\mu)$, reflecting the overlap with a specific reference state rather than the total coherence or entanglement.

This distinction has important consequences. Since fidelity depends on the projection of μ onto the real axis, it does not provide a complete characterization of entanglement. In particular, states with the same concurrence may exhibit different fidelities with respect to $|\Psi^+\rangle$. This behavior is clearly observed in the deterministic phase model of Experiment 1, where the parameter μ lies on the unit circle. In that case, the magnitude $|\mu| = 1$ remains constant, and therefore the concurrence is equal to one for all values of θ , while the fidelity varies as a function of the phase.

In contrast, the random-phase ensemble of Experiment 2 leads to a reduction of the magnitude $|\mu|$, reflecting the loss of phase coherence under statistical averaging. As a consequence, both purity and concurrence decrease, with purity quantified by $\gamma(\rho)$. In this regime, the reduction of fidelity is associated with a genuine loss of entanglement, rather than a simple change of phase relative to a fixed reference state.

A useful geometric interpretation can be obtained by representing μ as a point in the complex plane. The concurrence is proportional to the distance of this point from the origin, while the fidelity depends on its projection onto the real axis. Under deterministic phase evolution, μ moves along the unit circle, preserving its magnitude but changing its real part. Under ensemble averaging, μ contracts toward the origin, leading to a simultaneous reduction of coherence, purity (as measured by $\gamma(\rho)$), and entanglement.

The limiting cases illustrate these behaviors clearly. When

$$|\mu| = 1,$$

the ensemble retains full phase coherence, the global state is pure, and therefore

$$\gamma(\rho) = 1,$$

while the concurrence attains its maximal value. When instead

$$|\mu| = 0,$$

all off-diagonal coherence terms vanish, the state reduces to a classical mixture, for which

$$\gamma(\rho) = \frac{1}{2},$$

and the concurrence becomes zero. Intermediate values $0 < |\mu| < 1$ correspond to partially coherent ensembles, characterized by mixed states with reduced but non-vanishing entanglement.

It is important to emphasize that, while purity, fidelity, and concurrence are general concepts in quantum information theory, the simple relations above arise from the specific structure of the random-phase ensemble considered here. In more general situations, these quantities are not necessarily determined by a single parameter.

In summary, the coherence parameter μ provides a unifying description of the ensemble states studied in this work. Its magnitude controls both mixedness and entanglement, while its phase determines the observable correlations and the overlap with a chosen reference state. This interplay clarifies the distinct roles of purity, fidelity, and concurrence, and highlights the necessity of using a genuine entanglement measure to fully characterize bipartite quantum states.

5 Numerical Experiments and Results

5.1 Experiment 1: Deterministic Phase Sweep

5.1.1 Objective

The first experiment investigates the effect of a controlled relative phase θ on a maximally entangled two-qubit state. The objective is to characterize how both correlation observables and state-based quantities evolve under the reduced unitary fiber model.

5.1.2 Model

The initial Bell state is chosen as

$$|\Psi^+\rangle = \frac{1}{\sqrt{2}} (|01\rangle + |10\rangle).$$

After propagation through the effective phase model, the state becomes

$$|\psi(\theta)\rangle = \frac{1}{\sqrt{2}} (|01\rangle + e^{i\theta}|10\rangle).$$

The correlation observables evaluated in the simulator are

$$c_{xx} = \langle \sigma_x \otimes \sigma_x \rangle, \quad c_{yy} = \langle \sigma_y \otimes \sigma_y \rangle, \quad c_{zz} = \langle \sigma_z \otimes \sigma_z \rangle.$$

5.1.3 Analytical Predictions

For the state $|\psi(\theta)\rangle$, direct evaluation gives

$$c_{xx} = \cos \theta, \quad c_{yy} = \cos \theta, \quad c_{zz} = -1.$$

Since the evolution is generated by local unitary transformations, the global state remains pure, i.e.

$$\gamma(\rho(\theta)) = \text{Tr}(\rho(\theta)^2) = 1.$$

The reduced density operators remain maximally mixed:

$$\rho_A(\theta) = \rho_B(\theta) = \frac{I}{2}, \quad \gamma(\rho_A) = \gamma(\rho_B) = \text{Tr}(\rho_A^2) = \text{Tr}(\rho_B^2) = \frac{1}{2}.$$

The fidelity with respect to the reference Bell state $|\Psi^+\rangle$ is

$$F_{\Psi^+}(\rho(\theta)) = \frac{1 + \cos \theta}{2}.$$

Finally, the concurrence of the state is constant and equal to

$$C(\rho(\theta)) = 1,$$

reflecting the fact that bipartite entanglement is invariant under local unitary transformations.

5.1.4 Numerical Results

The numerical experiment performs a sweep over $\theta \in [0, 2\pi]$, computing both correlation observables and state-characterization quantities.




Figure 1: Comparison between numerical simulation and analytical predictions for the correlation observables c_{xx} , c_{yy} , and c_{zz} as functions of the relative phase θ . The agreement confirms the correctness of the reduced phase model implementation.




Figure 2: Numerical and analytical comparison of state-characterization quantities for the phase-evolved state. The figure shows global purity $\gamma(\rho_{AB})$, reduced purities $\gamma(\rho_A)$ and $\gamma(\rho_B)$, fidelity with respect to $|\Psi^+\rangle$, and concurrence. While fidelity varies with θ , concurrence remains constant and equal to one.

5.1.5 Discussion

The numerical results are in full agreement with the analytical predictions:

- c_{xx} and c_{yy} follow the expected cosine dependence,
- c_{zz} remains constant at -1 ,
- the global purity remains equal to $\gamma(\rho_{AB}) = 1$,
- the reduced purities remain equal to $\gamma(\rho_A) = \gamma(\rho_B) = 1/2$,
- the fidelity follows $F_{\Psi^+} = (1 + \cos \theta)/2$,
- the concurrence remains constant at $C = 1$.

These results confirm that the fiber acts as an effective relative phase shifter. While the global state remains pure and maximally entangled, both the observable correlations and the fidelity with respect to a fixed Bell reference vary continuously with the phase.

A key observation is that fidelity and concurrence exhibit fundamentally different behaviors. The fidelity depends explicitly on the phase θ , reflecting the overlap with a fixed reference state, whereas the concurrence remains invariant. This highlights that fidelity does not provide a direct measure of entanglement.

Therefore, although the phase evolution modifies observable correlations and the apparent similarity to a chosen Bell state, it does not alter the underlying entanglement structure. This demonstrates that purity (quantified by $\gamma(\rho)$) and fidelity alone are insufficient to characterize bipartite entanglement, and motivates the introduction of concurrence as a genuine entanglement measure.

5.2 Experiment 2: Random Phase Ensemble

5.2.1 Objective

The second experiment extends the deterministic phase model by introducing random phase realizations θ_k , modeling stochastic birefringence effects in realistic optical fibers. Instead of a single pure state, the system is described by an ensemble

$$\{|\psi(\theta_k)\rangle\}_{k=1}^N,$$

and by the corresponding averaged density operator.

The objective is to characterize how ensemble averaging modifies global coherence, correlation observables, purity, fidelity, and entanglement.