

1 Chapter 4: Exercises, Part II

Consider the Laguerre polynomials

$$L_n^{(\alpha)}(x) = \frac{(-1)^n}{n!} x^n + O(x^{n-1})$$

where $\alpha > -1$, which are orthogonal with respect to

$$\langle f, g \rangle_\alpha = \int_0^\infty f(x)g(x)x^\alpha e^{-x} dx.$$

1. Show that the Rodrigues formula holds:

$$L_n^{(\alpha)}(x) = \frac{x^{-\alpha} e^x}{n!} \frac{d^n}{dx^n} [x^{\alpha+n} e^{-x}].$$

In other words, prove that $L_n^{(\alpha)}(x)$ (defined by the Rodrigues formula)

- (i) is a polynomial of degree exactly n
- (ii) is orthogonal to all lower degree polynomials
- (iii) has leading coefficient $\frac{(-1)^n}{n!}$

Hints: For (i) and (iii), it may be helpful first to prove that

$$\frac{d}{dx} [x^{\alpha+1} e^{-x} L_n^{(\alpha+1)}(x)] = (n+1)x^\alpha e^{-x} L_{n+1}^{(\alpha)}(x)$$

For (ii), use integration by parts.

2. Show that

$$(n+1)L_{n+1}^{(\alpha)}(x) = (\alpha+n+1)L_n^{(\alpha)}(x) - xL_n^{(\alpha+1)}(x)$$

3. Show that

$$L_n^{(\alpha+1)}(x) = L_{n-1}^{(\alpha+1)}(x) + L_n^{(\alpha)}(x).$$

4. Show that the Laguerre polynomials satisfy the following three-term recurrence:

$$xL_n^{(\alpha)}(x) = -(n+\alpha)L_{n-1}^{(\alpha)}(x) + (2n+\alpha+1)L_n^{(\alpha)}(x) - (n+1)L_{n+1}^{(\alpha)}(x)$$

5. Prove that

$$\frac{dL_n^{(\alpha)}}{dx} = -L_{n-1}^{(\alpha+1)}(x)$$

6. Let

$$L^{(\alpha)}(x) = \left[L_0^{(\alpha)}(x) |L_1^{(\alpha)}(x)| \cdots \right].$$

Give matrices D_α and S_α such that

$$\frac{d}{dx} L^{(\alpha)}(x) = L^{(\alpha+1)}(x) \mathcal{D}_\alpha \quad \text{and} \quad L^{(\alpha)}(x) = L^{(\alpha+1)}(x) S_\alpha$$

7. Consider the advection equation on the half line:

$$u_t + u_x = 0, \quad x \in [0, \infty), \quad t \geq 0.$$

Suppose the solution has an expansion of the form

$$u(x, t) = e^{-x/2} \sum_{k=0}^{\infty} u_k(t) L_k^{(0)}(x) = e^{-x/2} L^{(0)}(x) \mathbf{u}(t)$$

where

$$\mathbf{u}(t) = \begin{bmatrix} u_0(t) \\ u_1(t) \\ \vdots \end{bmatrix}.$$

Show that

$$\mathbf{u}'(t) = A \mathbf{u}(t),$$

where A is a matrix that is expressible in terms of \mathcal{D}_0 and \mathcal{S}_0 (defined in question 6). Use software of your choice to build a 10×10 version of the matrix A .