

Renormalization Group Evolution of the Standard Model Dimension Six Operators II: Yukawa Dependence

Elizabeth E. Jenkins,^a Aneesh V. Manohar,^a Michael Trott^{b,1}

^a*Department of Physics, University of California at San Diego, 9500 Gilman Drive,
La Jolla, CA 92093-0319, USA*

^b*Theory Division, Physics Department, CERN, CH-1211 Geneva 23, Switzerland*

E-mail: ejenkins@ucsd.edu, amanohar@ucsd.edu, michael.trott@cern.ch

ABSTRACT: We calculate the complete order y^2 and y^4 terms of the 59×59 one-loop anomalous dimension matrix for the dimension-six operators of the Standard Model effective field theory, where y is a generic Yukawa coupling. These terms, together with the terms of order λ , λ^2 and λy^2 depending on the Standard Model Higgs self-coupling λ which were calculated in a previous work, yield the complete one-loop anomalous dimension matrix in the limit of vanishing gauge couplings. The Yukawa contributions result in non-trivial flavor mixing in the various operator sectors of the Standard Model effective theory.

¹Corresponding author.

Contents

1	Introduction	1
2	Formalism	3
3	Discussion	5
4	Conclusion	6
A	Results	7
A.1	X^3	8
A.2	H^6	8
A.3	$H^4 D^2$	8
A.4	$X^2 H^2$	8
A.5	$\psi^2 H^3$	9
A.6	$\psi^2 X H$	10
A.7	$\psi^2 H^2 D$	11
A.8	ψ^4	12
A.8.1	$(\overline{L}L)(\overline{L}L)$	12
A.8.2	$(\overline{R}R)(\overline{R}R)$	14
A.8.3	$(\overline{L}L)(\overline{R}R)$	15
A.8.4	$(\overline{L}R)(\overline{R}L)$	18
A.8.5	$(\overline{L}R)(\overline{L}R)$	18

1 Introduction

The LHC has discovered a Higgs-like boson with properties consistent with Standard Model (SM) predictions. In addition, the SM provides a successful description of experimental data up to energies $v = 246 \text{ GeV}$, the scale of electroweak symmetry breaking, and there is no evidence thus far for any additional particles beyond the SM. It is important to study the properties of the Higgs boson to high precision, and to increase the energy at the LHC to search for additional particles or phenomena at higher energy. A widely used approach to studying new physics in light of the current experimental situation is to generalize the SM to an effective field theory (EFT) by adding higher dimensional (non-renormalizable) operators constructed out of SM fields to the SM Lagrangian. This approach implicitly assumes that $SU(2)_L \times U(1)_Y$ is a linearly realized symmetry in the scalar sector, which is an assumption we adopt in this work. The non-renormalizable operators are suppressed by an energy scale

$\Lambda > v$, and they parametrize the low-energy effects of new physics at energies above Λ . In the effective field theory approach, higher dimensional operators yield effects which are ordered in a power series expansion in E/Λ . Operators of mass dimension d in the SM EFT yield effects which are order $(E/\Lambda)^{d-4}$. Thus, the largest effects for $E < \Lambda$ arise from the non-renormalizable operators of smallest mass dimension.

The leading operators which preserve lepton number first arise at dimension six, and have been classified in Refs. [1, 2]. There are 59 independent dimension-six operators which preserve baryon number after redundant operators have been eliminated by field redefinitions, or equivalently, by using the equations of motion (EOM). These 59 operators divide into eight operator classes, labelled by their field content and number of covariant derivatives. Denoting gauge field strengths by $X = G_{\mu\nu}, W_{\mu\nu}, B_{\mu\nu}$, the Higgs doublet scalar field by H , fermion fields by $\psi = q, u, d, l, e$, and covariant derivatives by D , the eight operator classes are 1 : X^3 , 2 : H^6 , 3 : $H^4 D^2$, 4 : $X^2 H^2$, 5 : $\psi^2 H^3$, 6 : $\psi^2 X H$, 7 : $\psi^2 H^2 D$ and 8 : ψ^4 . Since we make extensive use of these operators, we list them again in Table 1.

The anomalous dimensions of the dimension-six operators enter into Higgs phenomenology. In Ref. [3], we computed the anomalous dimension matrix of the eight $X^2 H^2$ operators which contribute to $h \rightarrow \gamma\gamma$, $h \rightarrow \gamma Z$ and $gg \rightarrow h$, which are crucial processes for precision Higgs experiments, and in [4] an exactly solvable model was constructed that generates precisely these operators. In Ref. [5], we embarked on the calculation of the complete one-loop anomalous dimension matrix for the 59 independent dimension-six operators. In addition, we calculated *all* contributions to the running of the $d \leq 4$ SM parameters from the 59 independent dimension-six operators. The contributions to the running of SM parameters from dimension-six operators is order v^2/Λ^2 , which is as important as the tree-level contribution of dimension-six operators.

The present paper continues our computation of the one-loop anomalous dimension matrix of the dimension-six operators. This matrix, with $59 \times 59 = 3481$ entries, can be broken into block submatrices γ_{ij} where $i, j = 1, \dots, 8$ label the eight operator classes. In this paper, we present the Yukawa terms, leaving the gauge terms for a subsequent publication. The Yukawa terms contribute to flavor-changing processes. The anomalous dimensions we compute can give flavor-changing Higgs couplings to fermions, and they can lead to rare decays such as $\mu \rightarrow e\gamma$. The anomalous dimension calculation involves a large number (~ 100) one-loop diagrams, a selection of which are shown in Fig. 1. Each graph is simple to compute, but computing the full flavor dependence is tedious. For example, a single graph in Fig. 1 for the $\psi^4 - \psi^4$ mixing of four-fermion operators into themselves leads to a set of anomalous dimensions that occupy seven pages of the appendix.

Some aspects of operator mixing of dimension-six operators due to Yukawa couplings has been previously calculated in Refs. [6–11]. In particular, Ref. [11] recently studied operator mixing from a different viewpoint, and calculated, for a single flavor, some entries in rows 3, 5 and 6 of the anomalous dimension matrix, including the gauge dependence.




Figure 1. A selection of graphs contributing to the one-loop anomalous dimension matrix. The solid square is a $\mathcal{L}^{(6)}$ vertex, and the dots are SM vertices. Some graphs containing gauge fields contribute to the anomalous dimension matrix in the limit $g^2 \rightarrow 0$ in the rescaled operator basis used for power counting. The first graph leads to anomalous dimension contributions seven pages long when all flavor combinations are evaluated.

2 Formalism

The Lagrangian of the SM EFT is given by

$$\mathcal{L} = \mathcal{L}_{SM} + \sum_{d>4} \mathcal{L}^{(d)}, \quad (2.1)$$

where

$$\begin{aligned} \mathcal{L}_{\text{SM}} = & -\frac{1}{4}G_{\mu\nu}^A G^{A\mu\nu} - \frac{1}{4}W_{\mu\nu}^I W^{I\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} + (D_\mu H^\dagger)(D^\mu H) + \sum_{\psi=q,u,d,l,e} \bar{\psi} i \not{D} \psi \\ & - \lambda \left(H^\dagger H - \frac{1}{2}v^2 \right)^2 - \left[H^{\dagger j} \bar{d} Y_d q_j + \tilde{H}^{\dagger j} \bar{u} Y_u q_j + H^{\dagger j} \bar{e} Y_e l_j + \text{h.c.} \right], \end{aligned} \quad (2.2)$$

and $\mathcal{L}^{(d)}$ denotes terms in the effective Lagrangian of mass dimension d . In this work, we restrict our attention to the dimension-six Lagrangian

$$\mathcal{L}^{(6)} = \sum_i C_i Q_i, \quad (2.3)$$

where the sum is over the 59 independent operators Q_i given in Table 1. Each independent operator Q_i appears in the sum with a corresponding operator coefficient C_i , which is proportional to $1/\Lambda^2$.

The one-loop anomalous dimension matrix of the dimension-six operators has entries which are proportional to gauge couplings g , the Higgs scalar doublet self-coupling λ and Yukawa couplings y . The coupling constant dependence of the anomalous dimension entries simplifies if the operators are normalized using naive dimensional analysis (NDA) [12], and a factor of g is absorbed into each field-strength tensor X and a factor of y is absorbed into the single chirality-flip operators $\psi^2 H^3$ and $\psi^2 X H$. In terms of these rescaled operators \tilde{Q}_i , the anomalous dimension entries have the form

$$\gamma \propto \left(\frac{g^2}{16\pi^2} \right)^{n_g} \left(\frac{\lambda}{16\pi^2} \right)^{n_\lambda} \left(\frac{y^2}{16\pi^2} \right)^{n_y}, \quad N = n_g + n_\lambda + n_y, \quad (2.4)$$

where N is the perturbative order of the anomalous dimension. The form Eq. (2.4), and a formula for N in terms of NDA weights w_i of the rescaled operators, was derived in Ref. [13]. At one-loop order, $N = 0, 1, 2, 3, 4$, so that entries in γ range from perturbative order $N = 0$, i.e. effectively “tree-level order,” to perturbative order $N = 4$, i.e. effectively “four-loop order.” The form of the one-loop anomalous dimension matrix for the rescaled operators is given in Table 1 of Ref. [13].

The rescaled operators \tilde{Q}_i give the simple form Eq. (2.4) for the coupling constant dependence of γ , which is useful to classify the terms in γ . However, the actual results given below are for the original unrescaled operator basis Q_i of Ref. [2]. Ref. [3] calculated the complete result for γ_{44} . Ref. [5] performed a complete classification of the entire anomalous dimension matrix, giving the allowed values of n_g, n_λ, n_y for all possible terms. There are non-zero entries in γ for which no one-particle irreducible graph exists, arising from operator conversions using the EOM. Ref. [5] calculated the order λ , λ^2 and λy^2 terms of the one-loop anomalous dimension matrix, i.e. entries with $n_\lambda \geq 1$ and $n_g = 0$. These anomalous dimensions had either $n_y = 0$ or $n_y = 1$. In this paper, we calculate the order y^2 and y^4 terms, which are the remaining terms with $n_g = 0$. The terms with $n_g \neq 0$ will be discussed in a subsequent publication.

3 Discussion

The anomalous dimensions are lengthy, and are given in the appendix. Here, we briefly comment on some of the results.

The dimension-six operators alter the formulae of SM observables at tree level. For example, the Higgs doublet coupling to leptons in the SM EFT involves the Lagrangian terms

$$\begin{aligned}\mathcal{L} &= -[Y_e]_{rs} H^{\dagger j} \bar{e}_r l_{sj} + C_{eH}^* (H^{\dagger} H) H^{\dagger j} \bar{e}_r l_{sj} + h.c. \\ &= -\frac{1}{\sqrt{2}} (v + h) \bar{e}_{Rr} Y_e e_{Ls} + \frac{1}{2\sqrt{2}} (v + h)^3 C_{eH}^* \bar{e}_{Rr} e_{Ls} + h.c.,\end{aligned}\quad (3.1)$$

where the second line has been evaluated in the spontaneously broken theory. Consequently, the charged lepton mass matrix

$$\frac{1}{\sqrt{2}} v \left([Y_e]_{rs} - \frac{1}{2} v^2 C_{eH}^* \right) \quad (3.2)$$

and the Higgs-charged lepton coupling matrix

$$\frac{1}{\sqrt{2}} \left([Y_e]_{rs} - \frac{3}{2} v^2 C_{eH}^* \right) \quad (3.3)$$

depend on different combinations of the usual SM $d = 4$ Yukawa matrix Y_e and the dimension-six terms $v^2 C_{eH}^*$. The two combinations are not simultaneously diagonalizable in general, which can lead to flavor-changing Higgs couplings to fermions. Keeping only the top-Yukawa coupling contributions to the C_{eH} anomalous dimension Eq. (A.18),

$$\dot{C}_{eH}^* = 4 C_{lequ}^{(1)} N_c [Y_u Y_u^\dagger Y_u]_{pt} + \dots, \quad (3.4)$$

where \dot{C} is defined in Eq. (A.1), we see that flavor violation from the four-fermion operator $Q_{lequ}^{(1)} = (\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)$ can feed into the lepton Yukawa couplings. Similar flavor-mixing terms are also present in the quark sector. These flavor-mixing effects, which depend on the flavor structure of the dimension-six operator coefficients, need not be suppressed by SM Yukawa couplings, and could be phenomenologically interesting. For some recent phenomenological studies of bounds on such Higgs related flavor violation, see Refs.[14, 15].

The lepton dipole operators also get contributions from ψ^4 operators,

$$\begin{aligned}\dot{C}_{eW}^* &= -2g_2 N_c C_{lequ}^{(3)} [Y_u]_{tp} + \dots, \\ \dot{C}_{eB}^* &= 4g_1 N_c (y_u + y_q) C_{lequ}^{(3)} [Y_u]_{tp} + \dots.\end{aligned}\quad (3.5)$$

Such a mixing of “tree-generated” into “loop-generated” operators was recently argued to vanish in general, but it is non-zero by explicit computation. (See Refs. [5, 16] for more discussion on related issues.) The linear combination

$$\mathcal{C}_{e\gamma} = \frac{1}{g_1} C_{eB}^* - \frac{1}{g_2} C_{eW}^* \quad (3.6)$$

is the coefficient of the electromagnetic dipole operator

$$e \mathcal{C}_{e\gamma} F_{\mu\nu} \bar{l}_r \sigma^{\mu\nu} e_s H, \quad (3.7)$$

which gives flavor transitions such as $\mu \rightarrow e\gamma$, when H is replaced by its vacuum expectation value $v/\sqrt{2}$. It has the anomalous dimension

$$\dot{\mathcal{C}}_{e\gamma} = [4N_c(y_u + y_q) + 2N_c] C_{lequ}^{(3)} [Y_u]_{tp} + \dots \quad (3.8)$$

The current bound on the $\mu \rightarrow e\gamma$ branching ratio from the MEG experiment [17] is 5.4×10^{-13} , with an order of magnitude improvement in sensitivity expected with a future MEG upgrade. As these sensitive probes of BSM flavor violation increase in precision, incorporating the RGE effects presented in this work will be crucial in correctly interpreting future limits or deviations from SM predictions. Furthermore, if deviations are not found, the non-trivial flavor mixing effects present in the RGEs puts strong constraints on the flavor structure of the SM EFT. One possible interpretation is that it implies the idea of a symmetry-based solution, such as minimal flavor violation [18–20], if BSM physics is not simply decoupled.

4 Conclusion

We have computed the Yukawa terms in the 59×59 one-loop anomalous dimension matrix of the baryon number conserving dimension-six operators in the Standard Model effective field theory. The anomalous dimension matrix of the dimension-six baryon number violating operators have been computed in Ref. [21]. The anomalous dimensions mix the eight different operator classes of dimension-six operators, and the resultant expressions are lengthy. Most of the complications arise from the 25 four-fermion operators (class 8 : ψ^4). There are interesting non-trivial flavor effects which can occur from the flavor structure of the renormalization group evolution equations. Incorporating these RGE effects refines the interpretation of the increasingly strong bounds derived from the many searches for flavor violation beyond the SM.

Acknowledgments

We would like to thank R. Alonso, G. Pruna and A. Signer [22], and B. Pecjak for pointing out some corrections, and C. Bobeth and U. Haisch [23] for confirming some of the equations. This work was supported in part by DOE grant DE-SC0009919.

A Results

The renormalization group equations are given in this appendix. These equations are to be added to Eqs. (4.3), (4.4), (6.1)–(6.4) of Ref. [5]. Eq. (5.6) of Ref. [5], which does not depend on the $\lambda(H^\dagger H)^2$ coupling, is included as part of the results of this paper in Sec. A.6, and should not be added again. The invariants $c_{F,3} = (N_c^2 - 1)/(2N_c)$ and $c_{A,3} = N_c$ are the $SU(3)$ quadratic Casimirs in the fundamental and adjoint representation, respectively; $N_c = 3$ is the number of colors, and $y_{q,l,u,d,e}$ denotes the $U(1)$ hypercharges of the fermions. We use the notation

$$\dot{C} \equiv 16\pi^2 \mu \frac{d}{d\mu} C \quad (\text{A.1})$$

in the renormalization group equations given below. The wavefunction renormalization contributions proportional to Yukawa couplings are written in terms of

$$\begin{aligned} \gamma_l^{(Y)} &= \frac{1}{2} [Y_e^\dagger Y_e]_{rs}, & \gamma_e^{(Y)} &= [Y_e Y_e^\dagger]_{rs}, & \gamma_H^{(Y)} &= \text{Tr} \left[N_c Y_u^\dagger Y_u + N_c Y_d^\dagger Y_d + Y_e^\dagger Y_e \right], \\ \gamma_q^{(Y)} &= \frac{1}{2} [Y_u^\dagger Y_u + Y_d^\dagger Y_d]_{rs}, & \gamma_u^{(Y)} &= [Y_u Y_u^\dagger]_{rs}, & \gamma_d^{(Y)} &= [Y_d Y_d^\dagger]_{rs}, \end{aligned} \quad (\text{A.2})$$

which are $16\pi^2$ times the field anomalous dimensions. The gauge contributions to wavefunction renormalization will be included with the gauge terms for the anomalous dimension matrix, since only the total combination is gauge invariant.

To simplify later expressions, it is useful to define the constants η_{1-5} and $\xi_{e,d,u}$. $\eta_{1,2}$ were already used in Ref. [5].

$$\begin{aligned} \eta_1 &= \frac{1}{2} N_c C_{dH} [Y_d]_{sr} + \frac{1}{2} N_c C_{dH}^* [Y_d^\dagger]_{rs} + \frac{1}{2} N_c C_{uH} [Y_u]_{sr} + \frac{1}{2} N_c C_{uH}^* [Y_u^\dagger]_{rs} + \frac{1}{2} C_{eH} [Y_e]_{sr} + \frac{1}{2} C_{eH}^* [Y_e^\dagger]_{rs}, \\ \eta_2 &= -2 N_c C_{Hq}^{(3)} [Y_u^\dagger Y_u + Y_d^\dagger Y_d]_{sr} + N_c C_{Hud} [Y_d Y_u^\dagger]_{sr} + N_c C_{Hud}^* [Y_u Y_d^\dagger]_{sr} - 2 C_{Hl}^{(3)} [Y_e^\dagger Y_e]_{sr}, \\ \eta_3 &= N_c C_{Hq}^{(1)} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{sr} + 3 N_c C_{Hq}^{(3)} [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{sr} + N_c C_{Hu} [Y_u Y_u^\dagger]_{sr} - N_c C_{Hd} [Y_d Y_d^\dagger]_{sr} \\ &\quad - N_c C_{Hud} [Y_d Y_u^\dagger]_{sr} - N_c C_{Hud}^* [Y_u Y_d^\dagger]_{sr} + (3 C_{Hl}^{(3)} + C_{Hl}^{(1)}) [Y_e^\dagger Y_e]_{sr} - C_{He} [Y_e Y_e^\dagger]_{sr}, \\ \eta_4 &= 4 N_c C_{Hq}^{(1)} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{sr} + 4 N_c C_{Hu} [Y_u Y_u^\dagger]_{sr} - 4 N_c C_{Hd} [Y_d Y_d^\dagger]_{sr} + 2 N_c C_{Hud} [Y_d Y_u^\dagger]_{sr} \\ &\quad + 2 N_c C_{Hud}^* [Y_u Y_d^\dagger]_{sr} + 4 C_{Hl}^{(1)} [Y_e^\dagger Y_e]_{sr} - 4 C_{He} [Y_e Y_e^\dagger]_{sr}, \\ \eta_5 &= -\frac{1}{2} i N_c C_{dH} [Y_d]_{sr} + \frac{1}{2} i N_c C_{dH}^* [Y_d^\dagger]_{rs} + \frac{1}{2} i N_c C_{uH} [Y_u]_{sr} - \frac{1}{2} i N_c C_{uH}^* [Y_u^\dagger]_{rs} - \frac{1}{2} i C_{eH} [Y_e]_{sr} + \frac{1}{2} i C_{eH}^* [Y_e^\dagger]_{rs} \end{aligned} \quad (\text{A.3})$$

$$\begin{aligned}
\xi_{pt}^e &= 2C_{prst}^{le} [Y_e^\dagger]_{rs} - N_c C_{ptsr}^{ledq} [Y_d^\dagger]_{rs} + C_{lequ}^{(1)} N_c [Y_u]_{rs} \\
\xi_d &= 2(C_{qd}^{(1)} + c_{F,3} C_{prst}^{(8)}) [Y_d^\dagger]_{rs} - (N_c C_{srpt}^{(1)} + \frac{1}{2} C_{prst}^{(1)} + \frac{1}{2} c_{F,3} C_{prst}^{(8)}) [Y_u]_{rs} - C_{srtp}^{*ledq} [Y_e^\dagger]_{sr} \\
\xi_u &= 2(C_{qu}^{(1)} + c_{F,3} C_{prst}^{(8)}) [Y_u^\dagger]_{rs} - (N_c C_{ptsr}^{(1)} + \frac{1}{2} C_{stpr}^{(1)} + \frac{1}{2} c_{F,3} C_{stpr}^{(8)}) [Y_d]_{rs} + C_{srtp}^{(1)} [Y_e]_{rs}
\end{aligned} \tag{A.4}$$

The Yukawa contributions to the one-loop renormalization group equations of the 59 dimension-six operator coefficients are listed by operator class in the following eight subsections.

A.1 X^3

$$\dot{C}_G = 0, \quad \dot{C}_{\tilde{G}} = 0, \quad \dot{C}_W = 0, \quad \dot{C}_{\tilde{W}} = 0. \tag{A.5}$$

A.2 H^6

$$\begin{aligned}
\dot{C}_H &= -4N_c C_{rs}^{uH} [Y_u Y_u^\dagger Y_u]_{sr} - 4N_c C_{rs}^{*uH} [Y_u^\dagger Y_u Y_u^\dagger]_{rs} - 4N_c C_{rs}^{dH} [Y_d Y_d^\dagger Y_d]_{sr} - 4N_c C_{rs}^{*dH} [Y_d^\dagger Y_d Y_d^\dagger]_{rs} \\
&\quad - 4C_{rs}^{eH} [Y_e Y_e^\dagger Y_e]_{sr} - 4C_{rs}^{*eH} [Y_e^\dagger Y_e Y_e^\dagger]_{rs} + 6\gamma_H^{(Y)} C_H
\end{aligned} \tag{A.6}$$

A.3 $H^4 D^2$

$$\dot{C}_{H\Box} = -2\eta_3 + 4\gamma_H^{(Y)} C_{H\Box}, \quad \dot{C}_{HD} = -2\eta_4 + 4\gamma_H^{(Y)} C_{HD} \tag{A.7}$$

A.4 $X^2 H^2$

$$\dot{C}_{HG} = -2g_3 (C_{rs}^{uG} [Y_u]_{sr} + [Y_u^\dagger]_{rs} C_{rs}^{*uG}) - 2g_3 (C_{rs}^{dG} [Y_d]_{sr} + [Y_d^\dagger]_{rs} C_{rs}^{*dG}) + 2\gamma_H^{(Y)} C_{HG} \tag{A.8}$$

$$\dot{C}_{H\tilde{G}} = 2g_3 (iC_{rs}^{uG} [Y_u]_{sr} - i[Y_u^\dagger]_{rs} C_{rs}^{*uG}) + 2g_3 (iC_{rs}^{dG} [Y_d]_{sr} - i[Y_d^\dagger]_{rs} C_{rs}^{*dG}) + 2\gamma_H^{(Y)} C_{H\tilde{G}} \tag{A.9}$$

$$\begin{aligned}
\dot{C}_{HW} &= -g_2 (C_{rs}^{eW} [Y_e]_{sr} + [Y_e^\dagger]_{rs} C_{rs}^{*eW}) - g_2 N_c (C_{rs}^{uW} [Y_u]_{sr} + [Y_u^\dagger]_{rs} C_{rs}^{*uW}) \\
&\quad - g_2 N_c (C_{rs}^{dW} [Y_d]_{sr} + [Y_d^\dagger]_{rs} C_{rs}^{*dW}) + 2\gamma_H^{(Y)} C_{HW}
\end{aligned} \tag{A.10}$$

$$\begin{aligned}
\dot{C}_{H\tilde{W}} &= g_2 (iC_{rs}^{eW} [Y_e]_{sr} - i[Y_e^\dagger]_{rs} C_{rs}^{*eW}) + g_2 N_c (iC_{rs}^{uW} [Y_u]_{sr} - i[Y_u^\dagger]_{rs} C_{rs}^{*uW}) \\
&\quad + g_2 N_c (iC_{rs}^{dW} [Y_d]_{sr} - i[Y_d^\dagger]_{rs} C_{rs}^{*dW}) + 2\gamma_H^{(Y)} C_{H\tilde{W}}
\end{aligned} \tag{A.11}$$

$$\begin{aligned}\dot{C}_{HB} = & \left(-2g_1(y_l + y_e)(C_{eB}[Y_e]_{sr} + [Y_e^\dagger]_{rs}C_{eB}^*) - 2g_1N_c(y_q + y_u)(C_{uB}[Y_u]_{sr} + [Y_u^\dagger]_{rs}C_{uB}^*) \right. \\ & \left. - 2g_1N_c(y_q + y_d)(C_{dB}[Y_d]_{sr} + [Y_d^\dagger]_{rs}C_{dB}^*) + 2\gamma_H^{(Y)}C_{HB} \right) \quad (A.12)\end{aligned}$$

$$\begin{aligned}\dot{C}_{H\tilde{B}} = & 2g_1(y_l + y_e)(iC_{eB}[Y_e]_{sr} - i[Y_e^\dagger]_{rs}C_{eB}^*) + 2g_1N_c(y_q + y_u)(iC_{uB}[Y_u]_{sr} - i[Y_u^\dagger]_{rs}C_{uB}^*) \\ & + 2g_1N_c(y_q + y_d)(iC_{dB}[Y_d]_{sr} - i[Y_d^\dagger]_{rs}C_{dB}^*) + 2\gamma_H^{(Y)}C_{H\tilde{B}} \quad (A.13)\end{aligned}$$

$$\begin{aligned}\dot{C}_{HWB} = & \left(-g_2(C_{eB}[Y_e]_{sr} + [Y_e^\dagger]_{rs}C_{eB}^*) + g_2N_c(C_{uB}[Y_u]_{sr} + [Y_u^\dagger]_{rs}C_{uB}^*) - g_2N_c(C_{dB}[Y_d]_{sr} + [Y_d^\dagger]_{rs}C_{dB}^*) \right. \\ & \left. - 2g_1(y_l + y_e)(C_{eW}[Y_e]_{sr} + [Y_e^\dagger]_{rs}C_{eW}^*) + 2g_1N_c(y_q + y_u)(C_{uW}[Y_u]_{sr} + [Y_u^\dagger]_{rs}C_{uW}^*) \right. \\ & \left. - 2g_1N_c(y_q + y_d)(C_{dW}[Y_d]_{sr} + [Y_d^\dagger]_{rs}C_{dW}^*) + 2\gamma_H^{(Y)}C_{HWB} \right) \quad (A.14)\end{aligned}$$

$$\begin{aligned}\dot{C}_{H\tilde{W}B} = & g_2(iC_{eB}[Y_e]_{sr} - i[Y_e^\dagger]_{rs}C_{eB}^*) - g_2N_c(iC_{uB}[Y_u]_{sr} - i[Y_u^\dagger]_{rs}C_{uB}^*) + g_2N_c(iC_{dB}[Y_d]_{sr} - i[Y_d^\dagger]_{rs}C_{dB}^*) \\ & + 2g_1(y_l + y_e)(iC_{eW}[Y_e]_{sr} - i[Y_e^\dagger]_{rs}C_{eW}^*) - 2g_1N_c(y_q + y_u)(iC_{uW}[Y_u]_{sr} - i[Y_u^\dagger]_{rs}C_{uW}^*) \\ & + 2g_1N_c(y_q + y_d)(iC_{dW}[Y_d]_{sr} - i[Y_d^\dagger]_{rs}C_{dW}^*) + 2\gamma_H^{(Y)}C_{H\tilde{W}B} \quad (A.15)\end{aligned}$$

A.5 $\psi^2 H^3$

$$\begin{aligned}\dot{C}_{uH} = & 2(\eta_1 + \eta_2 - i\eta_5)[Y_u^\dagger]_{rs} + [Y_u^\dagger Y_u Y_u^\dagger]_{rs}(C_{HD} - 6C_{H\Box}) - 2C_{Hq}^{(1)}[Y_u^\dagger Y_u Y_u^\dagger]_{ts} + 6C_{Hq}^{(3)}[Y_d^\dagger Y_d Y_u^\dagger]_{ts} \\ & + 2[Y_u^\dagger Y_u Y_u^\dagger]_{rt}C_{Hu} - 2[Y_d^\dagger Y_d Y_d^\dagger]_{rt}C_{Hud}^* + 8(C_{qd}^{(1)} + c_{F,3}C_{qd}^{(8)})[Y_u^\dagger Y_u Y_u^\dagger]_{pt} \\ & - 2(2N_c C_{quqd}^{(1)} + C_{quqd}^{(1)} + c_{F,3}C_{quqd}^{(8)})[Y_d Y_d^\dagger Y_d]_{pt} + 4C_{lequ}^{(1)}[Y_e^\dagger Y_e Y_e^\dagger]_{pt} + 4C_{uH}[Y_u Y_u^\dagger]_{ts} + 5[Y_u^\dagger Y_u]_{rt}C_{uH} \\ & - 2[Y_d^\dagger]_{rt}C_{dH}^*[Y_u^\dagger]_{us} - C_{dH}[Y_d Y_u^\dagger]_{ts} - 2[Y_d^\dagger Y_d]_{rt}C_{uH} + 3\gamma_H^{(Y)}C_{uH} + \gamma_q^{(Y)}C_{uH} + C_{uH}\gamma_u^{(Y)} \\ & \quad (A.16)\end{aligned}$$

$$\begin{aligned}\dot{C}_{dH} = & 2(\eta_1 + \eta_2 + i\eta_5)[Y_d^\dagger]_{rs} + [Y_d^\dagger Y_d Y_d^\dagger]_{rs}(C_{HD} - 6C_{H\Box}) + 2C_{Hq}^{(1)}[Y_d^\dagger Y_d Y_d^\dagger]_{ts} + 6C_{Hq}^{(3)}[Y_u^\dagger Y_u Y_d^\dagger]_{ts} \\ & - 2[Y_d^\dagger Y_d Y_d^\dagger]_{rt}C_{Hd} - 2[Y_u^\dagger Y_u Y_u^\dagger]_{rt}C_{Hud} + 8(C_{qd}^{(1)} + c_{F,3}C_{qd}^{(8)})[Y_d^\dagger Y_d Y_d^\dagger]_{pt} - 4C_{ledq}^*[Y_e^\dagger Y_e Y_e^\dagger]_{pt} \\ & - 2(2N_c C_{quqd}^{(1)} + C_{quqd}^{(1)} + c_{F,3}C_{quqd}^{(8)})[Y_u Y_u^\dagger Y_u]_{pt} + 4C_{dH}[Y_d Y_d^\dagger]_{ts} + 5[Y_d^\dagger Y_d]_{rt}C_{dH} \\ & - 2[Y_u^\dagger]_{rt}C_{uH}^*[Y_d^\dagger]_{us} - C_{uH}[Y_u Y_d^\dagger]_{ts} - 2[Y_u^\dagger Y_u]_{rt}C_{dH} + 3\gamma_H^{(Y)}C_{dH} + \gamma_q^{(Y)}C_{dH} + C_{dH}\gamma_d^{(Y)} \\ & \quad (A.17)\end{aligned}$$

$$\begin{aligned}
\dot{C}_{eH} = & 2(\eta_1 + \eta_2 + i\eta_5)[Y_e^\dagger]_{rs} + [Y_e^\dagger Y_e Y_e^\dagger]_{rs}(C_{HD} - 6C_{H\Box}) + 2C_{Hl}^{(1)}[Y_e^\dagger Y_e Y_e^\dagger]_{ts} - 2[Y_e^\dagger Y_e Y_e^\dagger]_{rt}C_{He} \\
& + 8C_{rpts}^{le}[Y_e^\dagger Y_e Y_e^\dagger]_{pt} - 4C_{rspt}^{ledq}N_c[Y_d^\dagger Y_d Y_d^\dagger]_{tp} + 4C_{rstp}^{(1)}N_c[Y_u Y_u^\dagger Y_u]_{pt} + 4C_{rt}^{eH}[Y_e Y_e^\dagger]_{ts} \\
& + 5[Y_e^\dagger Y_e]_{rt}C_{eH} + 3\gamma_H^{(Y)}C_{eH} + \gamma_l^{(Y)}C_{eH} + C_{eH}\gamma_e^{(Y)}
\end{aligned} \tag{A.18}$$

A.6 $\psi^2 XH$

$$\dot{C}_{eW} = -2g_2 N_c C_{lequ}^{(3)}[Y_u]_{tp} + C_{eW}[Y_e Y_e^\dagger]_{ts} + \gamma_H^{(Y)}C_{eW} + \gamma_l^{(Y)}C_{eW} + C_{eW}\gamma_e^{(Y)} \tag{A.19}$$

$$\dot{C}_{eB} = 4g_1 N_c (y_u + y_q) C_{lequ}^{(3)}[Y_u]_{tp} + C_{eB}[Y_e Y_e^\dagger]_{ts} + 2[Y_e^\dagger Y_e]_{rt}C_{eB} + \gamma_H^{(Y)}C_{eB} + \gamma_l^{(Y)}C_{eB} + C_{eB}\gamma_e^{(Y)} \tag{A.20}$$

$$\dot{C}_{uG} = 2[Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{rt}C_{uG} - C_{dG}[Y_d Y_u^\dagger]_{ts} + C_{uG}[Y_u Y_u^\dagger]_{ts} + \gamma_H^{(Y)}C_{uG} + \gamma_q^{(Y)}C_{uG} + C_{uG}\gamma_u^{(Y)} \tag{A.21}$$

$$\begin{aligned}
\dot{C}_{uW} = & -2g_2 C_{lequ}^{(3)}[Y_e]_{pt} + 2[Y_d^\dagger Y_d]_{rt}C_{uW} - C_{dW}[Y_d Y_u^\dagger]_{ts} + C_{uW}[Y_u Y_u^\dagger]_{ts} \\
& + \gamma_H^{(Y)}C_{uW} + \gamma_q^{(Y)}C_{uW} + C_{uW}\gamma_u^{(Y)}
\end{aligned} \tag{A.22}$$

$$\begin{aligned}
\dot{C}_{uB} = & 4g_1 (y_e + y_l) C_{lequ}^{(3)}[Y_e]_{pt} + 2[Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{rt}C_{uB} - C_{dB}[Y_d Y_u^\dagger]_{ts} + C_{uB}[Y_u Y_u^\dagger]_{ts} \\
& + \gamma_H^{(Y)}C_{uB} + \gamma_q^{(Y)}C_{uB} + C_{uB}\gamma_u^{(Y)}
\end{aligned} \tag{A.23}$$

$$\dot{C}_{dG} = -2[Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{rt}C_{dG} - C_{uG}[Y_u Y_d^\dagger]_{ts} + C_{dG}[Y_d Y_d^\dagger]_{ts} + \gamma_H^{(Y)}C_{dG} + \gamma_q^{(Y)}C_{dG} + C_{dG}\gamma_d^{(Y)} \tag{A.24}$$

$$\dot{C}_{dW} = 2[Y_u^\dagger Y_u]_{rt}C_{dW} - C_{uW}[Y_u Y_d^\dagger]_{ts} + C_{dW}[Y_d Y_d^\dagger]_{ts} + \gamma_H^{(Y)}C_{dW} + \gamma_q^{(Y)}C_{dW} + C_{dW}\gamma_d^{(Y)} \tag{A.25}$$

$$\dot{C}_{dB} = -2[Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{rt}C_{dB} - C_{uB}[Y_u Y_d^\dagger]_{ts} + C_{dB}[Y_d Y_d^\dagger]_{ts} + \gamma_H^{(Y)}C_{dB} + \gamma_q^{(Y)}C_{dB} + C_{dB}\gamma_d^{(Y)} \tag{A.26}$$

A.7 $\psi^2 H^2 D$

$$\begin{aligned}
\dot{C}_{Hq}^{(1)} = & \frac{1}{2} [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{pr} (C_{H\Box} + C_{HD}) - [Y_u^\dagger]_{ps} C_{Hu} [Y_u]_{tr} - [Y_d^\dagger]_{ps} C_{Hd} [Y_d]_{tr} + 2C_{prst}^{qe} [Y_e Y_e^\dagger]_{ts} - 2C_{stpr}^{(1)lq} [Y_e^\dagger Y_e]_{ts} \\
& + \frac{3}{2} [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{pt} C_{Hq}^{(1)tr} + \frac{3}{2} C_{Hq}^{(1)pt} [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{tr} + \frac{9}{2} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{pt} C_{Hq}^{(3)tr} + \frac{9}{2} C_{Hq}^{(3)pt} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{tr} \\
& - (2N_c C_{prst}^{(1)qq} + 2N_c C_{stpr}^{(1)qq} + C_{ptsr}^{(1)qq} + C_{srpt}^{(1)qq} + 3C_{ptsr}^{(3)qq} + 3C_{srpt}^{(3)qq}) [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{ts} \\
& - 2N_c C_{prst}^{(1)qu} [Y_u Y_u^\dagger]_{ts} + 2N_c C_{stpr}^{(1)qd} [Y_d Y_d^\dagger]_{ts} + 2\gamma_H^{(Y)} C_{pr}^{(1)Hq} + \gamma_{pt}^{(Y)} C_{pt}^{(1)Hq} + C_{tr}^{(1)Hq} \gamma_{tr}^{(Y)} \quad (A.27)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{Hq}^{(3)} = & -\frac{1}{2} [Y_u^\dagger Y_u + Y_d^\dagger Y_d]_{pr} C_{H\Box} + \frac{3}{2} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{pt} C_{Hq}^{(1)tr} + \frac{3}{2} C_{Hq}^{(1)pt} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{tr} \\
& + \frac{1}{2} [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{pt} C_{Hq}^{(3)tr} + \frac{1}{2} C_{Hq}^{(3)pt} [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{tr} \\
& - (2N_c C_{prst}^{(3)qq} + 2N_c C_{stpr}^{(3)qq} + C_{ptsr}^{(1)qq} + C_{srpt}^{(1)qq} - C_{ptsr}^{(3)qq} - C_{srpt}^{(3)qq}) [Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{ts} - 2C_{stpr}^{(3)lq} [Y_e^\dagger Y_e]_{ts} \\
& + 2\gamma_H^{(Y)} C_{pr}^{(3)Hq} + \gamma_{pt}^{(Y)} C_{pt}^{(3)Hq} + C_{tr}^{(3)Hq} \gamma_{tr}^{(Y)} \quad (A.28)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{Hd} = & [Y_d Y_d^\dagger]_{pr} (C_{H\Box} + C_{HD}) - 2[Y_d]_{ps} C_{Hq}^{(1)st} [Y_d^\dagger]_{tr} + 3[Y_d Y_d^\dagger]_{pt} C_{Hd} + 3C_{pt} [Y_d Y_d^\dagger]_{tr} - [Y_d Y_u^\dagger]_{pt} C_{Hud} \\
& - C_{tp}^* [Y_u Y_d^\dagger]_{tr} + 2(N_c C_{prst}^{dd} + N_c C_{stpr}^{dd} + C_{ptsr}^{dd} + C_{srpt}^{dd}) [Y_d Y_d^\dagger]_{ts} + 2C_{stpr}^{ed} [Y_e Y_e^\dagger]_{ts} - 2C_{stpr}^{ld} [Y_e^\dagger Y_e]_{ts} \\
& - 2N_c C_{stpr}^{(1)ud} [Y_u Y_u^\dagger]_{ts} - 2N_c C_{stpr}^{(1)qd} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{ts} + 2\gamma_H^{(Y)} C_{pr}^{Hd} + \gamma_{pt}^{(Y)} C_{pt}^{Hd} + C_{tr}^{Hd} \gamma_{tr}^{(Y)} \quad (A.29)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{Hu} = & -[Y_u Y_u^\dagger]_{pr} (C_{H\Box} + C_{HD}) - 2[Y_u]_{ps} C_{Hq}^{(1)st} [Y_u^\dagger]_{tr} + 3[Y_u Y_u^\dagger]_{pt} C_{Hu} + 3C_{pt} [Y_u Y_u^\dagger]_{tr} + [Y_u Y_d^\dagger]_{pt} C_{Hud}^* \\
& + C_{Hud} [Y_d Y_u^\dagger]_{tr} - 2(N_c C_{prst}^{uu} + N_c C_{stpr}^{uu} + C_{ptsr}^{uu} + C_{srpt}^{uu}) [Y_u Y_u^\dagger]_{ts} + 2C_{stpr}^{eu} [Y_e Y_e^\dagger]_{ts} - 2C_{stpr}^{lu} [Y_e^\dagger Y_e]_{ts} \\
& + 2N_c C_{prst}^{(1)ud} [Y_d Y_d^\dagger]_{ts} - 2N_c C_{stpr}^{(1)qu} [Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{ts} + 2\gamma_H^{(Y)} C_{pr}^{Hu} + \gamma_{pt}^{(Y)} C_{pt}^{Hu} + C_{tr}^{Hu} \gamma_{tr}^{(Y)} \quad (A.30)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{Hud} = & [Y_u Y_d^\dagger]_{pr} (2C_{H\Box} - C_{HD}) - 2[Y_u Y_d^\dagger]_{pt} C_{Hd} + 2C_{pt} [Y_u Y_d^\dagger]_{tr} + 4(C_{ptsr}^{(1)ud} + c_F 3C_{ptsr}^{(8)ud}) [Y_u Y_d^\dagger]_{ts} \\
& + 2[Y_u Y_u^\dagger]_{pt} C_{Hud} + 2C_{pt} [Y_d Y_d^\dagger]_{tr} + 2\gamma_H^{(Y)} C_{pr}^{Hud} + \gamma_{pt}^{(Y)} C_{pt}^{Hud} + C_{tr}^{Hud} \gamma_{tr}^{(Y)} \quad (A.31)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{Hl}^{(1)} = & -\frac{1}{2}[Y_e^\dagger Y_e]_{pr}(C_{H\Box} + C_{HD}) - [Y_e^\dagger]_{ps}C_{He}^{st}[Y_e]_{tr} + \frac{3}{2}[Y_e^\dagger Y_e]_{pt}(C_{Hl}^{(1)} + 3C_{Hl}^{(3)}) + \frac{3}{2}(C_{Hl}^{(1)} + 3C_{Hl}^{(3)})[Y_e^\dagger Y_e]_{tr} \\
& + 2C_{le}^{prst}[Y_e Y_e^\dagger]_{ts} + (-2C_{ll}^{prst} - 2C_{ll}^{stpr} - C_{ll}^{ptsr} - C_{ll}^{srpt})[Y_e^\dagger Y_e]_{ts} - 2N_c C_{lq}^{(1)}[Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{ts} \\
& - 2N_c C_{lu}^{prst}[Y_u Y_u^\dagger]_{ts} + 2N_c C_{ld}^{prst}[Y_d Y_d^\dagger]_{ts} + 2\gamma_H^{(Y)}C_{Hl}^{(1)} + \gamma_l^{(Y)}C_{Hl}^{(1)} + C_{Hl}^{(1)}\gamma_l^{(Y)} \quad (A.32)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{Hl}^{(3)} = & -\frac{1}{2}[Y_e^\dagger Y_e]_{pr}C_{H\Box} + \frac{1}{2}[Y_e^\dagger Y_e]_{pt}(3C_{Hl}^{(1)} + C_{Hl}^{(3)}) + \frac{1}{2}(3C_{Hl}^{(1)} + C_{Hl}^{(3)})[Y_e^\dagger Y_e]_{tr} \\
& - (C_{ll}^{ptsr} + C_{ll}^{srpt})[Y_e^\dagger Y_e]_{ts} - 2N_c C_{lq}^{(3)}[Y_d^\dagger Y_d + Y_u^\dagger Y_u]_{ts} + 2\gamma_H^{(Y)}C_{Hl}^{(3)} + \gamma_l^{(Y)}C_{Hl}^{(3)} + C_{Hl}^{(3)}\gamma_l^{(Y)} \quad (A.33)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{He} = & [Y_e Y_e^\dagger]_{pr}(C_{H\Box} + C_{HD}) - 2[Y_e]_{ps}C_{Hl}^{st}[Y_e^\dagger]_{tr} + 3[Y_e Y_e^\dagger]_{pt}C_{He}^{tr} + 3C_{He}^{pt}[Y_e Y_e^\dagger]_{tr} \\
& - 2C_{le}^{stpr}[Y_e^\dagger Y_e]_{ts} + 2(C_{ee}^{prst} + C_{ee}^{stpr} + C_{ee}^{ptsr} + C_{ee}^{srpt})[Y_e Y_e^\dagger]_{ts} - 2N_c C_{eu}^{prst}[Y_u Y_u^\dagger]_{ts} \\
& + 2N_c C_{ed}^{prst}[Y_d Y_d^\dagger]_{ts} - 2N_c C_{eq}^{stpr}[Y_d^\dagger Y_d - Y_u^\dagger Y_u]_{ts} + 2\gamma_H^{(Y)}C_{He}^{pr} + \gamma_e^{(Y)}C_{He}^{pt} + C_{He}^{tr}\gamma_e^{(Y)} \quad (A.34)
\end{aligned}$$

A.8 ψ^4

A.8.1 $(\bar{L}L)(\bar{L}L)$

$$\begin{aligned}
\dot{C}_{ll}^{prst} = & -\frac{1}{2}[Y_e^\dagger Y_e]_{pr}C_{Hl}^{(1)} - \frac{1}{2}[Y_e^\dagger Y_e]_{st}C_{Hl}^{(1)} + \frac{1}{2}[Y_e^\dagger Y_e]_{pr}C_{Hl}^{(3)} + \frac{1}{2}[Y_e^\dagger Y_e]_{st}C_{Hl}^{(3)} \\
& - [Y_e^\dagger Y_e]_{sr}C_{Hl}^{(3)} - [Y_e^\dagger Y_e]_{pt}C_{Hl}^{(3)} - \frac{1}{2}[Y_e^\dagger]_{sv}[Y_e]_{wt}C_{prvw}^{le} - \frac{1}{2}[Y_e^\dagger]_{pv}[Y_e]_{wr}C_{stvw}^{le} \\
& + \gamma_l^{(Y)}C_{pv}^{ll} + \gamma_l^{(Y)}C_{sv}^{ll} + C_{pv}^{ll}\gamma_l^{(Y)} + C_{prsv}^{ll}\gamma_l^{(Y)} \quad (A.35)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{prst}^{(1)qq} = & \frac{1}{2} [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{pr} C_{st}^{(1)Hq} + \frac{1}{2} [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{st} C_{pr}^{(1)Hq} \\
& + \frac{1}{4N_c} \left([Y_u^\dagger]_{pv} [Y_u]_{wr} C_{stvw}^{(8)qu} + [Y_u^\dagger]_{sv} [Y_u]_{wt} C_{prvw}^{(8)qu} \right) + \frac{1}{4N_c} \left([Y_d^\dagger]_{pv} [Y_d]_{wr} C_{stvw}^{(8)qd} + [Y_d^\dagger]_{sv} [Y_d]_{wt} C_{prvw}^{(8)qd} \right) \\
& - \frac{1}{8} \left([Y_u^\dagger]_{pv} [Y_u]_{wt} C_{srvw}^{(8)qu} + [Y_u^\dagger]_{sv} [Y_u]_{wr} C_{ptvw}^{(8)qu} \right) - \frac{1}{8} \left([Y_d^\dagger]_{pv} [Y_d]_{wt} C_{srvw}^{(8)qd} + [Y_d^\dagger]_{sv} [Y_d]_{wr} C_{ptvw}^{(8)qd} \right) \\
& + \frac{1}{16N_c} \left([Y_d]_{wt} [Y_u]_{vr} C_{quqd}^{(8)psvw} + [Y_d]_{wr} [Y_u]_{vt} C_{quqd}^{(8)supw} \right) + \frac{1}{16N_c} \left([Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{quqd}^{(8)*trvw} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{quqd}^{(8)*trvw} \right) \\
& + \frac{1}{16} \left([Y_d]_{wt} [Y_u]_{vr} C_{quqd}^{(8)psvw} + [Y_d]_{wr} [Y_u]_{vt} C_{quqd}^{(8)psvw} \right) + \frac{1}{16} \left([Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{quqd}^{(8)*trvw} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{quqd}^{(8)*trvw} \right) \\
& - \frac{1}{2} [Y_u^\dagger]_{pv} [Y_u]_{wr} C_{stvw}^{(1)qu} - \frac{1}{2} [Y_d^\dagger]_{pv} [Y_d]_{wr} C_{stvw}^{(1)qd} - \frac{1}{2} [Y_u^\dagger]_{sv} [Y_u]_{wt} C_{prvw}^{(1)qu} - \frac{1}{2} [Y_d^\dagger]_{sv} [Y_d]_{wt} C_{prvw}^{(1)qd} \\
& - \frac{1}{8} [Y_d]_{wt} [Y_u]_{vr} C_{quqd}^{(1)psvw} - \frac{1}{8} [Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{quqd}^{(1)*trvw} - \frac{1}{8} [Y_d]_{wr} [Y_u]_{vt} C_{quqd}^{(1)psvw} - \frac{1}{8} [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{quqd}^{(1)*trvw} \\
& + \gamma_q^{(Y)} C_{pv}^{(1)qq} + \gamma_q^{(Y)} C_{vrst}^{(1)qq} + \gamma_q^{(Y)} C_{sv}^{(1)qq} + \gamma_q^{(Y)} C_{prvt}^{(1)qq} + C_{pq}^{(1)qq} \gamma_q^{(Y)} + C_{pq}^{(1)qq} \gamma_q^{(Y)} \quad (A.36)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{prst}^{(3)qq} = & -\frac{1}{2} [Y_u^\dagger Y_u + Y_d^\dagger Y_d]_{pr} C_{st}^{(3)Hq} - \frac{1}{2} [Y_u^\dagger Y_u + Y_d^\dagger Y_d]_{st} C_{pr}^{(3)Hq} \\
& - \frac{1}{8} \left([Y_u^\dagger]_{pv} [Y_u]_{wt} C_{stvw}^{(8)qu} + [Y_u^\dagger]_{sv} [Y_u]_{wr} C_{ptvw}^{(8)qu} \right) - \frac{1}{8} \left([Y_d^\dagger]_{pv} [Y_d]_{wt} C_{stvw}^{(8)qd} + [Y_d^\dagger]_{sv} [Y_d]_{wr} C_{ptvw}^{(8)qd} \right) \\
& - \frac{1}{16N_c} \left([Y_d]_{wt} [Y_u]_{vr} C_{quqd}^{(8)psvw} + [Y_d]_{wr} [Y_u]_{vt} C_{quqd}^{(8)supw} \right) - \frac{1}{16N_c} \left([Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{quqd}^{(8)*trvw} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{quqd}^{(8)*trvw} \right) \\
& - \frac{1}{16} \left([Y_d]_{wt} [Y_u]_{vr} C_{quqd}^{(8)psvw} + [Y_d]_{wr} [Y_u]_{vt} C_{quqd}^{(8)psvw} \right) - \frac{1}{16} \left([Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{quqd}^{(8)*trvw} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{quqd}^{(8)*trvw} \right) \\
& + \frac{1}{8} [Y_d]_{wt} [Y_u]_{vr} C_{quqd}^{(1)psvw} + \frac{1}{8} [Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{quqd}^{(1)*trvw} + \frac{1}{8} [Y_d]_{wr} [Y_u]_{vt} C_{quqd}^{(1)psvw} + \frac{1}{8} [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{quqd}^{(1)*trvw} \\
& + \gamma_q^{(Y)} C_{pv}^{(3)qq} + \gamma_q^{(Y)} C_{vrst}^{(3)qq} + \gamma_q^{(Y)} C_{sv}^{(3)qq} + \gamma_q^{(Y)} C_{prvt}^{(3)qq} + C_{pq}^{(3)qq} \gamma_q^{(Y)} + C_{pq}^{(3)qq} \gamma_q^{(Y)} \quad (A.37)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{prst}^{(1)lq} = & -[Y_e^\dagger Y_e]_{pr} C_{st}^{(1)Hq} + [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{st} C_{pr}^{(1)Hl} + \frac{1}{4} [Y_u]_{wt} [Y_e]_{vr} C_{psvw}^{(1)lequ} + \frac{1}{4} [Y_u^\dagger]_{sw} [Y_e^\dagger]_{pv} C_{rvtw}^{(1)*lequ} \\
& - [Y_u^\dagger]_{sv} [Y_u]_{wt} C_{prvw}^{lu} - [Y_d^\dagger]_{sv} [Y_d]_{wt} C_{prvw}^{ld} - [Y_e^\dagger]_{pv} [Y_e]_{wr} C_{stvw}^{qe} \\
& + \frac{1}{4} \left([Y_d^\dagger]_{sw} [Y_e]_{vr} C_{pvwt}^{ledq} + [Y_e^\dagger]_{pv} [Y_d]_{wt} C_{rvws}^{ledq*} \right) - \frac{1}{4} \left([Y_e]_{vr} [Y_u]_{wt} C_{psvw}^{(3)lequ} + [Y_e^\dagger]_{pv} [Y_u^\dagger]_{sw} C_{rvtw}^{(3)*lequ} \right) \\
& + \gamma_l^{(Y)} C_{pv}^{(1)lq} + \gamma_l^{(Y)} C_{vrst}^{(1)lq} + \gamma_l^{(Y)} C_{sv}^{(1)lq} + \gamma_l^{(Y)} C_{prvt}^{(1)lq} + C_{pl}^{(1)lq} \gamma_l^{(Y)} + C_{pl}^{(1)lq} \gamma_l^{(Y)} \quad (A.38)
\end{aligned}$$

$$\begin{aligned} \dot{C}_{lq}^{(3)} = & -[Y_e^\dagger Y_e]_{pr} C_{Hq}^{(3)} - [Y_u^\dagger Y_u + Y_d^\dagger Y_d]_{st} C_{Hl}^{(3)} - \frac{1}{4} [Y_u]_{wt} [Y_e]_{vr} C_{lequ}^{(1)} - \frac{1}{4} [Y_u^\dagger]_{sw} [Y_e^\dagger]_{pv} C_{lequ}^{(1)*} \\ & + \frac{1}{4} \left([Y_d^\dagger]_{sw} [Y_e]_{vr} C_{ledq} + [Y_e^\dagger]_{pv} [Y_d]_{wt} C_{ledq}^* \right) + \textcolor{red}{3} \left([Y_e]_{vr} [Y_u]_{wt} C_{lequ}^{(3)} + [Y_e^\dagger]_{pv} [Y_u^\dagger]_{sw} C_{lequ}^{(3)*} \right) \\ & + \gamma_{pv}^{(Y)} C_{vrst}^{(3)} + \gamma_{sv}^{(Y)} C_{prvt}^{(3)} + C_{pvst}^{(3)} \gamma_{vr}^{(Y)} + C_{prsv}^{(3)} \gamma_{vt}^{(Y)} \end{aligned} \quad (\text{A.39})$$

A.8.2 $(\overline{R}R)(\overline{R}R)$

$$\begin{aligned} \hat{C}_{prst}^{ee} = & [Y_e Y_e^\dagger]_{pr} C_{st}^{He} + [Y_e Y_e^\dagger]_{st} C_{pr}^{He} - [Y_e^\dagger]_{wr} [Y_e]_{pv} C_{vwst}^{le} - [Y_e^\dagger]_{wt} [Y_e]_{sv} C_{vwpr}^{le} \\ & + \gamma_{pv}^{(Y)} C_{vrst}^{ee} + \gamma_{sv}^{(Y)} C_{prvt}^{ee} + C_{pvst}^{ee} \gamma_{vr}^{(Y)} + C_{prsv}^{ee} \gamma_{vt}^{(Y)} \end{aligned} \quad (\text{A.40})$$

$$\begin{aligned} \dot{C}_{prst}^{uu} = & -[Y_u Y_u^\dagger]_{pr} C_{st}^{Hu} - [Y_u Y_u^\dagger]_{st} C_{pr}^{Hu} - [Y_u^\dagger]_{wr} [Y_u]_{pv} C_{vwst}^{(1)qu} - [Y_u^\dagger]_{wt} [Y_u]_{sv} C_{vwpr}^{(1)qu} \\ & + \frac{1}{2N_c} [Y_u]_{pv} [Y_u^\dagger]_{wr} C_{vwst}^{(8)qu} + \frac{1}{2N_c} [Y_u]_{sv} [Y_u^\dagger]_{wt} C_{vwpr}^{(8)qu} - \frac{1}{2} [Y_u^\dagger]_{wr} [Y_u]_{sv} C_{vwpt}^{(8)qu} - \frac{1}{2} [Y_u^\dagger]_{wt} [Y_u]_{pv} C_{vwsr}^{(8)qu} \\ & + \gamma_u^{(Y)} C_{pv}^{uu} C_{vrst}^{uu} + \gamma_u^{(Y)} C_{sv}^{uu} C_{prvt}^{uu} + C_{pvst}^{uu} \gamma_u^{(Y)} C_{vr}^{uu} + C_{prsv}^{uu} \gamma_u^{(Y)} C_{vt}^{uu} \end{aligned} \quad (\text{A.41})$$

$$\begin{aligned} \dot{C}_{prst}^{dd} = & [Y_d Y_d^\dagger]_{pr} C_{st}^{Hd} + [Y_d Y_d^\dagger]_{st} C_{pr}^{Hd} - [Y_d^\dagger]_{wr} [Y_d]_{pv} C_{qd}^{(1)} - [Y_d^\dagger]_{wt} [Y_d]_{sv} C_{qd}^{(1)} \\ & + \frac{1}{2N_c} [Y_d]_{pv} [Y_d^\dagger]_{wr} C_{qvst}^{(8)} + \frac{1}{2N_c} [Y_d]_{sv} [Y_d^\dagger]_{wt} C_{qvpr}^{(8)} - \frac{1}{2} [Y_d^\dagger]_{wr} [Y_d]_{sv} C_{qvpt}^{(8)} - \frac{1}{2} [Y_d^\dagger]_{wt} [Y_d]_{pv} C_{qvrs}^{(8)} \\ & + \gamma_{pv}^{(Y)} C_{vrst}^{dd} + \gamma_{sv}^{(Y)} C_{prvt}^{dd} + C_{pvst}^{dd} \gamma_{vr}^{(Y)} + C_{prsv}^{dd} \gamma_{vt}^{(Y)} \end{aligned} \quad (\text{A.42})$$

$$\begin{aligned} \dot{C}_{prst}^{eu} = & 2[Y_e Y_e^\dagger]_{pr} C_{st}^{Hu} - 2[Y_u Y_u^\dagger]_{st} C_{pr}^{He} + \left([Y_e]_{pv} [Y_u]_{sw} C_{vrwt}^{lequ(1)} + [Y_e^\dagger]_{vr} [Y_u^\dagger]_{wt} C_{vpws}^{lequ(1)*} \right) \\ & - 2[Y_e]_{pv} [Y_e^\dagger]_{wr} C_{vwst}^{lu} - 12 \left([Y_e]_{pv} [Y_u]_{sw} C_{vrwt}^{lequ(3)} + [Y_e^\dagger]_{vr} [Y_u^\dagger]_{wt} C_{vpws}^{lequ(3)*} \right) \\ & - 2[Y_u]_{sv} [Y_u^\dagger]_{wt} C_{vwpr}^{qe} + \gamma_{pe}^{(Y)} C_{vrst}^{eu} + \gamma_{uv}^{(Y)} C_{prvt}^{eu} + C_{pvst}^{eu} \gamma_{vr}^{(Y)} + C_{prsv}^{eu} \gamma_{vt}^{(Y)} \end{aligned} \quad (\text{A.43})$$

$$\begin{aligned} \dot{C}_{prst}^{ed} = & 2[Y_e Y_e^\dagger]_{pr} C_{st}^{Hd} + 2[Y_d Y_d^\dagger]_{st} C_{pr}^{He} - 2[Y_e]_{pv} [Y_e^\dagger]_{wr} C_{vwt}^{ld} - 2[Y_d]_{sv} [Y_d^\dagger]_{wt} C_{vwp}^{qe} \\ & + [Y_e]_{pv} [Y_d^\dagger]_{wt} C_{vrsu}^{ledq} + [Y_e^\dagger]_{vr} [Y_d]_{sw} C_{vptw}^{ledq} + \gamma_{pv}^{(Y)} C_{vrst}^{ed} + \gamma_{sv}^{(Y)} C_{prvt}^{ed} + C_{pust}^{ed} \gamma_{vr}^{(Y)} + C_{prsv}^{ed} \gamma_{vt}^{(Y)} \end{aligned} \quad (\text{A.44})$$

$$\begin{aligned}
\dot{C}_{prst}^{(1)ud} = & -2[Y_u Y_u^\dagger]_{pr} C_{st}^{Hd} + 2[Y_d Y_d^\dagger]_{st} C_{pr}^{Hu} + \frac{2}{N_c} [Y_d Y_u^\dagger]_{sr} C_{pt}^{Hud} + \frac{2}{N_c} [Y_u Y_d^\dagger]_{pt} C_{rs}^{*Hud} \\
& + \frac{1}{N_c} \left([Y_d]_{sv} [Y_u]_{pw} C_{vrwt}^{(1)quqd} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{vpws}^{(1)*quqd} \right) - [Y_d]_{sw} [Y_u]_{pv} C_{vrwt}^{(1)quqd} - [Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{vpws}^{(1)*quqd} \\
& + \frac{N_c^2 - 1}{2N_c^2} \left([Y_d]_{sv} [Y_u]_{pw} C_{vrwt}^{(8)quqd} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{vpws}^{(8)*quqd} \right) - 2[Y_u]_{pv} [Y_u^\dagger]_{wr} C_{vwst}^{(1)quqd} - 2[Y_d]_{sv} [Y_d^\dagger]_{wt} C_{vwpr}^{(1)quqd} \\
& + \gamma_{pv}^{(Y)u} C_{vrst}^{(1)ud} + \gamma_{sv}^{(Y)d} C_{prvt}^{(1)ud} + C_{pvst}^{(1)ud} \gamma_{vr}^{(Y)u} + C_{prsv}^{(1)ud} \gamma_{vt}^{(Y)d}
\end{aligned} \tag{A.45}$$

$$\begin{aligned}
\dot{C}_{prst}^{(8)ud} = & 4[Y_d Y_u^\dagger]_{sr} C_{pt}^{Hud} + 4[Y_u Y_d^\dagger]_{pt} C_{rs}^{*Hud} + 2 \left([Y_d]_{sv} [Y_u]_{pw} C_{vrwt}^{(1)quqd} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{vpws}^{(1)*quqd} \right) \\
& - 2[Y_u]_{pv} [Y_u^\dagger]_{wr} C_{vwst}^{(8)quqd} - 2[Y_d]_{sv} [Y_d^\dagger]_{wt} C_{vwpr}^{(8)quqd} - \frac{1}{N_c} \left([Y_d]_{sv} [Y_u]_{pw} C_{vrwt}^{(8)quqd} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{vpws}^{(8)*quqd} \right) \\
& - \left([Y_d]_{sw} [Y_u]_{pv} C_{vrwt}^{(8)quqd} + [Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{vpws}^{(8)*quqd} \right) + \gamma_{pv}^{(Y)u} C_{vrst}^{(8)ud} + \gamma_{sv}^{(Y)d} C_{prvt}^{(8)ud} + C_{pvst}^{(8)ud} \gamma_{vr}^{(Y)u} + C_{prsv}^{(8)ud} \gamma_{vt}^{(Y)d}
\end{aligned} \tag{A.46}$$

A.8.3 $(\bar{L}L)(\bar{R}R)$

$$\begin{aligned}
\dot{C}_{prst}^{le} = & [Y_e]_{sr} \xi_{pt}^e + [Y_e^\dagger]_{pt} \xi_{rs}^{*e} - [Y_e^\dagger Y_e]_{pr} C_{st}^{He} + 2[Y_e Y_e^\dagger]_{st} C_{pr}^{Hl} - [Y_e^\dagger]_{pv} [Y_e]_{wr} C_{vtsw}^{ee} - [Y_e^\dagger]_{pw} [Y_e]_{vr} C_{wtsv}^{ee} \\
& - 2[Y_e^\dagger]_{pv} [Y_e]_{wr} C_{vwst}^{ee} + [Y_e^\dagger]_{pw} [Y_e]_{sv} C_{vrwt}^{le} - [Y_e^\dagger]_{wt} [Y_e]_{sv} C_{pvwr}^{ll} - [Y_e^\dagger]_{vt} [Y_e]_{sw} C_{pvwr}^{ll} \\
& - 4[Y_e^\dagger]_{wt} [Y_e]_{sv} C_{prvw}^{ll} + [Y_e^\dagger]_{vt} [Y_e]_{wr} C_{pvsw}^{le} + \gamma_l^{(Y)} C_{pv}^{le} + \gamma_{sv}^{(Y)e} C_{prvt}^{le} + C_{pvst}^{le} \gamma_{vr}^{(Y)l} + C_{prsv}^{le} \gamma_{vt}^{(Y)e}
\end{aligned} \tag{A.47}$$

$$\begin{aligned}
\dot{C}_{prst}^{lu} = & -[Y_e^\dagger Y_e]_{pr} C_{st}^{Hu} - 2[Y_u Y_u^\dagger]_{st} C_{pr}^{Hl} - \frac{1}{2} \left([Y_e]_{vr} [Y_u]_{sw} C_{pvwt}^{(1)lequ} + [Y_e^\dagger]_{pv} [Y_u^\dagger]_{wt} C_{rvws}^{(1)*lequ} \right) \\
& - 2[Y_u]_{sv} [Y_u^\dagger]_{wt} C_{prvw}^{(1)lq} - \left([Y_e]_{vr} [Y_u]_{sw} C_{pvwt}^{(3)lequ} + [Y_e^\dagger]_{pv} [Y_u^\dagger]_{wt} C_{rvws}^{(3)*lequ} \right) - [Y_e]_{wr} [Y_e^\dagger]_{pv} C_{vwst}^{eu} \\
& + \gamma_l^{(Y)} C_{pv}^{lu} + \gamma_{sv}^{(Y)u} C_{prvt}^{lu} + C_{pvst}^{lu} \gamma_{vr}^{(Y)l} + C_{prsv}^{lu} \gamma_{vt}^{(Y)u}
\end{aligned} \tag{A.48}$$

$$\begin{aligned}
\dot{C}_{prst}^{ld} = & -[Y_e^\dagger Y_e]_{pr} C_{st}^{Hd} + 2[Y_d Y_d^\dagger]_{st} C_{pr}^{Hl} - \frac{1}{2} \left([Y_e]_{vr} [Y_d^\dagger]_{wt} C_{pvsw}^{ledq} + [Y_e^\dagger]_{pv} [Y_d]_{sw} C_{rvtw}^{*ledq} \right) \\
& - 2[Y_d]_{sv} [Y_d^\dagger]_{wt} C_{prvw}^{(1)lq} - [Y_e]_{wr} [Y_e^\dagger]_{pv} C_{vwst}^{ed} + \gamma_l^{(Y)} C_{pv}^{ld} + \gamma_{sv}^{(Y)d} C_{prvt}^{ld} + C_{pvst}^{ld} \gamma_{vr}^{(Y)l} + C_{prsv}^{ld} \gamma_{vt}^{(Y)d}
\end{aligned} \tag{A.49}$$

$$\begin{aligned}
\dot{C}_{prst}^{qe} = & [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{pr} C_{st}^{He} + 2[Y_e Y_e^\dagger]_{st} C_{pr}^{(1)Hq} - \frac{1}{2} \left([Y_d^\dagger]_{pw} [Y_e]_{sv} C_{vtwr}^{ledq} + [Y_e^\dagger]_{vt} [Y_d]_{wr} C_{vswp}^{ledq*} \right) \\
& - 2[Y_e]_{sv} [Y_e^\dagger]_{wt} C_{vwpr}^{(1)lq} - \frac{1}{2} \left([Y_u]_{wr} [Y_e]_{sv} C_{vtpw}^{(1)lequ} + [Y_e^\dagger]_{vt} [Y_u^\dagger]_{pw} C_{vsrw}^{(1)*lequ} \right) - [Y_d]_{wr} [Y_d^\dagger]_{pv} C_{stvw}^{ed} \\
& - 6 \left([Y_e]_{sv} [Y_u]_{wr} C_{vtpw}^{(3)lequ} + [Y_e^\dagger]_{vt} [Y_u^\dagger]_{pw} C_{vsrw}^{(3)*lequ} \right) - [Y_u]_{wr} [Y_u^\dagger]_{pv} C_{stvw}^{eu} \\
& + \gamma_q^{(Y)} C_{pv}^{qe} + \gamma_e^{(Y)} C_{sv}^{qe} + C_{prvt}^{qe} \gamma_q^{(Y)} + C_{pvst}^{qe} \gamma_e^{(Y)} + C_{prsv}^{qe} \gamma_v^{(Y)} + C_{vrt}^{qe} \gamma_e^{(Y)} \quad (A.50)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{prst}^{(1)qu} = & \frac{1}{N_c} [Y_u]_{sr} \xi_{pt}^u + \frac{1}{N_c} [Y_u^\dagger]_{pt} \xi_{rs}^{*u} + [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{pr} C_{st}^{Hu} - 2[Y_u Y_u^\dagger]_{st} C_{pr}^{(1)Hq} \\
& + \frac{1}{N_c} \left([Y_u^\dagger]_{pw} [Y_u]_{sv} C_{vrwt}^{(1)qu} + [Y_u^\dagger]_{vt} [Y_u]_{wr} C_{pvsw}^{(1)qu} + [Y_d]_{wr} [Y_u]_{sv} C_{ptvw}^{(1)quqd} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{rsvw}^{(1)*quqd} \right) \\
& - \frac{1}{2N_c^2} \left([Y_u^\dagger]_{pw} [Y_u]_{sv} C_{vrwt}^{(8)qu} + [Y_u^\dagger]_{vt} [Y_u]_{wr} C_{pvsw}^{(8)qu} + [Y_d]_{wr} [Y_u]_{sv} C_{ptvw}^{(8)quqd} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{rsvw}^{(8)*quqd} \right) \\
& - \frac{1}{N_c} \left([Y_u^\dagger]_{vt} [Y_u]_{sw} C_{pvwr}^{(1)qq} + [Y_u^\dagger]_{wt} [Y_u]_{sv} C_{pvwr}^{(1)qq} + [Y_u^\dagger]_{pv} [Y_u]_{wr} C_{vtsw}^{uu} + [Y_u^\dagger]_{pw} [Y_u]_{vr} C_{wtsw}^{uu} \right) \\
& - \frac{3}{N_c} \left([Y_u^\dagger]_{vt} [Y_u]_{sw} C_{pvwr}^{(3)qq} + [Y_u^\dagger]_{wt} [Y_u]_{sv} C_{pvwr}^{(3)qq} \right) + \frac{1}{2} \left([Y_u^\dagger]_{pw} [Y_u]_{sv} C_{vrwt}^{(8)qu} + [Y_u^\dagger]_{vt} [Y_u]_{wr} C_{pvsw}^{(8)qu} \right) \\
& + \frac{1}{2} \left([Y_u]_{sv} [Y_d]_{wr} C_{vtpw}^{(1)quqd} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{vsrw}^{(1)*quqd} + [Y_u]_{sv} [Y_d]_{wr} C_{ptvw}^{(8)quqd} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{rsvw}^{(8)*quqd} \right) \\
& - 4[Y_u^\dagger]_{wt} [Y_u]_{sv} C_{prvw}^{(1)qq} - 2[Y_u^\dagger]_{pv} [Y_u]_{wr} C_{vwst}^{uu} - [Y_d^\dagger]_{pv} [Y_d]_{wr} C_{stvw}^{ud} \\
& + \gamma_q^{(Y)} C_{pv}^{qu} + \gamma_u^{(Y)} C_{sv}^{qu} + C_{prvt}^{qu} \gamma_q^{(Y)} + C_{pvst}^{qu} \gamma_u^{(Y)} + C_{prsv}^{qu} \gamma_v^{(Y)} + C_{vrt}^{qu} \gamma_e^{(Y)} \quad (A.51)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{prst}^{(1)qd} = & \frac{1}{N_c} [Y_d]_{sr} \xi_{pt}^d + \frac{1}{N_c} [Y_d^\dagger]_{pt} \xi_{rs}^* + [Y_u^\dagger Y_u - Y_d^\dagger Y_d]_{pr} C_{st}^{Hd} + 2[Y_d Y_d^\dagger]_{st} C_{pr}^{(1)Hq} \\
& + \frac{1}{N_c} \left([Y_d^\dagger]_{pw} [Y_d]_{sv} C_{vrwt}^{(1)qd} + [Y_d^\dagger]_{vt} [Y_d]_{wr} C_{pvsu}^{(1)qd} + [Y_u]_{wr} [Y_d]_{sv} C_{vwpt}^{(1)quqd} + [Y_u^\dagger]_{pw} [Y_d^\dagger]_{vt} C_{vurs}^{(1)*quqd} \right) \\
& - \frac{1}{2N_c^2} \left([Y_d^\dagger]_{pw} [Y_d]_{sv} C_{vrwt}^{(8)qd} + [Y_d^\dagger]_{vt} [Y_d]_{wr} C_{pvsu}^{(8)qd} + [Y_u]_{wr} [Y_d]_{sv} C_{vwpt}^{(8)quqd} + [Y_u^\dagger]_{pw} [Y_d^\dagger]_{vt} C_{vurs}^{(8)*quqd} \right) \\
& - \frac{1}{N_c} \left([Y_d^\dagger]_{vt} [Y_d]_{sw} C_{pvwr}^{(1)qq} + [Y_d^\dagger]_{wt} [Y_d]_{sv} C_{pvsu}^{(1)qq} + [Y_d^\dagger]_{pv} [Y_d]_{wr} C_{vtsu}^{dd} + [Y_d^\dagger]_{pw} [Y_d]_{vr} C_{vtsu}^{dd} \right) \\
& - \frac{3}{N_c} \left([Y_d^\dagger]_{vt} [Y_d]_{sw} C_{pvwr}^{(3)qq} + [Y_d^\dagger]_{wt} [Y_d]_{sv} C_{pvsu}^{(3)qq} \right) + \frac{1}{2} \left([Y_d^\dagger]_{pw} [Y_d]_{sv} C_{vrwt}^{(8)qd} + [Y_d^\dagger]_{vt} [Y_d]_{wr} C_{pvsu}^{(8)qd} \right) \\
& + \frac{1}{2} \left([Y_d]_{sw} [Y_u]_{vr} C_{pvwt}^{(1)quqd} + [Y_u^\dagger]_{pv} [Y_d^\dagger]_{wt} C_{rvus}^{(1)*quqd} + [Y_d]_{sv} [Y_u]_{wr} C_{vwpt}^{(8)quqd} + [Y_u^\dagger]_{pw} [Y_d^\dagger]_{vt} C_{vurs}^{(8)*quqd} \right) \\
& - 4[Y_d^\dagger]_{wt} [Y_d]_{sv} C_{pruv}^{(1)qq} - 2[Y_d^\dagger]_{pv} [Y_d]_{wr} C_{vust}^{dd} - [Y_u^\dagger]_{pv} [Y_u]_{wr} C_{vust}^{ud} \\
& + \gamma_{pv}^{(Y)q} C_{qrst}^{(1)qd} + \gamma_{sv}^{(Y)d} C_{qrst}^{(1)qd} + C_{qrst}^{(1)qd} \gamma_{vr}^{(Y)q} + C_{qrst}^{(1)qd} \gamma_{vt}^{(Y)d}
\end{aligned} \tag{A.52}$$

$$\begin{aligned}
\dot{C}_{prst}^{(8)qu} = & 2[Y_u]_{sr} \xi_{pt}^u + 2[Y_u^\dagger]_{pt} \xi_{rs}^* \\
& - \frac{1}{N_c} \left([Y_u^\dagger]_{pw} [Y_u]_{sv} C_{vrwt}^{(8)qu} + [Y_u^\dagger]_{vt} [Y_u]_{wr} C_{pvsu}^{(8)qu} + [Y_d]_{wr} [Y_u]_{sv} C_{vwpt}^{(8)quqd} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{vurs}^{(8)*quqd} \right) \\
& + 2 \left([Y_u]_{sv} [Y_d]_{wr} C_{ptvw}^{(1)quqd} + [Y_u^\dagger]_{vt} [Y_d^\dagger]_{pw} C_{rsuv}^{(1)*quqd} + \frac{1}{4} [Y_u]_{sv} [Y_d]_{wr} C_{vtpw}^{(8)quqd} + \frac{1}{4} [Y_u^\dagger]_{vt} [Y_d^\dagger]_{pw} C_{vsru}^{(8)*quqd} \right) \\
& - 2 \left([Y_u^\dagger]_{vt} [Y_u]_{sw} C_{pvwr}^{(1)qq} + [Y_u^\dagger]_{wt} [Y_u]_{sv} C_{pvsu}^{(1)qq} - [Y_u^\dagger]_{pw} [Y_u]_{sv} C_{vrwt}^{(1)qu} - [Y_u^\dagger]_{vt} [Y_u]_{wr} C_{pvsu}^{(1)qu} \right) \\
& - 2 \left([Y_u^\dagger]_{pv} [Y_u]_{wr} C_{vtsu}^{uu} + [Y_u^\dagger]_{pw} [Y_u]_{vr} C_{vtsu}^{uu} \right) - 6 \left([Y_u^\dagger]_{vt} [Y_u]_{sw} C_{pvwr}^{(3)qq} + [Y_u^\dagger]_{wt} [Y_u]_{sv} C_{pvsu}^{(3)qq} \right) \\
& - [Y_d^\dagger]_{pv} [Y_d]_{wr} C_{stuv}^{(8)ud} + \gamma_{pv}^{(Y)q} C_{qrst}^{(8)qu} + \gamma_{sv}^{(Y)u} C_{qrst}^{(8)qu} + C_{qrst}^{(8)qu} \gamma_{vr}^{(Y)q} + C_{qrst}^{(8)qu} \gamma_{vt}^{(Y)u}
\end{aligned} \tag{A.53}$$

$$\begin{aligned}
\dot{C}_{\substack{qd \\ prst}}^{(8)} &= 2[Y_d]_{sr}\xi_d + 2[Y_d^\dagger]_{pt}\xi_{rs}^* \\
&- \frac{1}{N_c} \left([Y_d^\dagger]_{pw} [Y_d]_{sv} C_{\substack{qd \\ vrw t}}^{(8)} + [Y_d^\dagger]_{vt} [Y_d]_{wr} C_{\substack{qd \\ pvs w}}^{(8)} + [Y_u]_{wr} [Y_d]_{sv} C_{\substack{quqd \\ vpw t}}^{(8)} + [Y_u^\dagger]_{pw} [Y_d^\dagger]_{vt} C_{\substack{quqd \\ vwr s}}^{(8)*} \right) \\
&+ 2 \left([Y_d]_{sv} [Y_u]_{wr} C_{\substack{quqd \\ vpw t}}^{(1)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{pw} C_{\substack{quqd \\ vwr s}}^{(1)*} + \frac{1}{4} [Y_u]_{vr} [Y_d]_{sw} C_{\substack{quqd \\ pvw t}}^{(8)} + \frac{1}{4} [Y_u^\dagger]_{pv} [Y_d^\dagger]_{wt} C_{\substack{quqd \\ rvws}}^{(8)*} \right) \\
&- 2 \left([Y_d^\dagger]_{vt} [Y_d]_{sw} C_{\substack{qq \\ pvwr}}^{(1)} + [Y_d^\dagger]_{wt} [Y_d]_{sv} C_{\substack{qq \\ pwr v}}^{(1)} - [Y_d^\dagger]_{pw} [Y_d]_{sv} C_{\substack{qd \\ vrw t}}^{(1)} - [Y_d^\dagger]_{vt} [Y_d]_{wr} C_{\substack{qd \\ pvs w}}^{(1)} \right) \\
&- 2 \left([Y_d^\dagger]_{pv} [Y_d]_{wr} C_{\substack{dd \\ vts w}} + [Y_d^\dagger]_{pw} [Y_d]_{vr} C_{\substack{dd \\ wts v}} \right) - 6 \left([Y_d^\dagger]_{vt} [Y_d]_{sw} C_{\substack{qq \\ pvs w}}^{(3)} + [Y_d^\dagger]_{wt} [Y_d]_{sv} C_{\substack{qq \\ pwr v}}^{(3)} \right) \\
&- [Y_u^\dagger]_{pv} [Y_u]_{wr} C_{\substack{ud \\ vws t}}^{(8)} + \gamma_{\substack{q \\ pv}}^{(Y)} C_{\substack{qd \\ vrs t}}^{(8)} + \gamma_{\substack{d \\ sv}}^{(Y)} C_{\substack{qd \\ prvt}}^{(8)} + C_{\substack{qd \\ pvs t}}^{(8)} \gamma_{\substack{q \\ vr}}^{(Y)} + C_{\substack{qd \\ prsv}}^{(8)} \gamma_{\substack{d \\ vt}}^{(Y)} \quad (A.54)
\end{aligned}$$

A.8.4 $(\bar{L}R)(\bar{R}L)$

$$\begin{aligned}
\dot{C}_{\substack{ledq \\ prst}} &= -2[Y_d]_{st}\xi_{pr}^e - 2[Y_e]_{pr}^* \xi_{st}^* + 2[Y_e^\dagger]_{pv} [Y_d]_{wt} C_{\substack{ed \\ vrs w}} - 2[Y_e^\dagger]_{vr} [Y_d]_{wt} C_{\substack{ld \\ pvs w}} + 2[Y_e^\dagger]_{vr} [Y_d]_{sw} C_{\substack{lq \\ pvwt}}^{(1)} \\
&+ 6[Y_e^\dagger]_{vr} [Y_d]_{sw} C_{\substack{lq \\ pvwt}}^{(3)} - 2[Y_e^\dagger]_{pw} [Y_d]_{sv} C_{\substack{qe \\ vtwr}} + 2[Y_d]_{sv} [Y_u]_{wt} C_{\substack{lequ \\ prvw}}^{(1)} \\
&+ \gamma_{\substack{l \\ pv}}^{(Y)} C_{\substack{ledq \\ vrs t}} + \gamma_{\substack{q \\ sv}}^{(Y)} C_{\substack{ledq \\ prvt}} + C_{\substack{ledq \\ pvs t}} \gamma_{\substack{e \\ vr}}^{(Y)} + C_{\substack{ledq \\ prsv}} \gamma_{\substack{d \\ vt}}^{(Y)} \quad (A.55)
\end{aligned}$$

A.8.5 $(\bar{L}R)(\bar{L}R)$

$$\begin{aligned}
\dot{C}_{\substack{lequ \\ prst}}^{(1)} &= 2[Y_u^\dagger]_{st}\xi_{pr}^e + 2[Y_e^\dagger]_{pr}\xi_{st}^u + 2[Y_d^\dagger]_{sv} [Y_u^\dagger]_{wt} C_{\substack{ledq \\ prvw}} + 2[Y_e^\dagger]_{pv} [Y_u^\dagger]_{sw} C_{\substack{eu \\ vrw t}} + 2[Y_e^\dagger]_{vr} [Y_u^\dagger]_{wt} C_{\substack{lq \\ pvs w}}^{(1)} \\
&- 6[Y_e^\dagger]_{vr} [Y_u^\dagger]_{wt} C_{\substack{lq \\ pvs w}}^{(3)} - 2[Y_e^\dagger]_{vr} [Y_u^\dagger]_{sw} C_{\substack{lu \\ pvwt}} - 2[Y_e^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{\substack{qe \\ svwr}} \\
&+ \gamma_{\substack{l \\ pv}}^{(Y)} C_{\substack{lequ \\ vrs t}}^{(1)} + \gamma_{\substack{q \\ sv}}^{(Y)} C_{\substack{lequ \\ prvt}}^{(1)} + C_{\substack{lequ \\ pvs t}}^{(1)} \gamma_{\substack{e \\ vr}}^{(Y)} + C_{\substack{lequ \\ prsv}}^{(1)} \gamma_{\substack{u \\ vt}}^{(Y)} \quad (A.56)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{\substack{lequ \\ prst}}^{(3)} &= -\frac{1}{2} [Y_u^\dagger]_{sw} [Y_e^\dagger]_{pv} C_{\substack{eu \\ vrw t}} - \frac{1}{2} [Y_e^\dagger]_{vr} [Y_u^\dagger]_{wt} C_{\substack{lq \\ pvs w}}^{(1)} + \frac{3}{2} [Y_e^\dagger]_{vr} [Y_u^\dagger]_{wt} C_{\substack{lq \\ pvs w}}^{(3)} \\
&- \frac{1}{2} [Y_e^\dagger]_{vr} [Y_u^\dagger]_{sw} C_{\substack{lu \\ pvwt}} - \frac{1}{2} [Y_e^\dagger]_{pw} [Y_u^\dagger]_{vt} C_{\substack{qe \\ svwr}} \\
&+ \gamma_{\substack{l \\ pv}}^{(Y)} C_{\substack{lequ \\ vrs t}}^{(3)} + \gamma_{\substack{q \\ sv}}^{(Y)} C_{\substack{lequ \\ prvt}}^{(3)} + C_{\substack{lequ \\ pvs t}}^{(3)} \gamma_{\substack{e \\ vr}}^{(Y)} + C_{\substack{lequ \\ prsv}}^{(3)} \gamma_{\substack{u \\ vt}}^{(Y)} \quad (A.57)
\end{aligned}$$

$$\begin{aligned}
\dot{C}_{quqd}^{(1)} = & -2[Y_u^\dagger]_{pr}\xi_d - 2[Y_d^\dagger]_{st}\xi_u \\
& - \frac{2}{N_c^2} \left([Y_u^\dagger]_{vr} [Y_d^\dagger]_{pw} C_{svwt}^{(8)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{sw} C_{pvwr}^{(8)} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{vrwt}^{(8)} \right) \\
& + \frac{4}{N_c} \left([Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{svpw}^{(1)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{pvsu}^{(1)} - 3[Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{svpw}^{(3)} - 3[Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{pvsu}^{(3)} \right) \\
& + \frac{4}{N_c} \left([Y_d^\dagger]_{pw} [Y_u^\dagger]_{vr} C_{svwt}^{(1)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{sw} C_{pvwr}^{(1)} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{vrwt}^{(1)} \right) \\
& - 4 \left([Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{pvsu}^{(1)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{svpw}^{(1)} \right) + 12 \left([Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{pvsu}^{(3)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{svpw}^{(3)} \right) \\
& + 2 \left([Y_d^\dagger]_{pw} [Y_u^\dagger]_{vr} C_{svwt}^{(8)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{sw} C_{pvwr}^{(8)} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{vrwt}^{(8)} \right) - 4 [Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{vrwt}^{(1)} \\
& + \gamma_{pv}^{(q)} C_{quqd}^{(1)} + \gamma_{sv}^{(Y)} C_{quqd}^{(1)} + C_{quqd}^{(1)} \gamma_{prvt}^{(Y)} + C_{quqd}^{(1)} \gamma_{pvsu}^{(Y)} + C_{quqd}^{(1)} \gamma_{vrwt}^{(Y)}
\end{aligned} \tag{A.58}$$

$$\begin{aligned}
\dot{C}_{quqd}^{(8)} = & - \frac{4}{N_c} \left([Y_d^\dagger]_{pw} [Y_u^\dagger]_{vr} C_{svwt}^{(8)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{sw} C_{pvwr}^{(8)} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{vrwt}^{(8)} \right) \\
& + 8 \left([Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{svpw}^{(1)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{pvsu}^{(1)} \right) - 24 \left([Y_d^\dagger]_{wt} [Y_u^\dagger]_{vr} C_{svpw}^{(3)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{wr} C_{pvsu}^{(3)} \right) \\
& + 8 \left([Y_d^\dagger]_{pw} [Y_u^\dagger]_{vr} C_{svwt}^{(1)} + [Y_d^\dagger]_{vt} [Y_u^\dagger]_{sw} C_{pvwr}^{(1)} + [Y_d^\dagger]_{pw} [Y_u^\dagger]_{sv} C_{vrwt}^{(1)} \right) - 4 [Y_d^\dagger]_{sw} [Y_u^\dagger]_{pv} C_{vrwt}^{(8)} \\
& + \gamma_{pv}^{(q)} C_{quqd}^{(8)} + \gamma_{sv}^{(Y)} C_{quqd}^{(8)} + C_{quqd}^{(8)} \gamma_{prvt}^{(Y)} + C_{quqd}^{(8)} \gamma_{pvsu}^{(Y)} + C_{quqd}^{(8)} \gamma_{vrwt}^{(Y)}
\end{aligned} \tag{A.59}$$

References

- [1] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl.Phys.* **B268** (1986) 621.
- [2] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **1010** (2010) 085, [[arXiv:1008.4884](#)].
- [3] C. Grojean, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Scaling of Higgs Operators and $h \rightarrow \gamma\gamma$ Decay*, [arXiv:1301.2588](#).
- [4] A. V. Manohar, *An Exactly Solvable Model for Dimension Six Higgs Operators and $h \rightarrow \gamma\gamma$* , [arXiv:1305.3927](#).
- [5] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators I: Formalism and λ Dependence*, [arXiv:1308.2627](#).
- [6] A. Brignole and A. Rossi, *Anatomy and phenomenology of mu-tau lepton flavor violation in the MSSM*, *Nucl.Phys.* **B701** (2004) 3–53, [[hep-ph/0404211](#)].
- [7] S. Khan, *Renormalization Group Evolution of the Non-Unitary operator*, *Nucl.Phys.* **B864** (2012) 38–51, [[arXiv:1203.5043](#)].
- [8] W. Dekens and J. de Vries, *Renormalization Group Running of Dimension-Six Sources of Parity and Time-Reversal Violation*, *JHEP* **1305** (2013) 149, [[arXiv:1303.3156](#)].
- [9] M. Raidal, A. van der Schaaf, I. Bigi, M. Mangano, Y. K. Semertzidis, et al., *Flavour physics of leptons and dipole moments*, *Eur.Phys.J.* **C57** (2008) 13–182, [[arXiv:0801.1826](#)].
- [10] J. Elias-Miro, J. Espinosa, E. Masso, and A. Pomarol, *Renormalization of dimension-six operators relevant for the Higgs decay h to $\gamma\gamma$* , [arXiv:1302.5661](#).
- [11] J. Elias-Miro, J. Espinosa, E. Masso, and A. Pomarol, *Higgs windows to new physics through $d = 6$ operators: Constraints and one-loop anomalous dimensions*, [arXiv:1308.1879](#).
- [12] A. Manohar and H. Georgi, *Chiral Quarks and the Nonrelativistic Quark Model*, *Nucl.Phys.* **B234** (1984) 189.
- [13] E. E. Jenkins, A. V. Manohar, and M. Trott, *Naive Dimensional Analysis Counting of Gauge Theory Amplitudes and Anomalous Dimensions*, [arXiv:1309.0819](#).
- [14] G. Blankenburg, J. Ellis, and G. Isidori, *Flavour-Changing Decays of a 125 GeV Higgs-like Particle*, *Phys.Lett.* **B712** (2012) 386–390, [[arXiv:1202.5704](#)].
- [15] R. Harnik, J. Kopp, and J. Zupan, *Flavor Violating Higgs Decays*, *JHEP* **1303** (2013) 026, [[arXiv:1209.1397](#)].
- [16] E. E. Jenkins, A. V. Manohar, and M. Trott, *On Gauge Invariance and Minimal Coupling*, [arXiv:1305.0017](#).
- [17] **MEG Collaboration** Collaboration, J. Adam et al., *New constraint on the existence of the μ to $e+\gamma$ decay*, [arXiv:1303.0754](#).
- [18] R. S. Chivukula and H. Georgi, *Composite Technicolor Standard Model*, *Phys.Lett.* **B188** (1987) 99.
- [19] G. D’Ambrosio, G. Giudice, G. Isidori, and A. Strumia, *Minimal flavor violation: An Effective field theory approach*, *Nucl.Phys.* **B645** (2002) 155–187, [[hep-ph/0207036](#)].

- [20] A. J. Buras, *Minimal flavor violation*, *Acta Phys.Polon.* **B34** (2003) 5615–5668, [[hep-ph/0310208](#)].
- [21] R. Alonso, H.-M. Chang, E. E. Jenkins, A. V. Manohar, and B. Shotwell, *Renormalization group evolution of dimension-six baryon number violating operators*, *Phys. Lett.* **B734** (2014) 302–307, [[arXiv:1405.0486](#)].
- [22] G. M. Pruna and A. Signer, *The $\mu \rightarrow e\gamma$ decay in a systematic effective field theory approach with dimension 6 operators*, *JHEP* **1410** (2014) 14, [[arXiv:1408.3565](#)].
- [23] C. Bobeth and U. Haisch, *Anomalous triple gauge couplings from B -meson and kaon observables*, [arXiv:1503.04829](#).

1 : X^3		2 : H^6		3 : $H^4 D^2$		5 : $\psi^2 H^3 + \text{h.c.}$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H	$(H^\dagger H)^3$	$Q_{H\Box}$	$(H^\dagger H)\Box(H^\dagger H)$	Q_{eH}	$(H^\dagger H)(\bar{l}_p e_r H)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$			Q_{HD}	$(H^\dagger D_\mu H)^* (H^\dagger D_\mu H)$	Q_{uH}	$(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W	$\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$					Q_{dH}	$(H^\dagger H)(\bar{q}_p d_r H)$
$Q_{\tilde{W}}$	$\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$						
4 : $X^2 H^2$		6 : $\psi^2 XH + \text{h.c.}$		7 : $\psi^2 H^2 D$			
Q_{HG}	$H^\dagger H G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{Hl}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$		
$Q_{H\tilde{G}}$	$H^\dagger H \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{Hl}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_p \tau^I \gamma^\mu l_r)$		
Q_{HW}	$H^\dagger H W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{He}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$		
$Q_{H\tilde{W}}$	$H^\dagger H \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$		
Q_{HB}	$H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$		
$Q_{H\tilde{B}}$	$H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	Q_{Hu}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$		
Q_{HWB}	$H^\dagger \tau^I H W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	Q_{Hd}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$		
$Q_{H\tilde{W}B}$	$H^\dagger \tau^I H \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$	$i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$		
8 : $(\bar{L}L)(\bar{L}L)$		8 : $(\bar{R}R)(\bar{R}R)$		8 : $(\bar{L}L)(\bar{R}R)$			
Q_{ll}	$(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee}	$(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le}	$(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$		
$Q_{qq}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu}	$(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu}	$(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$		
$Q_{qq}^{(3)}$	$(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd}	$(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld}	$(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$		
$Q_{lq}^{(1)}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu}	$(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe}	$(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$		
$Q_{lq}^{(3)}$	$(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed}	$(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$		
		$Q_{ud}^{(1)}$	$(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$		
		$Q_{ud}^{(8)}$	$(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$		
				$Q_{qd}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$		
8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$		8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$					
Q_{ledq}	$(\bar{l}_p^j e_r)(\bar{d}_s q_{tj})$	$Q_{quqd}^{(1)}$	$(\bar{q}_p^j u_r) \epsilon_{jk} (\bar{q}_s^k d_t)$				
		$Q_{quqd}^{(8)}$	$(\bar{q}_p^j T^A u_r) \epsilon_{jk} (\bar{q}_s^k T^A d_t)$				
		$Q_{lequ}^{(1)}$	$(\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)$				
		$Q_{lequ}^{(3)}$	$(\bar{l}_p^j \sigma_{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$				

Table 1. The 59 independent dimension-six operators built from Standard Model fields which conserve baryon number, as given in Ref. [2]. The operators are divided into eight classes: X^3 , H^6 , etc. Operators with +h.c. in the table heading also have hermitian conjugates, as does the $\psi^2 H^2 D$ operator Q_{Hud} . The subscripts p, r, s, t are flavor indices.