

UNIVERSITÀ DEGLI STUDI DI ROMA TOR VERGATA  
MACROAREA DI SCIENZE MATEMATICHE, FISICHE E NATURALI



LAUREA MAGISTRALE IN MATEMATICA PURA E APPLICATA

TITOLO

TO BE CONFIRMED WHEN IT IS DONE

**Relatore:**

Prof. Carlo Sinestrari

**Laureando:**

matricola: 0318523

Marco Tamburro

**Anno Accademico 2023/2024**



# Contents

<b>Introduction</b>	<b>v</b>
<b>1 The Alexandrov Reflection Method</b>	<b>1</b>
1.1 Geometry of immersed hypersurfaces (da espandere/sistemare) . . .	1
1.2 Local representation as a graph . . . . .	3
1.3 Some well established results from analysis . . . . .	4
1.4 Applying the maximum principle to non-linear PDEs . . . . .	8
1.5 The Alexandrov Reflection Method (da scrivere) . . . . .	10
1.6 The Alexandrov soap-bubble theorem (da scrivere) . . . . .	10
<b>2 The Chow Critical Planes Result</b>	<b>11</b>
2.1 Class of Equations we analyze . . . . .	11
2.2 Parabolicity of the differential equation (2.2) . . . . .	12
2.3 An existence result . . . . .	14
2.4 Local representation as a graph of a solution of (2.2) . . . . .	14
2.5 The Alexandrov Reflection Method and the Chow result . . . . .	15
2.6 Theorem 2.2 Chow [RENAME] . . . . .	16
<b>3 Third Chapter Title</b>	<b>19</b>
3.1 Reflections on spheres and hyperbolic spaces (da scrivere) . . . . .	19
3.2 The equation in constant curvature spaces (da scrivere) . . . . .	20
3.3 Extension of the result(da scrivere) . . . . .	21
3.4 Extension to constant curvature (da scrivere) . . . . .	22
<b>Bibliography</b>	<b>23</b>



# Introduction

To be confirmed



# Chapter 1

## The Alexandrov Reflection Method

### 1.1 Geometry of immersed hypersurfaces (da espandere/sistemare)

The study of immersed hypersurfaces is a fundamental topic in differential geometry, and is especially important in the field of geometric analysis. An immersed hypersurface  $X : M^n \rightarrow \overline{M}^{n+1}$  is a submanifold of a higher-dimensional space that is embedded in that space in such a way that the submanifold has the same dimension as the space in which it is embedded minus one. In other words, it is a submanifold of codimension one. We will always assume the embedding to be smooth. The most common case is  $\overline{M}^{n+1} = \mathbb{R}^{n+1}$ , usually extensively studied in undergraduate courses when  $n = 2$ . We will be assuming that the embedding is also isometric, i.e. the metric  $g$  on  $M^n$  is the one induced by  $(\overline{M}^{n+1}, \bar{g})$ . The pullback of the tangent bundle of  $\overline{M}^{n+1}$  to  $M^n$  is a smooth vector bundle on  $M^n$ :

$$X^*T\overline{M}^{n+1} = T\overline{M}^{n+1}|_{M^n} = \coprod_{p \in M^n} T_p\overline{M}^{n+1}$$

One of the most important concepts in the geometry of immersed hypersurfaces is the concept of a normal vector field. The normal vector field  $\nu$  is a vector field on the pullback vector bundle  $X^*TE$  on the manifold  $M^n$  that is perpendicular to the tangent space of the hypersurface at each point. At each point  $p$ :

$$T_p\overline{M}^{n+1} = T_pM^n \oplus N_pM^n$$

where  $N_pM^n$  is the normal vector bundle generated by the normal vector. This also allows us to define the tangent and normal projection on  $T\overline{M}^{n+1}|_{M^n}$  by taking the two respective components.

Clearly, taking  $\bar{\nabla}$  to be the Levi-Civita connection on  $(\bar{M}^{n+1}, \bar{g})$ , we can decompose it as:

$$\bar{\nabla}_v w = (\bar{\nabla}_v w)^\top + (\bar{\nabla}_v w)^\perp$$

**Definition 1.1.** *The second fundamental form is then defined as:*

$$\mathbb{I}(v, w) = (\bar{\nabla}_v w)^\perp$$

It is a bilinear symmetric tensor because  $TM^n$  is involutive in  $T\bar{M}^{n+1}$  and depends only on the local value of  $v$  and  $w$  by symmetry. we can therefore write it as

$$\mathbb{I}(v, w) = v^T A w = h_{ij} v^i w^j$$

for some matrix  $A = \{h_{ij}\}$ . we can define the principal curvatures of the hypersurface, as the eigenvalues of the second fundamental form.

It is also possible to check that  $(\bar{\nabla}_v w)^\top$  satisfies the definition the Levi-Civita connection on  $(M^n, g)$ , therefore from uniqueness:

$$\begin{aligned} \nabla_v w &= (\bar{\nabla}_v w)^\top \\ \bar{\nabla}_v w &= \nabla_v w + \mathbb{I}(v, w) \end{aligned}$$

This result is known as the *Gauss Formula*.

**Proposition 1.2. (The Weingarten Equations)** *If  $v, w \in TM^n$  and  $N \in NM^n$ , if one considers the corresponding derivations in  $T\bar{M}^{n+1}$  the following equation holds:*

$$\langle \bar{\nabla}_v N, w \rangle_{\bar{g}} = -\langle N, \mathbb{I}(v, w) \rangle_{\bar{g}}$$

*Proof.* As  $\langle N, w \rangle_{\bar{g}} \equiv 0$  on  $M$ ,

$$\begin{aligned} 0 &= v \langle N, w \rangle_{\bar{g}} \\ &= \langle \bar{\nabla}_v N, w \rangle_{\bar{g}} + \langle N, \bar{\nabla}_v w \rangle_{\bar{g}} \\ &= \langle \bar{\nabla}_v N, w \rangle_{\bar{g}} + \langle N, \mathbb{I}(v, w) \rangle_{\bar{g}} \end{aligned}$$

applying the Gauss Formula and the fact that  $\nabla_v w \in TM^n$  in the last step □

We are going to use these equations in local coordinates, in the form shown below.



**Proposition 1.3.** *The Weingarten Equations can be expressed in local coordinates as:*

$$\frac{\partial^2 X^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial X^\alpha}{\partial x^k} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X^\beta}{\partial x^i} \frac{\partial X^\delta}{\partial x^j} = -h_{ij} \nu^\alpha \quad (1.1)$$

$$\frac{\partial \nu^\alpha}{\partial x^i} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X^\beta}{\partial x^i} \nu^\delta = h_{ij} g^{jl} \frac{\partial X^\alpha}{\partial x^l} \quad (1.2)$$

where  $\nu$  is the normal unit vector at the point and  $A = \{h_{ij}\}$  is the second fundamental form, thus  $h_{ij} = \langle \nabla_i \nu, e_j \rangle$

*Proof.* **[DA SCRIVERE]** Vedi qui □

There exist many important partial differential equations (PDEs) which stem from the study of immersed hypersurfaces, for example, the minimal surface equation, which is a PDE that describes surfaces with minimal area, can be seen as a geometric condition on the mean curvature of an immersed hypersurface. Likewise, the mean curvature flow is a geometric flow that deforms an immersed hypersurface by moving each point along the normal vector field at that point by an amount proportional to the mean curvature at that point. This flow is important in the study of geometric PDEs and is used to obtain a lot of important results in differential geometry.

**[Da espandere più avanti, in base a quello che usiamo - forse]**

## 1.2 Local representation as a graph

A well known result, Dini's Theorem<sup>1</sup>, states that, given a smooth function  $F$  defined on an open subset of the product space  $R^n \times R^m$ , if  $F(x, y) = 0$  and the partial derivative of  $F$  with respect to  $y$  is nonzero at a point  $(x_0, y_0)$ , then there exists an open neighborhood of  $x_0$  in  $R^n$  and a unique smooth function  $y = g(x)$  defined on that neighborhood such that  $y_0$  is a regular value of  $g$  and  $(x, g(x))$  is a smooth solution to the equation  $F(x, y) = 0$ .

Consequence of the Dini's theorem is a powerful result that allows one to locally represent a submanifold of  $\mathbb{R}^n \times \mathbb{R}^m$  as the graph of a smooth function. This theorem is widely used in differential geometry, geometric analysis, and many other fields of mathematics and physics. We provide a version of this theorem below:

---

<sup>1</sup>also known as Implicit Function Theorem

**Theorem 1.4** (Local representation as a graph). *Let  $X^n$  be a submanifold  $X^n \subset \mathbb{R}^{n+1}$  and let  $x_0 \in X$ . Then there exists a neighbourhood of  $x_0$ ,  $U \subset X^n$ , such that  $U$  is the graph of a function. Moreover, this function can be of the form*

$$f : \pi(U) \subset \mathbb{R}^n \rightarrow \mathbb{R}$$

$$U = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} | x_0 = f(x_1, \dots, x_n)\}$$

for any of the possible orders of the usual basis for  $\mathbb{R}^n$ ,  $(e_0, \dots, e_n)$ , as long as  $e_0 \notin T_x M$ , where  $\pi(U)$  is the projection on the last  $n$  coordinates  $((x_0, \dots, x_n) \mapsto (x_1, \dots, x_n))$ .

A proof of the 2D-case of the version of the theorem can be found in [5] which extends naturally to the  $n$  dimensional case, with almost no changes. This immediately extends to:

**Corollary 1.5** (Local representation as a graph on the tangent). *Let  $X^n$  be a submanifold  $X^n \subset \mathbb{R}^{n+1}$  and let  $x_0 \in X$ . Then there exists a neighbourhood of  $x_0$   $U \subset X^n$  and a smooth function  $f : T_x X^n \rightarrow \mathbb{R}$  such that any  $x_0 \in U$  can be expressed as*

$$x_0 = p + f(p)\nu$$

where  $\nu$  is the vector normal to  $T_{x_0} X^n$ , for an appropriate point  $p \in T_{x_0} X^n$ . In other words, every submanifold  $X^n \subset \mathbb{R}^{n+1}$  is locally expressible as a graph on its tangent space.

*Proof.* By rotation, we may assume  $T_x X^n$  orthogonal to  $e_1$ . Then one can just apply the previous theorem.  $\square$

We will use this later to prove Theorem 2.3.

### 1.3 Some well established results from analysis

We now introduce some well known results from analysis. The first result we introduce is the maximum principle.

The maximum principle is a classical result of mathematical analysis, and it is usually introduced in a first course on partial differential equations. It is a fundamental tool in the theory of partial differential equations. It is a statement about the behavior of solutions to certain types of PDEs and provides a method for obtaining upper and lower bounds on the solutions. The principle states that the maximum and minimum values of a solution to elliptic or parabolic PDE occur on the boundary of the domain unless the function is constant.

The maximum principle can be used to prove the existence, uniqueness, and regularity of solutions to elliptic and parabolic PDEs. It can also be used to obtain estimates on the behavior of solutions and to study the asymptotic behavior of solutions as the domain becomes large. The principle is widely used in many fields of mathematics and physics, such as geometric analysis, mathematical physics, and fluid dynamics. One of the many versions of this well know theorem is this:

**Theorem 1.6** (Maximum principle for parabolic equations). *Let  $\Omega$  be an open, bounded, connected set. Assume  $u \in C_1^2(\Omega \times [0, T]) \cap C^1(\overline{\Omega} \times [0, T])$ . Suppose  $u$  satisfies:*

$$-\frac{\partial u}{\partial t} + \left( \sum_{i,j=1}^n a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_i b_i \frac{\partial}{\partial x_i} + c \right) u = -u_t + Lu \geq 0 \quad (1.3)$$

where  $L$  is an elliptic differential operator, i.e. there exists  $\theta > 0$  such that  $\sum_{i,j=1}^n a_{ij}(x, t) \xi_i \xi_j \geq \theta |\xi|^2$  for all  $\xi \in \mathbb{R}^n$  and  $(x, t) \in \Omega \times [0, T]$ . Suppose also that  $c \equiv 0$  in  $\Omega$ . Then:

- if  $u$  attains its maximum in an interior point  $(x_0, t_0) \in \Omega \times [0, T]$ , then  $u$  is constant in  $\Omega \times [0, t_0]$ .
- If, instead, under the same conditions,  $u_t + Lu \geq 0$  and attains its minimum in an interior point of  $\Omega \times [0, T]$ , then  $u$  is constant in  $\Omega \times [0, t_0]$

A proof of this result can be found, for example, in [4]. The theorem extends also to situations where the condition holds in a convex bounded connected region  $R \subseteq \Omega \times [0, T]$ : in that case, if  $u$  attains its maximum in an interior point then  $u$  has the same value at any point in  $R$  that can be connected to it through a segment going in the backwards direction of time and a "horizontal" line contained in  $\Omega$ . This version of the theorem can be found for example in [3]:

**Theorem 1.7.** *Let  $u$  satisfy the uniformly parabolic differential inequality (1.3) with  $c(x) \leq 0$  in a region  $R_T = \{(x_1, x_2, \dots, x_n, t) | t \leq T\}$  where  $R$  is a non-empty connected open set, and suppose that the coefficients of  $L$  are bounded. Suppose that the maximum of  $u$  in  $R_T$  is  $M$  and that it is attained at a point  $(x, t)$  of  $R_T$ . Thus if  $(y, s)$  is a point of  $R$  which can be connected to  $(x, t)$  by a path in  $R$  consisting only of horizontal segments and upward vertical segments, then  $u(y, s) = M$ .*

Hopf's boundary point lemma is another important classical tool in the study of PDEs that provides a criterion for determining the behavior of solutions to certain types of elliptic or parabolic PDEs near the boundary of the domain. The lemma states that if one has a solutions to some kinds of partial differential inequalities, then the normal derivative of the solution at that point is strictly positive.

It is often used to obtain estimates on the behavior of solutions near the boundary, and to prove the existence and uniqueness of solutions to boundary value problems. The lemma is named after the German mathematician Eberhard Hopf, who first formulated it in the 1950s. In [3] we find the following version of the Hopf's boundary point lemma:

**Theorem 1.8.** *Let  $u$  be a solution to the parabolic inequality*

$$-u_t + Lu \geq 0$$

*with  $L$  an elliptic linear differential operator with bounded coefficients such that  $c(x) \leq 0$ , in a domain  $E$ , and let  $E_t = \{(x, s) \in E \mid s \leq t\}$ . Suppose the maximum  $M$  of  $u$  is attained at a point  $P = (x, t)$  on the boundary  $\partial E$ .*

*Assume that a sphere through  $P$  can be constructed which is in  $E_s$  such that*

- *tangent to  $\partial E$  at  $P$*
- *the set of point of its interior  $(y, s)$  such that  $s \leq t$  lies in  $E_s$ ,*
- *$u < M$  in its interior.*

*Also, suppose that the radial direction from the centre of the sphere to  $P$  is not parallel to the  $t$ -axis.*

*Then, if  $\frac{\partial}{\partial \nu}$  denotes any directional derivative in an outward direction from  $E_s$ , we have*

$$\frac{\partial u}{\partial \nu} > 0$$

*at  $P$ .*

*Proof.* Let the sphere through  $P$  be  $B_1$ . We may construct a smaller sphere  $B_2$  centred at  $P$ . Let now:

$$S_1 = \partial B_1 \cap B_2 \cap E_t,$$

$$S_2 = B_1 \cap \partial B_2 \cap E_t, \text{ and}$$

$$S_3 = B_1 \cap B_2 \cap \partial E_t = B_1 \cap B_2 \cap \{(x, s) \in E \mid s = t\}.$$

The three sets satisfy  $S_1 \cup S_2 \cup S_3 = \partial(B_1 \cap B_2 \cap E_t)$ , we may call this region  $R = B_1 \cap B_2 \cap E_t$ . Without loss of generality, potentially taking a smaller sphere  $B_1$ , we may assume that  $u < M$  on  $B_1$  except at  $P$ . As  $R \subset B_1$ , we also get  $u < M$  on  $R$ . We may thus conclude that:

- $u < M$  on  $R$  except at  $P$
- $u \leq M - \delta$  on  $S_2$  for a sufficiently small  $\delta > 0$

- $u = M$  at  $P$ .

Now, let the centre of  $B_1$  be  $Q = (z, t_0)$  and let  $r$  be its radius. we can now introduce the function

$$v(y, s) = \exp \left( -\alpha(s - t_0)^2 - \sum_{i=1}^n \alpha(y_i - z_i)^2 \right) - \exp(-\alpha r^2)$$

This function is such that  $v(y, s) = 0$  if  $(y, s) \in S_1$  - including  $v(x, t) = 0$ , as there the first term is  $e^{-\alpha r^2}$ , and  $v(y, s) > 0$  in the interior of  $B_1$ .

Thus, in the region  $R$ ,  $v(y, s) \geq 0$  and has a minimum point at the boundary on  $(x, t)$ , where  $v(x, t) = 0$ .

We can also compute  $Lv$ . After some calculation, we get that

$$\begin{aligned} Lv = 2\alpha e^{(-\alpha(s-t_0)^2 - \sum_{i=1}^n \alpha(y_i - z_i)^2)} [2\alpha(y - z)^t A(y - z) + \\ + \sum_i^n [b_i(y_i - z_i) + a_{i,i}] + (s - t)] \end{aligned}$$

where  $A$  is the matrix of the  $a_{i,j}$ . In particular, one can choose an  $\alpha$  large enough, so that  $Lv > 0$  in  $R \cup \partial R$ .

We can thus introduce  $w = u + \varepsilon v$ . As both  $Lu$  and  $Lv$  are positive in  $R$ ,  $Lw > 0$  in  $R$ . We can also choose  $\varepsilon$  small enough so that  $w < M$  on  $S_2$ . Also, as  $v = 0$  on  $S_1$ ,  $w < M$  on  $S_1$  except at  $P$ , and  $w = M$  at  $P$ .

Therefore, we can apply the Strong Maximum Principle 1.7 to the region  $R$  to conclude that the maximum of  $w$  in  $R$  is attained at  $P$ . Therefore:

$$\frac{\partial w}{\partial \nu} = \frac{\partial u}{\partial \nu} + \varepsilon \frac{\partial v}{\partial \nu} \geq 0$$

But:

$$\frac{\partial v}{\partial \nu} = \nu \cdot n \frac{\partial v}{\partial R} = -2\nu \cdot n \alpha R e^{-\alpha R} < 0$$

Where  $n$  is the vector orthogonal to the sphere  $S_1$ . Therefore, one must have:

$$\frac{\partial u}{\partial \nu} > 0$$

as we wanted. □

**Remark 1.9.** If  $c(x)$  is now just bounded, we can consider, instead of  $u$ ,  $v = ue^{-\lambda t}$ , thus, by change of variables

$$-v_t + Lv - \lambda v \geq 0$$

whenever  $-u_t + Lu \geq 0$ , and we can chose  $\lambda$  large enough such that  $c(x) - \lambda < 0$  and thus we can remove the hypothesis  $c(x) \leq 0$  in both theorems when  $c$  is bounded.

## 1.4 Applying the maximum principle to non-linear PDEs

We can now introduce an important observation shown in [3] that allows us to apply the maximum principle 1.7 and Hopf's boundary point lemma 1.8 in some non-linear settings. Firstly, we must clarify what we mean by parabolic non-linear problem.

**Definition 1.10.** *A differential non-linear problem in the form*

$$Lu = F\left(t, x, v, \frac{\partial v}{\partial x_i}, \frac{\partial^2 v}{\partial x_i \partial x_j}\right) - v_t = f(x, t) \quad (1.4)$$

*given a smooth  $F$  is parabolic if for any real vector  $\xi$*

$$\sum_{i,j=1}^n F_{ij} \xi_i \xi_j > 0$$

*where  $F_{ij}$  are the derivatives of  $F$  with respect to  $\frac{\partial^2 v}{\partial x_i \partial x_j}$ .*

Secondly, we remind the reader of the following generalized version of the theorem of the mean, a.k.a. Lagrange's theorem:

**Theorem 1.11** (Lagrange's theorem). *Given a convex open set  $U \subseteq \mathbb{R}^n$  and a real function  $F \in C^1(U)$ , and given to points  $x, y$  in  $U$ , there exists a point  $z$  in the segment connecting  $x$  and  $y$  such that*

$$F(y) - F(x) = \langle \nabla F(z), (y - x) \rangle$$

Suppose that we have a solution to a non-linear parabolic problem  $v$ , i.e.  $v$  solves (1.4):

$$Lu = F\left(t, x, v, \frac{\partial v}{\partial x_i}, \frac{\partial^2 v}{\partial x_i \partial x_j}\right) - v_t = f(x, t)$$

for a non-linear elliptic operator  $L$  in some region  $E$ , where we assume  $F(t, x, y_i, z_{i,j})$  to be a given  $C^1$  function. Suppose also that there is a  $w$  which is a solution of the corresponding differential inequality:

$$Lw = F\left(t, x, w, \frac{\partial w}{\partial x_i}, \frac{\partial^2 w}{\partial x_i \partial x_j}\right) - w_t \leq f(x, t)$$

One can then consider  $u = v - w$ , and by combining the above we get:

$$Lv = \left( F \left( t, x, v, \frac{\partial v}{\partial x_i}, \frac{\partial^2 v}{\partial x_i \partial x_j} \right) - F \left( t, x, w, \frac{\partial w}{\partial x_i}, \frac{\partial^2 w}{\partial x_i \partial x_j} \right) \right) - u_t \leq 0$$

Now, we can apply Lagrange's theorem to  $F$  to get

$$Lv = \left\langle \nabla F(\xi(x, t)), \left( t, x, u, \frac{\partial u}{\partial x_i}, \frac{\partial^2 u}{\partial x_i \partial x_j} \right) \right\rangle - u_t \leq 0$$

for a fixed  $\xi(x, t)$ .

Thus, the difference  $u$  of two sub-solutions to a non-linear differential problem is a sub-solution to a (different) *linear* parabolic problem, as the derivatives of  $F$  and  $\xi$  do not depend on  $u$  ( $\xi$  can be chosen a-priori).

We can thus see that this new problem must be parabolic, and apply the maximum principle and the Hopf's boundary point lemma to  $u$ . This can allow us to state the following two results which we will be using later:

**Proposition 1.12** (Maximum principle for parabolic non-linear differential equations). *Suppose we have two solution  $v$  and  $w$  in the interval  $[0, T]$  to the same parabolic non-linear differential equation (1.4) on an bounded open set  $\Omega$ , but with different start conditions. Suppose also that  $F$  is smooth on  $\bar{\Omega}$ . Then, if  $v > w$  in the interior of  $\Omega$  at  $t = 0$  and  $v \geq w$  on  $\partial\Omega$ ,  $v > w$  for all  $t \in [0, T]$  in the interior of  $\Omega$ .*

*Proof.*  $u = v - w \geq 0$  is a solution of a parabolic *linear* differential equation, where the term independent of  $u$  is bounded. Furthermore, if we take  $c(x, t)$  it must be bounded by compactness. At  $t = 0$ ,  $u > 0$  in the interior of  $\Omega$ . If, at an interior point  $x$ ,  $v = w$  at a certain time  $t = \tau$ ,  $u(\tau, x) = 0$ , and thus  $u$  is not constant. However, it attains minimum ( $u = 0$ ) at an interior point, thus by 1.6 it must be constant, a contradiction.  $\square$

**Proposition 1.13** (Hopf's boundary point lemma for parabolic non-linear differential equations). *Suppose we have two solution  $v$  and  $w$  to the same parabolic non-linear differential equation (1.4) in a region  $E$ , but with different start conditions. Suppose also that  $F$  is smooth on  $\bar{\Omega}$ . Let  $u = v - w$  and suppose that the maximum of  $u$  is attained at the point  $P$ . Furthermore, assume that the conditions on the shape of the region  $E$  from theorem 1.8 hold. Then,*

$$\frac{\partial u}{\partial \nu}(P) > 0$$

where we take  $\nu$  as the normal to  $\partial\Omega$ .

*Proof.*  $u$  is a solution of a parabolic *linear* differential equation, where  $c(x, t)$  is bounded by compactness. We can then apply 1.8 to  $v$  (see also remark 1.9).  $\square$

## 1.5 The Alexandrov Reflection Method (da scrivere)

[DA SCRIVERE]

## 1.6 The Alexandrov soap-bubble theorem (da scrivere)

[DA SCRIVERE]



# Chapter 2

## The Chow Critical Planes Result

**[FIX PARAGRAPH]** The main result we want to establish in this chapter is theorem 2.6, a result in Geometric Analysis and the main topic of the thesis. Before we can tackle it, one has to go first through a couple of classical results.

**[READ CHAPTER AND DOUBLE CHECK IT]**

### 2.1 Class of Equations we analyze

We consider manifolds  $M^n$  embedded in  $\mathbb{R}^{n+1}$ , i.e. there is an embedding  $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$  parametrizing the hypersurface  $X_0(M^n)$ .

Let  $F : \{(\kappa_1, \dots, \kappa_n) \in \mathbb{R}^n | \kappa_1 \leq \dots \leq \kappa_n\} \rightarrow \mathbb{R}$  be a  $C^1$  function satisfying:

$$\frac{\partial F}{\partial \kappa_i} > 0 \text{ for all } i = 1, \dots, n \quad (2.1)$$

and consider the evolution equation

$$\begin{cases} \frac{\partial X_t}{\partial t} = F(\kappa_1(x), \dots, \kappa_n(x))\nu \\ X(0) = X_0 \end{cases} \quad (2.2)$$

where  $\nu$  is the inward normal to  $X_t(M^n)$  at the point  $X_t(x)$  and  $\kappa_1 \leq \dots \leq \kappa_n$  are the principal curvatures at  $X_t(x)$ .

## 2.2 Parabolicity of the differential equation (2.2)

The condition (2.1) will guarantee that equation (2.2) is a parabolic equation. This may be confusing, as (2.2) does not make it obvious how to apply definition 1.10.

**[FIX PARAGRAPH]** In order to classify a non-linear partial differential equation one has to understand how it behaves “close to a solution” in the solutions space. We want to prove that very close to any solution, “moving in any direction”, the change in the equation is always a parabolic PDE. This will then tell us that our equation is parabolic, and that the theorems that apply to solutions of parabolic partial differential equations apply to our equation as well.

Like in [9], as  $F$  is a symmetric function in the principal curvatures, we may interchangeably take  $F$  to be a function of the Weingarten map tensor or of the second fundamental form, and thus we get:

$$\frac{\partial X_t}{\partial t} = F(h_{ij}(X_t))\nu$$

To understand the behaviour close to a solution, we can substitute in our equation  $X_t$  with a  $X_t + \varepsilon u_t$  to get:

$$\frac{\partial X_t}{\partial t} + \varepsilon \frac{\partial u_t}{\partial t} = F(h_{ij}(X_t + \varepsilon u_t))\nu_{(X_t + \varepsilon u_t)} \quad (2.3)$$

where we mean that  $\nu_{(X_t + \varepsilon u_t)}$  is the normal to the perturbed immersion. We are interested in the behaviour of this equation for a small  $\varepsilon$ . This equation when taking the limit for  $\varepsilon \rightarrow 0$  is the so-called linearisation of the PDE; we want this PDE to be a parabolic equation to apply our results.

We can use the Weingarten equation (1.1) to write the RHS explicitly:

$$\begin{aligned} h_{ij}(X_t + \varepsilon u_t) &= -\langle v, \nu_{(X_t + \varepsilon u_t)} \rangle \text{ where} \\ v^\alpha &= \frac{\partial^2 X_t^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial X_t^\alpha}{\partial x^k} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X_t^\beta}{\partial x^i} \frac{\partial X_t^\delta}{\partial x^k} + \\ &+ \varepsilon \left( \frac{\partial^2 u_t^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial u_t^\alpha}{\partial x^k} + \bar{\Gamma}_{\beta\delta}^\alpha \left( \frac{\partial X_t^\beta}{\partial x^i} \frac{\partial u_t^\delta}{\partial x^k} + \frac{\partial u_t^\beta}{\partial x^i} \frac{\partial X_t^\delta}{\partial x^k} \right) \right) + \\ &+ \varepsilon^2 \left( \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial u_t^\beta}{\partial x^i} \frac{\partial u_t^\delta}{\partial x^k} \right) \\ v^\alpha &= w + \varepsilon \left( \frac{\partial^2 u_t^\alpha}{\partial x^i \partial x^j} + \text{lower order terms} \right) + o(\varepsilon) \end{aligned}$$

where  $h_{ij}(X_t) = -\langle w, \nu_{(X_t+\varepsilon u_t)} \rangle$ . Putting it all together in the first line:

$$\begin{aligned} h_{ij}(X_t + \varepsilon u_t) &= -\left\langle w + \varepsilon \left( \frac{\partial^2 u_t}{\partial x^i \partial x^j} + \text{lower order terms} \right) + o(\varepsilon), \nu_{(X_t+\varepsilon u_t)} \right\rangle \\ &= h_{ij}(X_t) - \varepsilon \left\langle \left( \frac{\partial^2 u_t}{\partial x^i \partial x^j} + \text{lower order terms} \right), \nu_{(X_t+\varepsilon u_t)} \right\rangle + o(\varepsilon) \\ &= h_{ij}(X_t) - \varepsilon H + o(\varepsilon) \end{aligned}$$

We can then expand F in the RHS of the equation (2.3) to the first order, as it is a  $C^1$  function:

**[PROBABILMENTE HAI PERSO UN  
PRODOTTO SCALARE PER LA STRADA]**

$$\begin{aligned} \frac{\cancel{\partial X_t}}{\partial t} + \varepsilon \frac{\partial u_t}{\partial t} &= (\cancel{F(h_{ij}(X_t))} - \varepsilon DF(H) + o(\varepsilon)) \nu_{(X_t+\varepsilon u_t)} \\ \varepsilon \frac{\partial u_t}{\partial t} &= -\varepsilon \left( \frac{\partial F}{\partial h_{ij}} g^{ik} g^{jl} \frac{\partial^2 u_t}{\partial x^i \partial x^j} + \text{lower order terms} \right) \nu_{(X_t+\varepsilon u_t)} + o(\varepsilon) \\ \frac{\partial u_t}{\partial t} &= -\frac{\partial F}{\partial h_{ij}} g^{ik} g^{jl} \frac{\partial^2 u_t}{\partial x^i \partial x^j} \nu_{(X_t+\varepsilon u_t)} + \text{lower order terms} + o(1) \end{aligned}$$

Using a similar line of reasoning for  $\nu_{(X_t+\varepsilon u_t)}$  to only keep the highest order terms, using 1.2, and letting  $\varepsilon \rightarrow 0$ , we get to:

$$\frac{\partial u_t}{\partial t} = \frac{\partial F}{\partial h_{ij}} g^{ik} g^{jl} \left\langle \frac{\partial^2 u_t}{\partial x_k \partial x_l}, \nu \right\rangle \nu + \text{lower order terms}$$

For the second order term to be positive definite, then,

$$\left( \frac{\partial F}{\partial h_{ij}} \right)_{i,j}$$

must be positive definite. Or equivalently, as the principal curvatures are the eigenvalues of the matrix  $(h_{ij})_{i,j}$ ,

$$\frac{\partial F}{\partial \kappa_i} > 0$$

**Remark 2.1.** While in this chapter we are using the standard metric of  $\mathbb{R}^n$ , the proof above is valid using any other metric.

## 2.3 An existence result

The paper from Huisken and Polden [9] states the following comforting existence result for the class of equations we are analysing under very broad hypothesis. The same paper also includes a proof of the result with some more restrictive hypotheses. A complete proof is quite more involved. We will not use it, nor prove it, but it is included here for completeness and peace of mind:

**Theorem 2.2** (Short term existence of a solution for (2.2)). *Suppose  $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$  is a smooth, closed hypersurface in  $\mathbb{R}^{n+1}$ , such that (2.1) holds at all points in  $X_0$ , i.e. for all the values of the principal curvatures  $\kappa_i$  realized at some point on  $X_0$ . Then, (2.2) has a smooth solution, at least on some short time interval  $[0, T)$ ,  $T > 0$ .*

## 2.4 Local representation as a graph of a solution of (2.2)

As a first step, from the Corollary 1.5, we can establish the following:

**Theorem 2.3** (Local representation as a graph of a solution of (2.2)). *Let  $X^n$  be a submanifold  $X^n \subset \mathbb{R}^{n+1}$  and let  $F : X^n \times (0, T) \rightarrow \mathbb{R}^{n+1}$  be a solution of (2.2). Also let  $t \in (0, T)$  and  $x \in F(X^n, t)$ . Then there exists a neighbourhood of  $(x, t)$ ,  $U \subset F(X^n \times (0, T))$ , and a smooth function  $f : T_x F(X^n, t) \times (0, T) \rightarrow \mathbb{R}$  such that any  $(x_0, t_0) \in U$  can be expressed as*

$$x_0 = p + f(p, t_0)\nu$$

where  $\nu$  is a vector normal to  $T_x F(X^n, t)$ , for an appropriate point  $p \in T_x F(X^n, t)$ .

*Proof.* We can consider the image of  $F : X^n \times (0, T) \rightarrow \mathbb{R}^{n+1}$  as a manifold in  $\mathbb{R}^{n+2}$  by considering  $G := (F(x, t), t)$ . Moreover  $\frac{\partial G_i}{\partial e_j} \equiv 0$  for all possible vectors of the canonical basis of  $\mathbb{R}^{n+1} \times \mathbb{R}$  except for the one corresponding to the time coordinate, where it is 1. Also,  $\frac{\partial G_i}{\partial e_j} \equiv \frac{\partial F_i}{\partial e_j}$  and the first  $n$  coordinates of  $\frac{\partial G}{\partial t}$  form a vector normal to  $T_x X^n$  by (2.2). Thus, the tangent space of  $\text{Im}(G)$  is  $T_x X^n \times \{0\} \oplus \text{span}\langle(\nu, 1)\rangle$  and we can apply corollary 1.5 to  $\text{Im}(G)$  to get a function  $\tilde{f}$  such that

$$(x_0, t) = [(p, 0) + (s\nu, s)] + \tilde{f}(p, s)(\nu, -\|\nu\|)$$

as  $(\nu, -\|\nu\|)$  is the vector orthogonal to  $T_x X^n \times \{0\} \oplus \text{span}\langle(\nu, 1)\rangle$ . Let  $\sigma_p : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be the function that associates to  $t$  the appropriate  $s$  in the expression above.

Projecting to the first  $n + 1$  coordinates and calling  $f(p, t) = \sigma_p(t) + \tilde{f}(p, \sigma_p(t))$  one gets:

$$x_0 = p + f(p, t)\nu$$

which is our thesis, as long as  $\sigma_p(t)$  is smooth. This is indeed the case, as the graph function  $\Gamma_f : x \mapsto (x, f(x))$  is smooth for any smooth function  $f$  and has a smooth inverse (and thus, the inverse  $(x_0, t) \mapsto ((p, 0) + (s\nu, s))$  is smooth).  $\square$

One can show through direct calculation (see [10], Exercise 1.1.2) that, if an immersed hypersurface  $\phi : M \rightarrow \mathbb{R}^{n+1}$  is locally the graph of a function (i.e., locally,  $(x, f(x)) = \phi$ ), then:

$$\begin{aligned} g_{ij} &= \delta_{ij} + \frac{\partial f}{\partial x_i} \frac{\partial f}{\partial x_j} \\ \nu &= -\frac{(\nabla f, -1)}{\sqrt{1 + |\nabla f|^2}} \\ h_{ij} &= -\frac{H_{ij}}{\sqrt{1 + |\nabla f|^2}} \end{aligned}$$

Where the matrix  $(H_{ij})_{ij}$  is the hessian of  $f$ . Thus, if one considers the principal curvatures, they are closely related to the eigenvalues of the hessian of  $f$ .

## 2.5 The Alexandrov Reflection Method and the Chow result

The key idea of the main result in this chapter is as follows: suppose we have an embedded smooth hypersurface  $M^n$  evolving according to (2.2) and a fixed hyperplane  $\Pi$ , intersecting  $M^n$ . We can also consider  $M_\Pi^n$ , the reflection of  $M^n$  with respect to  $\Pi$ , which will also evolve according to (2.2). Suppose that, at some time  $t$ ,  $M^n$  and  $M_\Pi^n$  touch outside of  $\Pi$ . We can consider  $M^n$  and  $M_\Pi^n$  as local graphs over the same hyperplane  $\Pi$ , and we can show that these function evolve according to the same differential equation. Using the strong maximum principle and the Hopf boundary point lemma, then, one can conclude that the two functions coincide, and have been coinciding up until that point. We can then conclude that if  $M^n$  and  $M_\Pi^n$  only touch on  $M^n \cap \Pi$  at the beginning of the evolution, then they will never touch elsewhere. We shall make this more precise in the following paragraph, loosely based on paragraph 2 in [1].

## 2.6 Theorem 2.2 Chow [RENAME]

Let  $\Pi$  be a hyperplane in  $\mathbb{R}^{n+1}$ . We may assume  $\Pi$  orthogonal to a unit vector  $V \in \mathbb{R}^{n+1}$ , i.e.  $\langle \Pi, V \rangle = C$  for some constant  $C$ .

Then,  $\mathbb{R}^{n+1}$  is divided by  $\Pi$  into two half-spaces, which we will name  $H^+(\Pi) = \{x \in \mathbb{R} : \langle x, V \rangle > C\}$  and  $H^-(\Pi) = \{x \in \mathbb{R} : \langle x, V \rangle < C\}$ .

**Definition 2.4.** We say we can reflect  $M^n$  strictly with respect to  $\Pi$  if

$$M_\Pi^n \cap H^-(\Pi) \subset \text{int}(M^n) \cap H^-(\Pi)$$

and  $V \notin TM_x^n$  for all  $x \in M^n \cap \Pi$ , where  $\text{int}(M^n)$  is the region inside  $M^n$ .

This fundamentally means that the reflection of one of the halves of  $M^n$  on the other side of  $\Pi$  is contained in the region inside  $M^n$  and the tangent spaces of  $M^n$  and of the half-reflection do not form a ninety degree angle with  $\Pi$ , at all points on  $\Pi \cap M^n$ .

**Definition 2.5.** We say we can reflect  $M^n$  strictly up to  $(\Pi, V)$  if we can reflect  $M^n$  strictly with respect to  $\Pi_K$  for all hyperplanes  $\Pi_K = \{x \in \mathbb{R} : \langle x, V \rangle = K\}$  such that  $K < C$ .

The main theorem is the following:

**Theorem 2.6** (Chow). Let  $X : M^n \times [0, T] \rightarrow \mathbb{R}^{n+1}$  be a  $C^2$  solution to 2.2. Then, if we can reflect  $X(M^n, 0) = M_0$  strictly with respect to  $\Pi$ , then for all  $t \in [0, T]$  we can reflect  $X(M^n, t) = M_t$  strictly with respect to  $\Pi$ .

*Proof.* By contradiction, suppose that there is a time  $t$  such that the thesis is false, and that it is the smallest such  $t$ . Then, for all  $\tau \in [0, t)$ ,  $M_{\tau, \Pi} \cap H^-(\Pi) \subset \text{int}(M_\tau) \cap H^-(\Pi)$ ; the unit vector orthogonal to  $\Pi$ ,  $V$ , is such that  $V \notin T_x M_\tau$  for all  $x \in M_\tau \cap \Pi$  and  $\tau \in [0, t)$ ; and either of the conditions fails at  $t$ , i.e. either:

(i)  $M_{t, \Pi} \cap H^-(\Pi) \cap M_t \neq \emptyset$

(ii)  $V \in T_x M_t$  for some  $x \in \Pi$ .

(i) Suppose the first case is true. Then, there exists  $x_0 \in M_{t, \Pi} \cap H^-(\Pi) \cap M_t$  such that at  $x_0$  the two manifolds are tangent.

We can take a neighbourhood of  $(x_0, t) \in M_t \times \mathbb{R}$  such that both  $M_{t, \Pi}$  and  $M_t$  are graphs over  $T_{x_0} M_t$  by 2.3.

We can explicitly write the functions  $f : U \times (t - \varepsilon, t + \varepsilon) \rightarrow M_t$ , where  $U \subset T_{x_0} M_t$ , and the corresponding  $f_\Pi$  for  $M_{t, \Pi}$ . We can also write

$$f : (x, t) \mapsto x + \tilde{f}(x, t)\nu$$

$$f_\Pi : (x, t) \mapsto x + \tilde{f}_\Pi(x, t)\nu$$

for appropriate functions  $\tilde{f} : U \times (t - \varepsilon, t + \varepsilon) \rightarrow \mathbb{R}$  and  $\tilde{f}_\Pi : U \times (t - \varepsilon, t + \varepsilon) \rightarrow \mathbb{R}$ , where  $\nu$  is a fixed unit vector normal to  $T_{x_0}M_t$ .  $\tilde{f}$  and  $\tilde{f}_\Pi$  are solutions to the same second order PDE, which is parabolic by what was discussed in paragraph 2.2, hence we can apply Proposition 1.12 to conclude that  $\tilde{f} \equiv \tilde{f}_\Pi$ , and thus  $M_{t,\Pi}$  and  $M_t$  coincide in a neighbourhood of  $(x, t)$ , a contradiction as we assumed that  $t$  is the first  $t$  where the flows touch.

(ii) Suppose instead that  $V \in T_x M_t$  for some  $t \in [0, t)$  and some  $x \in M_t \cap \Pi$ . Then  $T_x M_t = T_x M_{t,\Pi}$  and in a neighbourhood of  $(x, t)$  both  $M_t$  and  $M_{t,\Pi}$  are graphs of two smooth functions over  $T_x M_t$  by 2.3, i.e. again

$$\begin{aligned} f &: (x, t) \mapsto x + \tilde{f}(x, t)\nu \\ f_\Pi &: (x, t) \mapsto x + \tilde{f}_\Pi(x, t)\nu \end{aligned}$$

Moreover, in  $\overline{H^-(\Pi)}$ ,  $f_\Pi \geq f$ , because  $M_\Pi^n \cap H^-(\Pi) \subset \text{int}(M^n) \cap H^-(\Pi)$ . Finally,  $f(x, t) = f_\Pi(x, t)$ , hence  $f_\Pi - f(x, t) = 0$ , and thus  $(x, t)$  is a minimum point on the boundary for  $f_\Pi - f$ . Also, we must have

$$\frac{\partial f}{\partial V}(x, t) = \frac{\partial f_\Pi}{\partial V}(x, t)$$

because the graphs are both tangent to  $T_x M_t$ , and  $V$  here is the outward pointing normal to the boundary by definition of the reflection. Thus,

$$\frac{\partial(f - f_\Pi)}{\partial V}(x, t) = 0$$

But we must have

$$\frac{\partial(f - f_\Pi)}{\partial V}(x, t) > 0$$

at a minimum on the boundary by Proposition 1.13, a contradiction.  $\square$

The following is an immediate consequence of the main result:

**Corollary 2.7.** *Let  $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a  $C^2$  solution to 2.2. Then, if we can reflect  $M_0$  strictly up to  $(\Pi, V)$ , then for all  $t \in [0, T)$  we can reflect  $X(M, t)$  strictly up to  $(\Pi, V)$ .*

*Proof.* The hypothesis of the theorem are true for each  $\Pi_K$  in the definition, thus we can reflect strictly with respect to each  $\Pi_K$  for all  $t \in [0, T)$ , and thus we can reflect  $X(M, t)$  strictly up to  $(\Pi, V)$ .  $\square$





## Chapter 3

### Third Chapter Title

[DA SCRIVERE]

3.1 Reflections on spheres and hyperbolic spaces  
(da scrivere)

[DA SCRIVERE]

## **3.2 The equation in constant curvature spaces (da scrivere)**

**[DA SCRIVERE]**

### **3.3 Extension of the result(da scrivere)**

**[DA SCRIVERE]**

### **3.4 Extension to constant curvature (da scrivere)**

**[DA SCRIVERE]**

# Bibliography

- [1] Chow B.; *Geometric Aspects of Aleksandrov Reflection and Gradient Estimates for Parabolic Equations*, Communications in Analysis and Geometry, Volume 5, Number 2, 389-409, 1997.
- [2] Gidas B., Wei-Ming Ni, L.Nirenberg; *Symmetry and Related Properties via the Maximum Principle*, Commun. Math. Phys. 68, 209-243, 1979.
- [3] Protter, M. H.; Weinberger, H. F.; *Maximum Principles in Differential Equations*, Springer, 1984.
- [4] Evans, L.C.; *Partial Differential Equations*, American Mathematical Society, 2010.
- [5] Do Carmo, M.P.; *Differential Geometry of Curves and Surfaces, Second Edition*, Dover Books on Mathematics, 2016.
- [6] Lee, J.M.; *Riemannian Manifolds*, Springer Graduate Texts in Mathematics 176, 1997.
- [7] Lee, J.M.; *Introduction to Smooth Manifolds, Second Edition*, Springer Graduate Texts in Mathematics 218, 2013.
- [8] J. Scheuer; *Isotropic functions revisited*, Arch. Math. 110, 591–604, 2018.
- [9] G. Huisken, A. Polden *Geometric evolution equations for hypersurfaces*
- [10] C.Mantegazza *Lecture Notes on Mean Curvature Flow*
- [11] R.S. Hamilton, *Three-Manifolds with Positive Ricci Curvature*