

UNIVERSITÀ DEGLI STUDI DI ROMA TOR VERGATA
MACROAREA DI SCIENZE MATEMATICHE, FISICHE E NATURALI



LAUREA MAGISTRALE IN MATEMATICA PURA E APPLICATA

TITOLO

TO BE CONFIRMED WHEN IT IS DONE

Relatore:

Prof. Carlo Sinestrari

Laureando:

matricola: 0318523

Marco Tamburro

Anno Accademico 2023/2024

Contents

Introduction	5
1 The Alexandrov Reflection Method	7
1.1 Geometry of immersed hypersurfaces	7
1.2 Local representation as a graph	11
1.3 Some well established results from analysis	12
1.4 Applying the maximum principle to non-linear PDEs	16
1.5 Reflections on spheres and hyperbolic spaces	19
1.6 The Method of Moving Planes	23
1.7 The Alexandrov soap-bubble theorem	24
2 The Chow-Gulliver Critical Planes Result	29
2.1 Class of Equations we analyze	29
2.2 Parabolicity of the differential equation (2.2)	30
2.3 An existence result	32
2.4 Local representation as a graph of a solution of (2.2)	32
2.5 The Moving Planes Method and the Chow result	34
2.6 The Chow-Gulliver result	35
2.7 Some corollaries of the result	38
2.8 Applying the result to find gradient estimates	41
2.9 Expansive flows and ancient solutions	46
3 Extension to constant curvature spaces	49
3.1 The equation in constant curvature spaces	49
3.2 Extension of the result	50
3.3 Extending Corollaries	51
3.4 Extending Ancient solutions section	52
Bibliography	55

Introduction

To be confirmed

Chapter 1

The Alexandrov Reflection Method

1.1 Geometry of immersed hypersurfaces

The study of immersed hypersurfaces is a fundamental topic in differential geometry, and is especially important in the field of geometric analysis. An immersed hypersurface $X : M^n \rightarrow \overline{M}^{n+1}$ is a submanifold of a higher-dimensional space that is embedded in that space in such a way that the submanifold has the same dimension as the space in which it is embedded minus one. In other words, it is a submanifold of codimension one. We will always assume the embedding to be smooth. The most common case is $\overline{M}^{n+1} = \mathbb{R}^{n+1}$, usually extensively studied in undergraduate courses when $n = 2$. We will be assuming that the embedding is also isometric, i.e. the metric g on M^n is the one induced by $(\overline{M}^{n+1}, \overline{g})$.



Figure 1.1: A (hyper)surface immersed in \mathbb{R}^3

Figure 1.2: The tangent and the normal bundle at a point p

Symbols referring to $(\overline{M}^{n+1}, \bar{g})$ will have a line on top, otherwise the symbol will refer to (M^n, g) .

The pullback of the tangent bundle of \overline{M}^{n+1} to M^n is a smooth vector bundle on M^n :

$$X^*T\overline{M}^{n+1} = T\overline{M}^{n+1}|_{M^n} = \coprod_{p \in M^n} T_p\overline{M}^{n+1}$$

One of the most important concepts in the geometry of immersed hypersurfaces is the concept of a normal vector field. The normal vector field ν is a section of the pullback vector bundle $X^*T\overline{M}^{n+1}$ on the manifold M^n that is perpendicular to the tangent space of M^n at each point. At each point p :

$$T_p\overline{M}^{n+1} = T_pM^n \oplus N_pM^n$$

where N_pM^n is the normal vector bundle generated by the normal vector. If the manifold is oriented, there is usually a choice we can make on the normal vector:

we will generally chose the outward pointing normal vector. This also allows us to define the tangent and normal projection on $T\overline{M}^{n+1}|_{M^n}$ by taking the two respective components.

Clearly, taking $\overline{\nabla}$ to be the Levi-Civita connection on $(\overline{M}^{n+1}, \overline{g})$, we can decompose it as:

$$\overline{\nabla}_v w = (\overline{\nabla}_v w)^\top + (\overline{\nabla}_v w)^\perp$$

Definition 1.1. *The second fundamental form is then defined as:*

$$\mathbb{I}(v, w) = (\overline{\nabla}_v w)^\perp$$

It is a bilinear symmetric tensor because TM^n is involutive in $T\overline{M}^{n+1}$ and depends only on the local value of v and w by symmetry. we can therefore write it as

$$\mathbb{I}(v, w) = -(h_{ij}v^i w^j)\nu$$

for some matrix $A(p) = \{h_{ij}\}$. we can define the principal curvatures of the hypersurface, as the eigenvalues of this matrix.

It is also possible to check that $(\overline{\nabla}_v w)^\top$ satisfies the definition the Levi-Civita connection on (M^n, g) , therefore, from its uniqueness:

$$\begin{aligned}\nabla_v w &= (\overline{\nabla}_v w)^\top \\ \overline{\nabla}_v w &= \nabla_v w + \mathbb{I}(v, w)\end{aligned}$$

where ∇ is the Levi-Civita connection on (M^n, g) . This result is known as the *Gauss Formula*. It has to be noted however that we are implicitly considering tangent vectors that are not in the same space. Indeed, making that more explicit, the formula should be:

$$\overline{\nabla}_{X_*v} X_*w = X_*(\nabla_v w) + \mathbb{I}(v, w)$$

Proposition 1.2. (The Weingarten Equation) *If $v, w \in TM^n$ and $\nu \in NM^n$, if one considers the corresponding derivations in $T\overline{M}^{n+1}$ the following equation holds:*

$$\langle \overline{\nabla}_v \nu, w \rangle_{\overline{g}} = -\langle \nu, \mathbb{I}(v, w) \rangle_{\overline{g}}$$

Proof. As $\langle \nu, w \rangle_{\overline{g}} \equiv 0$ on M ,

$$\begin{aligned}0 &= v \langle \nu, w \rangle_{\overline{g}} \\ &= \langle \overline{\nabla}_v \nu, w \rangle_{\overline{g}} + \langle \nu, \overline{\nabla}_v w \rangle_{\overline{g}} \\ &= \langle \overline{\nabla}_v \nu, w \rangle_{\overline{g}} + \langle \nu, \mathbb{I}(v, w) \rangle_{\overline{g}}\end{aligned}$$

applying the Gauss Formula and the fact that $\nabla_v w \in TM^n$ in the last step \square

It is also usual to define the associated Weingarten map, which is the linear map between sections of M $s : \Gamma(M) \rightarrow \Gamma(M)$ satisfying:

$$\langle s(v), w \rangle_g = \langle \nu, \mathbb{I}(v, w) \rangle_{\bar{g}}$$

the linear map s is also known as the *shape operator* of M .

Combining this with the formula above, taking into account that it holds for a generic $w \in TM$:

$$s(v) = -(\bar{\nabla}_v \nu)^\top$$

We are going to use these equations in local coordinates, in the form shown below.

Proposition 1.3. *The above equations in local coordinates are equivalent to the following equations:*

$$\frac{\partial^2 X^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial X^\alpha}{\partial x^k} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X^\beta}{\partial x^i} \frac{\partial X^\delta}{\partial x^j} = -h_{ij} \nu^\alpha \quad (1.1)$$

$$\frac{\partial \nu^\alpha}{\partial x^i} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X^\beta}{\partial x^i} \nu^\delta = h_{ij} g^{jl} \frac{\partial X^\alpha}{\partial x^l} \quad (1.2)$$

where ν is the normal unit vector at the point and $A = \{h_{ij}\}$ is the second fundamental form, thus $h_{ij} = \langle \nu, \bar{\nabla}_{\partial_i} \bar{\partial}_j \rangle_{\bar{g}}$

Proof. For any connection ∇ . and any derivations $v = v^i \partial_i$ and $w = w^j \partial_j$:

$$\nabla_v w = \nabla_{(v^i \partial_i)} (w^j \partial_j) = v(w^k) \partial_k + (v^i w^j \Gamma_{ij}^k) \partial_k$$

Let $\partial_1, \dots, \partial_n$ be a basis of TM^n at a point, and let $\bar{\partial}_i = X_* \partial_i$, $\bar{\partial}_{n+1} = \nu$. Let's consider the Gauss Formula for two generic ∂_i, ∂_j , using roman letters for indices varying between 1 and n and greek letters for indices varying between 1 and $n+1$, and δ^{ij} the Kronecker delta:

$$\begin{aligned} \bar{\nabla}_{X_* \partial_i} X_* \partial_j &= X_*(\nabla_{\partial_i} \partial_j) + \mathbb{I}(\partial_i, \partial_j) \\ (\bar{\partial}_i (X_* \partial_j)^\alpha) \bar{\partial}_\alpha + ((X_* \partial_i)^\beta (X_* \partial_j)^\delta \bar{\Gamma}_{\beta\delta}^\alpha) \bar{\partial}_\alpha &= X_*(\Gamma_{ij}^k \partial_k) - h_{ij} \nu^\alpha \bar{\partial}_\alpha \\ \frac{\partial^2 X^\alpha}{\partial x^i \partial x^j} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X^\beta}{\partial x^i} \frac{\partial X^\delta}{\partial x^j} &= \Gamma_{ij}^k \frac{\partial X^\alpha}{\partial x^k} - h_{ij} \nu^\alpha \end{aligned}$$

Which is the formula (1.1). To get the second formula, first note that $s(v) = -(\bar{\nabla}_v \nu)^\top$. We then compute $-\langle s(\partial_i), \bar{\partial}_\alpha \rangle_{\bar{g}}$:

$$\begin{aligned} \langle \bar{\nabla}_{\partial_i} \nu, \bar{\partial}_\alpha \rangle_{\bar{g}} &= -\langle s(\partial_i), \bar{\partial}_\alpha \rangle_{\bar{g}} \\ \left\langle \left(\frac{\partial \nu^\alpha}{\partial x^i} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X^\beta}{\partial x^i} \nu^\delta \right) \bar{\partial}_\alpha, \bar{\partial}_\alpha \right\rangle_{\bar{g}} &= h_{ij} g^{jl} \frac{\partial X^\alpha}{\partial x^l} \end{aligned}$$

leading to (1.2). □

1.2 Local representation as a graph

A well known result, Dini's Theorem¹, states that, given a smooth function F defined on an open subset of the product space $\mathbb{R}^n \times \mathbb{R}^m$, if $F(x, y) = 0$ and the partial derivative of F with respect to y is nonzero at a point (x_0, y_0) , then there exists an open neighborhood of x_0 in \mathbb{R}^n and a unique smooth function $y = g(x)$ defined on that neighborhood such that y_0 is a regular value of g and $(x, g(x))$ is a smooth solution to the equation $F(x, y) = 0$.

Consequence of the Dini's theorem is a powerful result that allows one to locally represent a submanifold of $\mathbb{R}^n \times \mathbb{R}^m$ as the graph of a smooth function. This theorem is widely used in differential geometry, geometric analysis, and many other fields of mathematics and physics. We provide a version of this theorem below:

Theorem 1.4 (Local representation as a graph). *Let X^n be a submanifold $X^n \subset \mathbb{R}^{n+1}$ and let $x_0 \in X$. Then there exists a neighbourhood of x_0 , $U \subset X^n$, such that U is the graph of a function. Moreover, this function can be of the form*

$$f : \pi(U) \subset \mathbb{R}^n \rightarrow \mathbb{R}$$

$$U = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} | x_0 = f(x_1, \dots, x_n)\}$$

for any of the possible orders of the usual basis for \mathbb{R}^n , (e_0, \dots, e_n) , as long as $e_0 \notin T_x M$, where $\pi(U)$ is the projection on the last n coordinates $((x_0, \dots, x_n) \mapsto (x_1, \dots, x_n))$.

A proof of the 2D-case of the version of the theorem can be found in [13] which extends naturally to the n dimensional case, with almost no changes. This immediately extends to:

Corollary 1.5 (Local representation as a graph on the tangent). *Let X^n be a submanifold $X^n \subset \mathbb{R}^{n+1}$ and let $x_0 \in X$. Then there exists a neighbourhood of x_0 $U \subset X^n$ and a smooth function $f : T_x X^n \rightarrow \mathbb{R}$ such that any $x_0 \in U$ can be expressed as*

$$x_0 = p + f(p)\nu$$

where ν is the vector normal to $T_{x_0} X^n$, for an appropriate point $p \in T_{x_0} X^n$. In other words, every submanifold $X^n \subset \mathbb{R}^{n+1}$ is locally expressible as a graph on its tangent space.

Proof. By rotation, we may assume $T_x X^n$ orthogonal to e_1 . Then one can just apply the previous theorem. \square

We will use this later to prove Theorem 2.3.

¹also known as Implicit Function Theorem

1.3 Some well established results from analysis

We now introduce some well known results from analysis. The first result we introduce is the maximum principle.

The maximum principle is a classical result of mathematical analysis, and it is usually introduced in a first course on partial differential equations. It is a fundamental tool in the theory of partial differential equations. It is a statement about the behavior of solutions to certain types of PDEs and provides a method for obtaining upper and lower bounds on the solutions. The principle states that the maximum and minimum values of a solution to elliptic or parabolic PDE occur on the boundary of the domain unless the function is constant.

The maximum principle can be used to prove the existence, uniqueness, and regularity of solutions to elliptic and parabolic PDEs. It can also be used to obtain estimates on the behavior of solutions and to study the asymptotic behavior of solutions as the domain becomes large. The principle is widely used in many fields of mathematics and physics, such as geometric analysis, mathematical physics, and fluid dynamics. One of the many versions of this well know theorem is this:

Theorem 1.6 (Maximum principle for parabolic equations). *Let Ω be an open, bounded, connected set. Assume $u \in C_1^2(\Omega \times [0, T]) \cap C^1(\bar{\Omega} \times [0, T])$. Suppose u satisfies:*

$$-\frac{\partial u}{\partial t} + \left(\sum_{i,j=1}^n a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_i b_i \frac{\partial}{\partial x_i} + c \right) u = -u_t + Lu \geq 0 \quad (1.3)$$

where L is an elliptic differential operator, i.e. there exists $\theta > 0$ such that $\sum_{i,j=1}^n a_{ij}(x, t) \xi_i \xi_j \geq \theta |\xi|^2$ for all $\xi \in \mathbb{R}^n$ and $(x, t) \in \Omega \times [0, T]$. Suppose also that $c \equiv 0$ in Ω . Then:

- if u attains its maximum in an interior point $(x_0, t_0) \in \Omega \times [0, T]$, then u is constant in $\Omega \times [0, t_0]$.
- If, instead, under the same conditions, $u_t + Lu \geq 0$ and u attains its minimum in an interior point of $\Omega \times [0, T]$, then u is constant in $\Omega \times [0, t_0]$

A proof of this result can be found, for example, in [11]. The theorem extends also to situations where the condition holds in a convex bounded connected region $R \subseteq \Omega \times [0, T]$: in that case, if u attains its maximum in an interior point then u has the same value at any point in R that can be connected to it through a segment going in the backwards direction of time and a "horizontal" line contained in Ω . This version of the theorem can be found for example in [8]:

Theorem 1.7. *Let u satisfy the uniformly parabolic differential inequality (1.3) with $c(x) \leq 0$ in a region $R_T = \{(x_1, x_2, \dots, x_n, t) | t \leq T\}$ where R is a non-empty connected open set, and suppose that the coefficients of L are bounded. Suppose that the maximum of u in R_T is M and that it is attained at a point (x, t) of R_T . Thus if (y, s) is a point of R which can be connected to (x, t) by a path in R consisting only of horizontal segments and upward vertical segments, then $u(y, s) = M$.*

Hopf's boundary point lemma is another important classical tool in the study of PDEs that provides a criterion for determining the behavior of solutions to certain types of elliptic or parabolic PDEs near the boundary of the domain. The lemma states that if one has a solutions to some kinds of partial differential inequalities, then the normal derivative of the solution at that point is strictly positive.

It is often used to obtain estimates on the behavior of solutions near the boundary, and to prove the existence and uniqueness of solutions to boundary value problems. The lemma is named after the German mathematician Eberhard Hopf, who first formulated it in the 1950s. In [8] we find the following version of the Hopf's boundary point lemma:

Theorem 1.8. *Let u be a solution to the parabolic inequality*

$$-u_t + Lu \geq 0$$

with L an elliptic linear differential operator with bounded coefficients such that $c(x) \leq 0$, in a domain E , and let $E_t = \{(x, s) \in E | s \leq t\}$. Suppose the maximum M of u is attained at a point $P = (x, t)$ on the boundary ∂E .

Assume that a sphere through P can be constructed which is in E_s such that

- *tangent to ∂E at P*
- *the set of point of its interior (y, s) such that $s \leq t$ lies in E_s ,*
- *$u < M$ in its interior.*

Also, suppose that the radial direction from the centre of the sphere to P is not parallel to the t -axis.

Then, if $\frac{\partial}{\partial \nu}$ denotes any directional derivative in an outward direction from E_s , we have

$$\frac{\partial u}{\partial \nu} > 0$$

at P .




Figure 1.3: The setup in the proof at time t when there are two space dimensions. The region R also extends for times $s < t$ between the two spheres. The centre of B_1 can have time-coordinate different from t if the domain E is not “straight” in the time direction.

Proof. Let the sphere through P be B_1 . We may construct a smaller sphere B_2 centred at P . Let now:

$$\begin{aligned} S_1 &= \partial B_1 \cap B_2 \cap E_t, \\ S_2 &= B_1 \cap \partial B_2 \cap E_t, \text{ and} \\ S_3 &= B_1 \cap B_2 \cap \partial E_t = B_1 \cap B_2 \cap \{(x, s) \in E | s = t\}. \end{aligned}$$

The three sets satisfy $S_1 \cup S_2 \cup S_3 = \partial(B_1 \cap B_2 \cap E_t)$, we may call this region $R = B_1 \cap B_2 \cap E_t$. Without loss of generality, potentially taking a smaller sphere B_1 , we may assume that $u < M$ on B_1 except at P . As $R \subset B_1$, we also get $u < M$ on R . We may thus conclude that:

- $u < M$ on R except at P
- $u \leq M - \delta$ on S_2 for a sufficiently small $\delta > 0$
- $u = M$ at P .

Now, let the centre of B_1 be $Q = (z, t_0)$ and let r be its radius. we can now introduce the function

$$v(y, s) = \exp \left(-\alpha(s - t_0)^2 - \sum_{i=1}^n \alpha(y_i - z_i)^2 \right) - \exp(-\alpha r^2)$$

This function is such that $v(y, s) = 0$ if $(y, s) \in S_1$ - including $v(x, t) = 0$, as there the first term is $e^{-\alpha r^2}$, and $v(y, s) > 0$ in the interior of B_1 .

Thus, in the region R , $v(y, s) \geq 0$ and has a minimum point at the boundary on (x, t) , where $v(x, t) = 0$.

We can also compute Lv . After some calculation, we get that

$$\begin{aligned} Lv &= 2\alpha e^{(-\alpha(s-t_0)^2 - \sum_{i=1}^n \alpha(y_i - z_i)^2)} [2\alpha(y - z)^t A(y - z) + \\ &\quad + \sum_i^n [b_i(y_i - z_i) + a_{i,i}] + (s - t)] \end{aligned}$$

where A is the matrix of the $a_{i,j}$. In particular, one can choose an α large enough, so that $Lv > 0$ in $R \cup \partial R$.

We can thus introduce $w = u + \varepsilon v$. As both Lu and Lv are positive in R , $Lw > 0$ in R . We can also choose ε small enough so that $w < M$ on S_2 . Also, as $v = 0$ on S_1 , $w < M$ on S_1 except at P , and $w = M$ at P .

Therefore, we can apply the Strong Maximum Principle 1.7 to the region R to conclude that the maximum of w in R is attained at P . Therefore:

$$\frac{\partial w}{\partial \nu} = \frac{\partial u}{\partial \nu} + \varepsilon \frac{\partial v}{\partial \nu} \geq 0$$

But:

$$\frac{\partial v}{\partial \nu} = \nu \cdot n \frac{\partial v}{\partial R} = -2\nu \cdot n \alpha R e^{-\alpha R} < 0$$

Where n is the vector orthogonal to the sphere S_1 . Therefore, one must have:

$$\frac{\partial u}{\partial \nu} > 0$$

as we wanted. □

Remark 1.9. If $c(x)$ is now just bounded, we can consider, instead of u , $v = ue^{-\lambda t}$, thus, by change of variables

$$-v_t + Lv - \lambda v \geq 0$$

whenever $-u_t + Lu \geq 0$, and we can chose λ large enough such that $c(x) - \lambda < 0$ and thus we can remove the hypothesis $c(x) \leq 0$ in both theorems when c is bounded.

1.4 Applying the maximum principle to non-linear PDEs

We can now introduce an important observation shown in [8] that allows us to apply the maximum principle 1.7 and Hopf's boundary point lemma 1.8 in some non-linear settings. Firstly, we must clarify what we mean by parabolic non-linear problem.

Definition 1.10. *A differential non-linear problem in the form*

$$Lu = F\left(t, x, v, \frac{\partial v}{\partial x_i}, \frac{\partial^2 v}{\partial x_i \partial x_j}\right) - v_t = f(x, t) \quad (1.4)$$

given a smooth F is parabolic if for any real vector ξ

$$\sum_{i,j=1}^n F_{ij} \xi_i \xi_j > 0$$

where F_{ij} are the derivatives of F with respect to $\frac{\partial^2 v}{\partial x_i \partial x_j}$.

Secondly, we remind the reader of the following generalized version of the theorem of the mean, a.k.a. Lagrange's theorem:



Figure 1.4: In one dimension, Lagrange's theorem states that given a smooth function f on an interval $[a, b]$, there exists a ξ such that $f(b) - f(a) = f'(\xi)(b - a)$.

Theorem 1.11 (Lagrange's theorem). *Given a convex open set $U \subseteq \mathbb{R}^n$ and a real function $F \in C^1(U)$, and given to points x, y in U , there exists a point z in the segment connecting x and y such that*

$$F(y) - F(x) = \langle \nabla F(z), (y - x) \rangle$$

Suppose that we have a solution to a non-linear parabolic problem v , i.e. v solves (1.4):

$$Lu = F\left(t, x, v, \frac{\partial v}{\partial x_i}, \frac{\partial^2 v}{\partial x_i \partial x_j}\right) - v_t = f(x, t)$$

for a non-linear elliptic operator L in some region E , where we assume $F(t, x, y_i, z_{i,j})$ to be a given C^1 function. Suppose also that there is a w which is a solution of the corresponding differential inequality:

$$Lw = F\left(t, x, w, \frac{\partial w}{\partial x_i}, \frac{\partial^2 w}{\partial x_i \partial x_j}\right) - w_t \leq f(x, t)$$

One can then consider $u = v - w$, and by combining the above we get:

$$Lv = \left(F\left(t, x, v, \frac{\partial v}{\partial x_i}, \frac{\partial^2 v}{\partial x_i \partial x_j}\right) - F\left(t, x, w, \frac{\partial w}{\partial x_i}, \frac{\partial^2 w}{\partial x_i \partial x_j}\right) \right) - u_t \leq 0$$

Now, we can apply Lagrange's theorem to F to get

$$Lv = \left\langle \nabla F(\xi(x, t)), \left(t, x, u, \frac{\partial u}{\partial x_i}, \frac{\partial^2 u}{\partial x_i \partial x_j} \right) \right\rangle - u_t \leq 0$$

for a fixed $\xi(x, t)$.

Thus, the difference u of two sub-solutions to a non-linear differential problem is a sub-solution to a (different) *linear* parabolic problem, as the derivatives of F and ξ do not depend on u (ξ can be chosen a-priori).

We can thus see that this new problem must be parabolic, and apply the maximum principle and the Hopf's boundary point lemma to u . This can allow us to state the following two results which we will be using later:

Proposition 1.12 (Maximum principle for parabolic non-linear differential equations). *Suppose we have two solution v and w in the interval $[0, T]$ to the same parabolic non-linear differential equation (1.4) on an bounded open set Ω , but with different start conditions. Suppose also that F is smooth on $\bar{\Omega}$. Then, if $v > w$ in the interior of Ω at $t = 0$ and $v \geq w$ on $\partial\Omega$, $v > w$ for all $t \in [0, T]$ in the interior of Ω .*

Proof. $u = v - w \geq 0$ is a solution of a parabolic *linear* differential equation, where the term independent of u is bounded. Furthermore, if we take $c(x, t)$ it must be bounded by compactness. At $t = 0$, $u > 0$ in the interior of Ω . If, at an interior point x , $v = w$ at a certain time $t = \tau$, $u(\tau, x) = 0$, and thus u is not constant. However, it attains minimum ($u = 0$) at an interior point, thus by 1.6 it must be constant, a contradiction. \square

Proposition 1.13 (Hopf's boundary point lemma for parabolic non-linear differential equations). *Suppose we have two solution v and w to the same parabolic non-linear differential equation (1.4) in a region E , but with different start conditions. Suppose also that F is smooth on $\bar{\Omega}$. Let $u = v - w$ and suppose that the maximum of u is attained at the point P . Furthermore, assume that the conditions on the shape of the region E from theorem 1.8 hold. Then,*

$$\frac{\partial u}{\partial \nu}(P) > 0$$

where we take ν as the normal to $\partial\Omega$.

Proof. u is a solution of a parabolic *linear* differential equation, where $c(x, t)$ is bounded by compactness. We can then apply 1.8 to v (see also remark 1.9). \square

1.5 Reflections on spheres and hyperbolic spaces

In what follows, we will focus on manifolds embedded in spaces which have constant sectional curvature. Constant curvature manifolds are classified into three types based on the sign of the curvature:

- **Positive curvature:** *Spherical geometry*, where the curvature is positive and the manifold locally resembles a sphere (e.g., the standard sphere \mathbb{S}^n).
- **Zero curvature:** *Flat geometry*, where the curvature is zero and the manifold locally resembles Euclidean space \mathbb{R}^n .
- **Negative curvature:** *Hyperbolic geometry*, where the curvature is negative and the manifold locally resembles hyperbolic space \mathbb{H}^n .

As in [3], we will use the symbol \mathbb{M}^n to indicate a Riemannian manifold that can be replaced by any one of \mathbb{S}^n , \mathbb{R}^n or \mathbb{H}^n : the n -dimensional sphere, Euclidean plane or Hyperbolic space respectively.

We will also use the symbol \mathbb{M}_+^n to indicate a Riemannian manifold that can be replaced by \mathbb{S}_+^n , \mathbb{R}^n or \mathbb{H}^n : the n -dimensional hemisphere, Euclidean plane or Hyperbolic space, respectively.

Definition 1.14. \mathbb{H}^n is the n -dimensional hyperbolic plane. We can define it as the half space $\{x \in \mathbb{R}^n | x_n > 0\}$ with the Riemannian metric

$$g_x = \frac{1}{x_n^2} \langle \cdot, \cdot \rangle$$

where $\langle \cdot, \cdot \rangle$ is the Euclidean dot product on \mathbb{R}^n

Remark 1.15. This is not the only way we could define the hyperbolic space \mathbb{H}^n . Another alternative is taking the n -dimensional disc $D^n = \{x \in \mathbb{R}^n | \|x\| < 1\}$ with the Riemannian metric

$$g_x = \frac{4}{(1 - \|x\|^2)^2} \langle \cdot, \cdot \rangle$$

where $\langle \cdot, \cdot \rangle$ is the Euclidean dot product on \mathbb{R}^n .

Remark 1.16. Let $\mathbb{S}^n \setminus \{P\}$ be the standard n -dimensional unitary sphere minus a point, with the standard induced Euclidean metric. Through stereographic projection it is isometric to \mathbb{R}^n with metric:

$$g_x = \frac{4}{(1 + \|x\|^2)^2} \langle \cdot, \cdot \rangle$$

where $\langle \cdot, \cdot \rangle$ is again the Euclidean dot product on \mathbb{R}^n .




Figure 1.5: On the half-space model of \mathbb{H}^n , geodesics are half lines parallel to the x_n axis and circumferences with centre on the $x_n = 0$ hyperplane (see [9]).

Remark 1.17. The models above have curvature ± 1 . For other values of the curvature one has to consider the following Riemannian metrics

- for the half-space model of hyperbolic space:

$$g_x = \frac{R^2}{x_n^2} \langle \cdot, \cdot \rangle$$

- for the Poincaré ball model of hyperbolic space one has to take the disk with radius R and:

$$g_x = \frac{4R^4}{(R^2 - \|x\|^2)^2} \langle \cdot, \cdot \rangle$$

- For the stereographic projection of a sphere with radius R :

$$g_x = \frac{4R^4}{(R^2 + \|x\|^2)^2} \langle \cdot, \cdot \rangle$$

We will now define reflections in \mathbb{M}^n :

- On \mathbb{R}^n we choose an hyperplane and use the usual reflections.

- On S^n , reflections are those induced by \mathbb{R}^{n+1} when the fixed plane passes through the origin. Each one can be identified by vector orthogonal to the plane we chose. Each hyperplane through the origin in \mathbb{R}^{n+1} defines a $(n-1)$ -sphere through intersection with S^n , which can all be rotated into each other. We can parametrize each point in S_+^n the following way:

- choose a point O and a direction $v \in T_O S^n$, and consider the geodesic γ_v such that $\gamma_v(0) = O$ and $\dot{\gamma}_v(0) = v$. Assume that γ is parametrised by arc-length.
- Consider the $(n-1)$ -sphere π_0 passing through $\gamma_v(0)$ and orthogonal to $\dot{\gamma}_v(0)$. One can rotate that sphere along the geodesic γ_v so that it touches each point in S_+^n . We will call the sphere passing through $\gamma_v(t)$ $\pi_{v,t}$. Each point x is in a unique $\pi_{v,t}$.
- We can then assign to each point in S_+^n a unique couple of coordinates (x, t) , where x is the coordinate in $S^{n-1} \cap S_+^n$ when rotating it back to π_0 , and t is the unique t such that the point is in $\pi_{v,t}$.

Observe that a reflection about $\pi_{v,s}$ is $(x, t) \mapsto (x, 2s - t)$. This assignment can actually be extended to any point on S_-^n as well (the other hemisphere) minus any point on the plane orthogonal to v passing through O .

- On \mathbb{H}^n we can also construct a similarly constructed foliation:
 - As a first step, choose a point $O \in \mathbb{H}^n$. \mathbb{H}^n is a homogeneous space, so the construction does not depend on the choice we make.
 - Choose any direction in $v \in T_O \mathbb{H}^n$ and consider the geodesic $\gamma_v : \mathbb{R} \rightarrow \mathbb{H}^n$ satisfying $\gamma_v(0) = O$ and $\dot{\gamma}_v(0) = v$. Assume that γ is parametrised by arc-length.
 - Consider the hyperplane π_0 passing through $\gamma_v(0)$ and orthogonal to $\dot{\gamma}_v(0)$. Then consider the 1-parameter group of isometries of \mathbb{H}^n such that $g_t(\gamma_v(0)) = \gamma_v(t)$ and such that the curves $t \mapsto g_t(x)$ are orthogonal to π_0 for each $x \in \pi_0$. This allows us to assign to each point in \mathbb{H}^n coordinates (x, t) where $x \in \pi_0$ and $t \in \mathbb{R}$.
 - consider now any hyperplane π_t passing through $\gamma_v(t)$ and orthogonal to $\dot{\gamma}_v(t)$. The reflection fixing π_t will be the one given by the formula $(x, t) \mapsto (x, 2s - t)$.

One may find this insufficient, as there is no guarantee that this construction exists, but it is easy to convince oneself that it does: through rotations and translations it is possible to assume without loss of generality that the point we chose is e_n and the direction is e_1 (assuming the half-space model in the



Figure 1.6: The moving hyperplanes in \mathbb{H}^n when the point we chose is e_n and the direction is e_1 , projected onto the plane spanned by these two vectors. The centres of the hemispheres are the intersection of the $x_n = 0$ plane with the line spanned by the tangent vector.

definition above). We can do this because \mathbb{H}^n is frame-homogenous. Then the geodesic is the unit circle in the plane spanned by e_1 and e_n , and the (euclidean) hemispheres centred on the $x_n = 0$ plane and normal to the geodesic at one of its points are the hyperplanes (see figure 1.6). For a proof that \mathbb{H}^n is frame-homogenous, see proposition 3.9 in [9].

Remark 1.18. Another choice of direction on the hyperbolic plane allows one to write an explicit formula for the reflection about each hyperplanes, which may be more enlightening than the construction above. One can repeat the construction, taking the point e_n and direction e_n . The geodesic is then the vertical x_n axis and the totally geodesic hyperplanes are therefore the euclidean spheres centred at the origin. Assume without loss of generality that curvature is ± 1 , and take the sphere of radius r . We claim that the reflection about the sphere of radius r is the spherical inversion about the sphere. We remind the reader that a spherical inversion about a sphere of centre O is a transformation mapping a point P to the point P' on the (euclidean) line through O and P such that $\overline{OP} \cdot \overline{OP'} = r^2$. Therefore the point $(0, a)$ on the geodesic is mapped to $(0, \frac{r^2}{a})$ by the inversion.

Computing the geodesic distance of these two points to $(\underline{0}, r)$:

$$\begin{aligned}\int_r^a \frac{1}{x} dx &= [\ln x]_r^a = \ln a - \ln r \\ \int_{\frac{r^2}{a}}^r \frac{1}{x} dx &= [\ln x]_{\frac{r^2}{a}}^r = \ln r - 2 \ln r + \ln a - \ln r = \ln a - \ln r\end{aligned}$$

we see that the points are the equidistant from $(\underline{0}, r)$ and therefore the corresponding spheres/hyperplanes have to map to each other. On the other hand, it is a standard result that inversions are orientation-reversing isometries in hyperbolic space. Please note however that, if we consider $I_r(P)$ fix a point $P \in H^n$ and let r vary, it does not move along a geodesic unless it is on the geodesic generating the motion. Therefore, as the geodesic is the shortest path, taking r to be varying between R_0 and R_1

$$\text{dist}(I_{R_0}(P), I_{R_1}(P)) \leq \text{length}(I_r(P))$$

in general, with a strict inequality unless P is on the geodesic. This is unlike the situation on the plane, where equality holds everywhere.

Remark 1.19. The inequality

$$\text{dist}(I_{R_0}(P), I_{R_1}(P)) \leq \text{length}(I_r(P)) \quad (1.5)$$

taking corresponding objects, holds also on S^n .

1.6 The Method of Moving Planes

To provide some justification and context to the next chapters, we describe Alexandrov reflection, also known as Method of the Moving Planes in \mathbb{M}_+^{n+1} .

Let $X : M^n \rightarrow \mathbb{M}_+^{n+1}$ be a hypersurface in a constant curvature ambient space. If the ambient space is a sphere, X must be contained in a hemisphere to avoid issues with multiple self-intersections. Assume also that $X = \partial\Omega$ for a bounded domain Ω in \mathbb{M}_+^{n+1} .

- As a first step, choose a point $O \in \mathbb{M}^n$. As \mathbb{M}^n is a homogeneous space, so the construction does not depend on the choice we make. Without loss of generality, we choose the origin in \mathbb{R}^n , e_n in \mathbb{H}^n and the north pole in S^n .
- Choose any direction in $v \in T_O \mathbb{M}^n$ and consider the geodesic $\gamma_v : I \rightarrow \mathbb{M}^n$ satisfying $\gamma_v(0) = O$ and $\dot{\gamma}_v(0) = v$. Assume that γ is parametrised by arc-length. Here $I = \mathbb{R}$ if \mathbb{M}_+^n is flat or hyperbolic, and $I = (-\frac{\pi}{2}, \frac{\pi}{2})$ if \mathbb{M}_+^n is a hemisphere.

- Consider the hyperplanes $\pi_{v,s}$ passing through $\gamma_v(s)$ and orthogonal to $\dot{\gamma}_v(s)$.

The method consists of reflecting the part of X “below” the hyperplane into the top part, and using properties of both halves of the hypersurface together to prove some statement about the non-reflected hypersurface.

To make this more precise, we can define:

$$X_{v,s} = \{p \in X \mid p \in \pi_{v,t} \text{ for some } t < s\}$$

Let $X_{v,s}^\pi$ be the reflection of $X_{v,s}$ about $\pi_{v,s}$. We can then define:

$$\begin{aligned} m_v &= \sup \{s \in I \mid X_{v,s}^\pi \subset \Omega \text{ for every } t < s\} \\ &= \sup \{s \in I \mid X \cap X_{v,s}^\pi = \emptyset \text{ for every } t < s\} \end{aligned}$$

the last time at which the hypersurface and its reflection do not touch internally. Please note that at m_v the two surfaces are tangent at some point, which can be either in the interior of X , or on the boundary. The hyperplane π_{v,m_v} is the critical hyperplane.

1.7 The Alexandrov soap-bubble theorem

To give some justification to the method we described, we will outline the proof of an important result that uses this method, the so-called Alexandrov soap bubble theorem. For more details, see [3].

Theorem 1.20. *The only C^2 -regular connected hypersurfaces embedded in \mathbb{M}_+^{n+1} and such that the mean curvature is constant are the distance spheres.*

In [3] this theorem is proved more generally: let H_X be a C^2 function of the ordered principal curvatures $H_X = f(\kappa_1, \dots, \kappa_n)$, and

$$f : \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 \leq x_2 \leq \dots \leq x_n\} \rightarrow \mathbb{R}$$

is such that

$$f(x) > 0 \text{ if } x_i > 0 \text{ for every } i = 1, \dots, n$$

and it is concave on the component of $\{x \in \mathbb{R}^n \mid f(x) > 0\}$ containing $\{x \in \mathbb{R}^n \mid x_i > 0\}$. Then the following more general theorem holds:

Theorem 1.21. *The only C^2 -regular connected hypersurfaces embedded in \mathbb{M}_+^{n+1} and such that H_X is constant are the distance spheres.*




Figure 1.7: The method of the moving planes in \mathbb{R}^2 . In the figure, the “plane” is centred for easier pagination, but the hypersurface is fixed and the plane varies. The third figure shows the critical time m_v when the hypersurface and its reflection touch internally.

The following proposition holds:

Proposition 1.22. *Let $X = \partial\Omega$ be a C^2 -regular, connected, closed hypersurface embedded in \mathbb{M}_+^{n+1} , where Ω is a bounded domain. Assume that for every geodesic $\gamma : \mathbb{R} \rightarrow \mathbb{M}^{n+1}$ there exists a hyperplane orthogonal to γ such that X is symmetric about π . Then X is a distance sphere about its center of mass O , i.e. the unique minimum of*

$$P_\Omega(x) = \int_\Omega d(x, a)^2 da$$

The proof of this proposition is in [3] and is omitted here. We now move on to the proof of Theorem 1.20:

Proof. Assume that X is a manifold with constant mean curvature. We want to show that, for any point O and for every direction $v \in T_O\mathbb{M}^{n+1}$, X is symmetric about a plane perpendicular to the geodesic $\exp_O(tv)$. We put ourselves in the hypothesis of the method of the moving planes described in section 1.6. At m_v , $X \cap X_{v, m_v}^\pi$ is non-empty by definition of m_v and therefore it is closed in X_{v, m_v}^π . We want to show that $X \cap X_{v, m_v}^\pi$ is an open set in X_{v, m_v}^π .

Let $p \in X \cap X_{v, m_v}^\pi$. As the two manifolds are tangent

$$T_p X = T_p X_{v, m_v}^\pi.$$

By Theorem 1.5, we can represent both manifold as the Euclidean graph of two functions C^2 functions u and \tilde{u} defined in a neighbourhood of p inside the tangent space. Consider now the differential equation:

$$K(u(x)) = K(\tilde{u}(x)) = \text{constant}$$

where K is the mean-curvature operator. As X is a manifold with constant mean curvature, it holds everywhere. Looking at equation (1.1), we see that, given that the principal curvature are a function of the h_{ij} , the operator K is an elliptic operator. Reasoning exactly like we did in section 1.4 for parabolic differential equations, we see that $u - \tilde{u}$ is the solution of a linear elliptic differential equation of the form $L(u - \tilde{u}) = 0$, with $u(p) = \tilde{u}(p) = 0$.

Without loss of generality, we can assume that, in the neighbourhood where we defined u and \tilde{u} , $u - \tilde{u} \geq 0$.

If p is an interior point in X_{v, m_v}^π , we can then apply the maximum principle for elliptic equations (see section 6.4 in [11]) and obtain that $u = \tilde{u}$ in the neighbourhood.

Otherwise, if $p \in \pi_{v, s}$, $\nabla u(p) = \nabla \tilde{u}(p) = 0$ and one can apply Hopf's boundary point lemma for elliptic equations (see section 6.4 in [11]) to conclude again that $u = \tilde{u}$ in the neighbourhood.

Therefore, the whole neighbourhood is in $X \cap X_{v,m_v}^\pi$, hence the intersection is open, as every point p is contained in an open ball. This proves that X is symmetric about π_{v,m_v} . The theorem is then consequence of Proposition 1.22 \square

Remark 1.23. To prove the theorem in the aforementioned more general setting, one would need to only prove that the differential equation $H_X(u(x)) = \text{constant}$ is an elliptic equation. Its ellipticity is a standard result in Geometric Analysis, and can be found in the literature. It is an immediate consequence of the fact that equation (2.2) is parabolic (see section 2.2).

Chapter 2

The Chow-Gulliver Critical Planes Result

The main result we want to establish in this chapter is theorem 2.6, a result about critical hyperplanes when applying the method of the moving planes to solution of a large class of non-linear parabolic partial differential equations, and whose proof is somewhat similar to Theorem 1.20.

We will first describe the differential equations we are analysing, then prove that they are parabolic, then prove the theorem.

After proving the theorem we will include some corollaries of the result and use it to find some estimates for the gradient of the support function and the gradient radial function.

Finally, an application of the result is presented, showing that the condition of *coming out of a point* is particularly rigid on expansive flows.

2.1 Class of Equations we analyze

We consider manifolds M^n embedded in \mathbb{R}^{n+1} , i.e. there is an embedding $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$ parametrizing the hypersurface $X_0(M^n)$.

Let $F : \{(\kappa_1, \dots, \kappa_n) \in \mathbb{R}^n | \kappa_1 \leq \dots \leq \kappa_n\} \rightarrow \mathbb{R}$ be a C^1 function satisfying:

$$\frac{\partial F}{\partial \kappa_i} > 0 \text{ for all } i = 1, \dots, n \quad (2.1)$$

and consider the evolution equation

$$\begin{cases} \frac{\partial X_t}{\partial t} = -F(\kappa_1(x), \dots, \kappa_n(x))\nu \\ X(0) = X_0 \end{cases} \quad (2.2)$$

where ν is the outward normal to $X_t(M^n)$ at the point $X_t(x)$ and $\kappa_1 \leq \dots \leq \kappa_n$ are the principal curvatures at $X_t(x)$.

2.2 Parabolicity of the differential equation (2.2)

The condition (2.1) will guarantee that equation (2.2) is a parabolic equation. This may be confusing, as (2.2) does not make it obvious how to apply definition 1.10.

In order to classify a non-linear partial differential equation one has to understand how it behaves “close to a solution” in the solutions space. We want to prove that very close to any solution, “moving in any direction”, the change in the equation is always a parabolic PDE. This will then tell us that our equation is parabolic, and that the theorems that apply to solutions of parabolic partial differential equations apply to our equation as well. To do so, we are going to “linearise” the differential equation about a solution.

Like in [7], as F is a symmetric function in the principal curvatures, we may interchangeably take F to be a function of the Weingarten map tensor or of the second fundamental form, and thus we get:

$$\frac{\partial X_t}{\partial t} = -F(h_{ij}(X_t))\nu$$

To understand the behaviour close to a solution, we can substitute in our equation X_t with a $X_t + \varepsilon u_t$ to get:

$$\frac{\partial X_t}{\partial t} + \varepsilon \frac{\partial u_t}{\partial t} = -F(h_{ij}(X_t + \varepsilon u_t))\nu_{(X_t + \varepsilon u_t)} \quad (2.3)$$

where we mean that $\nu_{(X_t + \varepsilon u_t)}$ is the normal to the perturbed immersion. We are interested in the behaviour of this equation for a small ε . This equation when taking the limit for $\varepsilon \rightarrow 0$ is the so-called linearisation of the PDE; we want this PDE to be a parabolic equation to apply our results.

We can use the Weingarten equation (1.1) to write the RHS explicitly:

$$\begin{aligned}
h_{ij}(X_t + \varepsilon u_t) &= -\langle v, \nu_{(X_t + \varepsilon u_t)} \rangle \text{ where} \\
v^\alpha &= \frac{\partial^2 X_t^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial X_t^\alpha}{\partial x^k} + \bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial X_t^\beta}{\partial x^i} \frac{\partial X_t^\delta}{\partial x^k} + \\
&\quad + \varepsilon \left(\frac{\partial^2 u_t^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial u_t^\alpha}{\partial x^k} + \bar{\Gamma}_{\beta\delta}^\alpha \left(\frac{\partial X_t^\beta}{\partial x^i} \frac{\partial u_t^\delta}{\partial x^k} + \frac{\partial u_t^\beta}{\partial x^i} \frac{\partial X_t^\delta}{\partial x^k} \right) \right) + \\
&\quad + \varepsilon^2 \left(\bar{\Gamma}_{\beta\delta}^\alpha \frac{\partial u_t^\beta}{\partial x^i} \frac{\partial u_t^\delta}{\partial x^k} \right) \\
v^\alpha &= w + \varepsilon \left(\frac{\partial^2 u_t^\alpha}{\partial x^i \partial x^j} + \text{lower order terms} \right) + o(\varepsilon)
\end{aligned}$$

where $h_{ij}(X_t) = -\langle w, \nu_{X_t} \rangle$.

Putting it all together in the first line:

$$\begin{aligned}
h_{ij}(X_t + \varepsilon u_t) &= -\left\langle w + \varepsilon \left(\frac{\partial^2 u_t}{\partial x^i \partial x^j} + \text{lower order terms} \right) + o(\varepsilon), \nu_{(X_t + \varepsilon u_t)} \right\rangle \\
&= \langle w, \nu_{(X_t + \varepsilon u_t)} \rangle - \varepsilon \left\langle \left(\frac{\partial^2 u_t}{\partial x^i \partial x^j} + \text{lower order terms} \right), \nu_{(X_t + \varepsilon u_t)} \right\rangle + o(\varepsilon) \\
&= h_{ij}(X_t) + \langle w, \nu_{(X_t + \varepsilon u_t)} - \nu_{X_t} \rangle - \varepsilon H_{ij} + o(\varepsilon) \\
&= h_{ij}(X_t) - \varepsilon H_{ij} + o(\varepsilon)
\end{aligned}$$

Were on the last step we are using the fact that $\nu_{(X_t + \varepsilon u_t)} - \nu_{X_t} = O(\varepsilon)$, and as this gets smaller the component of w parallel to the difference also is $O(\varepsilon)$, as w is parallel to ν_{X_t} . We can then expand F in the RHS of the equation (2.3) to the first order, as it is a C^1 function:

$$\begin{aligned}
\frac{\partial X_t}{\partial t} + \varepsilon \frac{\partial u_t}{\partial t} &= - \left(F(h_{ij}(X_t)) + \varepsilon \langle DF, H \rangle + o(\varepsilon) \right) \nu_{(X_t + \varepsilon u_t)} \\
\varepsilon \frac{\partial u_t}{\partial t} &= \varepsilon \left(\frac{\partial F}{\partial h_{ij}} g^{ik} g^{jl} \left\langle \frac{\partial^2 u_t}{\partial x^k \partial x^l}, \nu_{(X_t + \varepsilon u_t)} \right\rangle + \text{lower order terms} \right) \nu_{(X_t + \varepsilon u_t)} + o(\varepsilon) \\
\frac{\partial u_t}{\partial t} &= \frac{\partial F}{\partial h_{ij}} g^{ik} g^{jl} \left\langle \frac{\partial^2 u_t}{\partial x^k \partial x^l}, \nu_{(X_t + \varepsilon u_t)} \right\rangle \nu_{(X_t + \varepsilon u_t)} + \text{lower order terms} + o(1)
\end{aligned}$$

Letting $\varepsilon \rightarrow 0$, we get to:

$$\frac{\partial u_t}{\partial t} = \frac{\partial F}{\partial h_{ij}} g^{ik} g^{jl} \left\langle \frac{\partial^2 u_t}{\partial x_k \partial x_l}, \nu \right\rangle \nu + \text{lower order terms}$$

For the second order term to be positive definite, then,

$$\left(\frac{\partial F}{\partial h_{ij}} \right)_{i,j}$$

must be positive definite. Or equivalently, as the principal curvatures are the eigenvalues of the matrix $(h_{ij})_{i,j}$,

$$\frac{\partial F}{\partial \kappa_i} > 0$$

A more general approach proving that the differential equation is parabolic can be found in [12].

Remark 2.1. While in this chapter we are using the standard metric of \mathbb{R}^n , the proof above is valid using any other metric.

2.3 An existence result

In [7] one finds the following comforting existence result for the class of equations we are analysing under very broad hypothesis. The same paper also includes a proof of the result with some more restrictive hypotheses. A complete proof is quite more involved. We will not use it, nor prove it, but it is included here for completeness and peace of mind:

Theorem 2.2 (Short term existence of a solution for (2.2)). *Suppose $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$ is a smooth, closed hypersurface in \mathbb{R}^{n+1} , such that (2.1) holds at all points in X_0 , i.e. for all the values of the principal curvatures κ_i realized at some point on X_0 . Then, (2.2) has a smooth solution, at least on some short time interval $[0, T)$, $T > 0$.*

A much more extended analysis of the problem of the existence of solution to the equation can be found in [12].

2.4 Local representation as a graph of a solution of (2.2)

As a first step, from the Corollary 1.5, we can establish the following:

Theorem 2.3 (Local representation as a graph of a solution of (2.2)). *Let X^n be a submanifold $X^n \subset \mathbb{R}^{n+1}$ and let $F : X^n \times (0, T) \rightarrow \mathbb{R}^{n+1}$ be a solution of (2.2). Also let $t \in (0, T)$ and $x \in F(X^n, t)$. Then there exists a neighbourhood of (x, t) ,*

2.4. LOCAL REPRESENTATION AS A GRAPH OF A SOLUTION OF (2.2)33

$U \subset F(X^n \times (0, T))$, and a smooth function $f : T_x F(X^n, t) \times (0, T) \rightarrow \mathbb{R}$ such that any $(x_0, t_0) \in U$ can be expressed as

$$x_0 = p + f(p, t_0)\nu$$

where ν is a vector normal to $T_x F(X^n, t)$, for an appropriate point $p \in T_x F(X^n, t)$.

Proof. We can consider the image of $F : X^n \times (0, T) \rightarrow \mathbb{R}^{n+1}$ as a manifold in \mathbb{R}^{n+2} by considering $G := (F(x, t), t)$. Moreover $\frac{\partial G_t}{\partial e_j} \equiv 0$ for all possible vectors of the canonical basis of $\mathbb{R}^{n+1} \times \mathbb{R}$ except for the one corresponding to the time coordinate, where it is 1. Also, $\frac{\partial G_i}{\partial e_j} \equiv \frac{\partial F_i}{\partial e_j}$ and the first n coordinates of $\frac{\partial G}{\partial t}$ form a vector normal to $T_x X^n$ by (2.2). Thus, the tangent space of $\text{Im}(G)$ is $T_x X^n \times \{0\} \oplus \text{span}\langle(\nu, 1)\rangle$ and we can apply corollary 1.5 to $\text{Im}(G)$ to get a function \tilde{f} such that

$$(x_0, t) = [(p, 0) + (s\nu, s)] + \tilde{f}(p, s)(\nu, -\|\nu\|)$$

as $(\nu, -\|\nu\|)$ is the vector orthogonal to $T_x X^n \times \{0\} \oplus \text{span}\langle(\nu, 1)\rangle$. Let $\sigma_p : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be the function that associates to t the appropriate s in the expression above. Projecting to the first $n + 1$ coordinates and calling $f(p, t) = \sigma_p(t) + \tilde{f}(p, \sigma_p(t))$ one gets:

$$x_0 = p + f(p, t)\nu$$

which is our thesis, as long as $\sigma_p(t)$ is smooth. This is indeed the case, as the graph function $\Gamma_f : x \mapsto (x, f(x))$ is smooth for any smooth function f and has a smooth inverse (and thus, the inverse $(x_0, t) \mapsto ((p, 0) + (s\nu, s))$ is smooth). \square

One can show through direct calculation (see [14], Exercise 1.1.2) that, if an immersed hypersurface $\phi : M \rightarrow \mathbb{R}^{n+1}$ is locally the graph of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ (i.e., locally, $(x, f(x)) = \phi$), then:

$$\begin{aligned} g_{ij} &= \delta_{ij} + \frac{\partial f}{\partial x_i} \frac{\partial f}{\partial x_j} \\ \nu &= -\frac{(\nabla f, -1)}{\sqrt{1 + |\nabla f|^2}} \\ h_{ij} &= -\frac{H_{ij}}{\sqrt{1 + |\nabla f|^2}} \end{aligned}$$

Where the matrix $(H_{ij})_{ij}$ is the hessian of f . Thus, if one considers the principal curvatures, they are closely related to the eigenvalues of the hessian of f .

2.5 The Moving Planes Method and the Chow result

We will now present the Chow-Gulliver result, presented more or less as in paragraph 2 in [1]. Suppose that we have a hypersurface embedded in \mathbb{R}^{n+1} evolving according to equation (2.2). For a fixed time t , we can apply the method of the moving planes as described in section 1.6: we can take parallel hyperplanes $\pi_{v,s}$ orthogonal to v intersecting X , and consider the reflection $X_{v,s}^\pi$. There will be a hyperplane π_{v,m_v} where X and $X_{v,s}^\pi$ are tangent. As the hypersurface evolves, we may wonder how this critical threshold changes over time. If it behaves in a predictable way, we can hope to use the technique on the evolving manifold, otherwise, it may be hopeless if the critical plane moves back and forth multiple times. In the next section, we are going to prove a result in this general direction, to show a form of "regularity" in this sense. First, however, we need to introduce a marginally stricter definition for the concept of "reflecting inside itself".

Let π be a hyperplane in \mathbb{R}^{n+1} . We may assume π orthogonal to a unit vector $v \in \mathbb{R}^{n+1}$, i.e. $\langle x, v \rangle = C$ for all $x \in \pi$ for some constant C . In our notation for the method of moving planes, assuming we pick the origin as a starting point, this means that $\pi = \pi_{v,C}$.

Then, \mathbb{R}^{n+1} is divided by π into two half-spaces, which we will name

$$H^+(\pi) = \{x \in \mathbb{R}^{n+1} : \langle x, v \rangle > C\} = \bigcup_{s>C} \pi_{v,s} \text{ and}$$

$$H^-(\pi) = \{x \in \mathbb{R}^{n+1} : \langle x, v \rangle < C\} = \bigcup_{s<C} \pi_{v,s}.$$

Definition 2.4. We say we can reflect $X : M^n \rightarrow \mathbb{R}^{n+1}$ strictly with respect to π if both:

- $X^\pi \cap H^-(\pi) \subset \text{int}(X) \cap H^-(\pi)$ where X^π is the reflection of X about π and $\text{int}(X)$ is the region inside X .
- $V \notin T_x M$ for all $x \in M^n \cap \pi$

This fundamentally means that the reflection of one of the halves of X on the other side of π is contained in the region inside M^n and the tangent spaces of X and of the half-reflection do not form a ninety degree angle with π , at all points on $\pi \cap X$. As the two tangent spaces are one the reflection of the other, this means that they do not coincide.

Definition 2.5. We say we can reflect $X : M^n \rightarrow \mathbb{R}^{n+1}$ strictly up to (π, v) if we can reflect M^n strictly with respect to $\pi_{v,s}$ for all hyperplanes $\pi_{v,s}$ such that $s < C$.




Figure 2.1: Example: We can reflect $X : M^n \rightarrow \mathbb{R}^{n+1}$ strictly with respect to π

The key idea of the main result in the next section is as follows: suppose we have an embedded smooth hypersurface X evolving according to (2.2) and a fixed hyperplane π , intersecting X . Suppose that, at some time t , X and X_π touch outside of π . We can consider X and X_π as local graphs over the same hyperplane π , and we can show that these function evolve according to the same differential equation. Using the strong maximum principle and the Hopf boundary point lemma, then, one can conclude that the two functions coincide, and have been coinciding up until that point. We can then conclude that if X and X_π only touch in $X \cap \pi$ at the beginning of the evolution, then they will never touch elsewhere.

2.6 The Chow-Gulliver result

The main theorem is the following:

Theorem 2.6 (Chow-Gulliver). *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 solution to equation (2.2). Then, if we can reflect $X(M^n, 0) = X_0$ strictly with respect to π , then for all $t \in [0, T)$ we can reflect $X(M^n, t) = X_t$ strictly with respect to π .*

Proof. By contradiction, suppose that there is a time t such that the thesis is false, and that it is the smallest such t . Then, for all $\tau \in [0, t)$, $X_{\tau, \pi} \cap H^-(\pi) \subset \text{int}(X_\tau) \cap H^-(\pi)$; the unit vector orthogonal to π , V , is such that $V \notin T_x X_\tau$ for all $x \in X_\tau \cap \pi$ and $\tau \in [0, t)$; and either of the conditions fails at t , i.e. either:




Figure 2.2: Example: We cannot reflect $X : M^n \rightarrow \mathbb{R}^{n+1}$ strictly with respect to π , because there is an interior contact




Figure 2.3: Example: We cannot reflect $X : M^n \rightarrow \mathbb{R}^{n+1}$ strictly with respect to π , because the tangent spaces coincide at a point about which we are reflecting

- (i) $X_{t,\pi} \cap H^-(\pi) \cap X_t \neq \emptyset$
- (ii) $V \in T_x X_t$ for some $x \in \pi$.

(i) Suppose the first case is true. Then, there exists $x_0 \in X_{t,\pi} \cap H^-(\pi) \cap X_t$ such that at x_0 the two manifolds are tangent.

We can take a neighbourhood of $(x_0, t) \in X_t \times \mathbb{R}$ such that both $X_{t,\pi}$ and X_t are graphs over $T_{x_0} X_t$ by 2.3.

We can explicitly write the functions $f : U \times (t - \varepsilon, t + \varepsilon) \rightarrow X_t$, where $U \subset T_{x_0} X_t$, and the corresponding f_π for $X_{t,\pi}$. We can also write

$$\begin{aligned} f &: (x, t) \mapsto x + \tilde{f}(x, t)\nu \\ f_\pi &: (x, t) \mapsto x + \tilde{f}_\pi(x, t)\nu \end{aligned}$$

for appropriate functions $\tilde{f} : U \times (t - \varepsilon, t + \varepsilon) \rightarrow \mathbb{R}$ and $\tilde{f}_\pi : U \times (t - \varepsilon, t + \varepsilon) \rightarrow \mathbb{R}$, where ν is a fixed unit vector normal to $T_{x_0} X_t$. \tilde{f} and \tilde{f}_π are solutions to the same second order PDE, which is parabolic by what was discussed in paragraph 2.2, hence we can apply Proposition 1.12 to conclude that $\tilde{f} \equiv \tilde{f}_\pi$, and thus $X_{t,\pi}$ and X_t coincide in a neighbourhood of (x, t) , a contradiction as we assumed that t is the first t where the flows touch.

(ii) Suppose instead that $V \in T_x X_t$ for some $t \in [0, t)$ and some $x \in X_t \cap \pi$. Then $T_x X_t = T_x X_{t,\pi}$ and in a neighbourhood of (x, t) both X_t and $X_{t,\pi}$ are graphs of two smooth functions over $T_x X_t$ by 2.3, i.e. again

$$\begin{aligned} f &: (x, t) \mapsto x + \tilde{f}(x, t)\nu \\ f_\pi &: (x, t) \mapsto x + \tilde{f}_\pi(x, t)\nu \end{aligned}$$

Moreover, in $\overline{H^-(\pi)}$, $f_\pi \geq f$, because $M_\pi^n \cap H^-(\pi) \subset \text{int}(M^n) \cap H^-(\pi)$. Finally, $f(x, t) = f_\pi(x, t)$, hence $f_\pi - f(x, t) = 0$, and thus (x, t) is a minimum point on the boundary for $f_\pi - f$. Also, we must have

$$\frac{\partial f}{\partial V}(x, t) = \frac{\partial f_\pi}{\partial V}(x, t)$$

because the graphs are both tangent to $T_x X_t$, and V here is the outward pointing normal to the boundary by definition of the reflection. Thus,

$$\frac{\partial(f - f_\pi)}{\partial V}(x, t) = 0$$

But we must have

$$\frac{\partial(f - f_\pi)}{\partial V}(x, t) > 0$$

at a minimum on the boundary by Proposition 1.13, a contradiction. \square

2.7 Some corollaries of the result

In this section we collect a number of corollaries to the main result above. This first one is an immediate consequence of the main result:

Corollary 2.7. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 embedded solution to equation (2.2). Then, if we can reflect X_0 strictly up to $(\pi_{v,C}, v)$, for all $t \in [0, T)$ $v \notin T_x X_t$ for all $x \in X_t \cap \overline{H^+(\pi)}$. In particular, $X_t \cap \overline{H^+(\pi)}$ is a graph over π for all $t \in [0, T)$.*

Another immediate consequence of theorem 2.6 is the following:

Corollary 2.8. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 solution to equation (2.2). Then, if we can reflect X_0 strictly up to $(\pi_{v,C}, v)$, for all $t \in [0, T)$ we can reflect $X(M, t)$ strictly up to $(\pi_{v,C}, v)$.*

Proof. The hypothesis of the theorem are true for each π_K in the definition, thus we can reflect strictly with respect to each π_K for all $t \in [0, T)$, and thus we can reflect $X(M, t)$ strictly up to $(\pi_{v,C}, v)$. \square

Furthermore, it is clear that for every direction v there exists a hyperplane Π perpendicular to v such that we can reflect X_0 up to (Π, v) and Π intersects the interior of X_0 . To be more precise, for every direction v there exists a hyperplane Π_0^v tangent to X_0 and such that $X_0 \cap H^+(\Pi_0^v) = \emptyset$. Suppose that for every $\varepsilon > 0$ we can find a plane π_ε such $H^+(\pi_\varepsilon) \cap B_{R-\varepsilon}(C) = \emptyset$ and we cannot reflect X_0 strictly at π_ε , where R is such that $X_0 \subset B_R(C)$. We can take the corresponding plane Π_0^ε parallel to π_ε and tangent to X_0 at the point p_ε . Taking a sequence $\varepsilon_n \rightarrow 0$ we find a corresponding limited sequence p_{ε_n} which, by compactness, converges to a point $p \in X_0$. This point p is such that arbitrarily close to it there is a point such that we cannot reflect by more than any chosen $\varepsilon > 0$ in the direction of its normal. As we can always represent X_0 as the graph of a function in a neighbourhood of p , we get a contradiction, because strict reflection in the direction of the normal by at least a fixed uniform amount ε at each point is always possible locally for any graph of a smooth function. Thus, we obtain the following:

Corollary 2.9. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 embedded solution to equation (2.2). There exists $\varepsilon > 0$ depending only on X_0 such that for all $t \in [0, T)$ we can reflect X_t up to $(\Pi_0^v + \varepsilon v, v)$ for every $v \in S^n$. In particular, if $X_0 \subset B_R(C)$, then we can always reflect X_t up to (Π, v) whenever $H^+(\Pi) \cap B_{R-\varepsilon}(C) = \emptyset$.*

In other words, we can always reflect a little ε in any direction, uniformly. We can use the fact that we can always reflect about a plane outside a sphere containing X_0 to prove the following estimate:

Corollary 2.10. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 embedded solution to equation (2.2). There exists $C > 0$ depending only on X_0 such that for all $t \in [0, T)$:*

$$\max_{x \in X_t} |x| - \min_{x \in X_t} |x| < C$$

Proof. We can reflect $X_0 \subset B_R(0)$ up to any plane tangent to $B_R(0)$, i.e. any plane $\pi_{v,K}$ for any $K \geq R$ and v unit vector, i.e. $\pi_{v,K} = \{p \in \mathbb{R}^{n+1} : \langle p, v \rangle = K\}$. Let $x_1, x_2 \in X_t$ such that $|x_1| = \min_{x \in X_t} |x|$ and $|x_2| = \max_{x \in X_t} |x|$. Let $v = \frac{x_2 - x_1}{|x_2 - x_1|}$: we can reflect X_t up to $(\pi_{v,R}, v)$ by theorem 2.6, therefore $\text{dist}(x_2, \pi_{v,R}) <$

$\text{dist}(x_1, \pi_{v,R})$, or in other words:

$$\begin{aligned}
\left\langle x_2, \frac{x_2 - x_1}{|x_2 - x_1|} \right\rangle - R &< \left\langle x_1, \frac{x_1 - x_2}{|x_2 - x_1|} \right\rangle + R \\
\frac{|x_2|^2}{|x_2 - x_1|} - \frac{\langle x_2, x_1 \rangle}{|x_2 - x_1|} - R &< \frac{|x_1|^2}{|x_2 - x_1|} - \frac{\langle x_2, x_1 \rangle}{|x_2 - x_1|} + R \\
\frac{|x_2|^2 - |x_1|^2}{|x_2 - x_1|} &< 2R \\
|x_2|^2 - |x_1|^2 &< 2R|x_2 - x_1| \leq 4R|x_2| \\
|x_2|^2 &< |x_1|^2 + 4R|x_2| \\
|x_2| &< |x_1| \frac{|x_1|}{|x_2|} + 4R < |x_1| + 4R \\
|x_2| - |x_1| &< 4R
\end{aligned}$$

□

Lastly:

Corollary 2.11. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 embedded solution to equation (2.2). Let $s_v : [0, T) \rightarrow I$ be such that*

$$s_v(t) = \sup \{s \in I \mid \text{we can reflect } X_t \text{ strictly up to } (\pi_{v,s}, v)\}.$$

Then $s_v(t)$ is a non-decreasing function. Also, if X_0 is compact, the limit

$$\lim_{t \rightarrow T^-} s_v(t)$$

exists and is finite.

Proof. When taking X_t as the starting manifold, the hypothesis of the theorem are still true, therefore we can reflect about $\pi_{c,v}$ for all $c < s_v(t)$ at all subsequent times. Thus $s_v(t)$ is non-decreasing. $\lim_{t \rightarrow T^-} s_v(t) = \sup s_v(t)$, therefore the limit exists. Also, if X_0 is bounded, there exists $R > 0$ such that $X_0 \subset B_R(0)$, therefore we can reflect X_0 strictly about any hyperplane non intersecting $B_R(0)$, as it does not touch X_0 , and therefore we can also reflect X_t strictly about the same hyperplanes by theorem 2.6. At the same time, there exists a hyperplane such that X_t cannot be reflected strictly about it, because there will always be a straight line parallel to v intersecting X_t in multiple points, and we can always consider the hyperplane orthogonal to v passing through their midpoint, about which X_t cannot be reflected strictly, and $s_v(t) \neq +\infty$. Therefore $s_v(t) \in [-R, R]$, and the limit above is finite. □

2.8 Applying the result to find gradient estimates

In this section we collect some applications of theorem 2.6 to gradient estimates for the support function and the radial function from [1] (a more in-depth definition these two functions can be found in the introduction of [15]). Central in what will follow is this corollary providing an estimate for the tangent component of the position vector x of a point on the hypersurface in its own tangent space:

Corollary 2.12. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be an embedded solution to equation (2.2). There exists a constant C , depending only on the initial hypersurface X_0 , such that for all points $x \in X_t$ and $t \in [0, T)$, the following inequality holds:*

$$|x - \langle x \cdot \nu \rangle \nu| \leq C,$$

where ν is the unit normal to X_t at the point x .

Proof. : Choose $C > 0$ such that $X_0 \subset B_C(0)$. By Theorem 2.6, we can reflect X_t strictly up to any plane tangent to the ball, like in the previous corollaries. Thus, for any point $x \in X_t$ and outside the ball, we know that whenever $(x, V) > C$, then $V \notin T_x X_t$. This is equivalent to saying that for all $W \in T_x X_t$, we have:

$$(x, W) \leq C.$$

If we take now the projection of x on $T_x X_t$, $W = \frac{x - (x \cdot \nu)\nu}{|x - (x \cdot \nu)\nu|} \in T_x X_t$, then we obtain:

$$C \geq (x, W) = \frac{|x - (x \cdot \nu)\nu|}{|W|} = |x - (x \cdot \nu)\nu|.$$

Thus, the corollary is proved. \square

Let's now assume that the hypersurfaces in the solution to the equation are convex. We are going to show a gradient estimate for the support function of the hypersurface. Support functions are one of the most important concepts stemming from the study of convex sets.

Definition 2.13. *Let K be a non-empty compact convex set in \mathbb{R}^n . We define the support function $u_K : S^n \rightarrow \mathbb{R}$ as*

$$u_K(\nu) = \sup\{\langle x, \nu \rangle : x \in K\}$$

Remark 2.14. Many authors define the support function on the whole euclidean space. As its gradient scales linearly with ∇u_K , we will not be doing so out of simplicity. Notice that given two convex sets K_1 and K_2 , $K_1 \subseteq K_2$ if and only if $u_{K_1} \leq u_{K_2}$. In this sense, a convex set is determined by its support function. Also, the tangent plane at the point in ∂K which has ν as a normal vector is $h_\nu = \{x \in \mathbb{R}^n : x \cdot \nu = u_K(\nu)\}$.




Figure 2.4: A geometric interpretation of corollary 2.12: the normal line always intersects $B_C(0)$. $x^\top = x - (x \cdot \nu)\nu$.

Corollary 2.15. *Let $u : S^n \times [0, T) \rightarrow \mathbb{R}$ be the support function of convex hypersurfaces X_t , solving the equation (2.2). There exists a constant C , depending only on $u(0)$, such that:*

$$|\nabla u(\nu, t)| \leq C,$$

for all $(\nu, t) \in S^n \times [0, T)$.

Proof. For each unit normal vector $\nu \in S^n$, let $x_t \in X_t$ be the unique point such that ν is the outward unit normal to X_t at x_t . We compute $\nabla u(\nu)$: let R_θ be a rotation of an angle θ in the direction $\partial_i \in T_\nu S^n$:

$$\begin{aligned} \partial_i u(\nu) &= \lim_{\theta \rightarrow 0} [\langle x_t(R_\theta \nu), R_\theta \nu \rangle - \langle x_t(\nu), \nu \rangle] / \theta \\ &= \lim_{\theta \rightarrow 0} [\langle x_t(R_\theta \nu), \nu - \nu + R_\theta \nu \rangle - \langle x_t(\nu), \nu \rangle] / \theta \\ &= \lim_{\theta \rightarrow 0} [\langle x_t(R_\theta \nu), R_\theta \nu - \nu \rangle + \langle x_t(R_\theta \nu) - x_t(\nu), \nu \rangle] / \theta \\ &= \lim_{\theta \rightarrow 0} \left[\left\langle x_t(R_\theta \nu), \frac{R_\theta \nu - \nu}{\theta} \right\rangle + \left\langle \frac{x_t(R_\theta \nu) - x_t(\nu)}{\theta}, \nu \right\rangle \right] \\ &= \langle x_t(\nu), \partial_i \rangle + \langle (dx_t)_\nu(\partial_i), \nu \rangle \end{aligned}$$

where $\langle (dx_t)_\nu(\partial_i), \nu \rangle = 0$ because $(dx_t)_\nu(\partial_i) \in T_{x_t} X_t$. Therefore

$$\begin{aligned} \nabla u(\nu) &= (x_t)^\top = x_t - (x_t)^\perp \\ \nabla u(\nu) &= x_t - \langle x_t, \nu \rangle \nu = x_t - u(\nu) \nu \\ |\nabla u(\nu, t)| &= |x_t - \langle x_t, \nu \rangle \nu| \leq C \end{aligned}$$

by applying Corollary 2.12, which completes the proof. \square

Now, let's consider the case where the hypersurfaces X_t are starshaped for all $t \in [0, T)$. We obtain a gradient estimate for the radial function at points outside a certain compact starshaped region associated with the initial hypersurface X_0 .

Definition 2.16. *Suppose that $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ parametrizes starshaped hypersurfaces X_t with respect to the origin. The radial function $r : S^n \times [0, T) \rightarrow \mathbb{R}^+$ is defined so that for each $(z, t) \in S^n \times [0, T)$, the point $r(z, t)z$ belongs to X_t .*

We will need the following lemma:

Lemma 2.17. *In the hypothesis of the definition above, there exists a constant C , depending only on X_0 , such that for all points $(v, t) \in S^n \times [0, T)$, we have:*

$$r^2 |\nabla r|^2 \leq C(r^2 + |\nabla r|^2).$$

In particular, if $r^2 > C$, then:

$$|\nabla r|^2 \leq \frac{Cr^2}{r^2 - C}.$$

Proof. As $x = r(z)z$, let $\partial_i \in T_z S^n$ and $\bar{\partial}_i = \partial_i x \in T_z X_t$ for $i = 1 \dots n$ be corresponding bases in the tangent spaces. Computing this explicitly in \mathbb{R}^{n+1} yields:

$$\begin{aligned}\bar{\partial}_i &= \partial_i x \\ &= \partial_i(r(z)z) \\ &= (\partial_i r(z))z + r(z)\partial_i\end{aligned}$$

Notice here that z and the ∂_i are orthogonal, therefore $|az + b^i \partial_i|^2 = a^2 + \sum_i (b^i)^2$. Computing this vector's scalar product with $r(z)z - \nabla r(z)$ yields:

$$\begin{aligned}\langle r(z)z - \nabla r(z), \bar{\partial}_i \rangle &= \langle r(z)z - \sum_j \partial_j r(z) \partial_j, \bar{\partial}_i \rangle \\ &= \langle r(z)z, \bar{\partial}_i \rangle - \langle \sum_j \partial_j r(z) \partial_j, \bar{\partial}_i \rangle \\ &= r(z) \langle z, \bar{\partial}_i \rangle - \sum_j \partial_j r(z) \langle \partial_j, \bar{\partial}_i \rangle \\ &= r(z) (\partial_i r(z)) - \partial_i r(z) r(z) \\ &= 0\end{aligned}$$

Moreover, notice that $\nabla r(z) \in T_z X_t$, which is orthogonal to z . Therefore, the following formula holds for the normal at a point with respect to the radial function:

$$\nu = \frac{r(z)z - \nabla r(z)}{\sqrt{r(z)^2 + |\nabla r(z)|^2}}$$

when taking $x = r(z)z$. Substituting the expressions above into the inequality from Corollary 2.12, we obtain:

$$\begin{aligned}C &\geq \left| r(z)z - \langle r(z)z, r(z)z - \nabla r(z) \rangle \frac{r(z)z - \nabla r(z)}{r(z)^2 + |\nabla r(z)|^2} \right| \\ &\geq \left| r(z)z - \frac{r(z)^2(r(z)z - \nabla r(z))}{r(z)^2 + |\nabla r(z)|^2} \right| \\ &\geq \left| \frac{r(z)z(r(z)^2 + |\nabla r(z)|^2) - r(z)^2(r(z)z - \nabla r(z))}{r(z)^2 + |\nabla r(z)|^2} \right| \\ &\geq \left| \frac{r(z)|\nabla r(z)|^2 z + r(z)^2 \nabla r(z)}{r(z)^2 + |\nabla r(z)|^2} \right|\end{aligned}$$

Multiplying by $r(z)^2 + |\nabla r(z)|^2$ and squaring (still using the fact that z and $\nabla r(z)$

are orthogonal):

$$\begin{aligned} |r(z)|^2 |\nabla r(z)|^2 z + r(z)^2 |\nabla r(z)|^2 &\leq C(r(z)^2 + |\nabla r(z)|^2) \\ r(z)^2 |\nabla r(z)|^4 + r(z)^4 |\nabla r(z)|^2 &\leq C^2(r(z)^2 + |\nabla r(z)|^2)^2 \\ r^2 |\nabla r|^4 + r^4 |\nabla r|^2 &\leq C^2(r^2 + |\nabla r|^2)^2. \end{aligned}$$

From here, the lemma follows. \square

The estimate is:

Proposition 2.18. *With the same hypotheses as the lemma above, there exists a constant C , depending only on X_0 , such that for all $(v, t) \in S^n \times [0, T)$, taking Π_0^v as in corollary 2.9, if $r(v, t)v \in \overline{H^+(\Pi_0^v)}$, then:*

$$|\nabla r(v, t)| \leq K.$$

Proof. By Corollary 2.9, there exists a constant $\epsilon > 0$, depending only on X_0 , such that for all $(z, t) \in S^n \times [0, T)$ with $r(z, t)z \in H^+(\Pi_0^v)$, and for all $w \in S^n$ with $\langle W, z \rangle > 1 - \epsilon$, we can reflect X_t up to the hyperplane $\Pi_w = \{r(z, t)z + p : \langle p, w \rangle = 0\}$. This means that $w \notin T_{r(z, t)z}X_t$, i.e., W is not tangent to X_t . In the proof of the previous lemma we showed that $r(z)z - \nabla r(z)$ is normal to the manifold's tangent plane at $r(z)z$, so, letting w_z be the projection of w in the z direction:

$$\begin{aligned} 0 &< \langle w, r(z)z - \nabla r(z) \rangle \\ \langle w, \nabla r(z) \rangle &< \langle w, r(z)z \rangle \\ \langle w - w_z, \nabla r(z) \rangle &< r(z) \langle w, z \rangle \\ |w - w_z| |\nabla r(z)| &< r(z) \langle w, z \rangle \\ |\nabla r(z)| &< \frac{r(z) \langle w, z \rangle}{|w - w_z|} \end{aligned}$$

by the fact that w is arbitrary, and estimating $|w - w_z|$ as $\sqrt{1^2 - (1 - \epsilon)^2}$:

$$\begin{aligned} |\nabla r(z)| &< \frac{r(z)(1 - \epsilon)}{\sqrt{1^2 - (1 - \epsilon)^2}} \\ |\nabla r(z)| &< r(z) \frac{(1 - \epsilon)}{\sqrt{\epsilon(2 - \epsilon)}} \end{aligned}$$

By the preceding lemma, if $r^2 > C$, then:

$$\begin{aligned} |\nabla r|^2 &\leq \frac{Cr^2}{r^2 - C} < C \\ |\nabla r| &\leq \sqrt{C} \end{aligned}$$

otherwise,

$$|\nabla r(z)| < C \frac{(1 - \epsilon)}{\sqrt{\epsilon(2 - \epsilon)}}$$

Hence:

$$|\nabla r(z)| < \max \left(\sqrt{C}, C \frac{(1 - \epsilon)}{\sqrt{\epsilon(2 - \epsilon)}} \right)$$

□

Finally, we include the following result about the part of X_t outside a ball:

Corollary 2.19. *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be an embedded solution to equation (2.2). Then, if, for a sphere B , $X_0 \subset B$, at all times $t \in [0, T)$ $X_t \setminus B$ is star-shaped with respect to the centre of B .*

Proof. $X_0 \subset B$ therefore we can reflect X_t about any hyperplane tangent to B by Corollary 2.9. By Corollary 2.7 $X_t \cap \overline{H^+(\pi)}$ is a graph over π , therefore, it is not possible that a normal line coming out of B and orthogonal to π intersects $X_t \cap \overline{H^+(\pi)}$ more than once. This implies that $X_t \setminus B$ is star-shaped with respect to the centre of B □

2.9 Expansive flows and ancient solutions

In this section, roughly following [5], we consider solutions to 2.2 which are defined on a larger interval (T_0, T_1) , with $-\infty \leq T_0 \leq 0 < T_1 \leq \infty$.

We will limit our analysis to a sub-class of solutions to 2.2:

Definition 2.20. *We say that a solution to 2.2 is expansive if $F < 0$.*

An example of such a flow is the inverse curvature flow, extensively studied in the literature.

Remark 2.21. This assumption on F implies that whenever $t < s$, $X_t \subset \text{int}(X_s)$. This is implied by the fact that, in the equation, the time derivative is always an outward pointing non-zero vector. In this sense, the hypersurface is expanding in the ambient space.

Definition 2.22. *Let $X : M^n \times (T_0, T_1) \rightarrow \mathbb{R}^{n+1}$ be an embedded expansive solution to equation (2.2). We say that X comes out of a point if there exists a point y_∞ such that for every $\varepsilon > 0$, there exists a time $\tau \in (T_0, T_1)$ such that $X_\tau \subset B_\varepsilon(y_\infty)$.*

Remark 2.23. If we take the optimal τ in the definition above, Remark 2.21 implies that the function $\tau(\varepsilon)$ mapping ε to the last time where X_τ is contained in the ball is non-increasing.

In particular, we will show that expansive solutions "coming out of a point" must be expanding spheres. It is easy to check that homothetically expanding spheres do satisfy the equation: Any spherical solution is completely determined by its radius at time t , because the equation is invariant at each point, being the principal curvatures constant: thus, solving the ordinary differential equation $r'(t) = \varphi(r(t))$ completely determines the flow, where $\varphi(r) = -F(\frac{1}{r}, \dots, \frac{1}{r})$.

We say that a solution to the equation is ancient if $T_0 = -\infty$. In the case of the spherical solution coming out of a point, there exists an ancient solution if and only if

$$+\infty = (T_1 - T_0) = \int_{T_0}^{T_1} 1 dt = \int_{T_0}^{T_1} \frac{r'(t)}{\varphi(r(t))} dt = \int_{r(T_0)}^{r(T_1)} \frac{1}{\varphi(r)} dr = \int_0^c \frac{1}{\varphi(r)} dr$$

$$\int_0^c \frac{1}{\varphi(r)} dr = +\infty$$

This equation allows us to determine, given F , if an ancient solution can exist (even if not coming out of a point). Solutions to (2.2) obey a comparison principle: if a solution is inside another one, it stays inside it at all times. This can be proven similarly to theorem 2.6, if at some (x, t) this fails, one can apply the maximum principle to show that they coincide in a neighbourhood, which is a contradiction if one takes the first or the last t where this happens. A generic solution, then, will be sandwiched between two expanding sphere solutions, one inside and one outside, and therefore cannot be ancient if no ancient expanding sphere solution exists.

The idea of ancient solutions was introduced by Richard Hamilton in his work on the Ricci flow. It has since been applied to other geometric flows as well as to other partial differential equations. Ancient solutions are significant because they capture key asymptotic features of the flow and often have unique, rigid properties that distinguish them from other solutions. For instance, in contractive flows, ancient solutions can form complex, non-spherical shapes, whereas in expansive flows, they tend to exhibit strong rigidity, commonly resulting in spherical shapes under certain conditions. These solutions thus help in understanding the geometry and topology of hypersurfaces as they evolve, especially in applications like mean curvature flow and inverse mean curvature flow in mathematics and physics.

Indeed, the property of coming out a point is particularly rigid, as shown in the following result:

Theorem 2.24. *Let $X : M^n \times (T_0, T_1) \rightarrow \mathbb{R}^{n+1}$ be a smooth, closed, embedded expansive solution to equation (2.2) coming out of a point. Then it is a family of expanding spheres.*

Previous results in geometric flows have shown multiple non-trivial examples of contracting flows sweeping the whole space. This could therefore be somewhat surprising, because it shows an opposite result, when running the equation in the opposite direction. It can however be explained intuitively by the idea that parabolic flows tend to *smooth things out*: it is thus possible to arrive to a point from a more irregular hypersurface, however the only thing that can “come out” of a point is something just as symmetric as a point, i.e. a sphere.

The proof is a relatively simple application of the reflection technique. The outline of the proof in [5] is the same as the one below, but using Corollary 2.9 makes it a bit simpler:

Proof. Fix any hyperplane π not passing through y_∞ . There is $R > 0$ such that $B_{2R}(y_\infty)$ does not intersect it. By definition, there is also a time $\tau \in (T_0, T_1)$ such that $X_\tau \subset B_R(y_\infty)$. By Corollary 2.9, then, we can reflect strictly X_t up to π for any $t > \tau$.

Now consider a sequence $\epsilon_n \rightarrow 0$. Up to a subsequence, we can then find a corresponding converging non-increasing sequence $\tau_n \rightarrow \bar{t} \in [T_0, T_1)$ (here note that \bar{t} can be $-\infty$) such that $X_{\tau_n} \subseteq B_{\epsilon_n}(y_\infty)$, therefore in (τ_n, T_1) I can reflect up to any hyperplane tangent to $B_{\epsilon_n}(y_\infty)$ in any direction, reasoning like we just did. At time \bar{t} , $X_{\bar{t}} \subseteq \cap_r B_r(y_\infty) = \{y_\infty\}$, thus we would have a singularity at \bar{t} if $\bar{t} \in (T_0, T_1)$ and therefore $\bar{t} = T_0$.

On the other hand, by construction, we can reflect X_t strictly about any hyperplane not intersecting $\cap_r B_r(y_\infty) = \{y_\infty\}$ at any time $t > \bar{t} = T_0$, hence we can reflect X_t strictly up to any hyperplane passing through y_∞ , in both directions, at any time $t \in (T_0, T_1)$. We observe that in the limit, the reflection property becomes non-strict, in the sense that we have to replace the interior of X in definition 2.4 with its closure, therefore X_t may touch its reflection at the limit plane, i.e. the one passing through y_∞ . Similarly, it cannot be that the other condition is the one causing the strict reflection definition to fail, as the other condition stays instead strict.

This implies that, taking any hyperplane passing through y_∞ and considering opposite directions for the reflection, X_t is symmetric about said hyperplane for any time $t \in (T_0, T_1)$. By Proposition 1.22, then, we conclude that X_t must be a ball. \square

Chapter 3

Extension to constant curvature spaces

In this chapter we want to extend the Chow-Gulliver result (theorem 2.6) to constant curvature spaces. We will use the same notation as in section 1.5 and 1.6. Throughout the chapter, by hyperplane we mean a totally geodesic hyperplane.

3.1 The equation in constant curvature spaces

As shown in [7], the equation we analysed in the previous chapter can be also considered in non-flat ambient spaces. In particular, we will analyse the case where the ambient space is one of those described in section 1.5: \mathbb{R}^{n+1} , \mathbb{H}^{n+1} , or \mathbb{S}_+^n . We again use the symbol \mathbb{M}_+^{n+1} to indicate any of these spaces. Let $X_0 : M^n \rightarrow \mathbb{M}_+^{n+1}$ be a manifold embedded in \mathbb{M}_+^{n+1} . Let $F : \{(\kappa_1, \dots, \kappa_n) \in \mathbb{R}^n | \kappa_1 \leq \dots \leq \kappa_n\} \rightarrow \mathbb{R}$ be a C^1 function satisfying:

$$\frac{\partial F}{\partial \kappa_i} > 0 \text{ for all } i = 1, \dots, n \quad (3.1)$$

and consider the evolution equation

$$\begin{cases} \frac{\partial X_t}{\partial t} = -F(\kappa_1(x), \dots, \kappa_n(x))\nu \\ X(0) = X_0 \end{cases} \quad (3.2)$$

where ν is the outward normal to $X_t(M^n)$ at the point $X_t(x)$ and $\kappa_1 \leq \dots \leq \kappa_n$ are the principal curvatures at $X_t(x)$.

As we saw in the previous chapter, it is a non-linear parabolic differential equation, as the calculation in chapter 2.2 is valid for any metric on $\mathbb{R}n + 1$, and in particular, for the metrics in our models for \mathbb{M}_+^{n+1} in section 1.5. The existence

result in [7] also holds as well in this case. Finally, the result in section 2.4 is not using the metric tensor, and therefore is valid in this setting as well, again because there are models on \mathbb{R}^{n+1} .

3.2 Extension of the result

Assume the hypothesis in section 1.5: $X : M^n \rightarrow \mathbb{M}_+^{n+1}$ is a hypersurface in a constant curvature ambient space, and we choose a point and a direction v to foliate the ambient space. Consider a hyperplane in the foliation, $\pi = \pi_{v,C}$. As in section 1.5 we can define the reflection about π . As in the previous chapter, let X^π be the reflection of X about π .

Then, \mathbb{M}_+^{n+1} is divided by π into two half-spaces:

$$H^+(\pi) = \bigcup_{s>C} \pi_{v,s} \text{ and } H^-(\pi) = \bigcup_{s<C} \pi_{v,s}.$$

Definition 3.1. We say we can reflect $X : M^n \rightarrow \mathbb{M}_+^{n+1}$ strictly with respect to π if both:

- $X^\pi \cap H^-(\pi) \subset \text{int}(X) \cap H^-(\pi)$
- The tangent spaces $T_x X$ and $T_x X^\pi$ do not coincide for each $x \in X(M^n) \cap \pi$ (when seen as subspaces of $T_x \mathbb{M}^{n+1}$)

This fundamentally means that the reflection of one of the halves of X on the other side of π is contained in the region inside M^n and the tangent spaces of X and of the half-reflection do not form a ninety degree angle with π , at all points on $\pi \cap X$. As the two tangent spaces are one the reflection of the other, this means that they do not coincide.

Definition 3.2. We say we can reflect $X : M^n \rightarrow \mathbb{R}^{n+1}$ strictly up to (π, v) if we can reflect M^n strictly with respect to $\pi_{v,s}$ for all hyperplanes $\pi_{v,s}$ such that $s < C$.

The result then becomes:

Theorem 3.3 (Extended Chow-Gulliver). *Let $X : M^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$ be a C^2 solution to equation (3.2). Then, if we can reflect $X(M^n, 0) = M_0$ strictly with respect to π , then for all $t \in [0, T)$ we can reflect $X(M^n, t) = M_t$ strictly with respect to π .*

Proof. As before, by contradiction, suppose that there is a time t such that the thesis is false, and that it is the smallest such t . Then, for all $\tau \in [0, t)$, $M_{\tau,\pi} \cap H^-(\pi) \subset \text{int}(M_\tau) \cap H^-(\pi)$; the unit vector orthogonal to π , V , is such that $V \notin T_x M_\tau$ for all $x \in M_\tau \cap \pi$ and $\tau \in [0, t)$; and either of the conditions fails at t , i.e. either:

(i) $M_{t,\pi} \cap H^-(\pi) \cap M_t \neq \emptyset$

(ii) The tangent spaces $T_x X$ and $T_x X^\pi$ coincide for some $x \in \pi$.

(i) Suppose the first case is true. Then, there exists $x_0 \in M_{t,\pi} \cap H^-(\pi) \cap M_t$ such that at x_0 the two manifolds are tangent.

We can then reason as in the proof of theorem 2.6.

(ii) Suppose instead that $T_x M_t = T_x M_t^\pi$ and in a neighbourhood of (x, t) both M_t and M_t^π are graphs of two smooth functions over $T_x M_t$ by 2.3, i.e. again

$$\begin{aligned} f &: (x, t) \mapsto x + \tilde{f}(x, t)\nu \\ f_\pi &: (x, t) \mapsto x + \tilde{f}_\pi(x, t)\nu \end{aligned}$$

We can do this because the theorem allowing us to do so is a theorem on smooth manifolds, and requires nothing on the metric, therefore the existence of models in \mathbb{R}^{n+1} of the ambient spaces allows us to do the same procedure. Reasoning again as in the proof of theorem 2.6, in $\overline{H^-(\pi)}$, $f_\pi \geq f$, because $M_\pi^n \cap H^-(\pi) \subset \text{int}(M^n) \cap H^-(\pi)$. Finally, $f(x, t) = f_\pi(x, t)$, hence $f_\pi - f(x, t) = 0$, and thus (x, t) is a minimum point on the boundary for $f_\pi - f$. Also, for the outward pointing normal to the boundary V

$$\frac{\partial f}{\partial V}(x, t) = \frac{\partial f_\pi}{\partial V}(x, t)$$

because the graphs are both tangent to $T_x M_t$. We note also that the models in section 1.5 are conformal, so a vector is orthogonal to the boundary if and only if it is orthogonal to the corresponding region when seen as a region in \mathbb{R}^{n+1} . Thus,

$$\frac{\partial(f - f_\pi)}{\partial V}(x, t) = 0$$

But we must have

$$\frac{\partial(f - f_\pi)}{\partial V}(x, t) > 0$$

at a minimum on the boundary by Proposition 1.13, a contradiction. \square

3.3 Extending Corollaries

We now shift the focus to extending the corollaries of theorem 2.6 in section 2.7.

The author is not aware of a standard definition of a support and a radial function in a curved setting, so we do not attempt to extend the results in section 2.8. **[DA RIMUOVERE SE CAMBIAMO IDEA]**

Clearly, corollary 2.8 has a direct equivalent in this setting:

Corollary 3.4. *Let $X : M^n \times [0, T) \rightarrow \mathbb{M}_+^{n+1}$ be a C^2 solution to equation (2.2). Then, if we can reflect X_0 strictly up to $(\pi_{v,C}, v)$, for all $t \in [0, T)$ we can reflect X_t strictly up to $(\pi_{v,C}, v)$.*

The second result that we can extend is Corollary 2.7, although the meaning of *graph* in curved spaces is ambiguous. We adapt the corollary as follows:

Corollary 3.5. *Let $X : M^n \times [0, T) \rightarrow \mathbb{M}_+^{n+1}$ be a C^2 embedded solution to equation (3.2). Then, if we can reflect X_0 strictly up to $(\pi_{v,C}, v)$, then for all $t \in [0, T)$, $X_t \cap H^+(\pi_{v,C})$ is such that the projection of the coordinates of its points onto $\pi_{v,C}$, $(p, \tau) \mapsto p$, is injective.*

This condition guarantees that we can build a map from $s : \overline{\text{int}(X_t \cap \pi_{v,C})} \rightarrow \mathbb{R}$ such that $(p, s(p)) \in X_t$, making $p \mapsto (p, s(p))$ act as a sort of curved graph.

Proof. By theorem 3.3 we can reflect up to $(\pi_{v,C}, v)$ at all times in $[0, T)$. Let $\gamma_p(s) = (p, s + C)$ be the path followed by $p \in \pi_{v,C}$ as the planes sweep through the ambient space. If two points exist with the same p coordinate, say (p, C_1) and (p, C_2) , the reflection about the hyperplane $\pi_{v, \frac{C_1+C_2}{2}}$ would map one onto the other, but as both C_1 and C_2 are greater than C , then we should also be able to reflect strictly about it, as we would have $\frac{C_1+C_2}{2} > C$. \square

Let $\tilde{B}_r(y) = \{p \in \mathbb{M}_+^{n+1} : \text{dist}(p, y) < r\}$. Reasoning exactly like in corollary 2.9, we also can extend it to this setting:

Corollary 3.6. *Let $X : M^n \times [0, T) \rightarrow \mathbb{M}_+^{n+1}$ be a C^2 embedded solution to equation (3.2). There exists $\varepsilon > 0$ depending only on X_0 such that for all $t \in [0, T)$ we can reflect X_t up to $(\Pi_0^v + \varepsilon v, v)$ for every $v \in S^n$. In particular, if $X_0 \subset \tilde{B}_R(C)$, then we can always reflect X_t up to (Π, v) whenever $H^+(\Pi) \cap \tilde{B}_{R-\varepsilon}(C) = \emptyset$.*

[Corollary 2.12 - If/when it can be done - DA SCRIVERE]

[Corollary 2.10 - If/when it can be done - DA SCRIVERE] I fear that this one cannot be done because the line mapping the minimum into the maximum in the reflection is not a geodesic anymore in the hyperbolic case

3.4 Extending Ancient solutions section

[DA SCRIVERE]

Like before:

Definition 3.7. We say that a solution to 3.2 is expansive if $F < 0$.

Definition 3.8. Let $X : M^n \times (T_0, T_1) \rightarrow \mathbb{M}_+^{n+1}$ be an embedded expansive solution to equation (3.2). We say that X comes out of a point if there exists a point y_∞ such that for every $\varepsilon > 0$, there exists a time $\tau \in (T_0, T_1)$ such that $X_\tau \subset \tilde{B}_\varepsilon(y_\infty)$, where $\tilde{B}_r(y) = \{p \in \mathbb{M}_+^{n+1} : \text{dist}(p, y) < r\}$.

[DA SCRIVERE]

Theorem 3.9. Let $X : M^n \times (T_0, T_1) \rightarrow \mathbb{M}_+^{n+1}$ be a smooth, closed, embedded expansive solution to equation (3.2) coming out of a point. Then it is a family of expanding distance spheres.

Proof. **[DA CONTROLLARE]**

Fix any hyperplane π not passing through y_∞ . There is $R > 0$ such that $\tilde{B}_{2R}(y_\infty)$ does not intersect it. By definition, there is also a time $\tau \in (T_0, T_1)$ such that $X_\tau \subset \tilde{B}_R(y_\infty)$. By Corollary 3.6, then, we can reflect strictly X_t up to π for any $t > \tau$.

Now consider a sequence $\epsilon_n \rightarrow 0$. Up to a subsequence, we can then find a corresponding converging non-increasing sequence $\tau_n \rightarrow \bar{t} \in [T_0, T_1]$ (here note that \bar{t} can be $-\infty$) such that $X_{\tau_n} \subseteq B_{\epsilon_n}(y_\infty)$, therefore in (τ_n, T_1) I can reflect up to any hyperplane outside $\tilde{B}_{\epsilon_n}(y_\infty)$ in any direction, reasoning like we just did. At time \bar{t} , $X_{\bar{t}} \subseteq \cap_r \tilde{B}_r(y_\infty) = \{y_\infty\}$, thus we would have a singularity at \bar{t} if $\bar{t} \in (T_0, T_1)$ and therefore $\bar{t} = T_0$.

On the other hand, by construction, we can reflect X_t strictly about any hyperplane not intersecting $\cap_r B_r(y_\infty) = \{y_\infty\}$ at any time $t > \bar{t} = T_0$, hence we can reflect X_t strictly up to any hyperplane passing through y_∞ , in both directions, at any time $t \in (T_0, T_1)$. We observe that in the limit, the reflection property becomes non-strict, in the sense that we have to replace the interior of X in definition 2.4 with its closure, therefore X_t may touch its reflection at the limit plane, i.e. the one passing through y_∞ . Similarly, it cannot be that the other condition is the one causing the strict reflection definition to fail, as the other condition stays instead strict.

This implies that, taking any hyperplane passing through y_∞ and considering opposite directions for the reflection, X_t is symmetric about said hyperplane for any time $t \in (T_0, T_1)$. By Proposition 1.22, then, we conclude that X_t must be a ball. \square

Bibliography

- [1] Chow B.; *Geometric Aspects of Aleksandrov Reflection and Gradient Estimates for Parabolic Equations*, Communications in Analysis and Geometry, Volume 5, Number 2, 389-409, 1997.
- [2] Chow, B. (2023). *Aleksandrov reflection for extrinsic geometric flows of Euclidean hypersurfaces*. Advanced Nonlinear Studies, Vol. 23 (Issue 1), pp. 20220034.
- [3] Ciraolo, G., Roncoroni, A., Vezzoni, L.; *Quantitative stability for hypersurfaces with almost constant curvature in space forms*, Annali di Matematica 200, 2043–2083 (2021).
- [4] Levitt, G.; Rosenberg, H.. (1984). *Symmetry of Constant Mean Curvature Hypersurfaces in Hyperbolic Space*. Retrieved from the University Digital Conservancy.
- [5] Sinestrari C., Risa S.; *Strong spherical rigidity of ancient solutions of expansive curvature flows*, Bull. London Math. Soc. 52, (2020) 94-99.
- [6] Gidas B., Wei-Ming Ni, L.Nirenberg; *Symmetry and Related Properties via the Maximum Principle*, Commun. Math. Phys. 68, 209-243, 1979.
- [7] Gerhard, H., Alexander, P. (1999). *Geometric evolution equations for hypersurfaces*. In: Hildebrandt, S., Struwe, M. (eds) Calculus of Variations and Geometric Evolution Problems. Lecture Notes in Mathematics, vol 1713. Springer, Berlin, Heidelberg.
- [8] Protter, M. H.; Weinberger, H. F.; *Maximum Principles in Differential Equations* , Springer, 1984.
- [9] Lee, J.M.; *Introduction to Riemannian Manifolds, Second Edition*, Springer Graduate Texts in Mathematics 176, 2018.
- [10] Lee, J.M.; *Introduction to Smooth Manifolds, Second Edition*, Springer Graduate Texts in Mathematics 218, 2013.

- [11] Evans, L.C.; *Partial Differential Equations*, American Mathematical Society, 2010.
- [12] Gerhardt, C.; *Curvature problems*, International Press, 2006.
- [13] Do Carmo, M.P.; *Differential Geometry of Curves and Surfaces, Second Edition*, Dover Books on Mathematics, 2016.
- [14] Mantegazza C. *Lecture Notes on Mean Curvature Flow*
- [15] Gardner R.J. *Geometric Tomography*. 2nd ed. Cambridge: Cambridge University Press; 2006
- [16] Gilardi, G.; *Analisi Matematica di Base*, Second Edition, McGraw-Hill, 2011.