



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)


Small Cell Lung Cancer

NCCN Evidence Blocks™

Version 1.2027 — July 1, 2026

NCCN.org

NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.





NCCN Guidelines Version 1.2024

Small Cell Lung Cancer

NCCN Evidence Blocks™

***Apar Kishor P. Ganti, MD, Chair †**
Fred & Pamela Buffett Cancer Center

***Billy W. Loo, Jr, MD, PhD/Vice Chair §**
Stanford Cancer Institute

Shahed Badiyan, MD §
UT Southwestern Simmons
Comprehensive Cancer Center

Michael Bassetti, MD §
University of Wisconsin Carbone Cancer Center

Anne Chiang, MD, PhD †
Yale Cancer Center/Smilow Cancer Hospital

Noura Choudhury, MD †
The UChicago Medicine
Comprehensive Cancer Center

Alissa Cooper, MD †
Dana-Farber Cancer Institute

Christopher A. D'Avella, MD †
Abramson Cancer Center
at the University of Pennsylvania

Afshin Dowlati, MD †
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center and
Cleveland Clinic Taussig Cancer Institute

Martin Edelman, MD †
Fox Chase Cancer Center

Charles Florsheim, JD ≠
Patient Advocate

Kathryn A. Gold, MD †
UC San Diego Moores Cancer Center

Jonathan W. Goldman, MD †
UCLA Jonsson Comprehensive Cancer Center

John C. Grecula, MD §
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Matthew Gumbleton, MD, PhD † ≠
Huntsman Cancer Institute
at the University of Utah

Christine Hann, MD, PhD †
Johns Hopkins Kimmel Cancer Center

Maya Khalil, MD † ≠
O'Neal Comprehensive Cancer Center at UAB

Robert Lee, MD ≡
Memorial Sloan Kettering Cancer Center

Jyoti Malhotra, MD †
City of Hope National Medical Center

Robert E. Merritt, MD ¶
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Nisha Mohindra, MD †
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Julian R. Molina, MD, PhD †
Mayo Clinic Comprehensive Cancer Center

Cesar Moran, MD ≠
The University of Texas
MD Anderson Cancer Center

Claire Mulvey, MD † ≠
UCSF Helen Diller Family
Comprehensive Cancer Center

Evan Osmundson, MD, PhD §
Vanderbilt-Ingram Cancer Center

Tejas Patil, MD †
University of Colorado Cancer Center

Saraswati Pokharel, MD ≠
Roswell Park Comprehensive Cancer Center

Sonam Puri, MD ≠ † ≠
Moffitt Cancer Center

Angel Qin, MD †
University of Michigan Rogel Cancer Center

Surbhi Singhal, MD †
UC Davis Comprehensive Cancer Center

Misty Shields, MD, PhD †
Indiana University Melvin and Bren Simon
Comprehensive Cancer Center

Tina D. Tailor, MD ϕ
Duke Cancer Institute

Saiana N. Waqar, MD † ≠
Siteman Cancer Center at Barnes-
Jewish Hospital and WashU Medicine

NCCN
Carly J. Cassara, MSc
Swathi Ramakrishnan, PhD
Victoria Reed, PhD

ϕ Diagnostic radiology
‡ Hematology/Hematology oncology
≡ Internal medicine
† Medical oncology
≠ Pathology
≠ Patient advocacy
≡ Pulmonary medicine
§ Radiotherapy/Radiation oncology
¶ Surgery/Surgical oncology
* Discussion writing committee member

[NCCN Guidelines Panel Disclosures](#)



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

[NCCN Small Cell Lung Cancer Panel Members](#)
[NCCN Evidence Blocks Definitions \(EB-1\)](#)

[Initial Evaluation and Staging \(SCL-1\)](#)
[Limited Stage, Workup and Treatment \(SCL-2\)](#)
[Extensive Stage, Primary Treatment \(SCL-5\)](#)
[Response Assessment Following Primary Treatment and Adjuvant Therapy \(SCL-6\)](#)
[Progressive Disease: Subsequent Therapy and Palliative Therapy \(SCL-8\)](#)

[Signs and Symptoms of Small Cell Lung Cancer \(SCL-A\)](#)
[Principles of Pathologic Review \(SCL-B\)](#)
[Principles of Surgical Resection \(SCL-C\)](#)
[Principles of Supportive Care \(SCL-D\)](#)
[Principles of Systemic Therapy \(SCL-E\)](#)
[Principles of Radiation Therapy \(SCL-F\)](#)
[Principles of Imaging \(SCL-G\)](#)

[Staging \(ST-1\)](#)

[Lung Neuroendocrine Tumors – See NCCN Guidelines for Neuroendocrine and Adrenal Tumors](#)
[Abbreviations \(ABBR-1\)](#)

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

[NCCN Guidelines Navigator™](#)
[NCCN Guidelines for Patients®](#)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS

5					
4					
3					
2					
1					

E S Q C A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

Example Evidence Block

5					
4					
3					
2					
1					

E S Q C A

E = 4
S = 4
Q = 3
C = 4
A = 3

Efficacy of Regimen/Agent

5	Highly effective: Cure likely and often provides long-term survival advantage
4	Very effective: Cure unlikely but sometimes provides long-term survival advantage
3	Moderately effective: Modest impact on survival, but often provides control of disease
2	Minimally effective: No, or unknown impact on survival, but sometimes provides control of disease
1	Palliative: Provides symptomatic benefit only

Safety of Regimen/Agent

5	Usually no meaningful toxicity: Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
4	Occasionally toxic: Rare significant toxicities or low-grade toxicities only; little interference with ADLs
3	Mildly toxic: Mild toxicity that interferes with ADLs
2	Moderately toxic: Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
1	Highly toxic: Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

Note: For significant chronic or long-term toxicities, score decreased by 1

Quality of Evidence

5	High quality: Multiple well-designed randomized trials and/or meta-analyses
4	Good quality: One or more well-designed randomized trials
3	Average quality: Low quality randomized trial(s) or well-designed non-randomized trial(s)
2	Low quality: Case reports or extensive clinical experience
1	Poor quality: Little or no evidence

Consistency of Evidence

5	Highly consistent: Multiple trials with similar outcomes
4	Mainly consistent: Multiple trials with some variability in outcome
3	May be consistent: Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
2	Inconsistent: Meaningful differences in direction of outcome between quality trials
1	Anecdotal evidence only: Evidence in humans based upon anecdotal experience

Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)

5	Very inexpensive
4	Inexpensive
3	Moderately expensive
2	Expensive
1	Very expensive



NCCN Guidelines Version 1.2027


Small Cell Lung Cancer

NCCN Evidence Blocks™

DIAGNOSIS

INITIAL EVALUATION^{a,b}

STAGE



^a If extensive stage is established, further staging evaluation is optional and dependent on the clinical situation. However, brain imaging—MRI (preferred) or CT with contrast—is recommended.

^b Workup of SCLC should be expedited, with studies done in parallel whenever possible.

^c [Signs and Symptoms of Small Cell Lung Cancer \(SCL-A\)](#).

^d [Principles of Pathologic Review \(SCL-B\)](#).

^e Brain MRI is more sensitive than CT for identifying brain metastases and is preferred over CT. See Principles of Imaging ([SCL-G](#)).

^f If FDG-PET/CT is not available, bone scan may be used to identify bone metastases. Pathologic confirmation is recommended for isolated or equivocal lesions if their involvement would change clinical management.

^g Biomarker testing via blood, tissue, or both can be considered in rare cases—particularly for patients with extensive-stage/relapsed SCLC who do not smoke tobacco, lightly smoke, have remote smoking history, or have diagnostic or therapeutic dilemma, or at time of relapse—if not previously done, because this may change management.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

STAGE

ADDITIONAL WORKUP^b



^b Workup of SCLC should be expedited, with studies done in parallel whenever possible.

^h While most pleural effusions in patients with lung cancer are due to tumor, there are a few patients in whom multiple cytopathologic examinations of pleural fluid are negative for tumor and fluid is non-bloody and not an exudate. When these elements and clinical judgment dictate that the effusion is not related to the tumor, the effusion should be excluded as a staging element. Pericardial effusion is classified using the same criteria.

ⁱ Selection criteria include: nucleated red blood cells (RBCs) on peripheral blood smear, neutropenia, or thrombocytopenia suggestive of bone marrow infiltration. Perform bone marrow biopsy only if it changes clinical management.

^j [Principles of Surgical Resection \(SCL-C\)](#).

^k Lymph node staging procedures include endobronchial ultrasound-guided biopsy (preferred, if expertise exists), mediastinoscopy, mediastinotomy, esophageal ultrasound-guided biopsy, and video-assisted thoracoscopy. If nodal involvement is established by one modality of mediastinal staging, additional modalities are not necessary, but staging biopsies of visible nodes in all accessible nodal stations is encouraged to inform radiotherapy targeting.


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™



^j [Principles of Surgical Resection \(SCL-C\)](#).

^k Lymph node staging procedures include endobronchial ultrasound-guided biopsy (preferred, if expertise exists), mediastinoscopy, mediastinotomy, esophageal ultrasound-guided biopsy, and video-assisted thoracoscopy. If nodal involvement is established by one modality of mediastinal staging, additional modalities are not necessary, but staging biopsies of visible nodes in all accessible nodal stations is encouraged to inform radiotherapy targeting.

^l Select patients may be treated with systemic therapy/RT as an alternative to surgical resection.

^m [Principles of Systemic Therapy \(SCL-E\)](#).

ⁿ [Principles of Radiation Therapy \(SCL-F\)](#).

^o Systemic therapy may be initiated first if time to initiation of stereotactic body radiotherapy (SABR) will be prolonged.

^p [Principles of Imaging \(SCL-G\)](#).

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.**




NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRIMARY TREATMENT



^m [Principles of Systemic Therapy \(SCL-E\)](#)..
ⁿ [Principles of Radiation Therapy \(SCL-F\)](#)..
^p [Principles of Imaging \(SCL-G\)](#)..
^q [Principles of Supportive Care \(SCL-D\)](#)..

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

STAGE

PRIMARY TREATMENT



^m [Principles of Systemic Therapy \(SCL-E\)](#).

ⁿ [Principles of Radiation Therapy \(SCL-F\)](#).

^p [Principles of Imaging \(SCL-G\)](#).

^q [Principles of Supportive Care \(SCL-D\)](#).

^r If brain metastases progress while on systemic therapy, it is recommended that brain RT is initiated before completion of systemic therapy.

^s For transformation to SCLC from NSCLC, consider referral to a center with expertise ([SCL-E 4 of 6](#)).

^t Initiate steroids for patients with symptomatic neurologic disease. Eg, dexamethasone 10 mg loading dose followed by 4–6 mg maintenance dose (IV or PO every 4–6 hours [or as appropriate]). Kumar A, et al. Clin Spine Surg 2017;4:156-163.

^u With neurologic symptoms, RT is preferred before systemic therapy. Systemic therapy may start first if RT cannot be started expeditiously or if controlling systemic symptoms is more urgent.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.**




NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

RESPONSE ASSESSMENT FOLLOWING PRIMARY TREATMENT

ADJUVANT THERAPY



[Surveillance \(SCL-7\)](#)

[Subsequent Therapy/Palliative Therapy \(SCL-8\)](#)

Footnotes

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

FOOTNOTES

^c [Signs and Symptoms of Small Cell Lung Cancer \(SCL-A\)](#).

^e Brain MRI is more sensitive than CT for identifying brain metastases and is preferred over CT. See [Principles of Imaging \(SCL-G\)](#).

^m [Principles of Systemic Therapy \(SCL-E\)](#).

ⁿ [Principles of Radiation Therapy \(SCL-F\)](#).

^v PCI is not recommended in patients with poor PS or impaired neurocognitive function. Increased cognitive decline after PCI has been observed in older adults (≥60 years) in prospective trials; the risks and benefits of PCI versus close brain surveillance, MRI (preferred), or CT with contrast should be carefully discussed with these patients. Edelman MJ, Am Soc Clin Oncol Educ Book 2020:40:24-28.

^w The benefit of PCI is unclear in patients who have undergone definitive therapy for pathologic stage I (T1–2a,N0,M0) SCLC. See [Principles of Radiation Therapy \(SCL-F\)](#).

^x Sequential RT to thorax in selected patients, especially with residual thoracic disease and low-bulk extrathoracic metastatic disease that has responded to systemic therapy.

^y If PCI is considered, it should be given prior to durvalumab.

^z For those with good PS who are medically inoperable or decision was made not to pursue surgical resection.

^{aa} Imaging every 8 weeks for the first 18 months and every 12 weeks after.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.**




NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

SURVEILLANCE



^c [Signs and Symptoms of Small Cell Lung Cancer \(SCL-A\)](#).

^{aa} [NCCN Guidelines for Survivorship](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PROGRESSIVE DISEASE

SUBSEQUENT THERAPY/PALLIATIVE THERAPY



^m [Principles of Systemic Therapy \(SCL-E\)](#).

ⁿ [Principles of Radiation Therapy \(SCL-F\)](#).

^p [Principles of Imaging \(SCL-G\)](#).

^q [Principles of Supportive Care \(SCL-D\)](#).

^{cc} Consider biomarker testing, if not previously done, to determine clinical trial eligibility.

^{dd} [NCCN Guidelines for Palliative Care](#).

^{ee} For central nervous system (CNS) progression only, continue systemic therapy and treat the brain metastases with RT. See [Principles of Radiation Therapy \(SCL-F\)](#).

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

SIGNS AND SYMPTOMS OF SMALL CELL LUNG CANCER

Signs and Symptoms Due to Local Primary Tumor Growth

- Cough – endobronchial irritation, bronchial compression
- Hemoptysis – usually central or cavitory lesion
- Wheezing – partially obstructing endobronchial lesion
- Fever – postoperative pneumonia
- Dyspnea – bronchial obstruction, pneumonia, pleural effusion

Signs and Symptoms Due to Primary Tumor Invasion or Regional Lymphatic Metastases

- Hoarseness – left vocal cord paralysis due to tumor invasion or lymphadenopathy in the aortopulmonary window
- Hemidiaphragm elevation – due to phrenic nerve compression
- Dysphagia – due to esophageal compression
- Chest pain – involvement of pleura or chest wall, often dull and non-localized
- SVC syndrome – due to local invasion into mediastinum or lymphadenopathy in right paratracheal region
- Pericardial effusion and tamponade
- Cervical or supraclavicular lymph node enlargement

Signs and Symptoms Due to Extrathoracic (Hematogenous) Metastases

- Brain metastases:
 - Headache, focal weakness or numbness, confusion, slurred speech, gait instability, incoordination
- Leptomeningeal carcinomatosis:
 - Headache, confusion, cranial nerve palsy, diplopia, slurred speech, radicular back pain, spinal cord compression
- Adrenal metastases:
 - Mid-back or flank pain, costovertebral angle tenderness
 - Adrenal insufficiency due to tumor involvement (rare)
- Liver metastases:
 - Right upper quadrant pain or tenderness, jaundice, fatigue, fever, hepatomegaly
- Bone metastases:
 - Bone pain
 - Spinal cord compression – back pain, muscle weakness, numbness, paresthesia, loss of bowel and bladder control
- Constitutional:
 - Anorexia/cachexia – weight loss
 - Fatigue

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



SIGNS AND SYMPTOMS OF SMALL CELL LUNG CANCER

Signs and Symptoms of Paraneoplastic Syndromes

- Presence does not imply metastases or incurability

Endocrine:

- Due to ectopic peptide hormone production
- Usually reversible with successful anti-tumor therapy
- Syndrome of inappropriate antidiuretic hormone secretion (SIADH)^a:
 - Ectopic vasopressin (antidiuretic hormone, ADH) secretion
 - Clinically significant hyponatremia in 5%–10% of SCLC
 - Malaise, weakness, confusion, obtundation, volume depletion, nausea
 - Hyponatremia, euolemia, low serum osmolality, inappropriately concentrated urine osmolality, normal thyroid and adrenal function
- Cushing syndrome^a:
 - Ectopic adrenocorticotrophic hormone (ACTH) secretion
 - Weight gain, moon facies, hypertension, hyperglycemia, generalized weakness
 - High serum cortisol and ACTH, hypernatremia, hypokalemia, alkalosis

Neurologic:

- All specific syndromes are rare
 - Subacute cerebellar degeneration – ataxia, dysarthria
 - Encephalomyelitis – confusion, obtundation, dementia
 - Sensory neuropathy – pain, sensory loss
 - Lambert-Eaton myasthenic syndrome (LEMS)^a – proximal muscle weakness, autonomic dysfunction
 - ◊ Neurologic workup, preferably in consultation with neurology, may include but not limited to tests for P/Q- and N-type voltage gated calcium channel (VGCC) antibodies.
 - Cancer-associated retinopathy – visual loss, photosensitivity
- Consider early subspecialty consultation for unusual paraneoplastic neurologic syndromes to ensure the most recent management is done
- If paraneoplastic neurologic syndrome is suspected, consider obtaining a neurologic consultation and/or comprehensive paraneoplastic antibody panel

Hematologic:

- Anemia
- Leukemoid reaction – leukocytosis
- Trousseau syndrome – migratory thrombophlebitis

^a [Principles of Supportive Care \(SCL-D\).](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2024

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGIC REVIEW

Pathologic Evaluation

- Pathologic evaluation is performed to determine the histologic classification of lung tumors and relevant staging parameters.
- The World Health Organization (WHO) tumor classification system provides the foundation for the classification of lung tumors, including histologic subtype, staging factors, clinical features, molecular characteristics, genetics, and epidemiology.¹⁻³
- SCLC is a poorly differentiated neuroendocrine carcinoma. Distinguishing SCLC from other neuroendocrine tumors, particularly typical and atypical carcinoids, is important due to significant differences in epidemiology, genetics, treatment, and prognosis.⁴⁻⁶
- SCLC can be diagnosed on good-quality histologic samples via high-quality hematoxylin and eosin (H&E)-stained sections or on well-preserved cytologic samples.
 - ▶ SCLC is characterized by small blue cells with scant cytoplasm, high nuclear-to-cytoplasmic ratio, granular chromatin, and absent or inconspicuous nucleoli.
 - ▶ SCLC cells are round, oval, or spindle-shaped with molding and high mitotic counts.⁷⁻⁹
 - ▶ The most useful characteristics for distinguishing SCLC from large-cell neuroendocrine carcinoma (LCNEC) are the high nuclear-to-cytoplasmic ratio and paucity of nucleoli in SCLC.
- Careful counting of mitoses is essential, because it is the most important histologic criterion for distinguishing SCLC from typical and atypical carcinoids. Strongly recommend a second opinion—with a pathologist specializing in the diagnosis of thoracic malignancies—for diagnostic dilemma, including carcinoid.
 - ▶ SCLC (>10 mitoses/2 mm² field); atypical carcinoid (2–10 mitoses/2 mm² field); typical carcinoid (0–1 mitoses/2 mm² field)
 - ▶ Mitoses should be counted in the areas of highest activity and per 2 mm² field, rather than per 10 high-power fields.
 - ▶ In tumors that are near the defined cutoffs of 2 or 10 mitoses per 2 mm², at least three 2-mm² fields should be counted and the calculated mean (rather than the single highest mitotic count) should be used to determine the overall mitotic rate.^{1,2}
- SCLC is often associated with necrosis. However, necrosis, usually punctate, is also seen in atypical carcinoid tumors. Counting mitotic figures helps to distinguish these two entities.
- Combined SCLC consists of both SCLC histology and NSCLC histology (squamous cell, adenocarcinoma, spindle/pleomorphic, and/or large cell). There is no minimal percentage of NSCLC histologic elements required; when any are present along with SCLC, this can be called combined SCLC, except in combination with LCNEC. At least 10% of the tumor should show LCNEC morphology to be classified as combined SCLC and LCNEC.¹
- Biomarker testing via blood, tissue, or both can be considered in rare cases—particularly for patients with extensive-stage/relapsed SCLC who do not smoke tobacco, lightly smoke, have remote smoking history, or have diagnostic or therapeutic dilemma, or at time of relapse—if not previously done, because this may change management.

Immunohistochemical Staining

- Immunohistochemistry can be very helpful in diagnosing SCLC in limited samples.^{5,7}
 - ▶ Nearly all SCLCs are positive for cytokeratin antibody mixtures with broad reactivity, such as AE1/AE3 and CAM5.2.^{1,10}
 - ▶ The majority of SCLCs are reactive to markers of neuroendocrine differentiation, including insulinoma-associated protein 1 (INSM1), CD56/NCAM, synaptophysin, and chromogranin A. Fewer than 5% of SCLCs are negative for all neuroendocrine markers.^{11,12} For cases with suspicious SCLC morphology without expression of neuroendocrine markers, POU2F3 immunohistochemical staining can be considered.^{13,14}
 - ▶ Thyroid transcription factor-1 (TTF-1) is positive in 85% to 90% of SCLCs.¹⁵⁻¹⁸
 - ▶ Additional immunohistochemical markers are useful in distinguishing small cell carcinoma from poorly differentiated non-small cell carcinoma and combined carcinoma using Napsin A as a marker of adenocarcinoma, and p40 or p63 as a marker of squamous differentiation.¹⁰ It should, however, be noted that p40 and p63 can be focally positive in small cell carcinoma.
- Ki-67 immunostaining can be very helpful in distinguishing SCLC from carcinoid tumors, especially in small biopsy samples with crushed or necrotic tumor cells in which counting mitotic figures is difficult.^{4,5}
 - ▶ The Ki-67 proliferative index in SCLC is typically 50% to 100%.¹
- Carcinoid tumors generally retain RB1 expression and show a wild-type p53 staining pattern. In contrast, most high-grade neuroendocrine carcinomas demonstrate loss of RB1 expression and/or aberrant (mutation-type) p53 expression, characterized by either diffuse strong nuclear staining (overexpression) or complete absence of staining (null pattern).

References

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.

PRINCIPLES OF PATHOLOGIC REVIEW - REFERENCES

- ¹ WHO Classification of Tumours Editorial Board. Thoracic Tumours. In: WHO classification of tumours series. 5th ed. Lyon, France. International Agency for Research on Cancer; 2021.
- ² Travis WD, Brambilla E, Burke AP, et al. Introduction to The 2015 World Health Organization Classification of Tumors of the Lung, Pleura, Thymus, and Heart. *J Thorac Oncol* 2015;10:1240-1242.
- ³ Travis WD, Brambilla E, Nicholson AG, et al, and WHO Panel. The 2015 World Health Organization Classification of Lung Tumors: Impact of Genetic, Clinical and Radiologic Advances Since the 2004 Classification. *J Thorac Oncol* 2015;10:1243-1260.
- ⁴ Pelosi G, Rindi G, Travis WD, Papotti M. Ki-67 antigen in lung neuroendocrine tumors: unraveling a role in clinical practice. *J Thorac Oncol* 2014;9:273-284.
- ⁵ Pelosi G, Rodriguez J, Viale G, Rosai J. Typical and atypical pulmonary carcinoid tumor overdiagnosed as small-cell carcinoma on biopsy specimens: a major pitfall in the management of lung cancer patients. *Am J Surg Pathol* 2005;29:179-187.
- ⁶ Rindi G, Klersy C, Inzani F, et al. Grading the neuroendocrine tumors of the lung: an evidence-based proposal. *Endocr Relat Cancer* 2013;21:1-16.
- ⁷ Travis WD. Advances in neuroendocrine lung tumors. *Ann Oncol* 2010;21 Suppl 7:vii65-vii71.
- ⁸ Zakowski MF. Pathology of small cell carcinoma of the lung. *Semin Oncol* 2003;30:3-8.
- ⁹ Nicholson SA, Beasley MB, Brambilla E, et al. Small cell lung carcinoma (SCLC): a clinicopathologic study of 100 cases with surgical specimens. *Am J Surg Pathol* 2002;26:1184-1197.
- ¹⁰ Masai K, Tsuta K, Kawago M, et al. Expression of squamous cell carcinoma markers and adenocarcinoma markers in primary pulmonary neuroendocrine carcinomas. *Appl Immunohistochem Mol Morphol* 2013;21:292-297.
- ¹¹ Rooper LM, Sharma R, Li QK, et al. INSM1 demonstrates superior performance to the individual and combined use of synaptophysin, chromogranin and CD56 for diagnosing neuroendocrine tumors of the thoracic cavity. *Am J Surg Pathol* 2017;41:1561-1569.
- ¹² Bellizzi AM. Immunohistochemistry in the diagnosis and classification of neuroendocrine neoplasms: what can brown do for you? *Hum Pathol* 2020;96:8-33.
- ¹³ Wang Y, Jin Y, Shen X, et al. POU2F3: A sensitive and specific diagnostic marker for neuroendocrine-low/negative small cell lung cancer. *Am J Surg Pathol* 2023;47:1059-1066.
- ¹⁴ Baine MK, Febres-Aldana CA, Chang JC, et al. POU2F3 in SCLC: clinicopathologic and genomic analysis with a focus on its diagnostic utility in neuroendocrine-low SCLC. *J Thorac Oncol* 2022;17:1109-1121.
- ¹⁵ Ordonez NG. Value of thyroid transcription factor-1 immunostaining in distinguishing small cell lung carcinomas from other small cell carcinomas. *Am J Surg Pathol* 2000;24:1217-1223.
- ¹⁶ Kaufmann O, Dietel M. Expression of thyroid transcription factor-1 in pulmonary and extrapulmonary small cell carcinomas and other neuroendocrine carcinomas of various primary sites. *Histopathology* 2000;36:415-420.
- ¹⁷ Lantuejoul S, Moro D, Michalides RJ, et al. Neural cell adhesion molecules (NCAM) and NCAM-PSA expression in neuroendocrine lung tumors. *Am J Surg Pathol* 1998;22:1267-1276.
- ¹⁸ Wick MR. Immunohistology of neuroendocrine and neuroectodermal tumors. *Semin Diagn Pathol* 2000;17:194-203.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2024

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SURGICAL RESECTION

- Stage I–IIA SCLC is diagnosed in less than 5% of patients with SCLC.
- Patients most likely to benefit from surgery are those with SCLC that is clinical stage I–IIA (T1–2,N0,M0) after standard staging evaluation (including CT of the chest and upper abdomen, brain imaging, FDG-PET/CT imaging, and lymph node staging).^{1,2}
 - ▶ Prior to resection, all patients should undergo mediastinoscopy or other surgical mediastinal staging to rule out occult nodal disease. This may also include an endoscopic staging procedure.
 - ▶ For patients undergoing definitive surgical resection, the preferred operation is lobectomy with mediastinal lymph node dissection or systematic lymph node sampling (eg, ≥3 N2 and ≥1 N1 stations).^{3–7}
- In patients who do not smoke, small lesions that are presumed to be small cell carcinoma on biopsy should be resected because they are likely carcinoids that have been misdiagnosed (See [NCCN Guidelines for Neuroendocrine and Adrenal Tumors](#)).
- Surgery may be considered for selected patients with T3 (based on size), N0 SCLC, if invasive mediastinal lymph node staging is negative.
- Intraoperative diagnosis of likely SCLC in a patient with no prior biopsy
 - ▶ Mediastinal lymph node dissection or systematic lymph node sampling with frozen section is recommended to assess extent of disease and overall burden of disease.
 - ▶ If primary site and lymph nodes appear resectable, perform anatomic resection, preferably lobectomy. Pneumonectomy should not be performed if needed to encompass nodal metastatic disease.
- Patients who undergo complete resection should be treated with postoperative systemic therapy.⁸ Patients without nodal metastases should be treated with systemic therapy alone. Patients with N2 or N3 nodal metastases should be treated with postoperative concurrent or sequential systemic therapy and mediastinal RT. Patients with N1 nodal metastases may be considered for postoperative mediastinal radiation.
- The benefit of PCI is unclear in patients who have undergone definitive therapy for pathologic stage I (T1–2a,N0,M0); see [SCL-F](#).

¹ Lad T, Piantadosi S, Thomas P, et al. A prospective randomized trial to determine the benefit of surgical resection of residual disease following response of small cell lung cancer to combination chemotherapy. Chest 1994;106:320S-323S.

² Yang CJ, Chan DY, Shah SA, et al. Long-term survival after surgery compared with concurrent chemoradiation for node-negative small cell lung cancer. Ann Surg 2018;268:1105-1112.

³ Katz MHG, Francescatti AB, Hunt KK; Cancer Surgery Standards Program of the American College of Surgeons. Technical Standards for Cancer Surgery: Commission on Cancer Standards 5.3-5.8. Ann Surg Oncol 2022;29:6549-6558.

⁴ Darling GE, Allen MS, Decker PA, et al. Randomized trial of mediastinal lymph node sampling versus complete lymphadenectomy during pulmonary resection in the patient with N0 or N1 (less than hilar) non-small cell carcinoma: results of the American College of Surgery Oncology Group Z0030 Trial. J Thorac Cardiovasc Surg 2011;141:662-670.

⁵ Darling GE, Allen MS, Decker PA, et al. Number of lymph nodes harvested from a mediastinal lymphadenectomy: results of the randomized, prospective American College of Surgeons Oncology Group Z0030 trial. Chest 2011;139:1124-1129.

⁶ Osarogiagbon RU, Decker PA, Ballman K, et al. Survival Implications of Variation in the Thoroughness of Pathologic Lymph Node Examination in American College of Surgeons Oncology Group Z0030 (Alliance). Ann Thorac Surg 2016;102:363-369.

⁷ Su S, Scott WJ, Allen MS, et al. Patterns of survival and recurrence after surgical treatment of early stage non-small cell lung carcinoma in the ACOSOG Z0030 (ALLIANCE) trial. J Thorac Cardiovasc Surg 2014;147:747-752: Discussion 752-753.

⁸ Yang CE, Chan DY, Speicher PJ, et al. Role of adjuvant therapy in a population-based cohort of patients with early-stage small-cell lung cancer. J Clin Oncol 2016;34:1057-1064.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SUPPORTIVE CARE

- **Smoking cessation advice, counseling, and pharmacotherapy**
 - Use the 5 A's Framework: Ask, Advise, Assess, Assist, Arrange (<https://www.ahrq.gov/prevention/guidelines/tobacco/5steps.html>)
 - See [NCCN Guidelines for Smoking Cessation](#)
- **Granulocyte-macrophage colony-stimulating factor (GM-CSF) or granulocyte colony-stimulating factor (G-CSF) are not recommended during concurrent systemic therapy plus RT (category 1 for not using GM-CSF).^{1,2}**
- **Trilaciclib or G-CSF may be used as prophylactic options to decrease the incidence of chemotherapy-induced myelosuppression when administering platinum/etoposide ± immune checkpoint inhibitor (ICI)-containing regimens or a topotecan-containing regimen for extensive stage SCLC (ES-SCLC).**
- **SIADH**
 - Fluid restriction
 - Saline infusion for symptomatic patients
 - Demeclocycline
 - Vasopressin receptor inhibitors (ie, conivaptan, tolvaptan) for refractory hyponatremia
- **Cushing syndrome**
 - Consider ketoconazole. If not effective, consider metyrapone.
 - Consider referral to an appropriate endocrinology subspecialist.
- **Leptomeningeal disease: See [NCCN Guidelines for Central Nervous System Cancers](#)**
- **Lambert-Eaton myasthenic syndrome**
 - Consider amifampridine or intravenous immunoglobulin (IVIG) in consultation with neurology
- **Pain management: See [NCCN Guidelines for Adult Cancer Pain](#)**
- **Nausea/vomiting: See [NCCN Guidelines for Antiemesis](#)**
- **Psychosocial distress: See [NCCN Guidelines for Distress Management](#)**
 - Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health
- **See [NCCN Guidelines for Palliative Care](#) as indicated**
- **Consider early subspecialty consultation for unusual paraneoplastic neurologic syndromes to ensure the most recent management is done**

¹ Bunn PA, Crowley J, Kelly K, et al. Chemoradiotherapy with or without granulocyte-macrophage colony-stimulating factor in the treatment of limited stage small-cell lung cancer: a prospective phase III randomized study of the Southwest Oncology Group. J Clin Oncol 1995;13:1632-1641.

² Wang C, Zhu S, Miao C, et al. Safety and efficacy of pegylated recombinant human granulocyte colony-stimulating factor during concurrent chemoradiotherapy for small-cell lung cancer: a retrospective, cohort-controlled trial. BMC Cancer 2022;22:542.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

PRIMARY OR ADJUVANT THERAPY FOR LIMITED-STAGE SCLC:

Four cycles of cytotoxic chemotherapy are recommended.

Planned cycle length should be every 21–28 days during concurrent RT.

The use of myeloid growth factors is not recommended during concurrent cytotoxic chemotherapy plus RT (category 1 for not using GM-CSF).¹

Preferred

- Cisplatin 75 mg/m² Day 1 and Etoposide 100 mg/m² Days 1, 2, 3²
- Cisplatin 60 mg/m² Day 1 and Etoposide 120 mg/m² Days 1, 2, 3³
- Carboplatin area under the curve (AUC) 5–6 Day 1 and Etoposide 100 mg/m² Days 1, 2, 3^{a,4}
- Consolidation Therapy
 - ▶ Durvalumab 1500 mg Day 1 every 28 Days for up to 24 months (category 1)^{b,5}

Other Recommended

- Cisplatin 25 mg/m² Days 1, 2, 3 and Etoposide 100 mg/m² Days 1, 2, 3²

PRIMARY THERAPY FOR EXTENSIVE-STAGE SCLC^c:

Four cycles of cytotoxic chemotherapy are recommended, but some patients may receive up to 6 cycles based on response and tolerability after 4 cycles.

Preferred

- Carboplatin AUC 5 Day 1 and Etoposide 100 mg/m² Days 1, 2, 3 and Atezolizumab 1200 mg Day 1 every 21 days x 4 cycles followed by maintenance Atezolizumab 1200 mg Day 1, every 21 days (category 1 for all)^{d,e,f,6}
- Carboplatin AUC 5 Day 1 and Etoposide 100 mg/m² Days 1, 2, 3 and Atezolizumab 1200 mg Day 1 every 21 days x 4 cycles followed by maintenance Atezolizumab 1680 mg Day 1, every 28 days^{d,e,f}
- Carboplatin AUC 5 Day 1 and Etoposide 100 mg/m² Days 1, 2, 3 and Atezolizumab 1200 mg Day 1 every 21 days x 4 cycles followed by maintenance Lurbinectedin 3.2 mg/m² and Atezolizumab 1200 mg Day 1, every 21 days (category 1 for all)^{g,h,7}
- Carboplatin AUC 5–6 Day 1 and Etoposide 80–100 mg/m² Days 1, 2, 3 and Durvalumab 1500 mg Day 1 every 21 days x 4 cycles followed by maintenance Durvalumab 1500 mg Day 1 every 28 days (category 1 for all)^{d,e,i,8}
- Cisplatin 75–80 mg/m² Day 1 and Etoposide 80–100 mg/m² Days 1, 2, 3 and Durvalumab 1500 mg Day 1 every 21 days x 4 cycles followed by maintenance Durvalumab 1500 mg Day 1 every 28 days (category 1 for all)^{d,e,i,8}

[See Evidence Blocks on EB-1](#)

Other Recommended

- Carboplatin AUC 5–6 Day 1 and Etoposide 100 mg/m² Days 1, 2, 3⁹
- Cisplatin 75 mg/m² Day 1 and Etoposide 100 mg/m² Days 1, 2, 3¹⁰
- Cisplatin 80 mg/m² Day 1 and Etoposide 80 mg/m² Days 1, 2, 3¹¹
- Cisplatin 25 mg/m² Days 1, 2, 3 and Etoposide 100 mg/m² Days 1, 2, 3¹²

Useful in Certain Circumstances

- Carboplatin AUC 5 Day 1 and Irinotecan 50 mg/m² Days 1, 8, 15¹³
- Cisplatin 60 mg/m² Day 1 and Irinotecan 60 mg/m² Days 1, 8, 15¹⁴
- Cisplatin 30 mg/m² Days 1, 8 and Irinotecan 65 mg/m² Days 1, 8¹⁵

[Footnotes \(SCL-E 2 of 6\)](#)
[Subsequent Systemic Therapy \(SCL-E 3 of 6\)](#)
[References \(SCL-E 5 of 6\)](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY - FOOTNOTES

- ^a Cisplatin contraindicated or not tolerated.
- ^b Those who did not experience disease progression after systemic therapy + concurrent RT may continue durvalumab until disease progression or intolerable toxicity, or for a maximum of 24 months.
- ^c For transformation to SCLC from NSCLC, consider referral to a center with expertise ([SCL-E 4 of 6](#)).
- ^d Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or concurrent use of immunosuppressive agents. If TKI is continued for transformed SCLC from NSCLC with oncogenic driver, ICI should be avoided due to known toxicity.
- ^e Maintenance immunotherapy with either atezolizumab or durvalumab should continue until progression or intolerable toxicity.
- ^f Atezolizumab and hyaluronidase-tqjs subcutaneous injection may be substituted for IV atezolizumab. Atezolizumab and hyaluronidase-tqjs has different dosing and administration instructions compared to atezolizumab for intravenous infusion.
- ^g Consider adding lurbinectedin to maintenance atezolizumab in patients who have achieved at least stable disease following four cycles of induction chemoimmunotherapy, have an ECOG PS of 0–1, and no history of brain metastases.
- ^h This regimen is not for use as re-treatment in subsequent-line therapy if lurbinectedin has been used previously.
- ⁱ Included patients with asymptomatic untreated brain metastases.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

SCLC SUBSEQUENT SYSTEMIC THERAPY (PS 0–2) ^j Consider dose reduction or growth factor support for patients with PS 2
Preferred • Tarlatamab-dlle^{k,l,16} (category 1) • Clinical trial enrollment • Irinotecan^{m,17,18} • Lurbinectedin (if not previously used)^{19,20} • If prolonged disease-free time, re-treatment with platinum-based doublet with or without immunotherapy²¹⁻²⁵ • Topotecan Oral (PO) or Intravenous (IV)^{17,23,26-28} Other Recommended • CAV (Cyclophosphamide/Doxorubicin/Vincristine)²⁶ • Docetaxel²⁹ • Gemcitabine^{30,31,32} • Nivolumabⁿ or Pembrolizumab^o (if not previously treated with an ICI)^{d,33-36} • Oral Etoposide^{37,38} • Paclitaxel^{39,40} • Temozolomide⁴⁰⁻⁴²

[See Evidence Blocks on EB-2](#)

^d Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or concurrent use of immunosuppressive agents. If TKI is continued for transformed SCLC from NSCLC with oncogenic driver, ICI should be avoided due to known toxicity.

^j Subsequent systemic therapy refers to second-line and beyond therapy.

^k For extensive stage with disease progression on or after platinum-based chemotherapy.

^l [NCCN Guidelines for Management of CAR T-Cell and Lymphocyte Engager-Related Toxicities](#).

^m For patients with CNS disease, consider using irinotecan.

ⁿ Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV nivolumab.

^o Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.

[References on \(SCL-E 5 of 6\)](#)

SCL-E
3 OF 6



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

- **Transformed SCLC from NSCLC with an Oncogenic Driver**

- ▶ This is a rare population of patients with very limited data to guide treatment.⁴³⁻⁴⁶
- ▶ Systemic cytotoxic chemotherapy is recommended using the NCCN Guidelines for Small Cell Lung Cancer.^{43,44}
- ▶ The role of immunotherapy in this setting is unclear based on limited data.⁴³⁻⁴⁶
- ▶ If tyrosine kinase inhibitor (TKI) is continued, ICI should be avoided, due to known toxicity.^{45,47,48}
- ▶ Consider referral to a center with experience managing transformed SCLC.

[References on \(SCL-E 5 of 6\)](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY – REFERENCES

- 1 Bunn PA, Crowley J, Kelly K, et al. Chemoradiotherapy with or without granulocyte-macrophage colony-stimulating factor in the treatment of limited stage small-cell lung cancer: a prospective phase III randomized study of the Southwest Oncology Group. *J Clin Oncol* 1995;13:1632-1641.
- 2 Faivre-Finn C, Snee M, Ashcroft L, et al. Concurrent once-daily versus twice-daily chemoradiotherapy in patients with limited stage small-cell lung cancer (CONVERT): an open-label, phase 3, randomised, superiority trial. *Lancet Oncol* 2017;18:1116-1125.
- 3 Turrisi AT 3rd, Kim K, Blum R, et al. Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with cisplatin and etoposide. *N Engl J Med* 1999;340:265-271.
- 4 Karlos DV, Samantas E, Briassoulis E, et al. Randomized comparison of early versus late hyperfractionated thoracic irradiation concurrently with chemotherapy in limited disease small-cell lung cancer: a randomized phase II study of the Hellenic Cooperative Oncology Group (HeCOG). *Ann Oncol* 2001;12:1231-1238.
- 5 Cheng Y, Spigel D, Cho BC, et al. Durvalumab after Chemoradiotherapy in Limited-Stage Small-Cell Lung Cancer. *N Engl J Med* 2024;39:1313-1327.
- 6 Horn L, Mansfield A, Szczesna A, et al. First-line atezolizumab plus chemotherapy in extensive stage small-cell lung cancer. *N Engl J Med* 2018;379:2220-2229.
- 7 Goldman JW, Dvorkin M, Chen Y, et al. Durvalumab, with or without tremelimumab, plus platinum-etoposide versus platinum-etoposide alone in first-line treatment of extensive stage small-cell lung cancer (CASPIAN): updated results from a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2021;22:51-65.
- 8 Okamoto H, Watanabe K, Nishiwaki Y, et al. Phase II study of area under the plasma-concentration-versus-time curve-based carboplatin plus standard-dose intravenous etoposide in elderly patients with small cell lung cancer. *J Clin Oncol* 1999;17:3540-3545.
- 9 Spigel DR, Townley PM, Waterhouse DM, et al. Randomized phase II study of bevacizumab in combination with chemotherapy in previously untreated extensive stage small-cell lung cancer: results from the SALUTE trial. *J Clin Oncol* 2011;29:2215-2222.
- 10 Niell HB, Herndon JE 2nd, Miller AA, et al. Randomized phase III Intergroup trial of etoposide and cisplatin with or without paclitaxel and granulocyte-colony stimulating factor in patients with extensive stage small-cell lung cancer: Cancer and Leukemia Group B trial 9732. *J Clin Oncol* 2005;23:3752-3759.
- 11 Evans WK, Shepherd FA, Feld R, et al. VP-16 and cisplatin as first-line therapy for small-cell lung cancer. *J Clin Oncol* 1985;3:1471-1477.
- 12 Schmittel A, Fischer von Weikersthal L, Sebastian M, et al. A randomized phase II trial of irinotecan plus carboplatin versus etoposide plus carboplatin treatment in patients with extended disease small-cell lung cancer. *Ann Oncol* 2006;17:663-667.
- 13 Noda K, Nishiwaki Y, Kawahara M, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med* 2002;346:85-91.
- 14 Hanna N, Bunn Jr, PA, Langer C, et al. Randomized phase III trial comparing irinotecan/cisplatin with etoposide/cisplatin in patients with previously untreated extensive stage disease small-cell lung cancer. *J Clin Oncol* 2006;24:2038-2043.
- 15 Postmus PE, Berendsen HH, van Zandwijk N, et al. Retreatment with the induction regimen in small cell lung cancer relapsing after an initial response to short term chemotherapy. *Eur J Cancer Clin Oncol* 1987;23:1409-1411.
- 16 Giaccone G, Ferrati P, Donadio M, et al. Reinduction chemotherapy in small cell lung cancer. *Eur J Cancer Clin Oncol* 1987;23:1697-1699.
- 17 Baize N, Monnet I, Greillier L, et al. Carboplatin plus etoposide versus topotecan as second-line treatment for patients with sensitive relapsed small-cell lung cancer: an open-label, multicentre, randomised, phase 3 trial. *Lancet Oncol* 2020;21:1224-1233.
- 18 Naito Y, Yamada K, Imamura Y, et al. Rechallenge treatment with a platinum-based regimen in patients with sensitive relapsed small-cell lung cancer. *Med Oncol* 2018;35:61.
- 19 Genestreti G, Tiseo M, Kenmotsu H, et al. Outcomes of platinum-sensitive small-cell lung cancer patients treated with platinum/etoposide rechallenge: a multi-institutional retrospective analysis. *Clin Lung Cancer* 2015;16:e223-e228.
- 20 Trigo J, Subbiah V, Besse B, et al. Lurbinectedin as second-line treatment for patients with small-cell lung cancer: a single-arm, open-label, phase 2 basket trial. *Lancet Oncol* 2020;21:645-654.
- 21 Subbiah V, Paz-Ares L, Besse B, et al. Antitumor activity of lurbinectedin in second-line small cell lung cancer patients who are candidates for re-challenge with the first-line treatment. *Lung Cancer* 2020;150:90-96.
- 22 von Pawel J, Schiller JH, Shepherd FA, et al. Topotecan versus cyclophosphamide, doxorubicin, and vincristine for the treatment of recurrent small-cell lung cancer. *J Clin Oncol* 1999;17:658-667.
- 23 O'Brien ME, Ciuleanu TE, Tsekov H, et al. Phase III trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J Clin Oncol* 2006;24:5441-5447.
- 24 Eckardt JR, von Pawel J, Pujol JL, et al. Phase III study of oral compared with intravenous topotecan as second-line therapy in small-cell lung cancer. *J Clin Oncol* 2007;25:2086-2092.
- 25 Edelman MJ, Dvorkin M, Laktionov K, et al. Randomized phase 3 study of the anti-disialoganglioside antibody dinutuximab and irinotecan vs irinotecan or topotecan for second-line treatment of small cell lung cancer. *Lung Cancer* 2022;166:135-142.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY – REFERENCES

- 26 Masuda N, Fukuoka M, Kusunoki Y, et al. CPT-11: a new derivative of camptothecin for the treatment of refractory or relapsed small-cell lung cancer. *J Clin Oncol* 1992;10:1225-1229.
- 27 Ahn MJ, Cho BC, Felip E, et al. Tarlatamab for patients with previously treated small-cell lung cancer. *N Engl J Med* 2023;389:2063-2075.
- 28 Antonia SJ, López-Martin JA, Bendell J, et al. Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (Checkmate 032): a multicentre, open-label phase 1/2 trial. *Lancet Oncol* 2016;17:883-895.
- 29 Ready NE, Ott PA, Hellmann MD, et al. Nivolumab monotherapy and nivolumab plus ipilimumab in recurrent small cell lung cancer: results from the CheckMate 032 randomized cohort. *J Thorac Oncol* 2020;15:426-435.
- 30 Chung HC, Piha-Paul SA, Lopez-Martin J, et al. Pembrolizumab After Two or More Lines of Previous Therapy in Patients With Recurrent or Metastatic SCLC: Results From the KEYNOTE-028 and KEYNOTE-158 Studies. *J Thorac Oncol* 2020;15:618-627.
- 31 Ott PA, Elez E, Hirt S, et al. Pembrolizumab in patients with extensive stage small-cell lung cancer: results from the phase Ib KEYNOTE-028 study. *J Clin Oncol* 2017;35:3823-3829.
- 32 Smit EF, Fokkema E, Biesma B, et al. A phase II study of paclitaxel in heavily pretreated patients with small-cell lung cancer. *Br J Cancer* 1998;77:347-351.
- 33 Yamamoto N, Tsurutani J, Yoshimura N, et al. Phase II study of weekly paclitaxel for relapsed and refractory small cell lung cancer. *Anticancer Res* 2006;26:777-781.
- 34 Pietanza MC, Kadota K, Huberman K, et al. Phase II trial of temozolomide with relapsed sensitive or refractory small cell lung cancer, with assessment of methylguanine-DNA methyltransferase as a potential biomarker. *Clin Cancer Res* 2012;18:1138-1145.
- 35 Zauderer MG, Drilon A, Kadota K, et al. Trial of a 5-day dosing regimen of temozolomide in patients with relapsed small cell lung cancers with assessment of methylguanine-DNA methyltransferase. *Lung Cancer* 2014;86:237-240.
- 36 Smyth JF, Smith IE, Sessa C, et al. Activity of docetaxel (Taxotere) in small cell lung cancer. The Early Clinical Trials Group of the EORTC. *Eur J Cancer* 1994;30A:1058-1060.
- 37 Masters GA, Declerck L, Blanke C, et al. Phase II trial of gemcitabine in refractory or relapsed small-cell lung cancer: Eastern Cooperative Oncology Group Trial 1597. *J Clin Oncol* 2003;21:1550-1555.
- 38 Van der Lee I, Smit EF, van Putten JW, et al. Single-agent gemcitabine in patients with resistant small-cell lung cancer. *Ann Oncol* 2001;12:557-561.
- 39 Hoang T, Kim K, Jaslawski A, et al. Phase II study of second-line gemcitabine in sensitive or refractory small cell lung cancer. *Lung Cancer* 2003;42:97-102.
- 40 Einhorn LH, Pennington K, McClean J. Phase II trial of daily oral VP-16 in refractory small cell lung cancer: a Hoosier Oncology Group study. *Semin Oncol* 1990;17:32-35.
- 41 Johnson DH, Greco FA, Strupp J, et al. Prolonged administration of oral etoposide in patients with relapsed or refractory small-cell lung cancer: a phase II trial. *J Clin Oncol* 1990;8:1613-1617.
- 42 Marcoux N, Gettinger SN, O'Kane G, et al. EGFR-mutant adenocarcinomas that transform to small-cell lung cancer and other neuroendocrine carcinomas: Clinical outcomes. *J Clin Oncol* 2019;37:278-285.
- 43 Ferrer L, Gaj Levra M, Brevet M, et al. A brief report of transformation from NSCLC to SCLC: Molecular and therapeutic characteristics. *J Thorac Oncol* 2019;14:130-134.
- 44 Chai X, Zhang X, Li W, Chai J. Small cell lung cancer transformation during antitumor therapies: A systematic review. *Open Med (Wars)* 2021;16:1160-1167.
- 45 Zhang CY, Sun H, Su JW, et al. A potential treatment option for transformed small-cell lung cancer on PD-L1 inhibitor-based combination therapy improved survival. *Lung Cancer* 2023;175:68-78.
- 46 Schoenfeld AJ, Arbour KC, Rizvi H, et al. Severe immune-related adverse events are common with sequential PD-(L)1 blockade and osimertinib. *Ann Oncol* 2019;30:839-844.
- 47 Oshima Y, Tanimoto T, Yuji K, Tojo A. EGFR-TKI-associated interstitial pneumonitis in nivolumab-treated patients with non-small cell lung cancer *JAMA Oncol* 2018;4:1112-1115.
- 48 Paz-Ares L, Borghaei H, Liu SV, et al. Efficacy and safety of first-line maintenance therapy with lurbinectedin plus atezolizumab in extensive-stage small-cell lung cancer (IMforte): a randomised, multicentre, open-label, phase 3 trial. *Lancet* 2025;405:2129-2143.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2024

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY

General Principles:

- General principles of RT for lung cancer—including commonly used abbreviations; standards for clinical and technologic expertise and quality assurance; and principles of RT simulation, planning, and delivery—are provided in the [NCCN Guidelines for Non-Small Cell Lung Cancer \(NSCL-C\)](#) and are applicable to RT for SCLC.
- RT has a potential role in all stages of SCLC, as part of either definitive or palliative therapy. Radiation oncology input, as part of a multidisciplinary evaluation or discussion, should be provided for all patients early in the determination of the treatment strategy.
- To maximize tumor control and to minimize treatment toxicity, critical components of modern RT include appropriate simulation, accurate target definition, conformal RT (CRT) planning, and ensuring accurate delivery of the planned treatment. A minimum standard is CT-planned 3D-CRT. Multiple fields should be used, with all fields treated daily.
- Use of more advanced technologies is appropriate when needed to deliver adequate tumor doses while respecting normal tissue dose constraints. Such technologies include (but are not limited to) 4D-CT and/or FDG-PET/CT simulation, intensity-modulated RT (IMRT)/volumetric modulated arc therapy (VMAT), image-guided RT (IGRT), and motion management strategies. IMRT is preferred over 3D-CRT on the basis of reduced toxicity in the setting of concurrent chemotherapy/RT.¹ Quality assurance measures are essential and are covered in the [NCCN Guidelines for Non-Small Cell Lung Cancer \(NSCL-C\)](#).
- Useful references include the ASTRO Guidelines and the American Radium Society.²⁻⁴

General Treatment Information:

Limited Stage:

- In patients with clinical stage I–IIA (T1–2,N0,M0) who have undergone lobectomy and are found to have regional nodal involvement on final pathology, postoperative RT is recommended in pathologic N2⁵ and may be considered in pathologic N1 stage, either sequentially or concurrently with chemotherapy. Principles of postoperative RT for NSCLC, including target volumes and doses, are recommended.
- Selected patients with stage I–IIA (T1–2,N0,M0) SCLC who are medically inoperable or in whom a decision is made not to pursue surgery may be candidates for stereotactic ablative radiotherapy (SABR), also known as stereotactic body RT (SBRT), to the primary tumor followed by adjuvant systemic therapy. Principles of SABR for SCLC are similar to those for NSCLC (see [NCCN Guidelines for Non-Small Cell Lung Cancer: NSCL-C](#)).⁶⁻⁸
- Timing: RT concurrent with systemic therapy is standard and preferred to sequential chemo/RT.⁹ RT should start early, with cycle 1 or 2 of systemic therapy (category 1).¹⁰ A shorter time from the start of any therapy to the end of RT (SER) is significantly associated with improved survival.¹¹
- Target definition: RT target volumes should be defined based on the pretreatment FDG-PET scan and CT scan obtained at the time of RT planning, as well as any positive biopsies. FDG-PET/CT is recommended, preferably within 4 weeks and no more than 8 weeks, before treatment. Ideally, FDG-PET/CT should be obtained in the treatment position.

Limited Stage (continued), Extensive Stage ([SCL-F 2 of 7](#))

Normal Tissue Dose Constraints, Prophylactic Cranial Irradiation ([SCL-F 3 of 7](#))

Brain Metastasis ([SCL-F 4 of 7](#))

[Continued](#)
[References on](#)
[\(SCL-F 5 of 7\)](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY

Limited Stage (continued):

- Historically, clinically uninvolved mediastinal nodes have been included in the RT target volume, whereas uninvolved supraclavicular nodes generally have not been included. Several more modern series, both retrospective and prospective, suggest that omission of elective nodal irradiation (ENI) results in low rates of isolated nodal recurrences (0%–11%, most <5%), particularly when incorporating FDG-PET staging/target definition (1.7%–3%).^{12–17} ENI has been omitted in recent prospective clinical trials (including CALGB 30610/RTOG 0538 and the EORTC 08072 [CONVERT] trial). Inclusion of the ipsilateral hilum in the target volume, even if not grossly involved, differs between these trials but may be reasonable.
- In patients who start systemic therapy before RT, the gross tumor volume (GTV) can be limited to the post-induction systemic therapy volume to avoid excessive toxicity. Initially involved nodal regions (but not their entire pre-systemic therapy volume) should be covered.^{14,18}
- Dose and schedule: For limited-stage SCLC, the optimal dose and schedule of RT have not been established.
 - ▶ Based on the randomized phase III trial, INT 0096, 45 Gy in 3 weeks (1.5 Gy twice daily [BID]) is superior (category 1) to 45 Gy in 5 weeks (1.8 Gy daily).^{19,20} When BID fractionation is used, there should be at least a 6-hour interfraction interval to allow for repair of normal tissue.
 - ▶ Retrospective and randomized phase II studies from Norway and Canada suggest that similarly accelerated doses of 40–42 Gy in 3 weeks but given in once-daily fractionation produce similar outcomes as 45 Gy in 3 weeks in BID fractionation, though regional practice between daily and twice daily fractionation has diverged between those countries after subsequent experience.^{21–23}
 - ▶ If using once-daily conventionally fractionated RT, higher doses of 66–70 Gy are preferred.^{24–27} Two randomized phase III trials did not demonstrate superiority of 66 Gy in 6.5 weeks/2 Gy daily (the European CONVERT trial) or 70 Gy in 7 weeks/2 Gy daily (CALGB 30610/RTOG 0538) over 45 Gy in 3 weeks/1.5 Gy BID, but overall survival and toxicity were similar.^{28–30}
 - ▶ Recent randomized phase II trials suggest that higher dose accelerated RT of 60–65 Gy in 4–5 weeks given in BID or daily fractionation may produce increased overall or progression-free survival compared to 45 Gy in 3 weeks in BID fractionation.^{31,32} Another randomized phase 3 trial demonstrated a survival advantage of high-dose accelerated RT given as 54 Gy simultaneous integrated boost (SIB) to the gross tumor volume in 30 BID fractions over 3 weeks compared to standard 45 Gy in 30 BID fractions over 3 weeks. These studies indicate an advantage of higher dose accelerated RT with concurrent chemotherapy.^{33,34}
 - ▶ Adjuvant immunotherapy with durvalumab in patients who have not progressed after the completion of chemoRT has been shown to provide a significant survival advantage. The optimal dose and schedule of RT in the context of immunotherapy is currently undefined.³⁴

Extensive Stage:

- Consolidative thoracic RT is beneficial for selected patients with ES-SCLC with complete response or good response to systemic therapy before immunotherapy, especially with residual thoracic disease and low-bulk extrathoracic metastatic disease. Studies have demonstrated that consolidative thoracic RT up to definitive doses is well-tolerated, results in fewer symptomatic chest recurrences, and improves long-term survival in some patients.^{35,36} The Dutch CREST randomized trial of modest-dose thoracic RT (30 Gy in 10 fractions) in patients with ES-SCLC that responded to chemotherapy (without immunotherapy) demonstrated significantly improved 2-year overall survival and 6-month progression-free survival, although the protocol-defined primary endpoint of 1-year overall survival was not significantly improved.³⁷ Subsequent exploratory analysis found the benefit of consolidative thoracic RT is limited to the majority of patients who had residual thoracic disease after systemic therapy.³⁸
- Dosing and fractionation of consolidative thoracic RT should be individualized within the range of 30 Gy in 10 daily fractions up to definitive dosing regimens in patients with a longer life expectancy.

Normal Tissue Dose Constraints, Prophylactic Cranial Irradiation ([SCL-F 3 of 7](#))

Brain Metastasis ([SCL-F 4 of 7](#))

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.

[Continued](#)

[References on](#)
[\(SCL-F 5 of 7\)](#)

SCL-F
2 OF 7



PRINCIPLES OF RADIATION THERAPY

Extensive Stage (continued):

Based on two randomized trials, immunotherapy during and after chemotherapy is a first-line approach,^{39,40} but these studies did not include consolidative thoracic RT. Nevertheless, consolidative thoracic RT after chemoimmunotherapy can be considered for selected patients as above, during or before maintenance immunotherapy (there are limited data on optimal sequencing or safety). The benefit of thoracic RT in the context of chemoimmunotherapy is under evaluation in the [RAPTOR/NRG LU007 trial](#).

Normal Tissue Dose Constraints:

- Normal tissue dose constraints depend on tumor size and location. For similar RT prescription doses, the normal tissue constraints used for NSCLC are appropriate (see [NSCLC-C](#)).
- When administering accelerated RT schedules (eg, BID) or lower total RT doses (eg, 45 Gy), more conservative constraints should be used. When using accelerated schedules (eg, 3–5 weeks), the spinal cord constraints from the CALGB 30610/RTOG 0538 protocol should be used as a guide: ie, the maximum spinal cord dose should be limited to ≤41 Gy (including scatter irradiation) for a prescription of 45 Gy BID in 3 weeks and limited to ≤50 Gy for more protracted schedules.

Prophylactic Cranial Irradiation:

- In patients with limited stage SCLC (LS-SCLC) who have a good response to initial therapy, PCI decreases brain metastases and increased overall survival^{41,42} in meta-analyses of past clinical trials. Of note, none of the past studies that have been used as the basis for PCI recommendations in LS-SCLC used MRI staging of the brain nor did any utilize FDG-PET scans for overall staging.
- The benefit of PCI is unclear in patients who have undergone definitive therapy for very early LS-SCLC, ie, pathologic stage I–IIA (T1–2,N0,M0).⁴³ These patients have a lower risk of developing brain metastases than patients with more advanced, LS-SCLC and may not benefit from PCI.⁴³ Brain MRI surveillance is recommended in patients not receiving PCI.⁴¹ However, PCI may have a benefit in patients who are found to have pathologic stage IIB or III SCLC after complete resection.^{42,43} This issue is being evaluated in the ongoing NCI cooperative group trial SWOG S1827/MAVERICK (brain MRI surveillance ± PCI), which includes the population undergoing surgical resection (<https://clinicaltrials.gov/ct2/show/NCT04155034>). In patients with ES-SCLC that have responded to systemic therapy, PCI decreases brain metastases. A randomized trial conducted by the European Organisation for Research and Treatment of Cancer (EORTC) found improved overall survival with PCI.⁴⁴ However, a Japanese randomized trial found that in patients who had no brain metastases on baseline MRI, PCI did not improve overall survival compared with routine surveillance MRI and treatment of asymptomatic brain metastases upon detection.⁴⁵ Surveillance imaging for brain metastases is recommended for all patients regardless of PCI status.
- The preferred dose for PCI to the whole brain is 25 Gy in 10 daily fractions. A shorter course (eg, 20 Gy in 5 fractions) may be appropriate in selected patients with extensive stage disease. In a large randomized trial (PCI 99-01), patients receiving a dose of 36 Gy had higher mortality and higher chronic neurotoxicity compared to patients treated with 25 Gy.^{46,47}
- Neurocognitive function: Increasing age and higher doses are the most predictive factors for development of chronic neurotoxicity. In trial RTOG 0212, 83% of patients >60 years experienced chronic neurotoxicity 12 months after PCI versus 56% of patients <60 years ($P = .009$).⁴⁷ PCI is not recommended in patients with poor PS or impaired neurocognitive function.⁴⁶ The role of PCI in MRI and FDG-PET staged SCLC in patients who are fit with normal neurocognitive function is the subject of ongoing debate, particularly in limited stage, and is being evaluated in the phase III SWOG S1827/MAVERICK trial comparing PCI (active comparator) to MRI surveillance (experimental) in both limited and extensive stage (<https://clinicaltrials.gov/ct2/show/NCT04155034>).

Brain Metastasis ([SCL-F 4 of 7](#))

[Continued](#)
[References on](#)
[\(SCL-F 5 of 7\)](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY

Prophylactic Cranial Irradiation (continued):

- Administer PCI after resolution of acute toxicities of initial therapy. PCI is not recommended in patients with poor PS or impaired neurocognitive functioning.
- When administering PCI, consider adding memantine during and after RT, which has been shown to decrease neurocognitive impairment following whole brain radiation therapy (WBRT) for brain metastases.⁴⁷ The dose of memantine used on RTOG 0614 was as follows: week 1 (starting on day 1 of WBRT), 5 mg each morning; week 2, 5 mg each morning and evening; week 3, 10 mg each morning and 5 mg each evening; and weeks 4–24, 10 mg each morning and evening (see [NCCN Guidelines for Central Nervous System Cancers](#)). Hippocampal-avoidance (HA) PCI using IMRT may be considered as a potential strategy to improve cognitive preservation. A phase III randomized trial of HA-WBRT versus conventional WBRT demonstrated improved cognitive preservation and patient-reported outcomes with HA-WBRT in patients with brain metastases from mixed histologies.⁴⁸ Conflicting data have been reported with HA-PCI versus conventional PCI in SCLC with one trial reporting no differences in cognition⁴⁹ and a separate trial reporting improved cognitive preservation with HA-PCI.⁵⁰ A larger randomized trial of HA-PCI versus conventional PCI, [NRG CC003](#),⁵¹ has completed accrual with results pending.⁵²
- An ongoing randomized trial, SWOG S1827/MAVERICK, is evaluating whether brain MRI surveillance alone is noninferior to MRI surveillance plus PCI with regard to overall survival for LS-SCLC and ES-SCLC.⁵³

Brain Metastases:

- Brain metastases have conventionally been treated with WBRT; however, selected patients with a small number of metastases may be appropriately treated with stereotactic radiotherapy (SRT)/radiosurgery (SRS).^{55,56,57} A current randomized trial, [NRG CC009](#), is comparing SRS to hippocampal-sparing WBRT plus memantine in this setting.
- Recommended dose for WBRT is 30 Gy in 10 daily fractions. Consider adding memantine during and after RT (see Prophylactic Cranial Irradiation for memantine dosing).⁵⁵
- In patients who develop brain metastases after PCI, repeat WBRT may be considered in carefully selected patients.^{56,57} SRS is preferred, if feasible.^{58,59}
- For patients with a better prognosis (eg, ≥ 4 months), hippocampal-sparing WBRT using IMRT plus memantine is preferred because it produces less cognitive function failure than conventional WBRT plus memantine.⁴⁸ However, patients with metastases within 5 mm of the hippocampi, leptomeningeal metastases, and other high-risk features were not eligible for hippocampal-sparing WBRT on [NRG CC001](#).⁴⁸ Although CC001 did not include patients with brain metastases from SCLC, it is reasonable to extrapolate the findings to SCLC.

Palliative Radiation for Extracranial Metastases:

- Common radiation dose-fractionation regimens (eg, 30 Gy in 10 fractions, 20 Gy in 5 fractions, 8 Gy in 1 fraction) used for palliation of other solid tumors are appropriate for palliation of SCLC metastases in most patients.
- Conformal techniques, such as IMRT, and/or higher dose intensity approaches, including SABR or SRS, may be appropriate in selected patients (eg, tumors with close proximity to organs at risk, reirradiation, or better prognosis).

[References on
\(SCL-F 5 of 7\)](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY – REFERENCES

- ¹ Chun SG, Hu C, Choy H, et al. Impact of intensity-modulated radiation therapy technique for locally advanced non-small-cell lung cancer: a secondary analysis of the NRG Oncology RTOG 0617 randomized clinical trial. *J Clin Oncol* 2017;35:56-62.
- ² Simone CB 2nd, Bogart JA, Cabrera AR, et al. Radiation Therapy for Small Cell Lung Cancer: An ASTRO Clinical Practice Guideline. *Pract Radiat Oncol* 2020;10:158-173.
- ³ Chun SG, Simone CB 2nd, Amini A, et al. American Radium Society Appropriate Use Criteria: Radiation Therapy for Limited-Stage SCLC 2020. *J Thorac Oncol* 2021;16:66-75.
- ⁴ Higgins KA, Simone CB 2nd, Amini A, et al. American Radium Society Appropriate Use Criteria on Radiation Therapy for Extensive-Stage SCLC. *J Thorac Oncol* 2021;16:54-65.
- ⁵ Zhang C, Zhao G, Wu H, et al. Application of postoperative adjuvant radiotherapy in limited stage small cell lung cancer: A systematic review and meta-analysis. *Radiother Oncol* 2024;193:110123.
- ⁶ Shioyama Y, Onishi H, Takayama K, et al. Clinical outcomes of stereotactic body radiotherapy for patients with stage I small-cell lung cancer: Analysis of a subset of the Japanese Radiological Society Multi-Institutional SBRT Study Group Database. *Technol Cancer Res Treat* 2018;17:1533033818783904.
- ⁷ Verma V, Simone CB 2nd, Allen PK, Lin SH. Outcomes of stereotactic body radiotherapy for T1-T2N0 small cell carcinoma according to addition of chemotherapy and prophylactic cranial irradiation: a multicenter analysis. *Clin Lung Cancer* 2017;18:675-681.e1.
- ⁸ Verma V, Simone CB 2nd, Allen PK, et al. Multi-institutional experience of stereotactic ablative radiation therapy for stage I small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2017;97:362-371.
- ⁹ Takada M, Fukuoka M, Kawahara M, et al. Phase III study of concurrent versus sequential thoracic radiotherapy in combination with cisplatin and etoposide for limited stage small-cell lung cancer: results of the Japan Clinical Oncology Group Study 9104. *J Clin Oncol* 2002;20:3054-3060.
- ¹⁰ Fried DB, Morris DE, Poole C, et al. Systematic review evaluating the timing of thoracic radiation therapy in combined modality therapy for limited stage small-cell lung cancer. *J Clin Oncol* 2004;22:4837-4845.
- ¹¹ De Ruyscher D, Pijls-Johannesma M, Bentzen SM, et al. Time between the first day of chemotherapy and the last day of chest radiation is the most important predictor of survival in limited-disease small-cell lung cancer. *J Clin Oncol* 2006;24:1057-1063.
- ¹² De Ruyscher D, Bremer RH, Koppe F, et al. Omission of elective node irradiation on basis of CT-scans in patients with limited disease small cell lung cancer: a phase II trial. *Radiother Oncol* 2006;80:307-312.
- ¹³ van Loon J, De Ruyscher D, Wanders R, et al. Selective nodal irradiation on basis of (18)FDG-PET scans in limited-disease small-cell lung cancer: a prospective study. *Int J Radiat Oncol Biol Phys* 2010;77:329-336.
- ¹⁴ Hu X, Bao Y, Xu YJ, et al. Final report of a prospective randomized study on thoracic radiotherapy target volume for limited stage small cell lung cancer with radiation dosimetric analyses. *Cancer* 2020;126:840-849.
- ¹⁵ Shirvani SM, Komaki R, Heymach JV, et al. Positron emission tomography/computed tomography-guided intensity-modulated radiotherapy for limited stage small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2012;82:e91-97.
- ¹⁶ Xia B, Chen GY, Cai XW, et al. Is involved-field radiotherapy based on CT safe for patients with limited stage small-cell lung cancer? *Radiother Oncol* 2012;102:258-262.
- ¹⁷ Colaco R, Sheikh H, Lorigan P, et al. Omitting elective nodal irradiation during thoracic irradiation in limited stage small cell lung cancer - Evidence from a phase II trial. *Lung Cancer* 2012;76:72-77.
- ¹⁸ Liengswangwong V, Bonner JA, Shaw EG, et al. Limited stage small-cell lung cancer: patterns of intrathoracic recurrence and the implications for thoracic radiotherapy. *J Clin Oncol* 1994;12:496-502.
- ¹⁹ Turrisi AT 3rd, Kim K, Blum R, et al. Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with cisplatin and etoposide. *N Engl J Med* 1999;340:265-271.
- ²⁰ Schild SE, Bonner JA, Shanahan TG, et al. Long-term results of a phase III trial comparing once-daily radiotherapy with twice-daily radiotherapy in limited stage small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2004;59:943-951.
- ²¹ Grønberg BH, Halvorsen TO, Fløtten Ø, et al. Randomized phase II trial comparing twice daily hyperfractionated with once daily hypofractionated thoracic radiotherapy in limited disease small cell lung cancer. *Acta Oncol* 2016;55:591-597.
- ²² Turgeon GA, Souhami L, Kopek N, et al. Thoracic irradiation in 3 weeks for limited stage small cell lung cancer: Is twice a day fractionation really needed? *Cancer Radiother* 2017;21:89-98.
- ²³ Graabak G, Grønberg BH, Sandvei MS, et al. Thoracic Radiotherapy in Limited stage SCLC-a Population-Based Study of Patterns of Care in Norway From 2000 Until 2018 *JTO Clin Res Rep* 2022;3:100270.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY – REFERENCES

- 24 Choi NC, Herndon JE 2nd, Rosenman J, et al. Phase I study to determine the maximum-tolerated dose of radiation in standard daily and hyperfractionated-accelerated twice-daily radiation schedules with concurrent chemotherapy for limited stage small-cell lung cancer. *J Clin Oncol* 1998;16:3528-3536.
- 25 Miller KL, Marks LB, Sibley GS, et al. Routine use of approximately 60 Gy once-daily thoracic irradiation for patients with limited stage small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2003;56:355-359.
- 26 Roof KS, Fidias P, Lynch TJ, et al. Radiation dose escalation in limited stage small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2003;57:701-708.
- 27 Bogart JA, Herndon JE, Lyss AP, et al. 70 Gy thoracic radiotherapy is feasible concurrent with chemotherapy for limited stage small-cell lung cancer: analysis of Cancer and Leukemia Group B study 39808. *Int J Radiat Oncol Biol Phys* 2004;59:460-468.
- 28 Faivre-Finn C, Snee M, Ashcroft L, et al. Concurrent once-daily versus twice-daily chemoradiotherapy in patients with limited stage small-cell lung cancer (CONVERT): an open-label, phase 3, randomised, superiority trial. *Lancet Oncol* 2017;18:1116-1125.
- 29 Bogart JA, Wang XF, Masters GA, et al. High-dose once-daily thoracic radiotherapy in limited stage small-cell lung cancer: CALGB 30610 (Alliance)/RTOG 0538. *J Clin Oncol* 2023;41:2394-2402.
- 30 Ganti A, Dueck AC, Fruth B, et al. Comparison of quality of life in patients randomized to high-dose once daily (QD) thoracic radiotherapy (TRT) with standard twice daily (BID) TRT in limited stage small cell lung cancer (LS-SCLC) on CALGB 30610 (Alliance, Sub-study CALGB 70702) [abstract]. *J Clin Oncol* 2022;40:8504.
- 31 Grønberg BH, Killingberg KT, Fløtten Ø, et al. High-dose versus standard dose twice-daily thoracic radiotherapy in limited stage small-cell lung cancer: final survival data, long-term toxicity and relapse patterns in a randomised, open-label, phase II trial. *J Thorac Oncol* 2025;S1556-0864;00696-3.
- 32 Qiu B, Li QW, Liu JL, et al. Moderately hypofractionated once-daily compared with twice-daily thoracic radiation therapy concurrently with etoposide and cisplatin in limited stage small-cell lung cancer: a multi-center, Phase II, randomized trial. *Int J Radiat Oncol Biol Phys* 2021;111:424-435.
- 33 Yu J, Jiang L, Zhao L, et al. High-dose hyperfractionated simultaneous integrated boost radiotherapy versus standard-dose radiotherapy for limited-stage small-cell lung cancer in China: a multicentre, open-label, randomised, phase 3 trial. *Lancet Respir Med* 2024;12:799-809.
- 34 Cheng Y, Spigel DR, Cho BC, et al; ADRIATIC Investigators. Durvalumab after chemoradiotherapy in limited-stage small-cell lung cancer. *N Engl J Med* 2024;391:1313-1327.
- 35 Jeremic B, Shibamoto Y, Nikolic N, et al. Role of radiation therapy in the combined-modality treatment of patients with extensive disease small-cell lung cancer: A randomized study. *J Clin Oncol* 1999;17:2092-2099.
- 36 Yee D, Butts C, Reiman A, et al. Clinical trial of post-chemotherapy consolidation thoracic radiotherapy for extensive stage small cell lung cancer. *Radiother Oncol* 2012;102:234-238.
- 37 Slotman BJ, van Tinteren H, Praag JO, et al. Use of thoracic radiotherapy for extensive stage small-cell lung cancer: a phase 3 randomised controlled trial. *Lancet* 2015;385:36-42.
- 38 Slotman BJ, van Tinteren H, Praag JO, et al. Radiotherapy for extensive stage small-cell lung cancer-Authors' reply. *Lancet* 2015;385:1292-1293.
- 39 Horn L, Mansfield A, Szczesna A, et al. First-line atezolizumab plus chemotherapy in extensive stage small-cell lung cancer. *N Engl J Med* 2018;379:2220-2229.
- 40 Paz-Ares L, Dvorkin M, Chen Y, et al. Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *Lancet* 2019;394:1929-1939.
- 41 Arriagada R, Le Chevalier T, Rivière A, et al. Patterns of failure after prophylactic cranial irradiation in small-cell lung cancer: analysis of 505 randomized patients. *Ann Oncol* 2002;13:748-754.
- 42 Aupérin A, Arriagada R, Pignon JP, et al. Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission. Prophylactic Cranial Irradiation Overview Collaborative Group. *N Engl J Med* 1999;341:476-484.
- 43 Yang Y, Zhang D, Zhou X, et al. Prophylactic cranial irradiation in resected small cell lung cancer: A systematic review with meta-analysis. *J Cancer* 2018;9:433-439.
- 44 Slotman B, Faivre-Finn C, Kramer G, et al. Prophylactic cranial irradiation in extensive small-cell lung cancer. *N Engl J Med* 2007;357:664-672.
- 45 Takahashi T, Yamanaka T, Seto T, et al. Prophylactic cranial irradiation versus observation in patients with extensive-disease small-cell lung cancer: a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2017;18:663-671.
- 46 Le Péchoux C, Dunant A, Senan S, et al. Standard-dose versus higher-dose prophylactic cranial irradiation (PCI) in patients with limited stage small-cell lung cancer in complete remission after chemotherapy and thoracic radiotherapy (PCI 99-01, EORTC 22003-08004, RTOG 0212, and IFCT 99-01): a randomised clinical trial. *Lancet Oncol* 2009;10:467-474.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY – REFERENCES

- ⁴⁷ Wolfson AH, Bae K, Komaki R, et al. Primary analysis of a phase II randomized trial Radiation Therapy Oncology Group (RTOG) 0212: Impact of different total doses and schedules of prophylactic cranial irradiation on chronic neurotoxicity and quality of life for patients with limited-disease small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2011;81:77-84.
- ⁴⁸ Brown P, Gondi V, Pugh S, et al. Hippocampal avoidance during whole-brain radiotherapy plus memantine for patients with brain metastases: Phase III trial NRG Oncology CC001. *J Clin Oncol* 2020;38:1019-1029.
- ⁴⁹ Belderbos JSA, De Ruyscher DKM, De Jaeger K, et al. Phase 3 randomized trial of prophylactic cranial irradiation with or without hippocampus avoidance in SCLC (NCT01780675). *J Thorac Oncol* 2021;16:840-849.
- ⁵⁰ Rodriguez De Dios N, Murcia M, Counago F, et al. Phase III trial of prophylactic cranial irradiation with or without hippocampal avoidance for small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2019;105:S35-S36.
- ⁵¹ Gondi V, Pugh S, Mehta MP, et al. Primary Endpoint Results of NRG CC003: Phase IIR/III Trial of Prophylactic Cranial Irradiation (PCI) with or without Hippocampal Avoidance (HA) for Small Cell Lung Cancer (SCLC) [abstract 2023]. *Int J Radiat Oncol Biol Phys* 2023;117:E3.
- ⁵² Gondi V, Pugh SL, Mehta MP, et al. NRG Oncology CC003: A randomized phase II/III trial of prophylactic cranial irradiation with or without hippocampal avoidance for small cell lung cancer. *J Clin Oncol* 2019;37:TPS 8578-TPS 8578.
- ⁵³ SWOG S1827 (MAVERICK) Testing whether the use of brain scans alone instead of brain scans plus preventive brain radiation affects lifespan in patients with small cell lung cancer. Available at: <https://clinicaltrials.gov/ct2/show/NCT04155034>.
- ⁵⁴ Rusthoven CG, Yamamoto M, Bernhardt D, et al. Evaluation of first-line radiosurgery vs whole-brain radiotherapy for small cell lung cancer brain metastases: the FIRE-SCLC cohort study. *JAMA Oncol* 2020;6:1028-1037.
- ⁵⁵ Brown PD, Pugh S, Laack NN, et al. Memantine for the prevention of cognitive dysfunction in patients receiving whole-brain radiotherapy: a randomized, double-blind, placebo-controlled trial. *Neuro Oncol* 2013;10:1429-1437.
- ⁵⁶ Sadikov E, Bezjak A, Yi QL, et al. Value of whole brain re-irradiation for brain metastases--single centre experience. *Clin Oncol (R Coll Radiol)* 2007;19:532-538.
- ⁵⁷ Son CH, Jimenez R, Niemierko A, et al. Outcomes after whole brain reirradiation in patients with brain metastases. *Int J Radiat Oncol Biol Phys* 2012;82:e167-172.
- ⁵⁸ Harris S, Chan MD, Lovato JF, et al. Gamma knife stereotactic radiosurgery as salvage therapy after failure of whole-brain radiotherapy in patients with small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2012;83:e53-e59.
- ⁵⁹ Wegner RE, Olson AC, Kondziolka D, et al. Stereotactic radiosurgery for patients with brain metastases from small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2011;81:e21-e27.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

PRINCIPLES OF IMAGING

General Principles:

- Both CT and MRI are performed with contrast, unless clinically contraindicated.
- Workup of SCLC should be expedited, with studies done in parallel whenever possible.
- If extensive stage is established, brain imaging MRI (preferred) or CT with contrast is recommended. Further evaluation is dependent on the clinical situation.
- Brain MRI is more sensitive than CT for identifying brain metastases and is preferred over CT.

Workup:

- Chest/abdomen/pelvis (C/A/P) CT with contrast
- Brain MRI (preferred) or CT with contrast
- FDG-PET/CT scan (skull base to mid-thigh), if needed to clarify extent of disease

Additional Workup:

- Bone imaging (radiographs or MRI) as appropriate, if FDG-PET/CT equivocal

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF IMAGING

Recommendations for scanning are based on patients without altered symptoms or with stable symptoms

Treatment Response Assessment

• Limited Stage

- ▶ For patients receiving adjuvant systemic therapy ± RT, response assessment is recommended only after completion of adjuvant therapy ([SCL-6](#)).
- ▶ Response assessment after adjuvant therapy involves C/A/P CT with contrast and brain MRI (preferred) with contrast or brain CT with contrast ([SCL-6](#)).
- ▶ For patients receiving systemic therapy + concurrent RT, response assessment is recommended only after completion of initial therapy ([SCL-6](#)).
- ▶ Repeating scans to assess response during adjuvant or initial treatment is not recommended in the absence of new symptoms.
- ▶ For patients receiving systemic therapy alone or sequential systemic therapy followed by RT, response assessment by C/A/P CT with contrast is recommended after every 2 cycles of systemic therapy and at completion of therapy ([SCL-6](#)).

• Extensive Stage

- ▶ In known brain metastases, if systemic therapy was initiated prior to brain RT, brain MRI (preferred) or CT with contrast is recommended to be repeated after every 2 cycles of systemic therapy until brain RT is initiated ([SCL-6](#)).
- ▶ During systemic therapy, response assessment by C/A/P CT with contrast is recommended after every 2–3 cycles of systemic therapy and at completion of therapy. For patients with known brain metastases, brain MRI (preferred) or brain CT with contrast should be obtained every 3–4 months, or at a frequency based on clinical indication.

Follow-up/Surveillance

- Most NCCN Member Institutions use chest CT ± abdomen/pelvis every 2–6 months (more frequently in years 1–2 and less frequently thereafter).
- Brain MRI or CT with contrast every 3–4 months during year 1, then every 6 months in year 2, then after year 2, as clinically indicated (regardless of PCI status).
 - ▶ Surveillance for all patients consists of:
 - ◊ CT Chest ± abdomen/pelvis
 - ◊ Brain MRI (preferred) or brain CT
- Imaging for known metastases (eg, CT neck or MRI spine) should be repeated for follow-up. Other imaging studies may be obtained based on individual clinical scenarios.
- New pulmonary nodule should initiate workup for potential new primary.
- FDG-PET/CT is not recommended for routine follow-up unless a contrast CT or MRI is contraindicated.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF IMAGING - REFERENCES

- Jett JR, Schild SE, Kesler KA, Kalemkerian GP. Treatment of small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e400S-e419S.
- Kalemkerian GP. Staging and imaging of small cell lung cancer. Cancer Imaging 2012;11:253-258.
- Seute T, Leffers P, ten Velde GP, Twijnstra A. Detection of brain metastases from small cell lung cancer: consequences of changing imaging techniques (CT versus MRI). Cancer 2008;112:1827-1834.
- Kalemkerian GP, Gadgeel SM. Modern staging of small cell lung cancer. J Natl Compr Canc Netw 2013;11:99-104.
- Podoloff DA, Ball DW, Ben-Josef E, et al. NCCN task force: clinical utility of PET in a variety of tumor types. J Natl Compr Canc Netw 2009;7 Suppl 2:S1-26.
- Bradley JD, Dehdashti F, Mintun MA, et al. Positron emission tomography in limited stage small-cell lung cancer: a prospective study. J Clin Oncol 2004;22:3248-3254.
- Fischer BM, Mortensen J, Langer SW, et al. A prospective study of PET/CT in initial staging of small-cell lung cancer: comparison with CT, bone scintigraphy and bone marrow analysis. Ann Oncol 2007;18:338-345.
- Wolfson AH, Bae K, Komaki R, et al. Primary analysis of a phase II randomized trial Radiation Therapy Oncology Group (RTOG) 0212: impact of different total doses and schedules of prophylactic cranial irradiation on chronic neurotoxicity and quality of life for patients with limited-disease small-cell lung cancer. Int J Radiat Oncol Biol Phys 2011;81:77-84.
- Le Pechoux C, Laplanche A, Faivre-Finn C, et al. Clinical neurological outcome and quality of life among patients with limited small-cell cancer treated with two different doses of prophylactic cranial irradiation in the intergroup phase III trial (PCI99-01, EORTC 22003-08004, RTOG 0212 and IFCT 99-01). Ann Oncol 2011;22:1154-1163.
- Farooqi AS, Holliday EB, Allen PK, et al. Prophylactic cranial irradiation after definitive chemoradiotherapy for limited stage small cell lung cancer: Do all patients benefit? Radiother Oncol 2017;122:307-312.
- Lok BH, Ma J, Foster A, et al. Factors influencing the utilization of prophylactic cranial irradiation in patients with limited stage small cell lung cancer. Adv Radiat Oncol 2017;2:548-554.
- Manapov F, Klockung S, Niyazi M, et al. Timing of failure in limited disease (stage I-III) small-cell lung cancer patients treated with chemoradiotherapy: a retrospective analysis. Tumori 2013;99:656-660.
- Johnson BE, Linnoila RI, Williams JP, et al. Risk of second aerodigestive cancers increases in patients who survive free of small-cell lung cancer for more than 2 years. J Clin Oncol 1995;13:101-111.
- Johnson BE. Second lung cancers in patients after treatment for an initial lung cancer. J Natl Cancer Inst 1998;90:1335-1345.
- Takahashi T, Yamanaka T, Seto T, et al. Prophylactic cranial irradiation versus observation in patients with extensive-disease small-cell lung cancer: a multicentre, randomised, open-label, phase 3 trial. Lancet Oncol 2017;18:663-671.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).
All recommendations are category 2A unless otherwise indicated.

RECOMMENDED FIRST-LINE SYSTEMIC THERAPY REGIMENS

First-Line Therapy for Limited-Stage SCLC in Excess of T1 2, N0

	PS 0-2	PS 3-4
Cisplatin (75 mg)/etoposide (100 mg) + RT		
Cisplatin (60 mg)/etoposide (120 mg) + RT		
Carboplatin/etoposide + RT		
Cisplatin (75 mg)/etoposide (100 mg)	—	
Cisplatin 25 mg/m ² and etoposide 100 mg/m ²		—
Cisplatin (60 mg)/etoposide (120 mg)	—	
Carboplatin/etoposide	—	

Adjuvant therapy following primary therapy for CR/PR/Stable Limited Stage disease

Consolidation Durvalumab	
--------------------------	---

First-Line Therapy for Extensive Stage SCLC

Carboplatin/etoposide/atezolizumab	
Carboplatin/etoposide/durvalumab	
Cisplatin/etoposide/durvalumab	
Carboplatin/etoposide	
Cisplatin (75 mg)/etoposide (100 mg)	
Cisplatin (80 mg)/etoposide (80 mg)	
Cisplatin (25 mg)/etoposide (100 mg)	
Carboplatin/irinotecan	
Cisplatin (60 mg)/irinotecan (60 mg)	
Cisplatin (30 mg)/irinotecan (65 mg)	

Maintenance Therapy

Atezolizumab [†]	
Durvalumab [‡]	
Lurbinectedin [†]	*

Postoperative Adjuvant Therapy for Limited Stage SCLC, T1 2

	N0
Cisplatin (75 mg)/etoposide (100 mg)	
Cisplatin (60 mg)/etoposide (120 mg)	
Carboplatin/etoposide	

	N+
Cisplatin (60 mg)/etoposide (120 mg) + RT	
Cisplatin (80 mg)/etoposide (100 mg) + RT	
Carboplatin/etoposide + RT	

*Evidence Block development in progress

[†]If atezolizumab was used with a first-line carboplatin/etoposide/atezolizumab regimen.

‡If durvalumab was used with a first-line carboplatin/etoposide/durvalumab regimen or a cisplatin/etoposide/durvalumab regimen.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.

RECOMMENDED SUBSEQUENT THERAPY REGIMENS FOR SCLC*

Preferred Regimens	
Tarlatamab-dlle	
Irinotecan	
Lurbinectedin	
Topotecan	
Other Recommended Regimens	
CAV (cyclophosphamide/doxorubicin/vincristine)	
Docetaxel	
Gemcitabine	
Nivolumab	
Pembrolizumab	
Oral etoposide	
Paclitaxel	
Temozolomide	

*Subsequent therapy refers to second-line and beyond therapy.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

Table 1 - Definition of small cell lung cancer consists of two stages:

- (1) Limited stage: Stage I-III (T any, N any, M0) that can be safely treated with definitive radiation doses. Excludes T3-4 due to multiple lung nodules that are too extensive or have tumor/nodal volume that is too large to be encompassed in a tolerable radiation plan.
- (2) Extensive stage: Stage IV (T any, N any, M 1a/b/c), or T3-4 due to multiple lung nodules that are too extensive or have tumor/nodal volume that is too large to be encompassed in a tolerable radiation plan.

Table 2 - American Joint Committee on Cancer (AJCC) Eighth ed., 2017 Definitions of TNM

T	Primary Tumor
TX	Primary tumor cannot be assessed, or tumor proven by the presence of malignant cells in sputum or bronchial washings but not visualized by imaging or bronchoscopy
T0	No evidence of primary tumor
Tis	Carcinoma <i>in situ</i> Squamous cell carcinoma <i>in situ</i> (SCIS) Adenocarcinoma <i>in situ</i> (AIS): adenocarcinoma with pure lepidic pattern, ≤3 cm in greatest dimension
T1	Tumor ≤3 cm in greatest dimension, surrounded by lung or visceral pleura, without bronchoscopic evidence of invasion more proximal than the lobar bronchus (i.e., not in the main bronchus)
T1mi	Minimally invasive adenocarcinoma: adenocarcinoma (≤3 cm in greatest dimension) with a predominantly lepidic pattern and ≤5 mm invasion in greatest dimension
T1a	Tumor ≤1 cm in greatest dimension. A superficial, spreading tumor of any size whose invasive component is limited to the bronchial wall and may extend proximal to the main bronchus also is classified as T1a, but these tumors are uncommon.
T1b	Tumor >1 cm but ≤2 cm in greatest dimension
T1c	Tumor >2 cm but ≤3 cm in greatest dimension
T2	Tumor >3 cm but ≤5 cm or having any of the following features: (1) Involves the main bronchus, regardless of distance to the carina, but without involvement of the carina; (2) Invades visceral pleura (PL1 or PL2); (3) Associated with atelectasis or obstructive pneumonitis that extends to the hilar region, involving part or all of the lung. T2 tumors with these features are classified as T2a if ≤4 cm or if the size cannot be determined and T2b if >4 cm but ≤5 cm.
T2a	Tumor >3 cm but ≤4 cm in greatest dimension
T2b	Tumor >4 cm but ≤5 cm in greatest dimension
T3	Tumor >5 cm but ≤7 cm in greatest dimension or directly invading any of the following: parietal pleura (PL3), chest wall (including superior sulcus tumors), phrenic nerve, parietal pericardium; or separate tumor nodule(s) in the same lobe as the primary
T4	Tumor >7 cm or tumor of any size invading one or more of the following: diaphragm, mediastinum, heart, great vessels, trachea, recurrent laryngeal nerve, esophagus, vertebral body, or carina; separate tumor nodule(s) in an ipsilateral lobe different from that of the primary

©, 2017, American College of Surgeons, All Rights Reserved. *Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging System.*

[Continued](#)



Table 2. Definitions for T, N, M (continued)

N	Regional Lymph Nodes
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in ipsilateral peribronchial and/or ipsilateral hilar lymph nodes and intrapulmonary nodes, including involvement by direct extension
N2	Metastasis in ipsilateral mediastinal and/or subcarinal lymph node(s)
N3	Metastasis in contralateral mediastinal, contralateral hilar, ipsilateral or contralateral scalene, or supraclavicular lymph node(s)
M	Distant Metastasis
MX	Distant metastasis cannot be assessed
M0	No distant metastasis
M1	Distant metastasis
M1a	Separate tumor nodule(s) in a contralateral lobe; tumor with pleural or pericardial nodules or malignant pleural or pericardial effusion ^a
M1b	Single extrathoracic metastasis in a single organ (including involvement of a single nonregional node)
M1c	Multiple extrathoracic metastases in a single organ or in multiple organs

Table 3. AJCC Prognostic Groups

	T	N	M
Occult carcinoma	TX	N0	M0
Stage 0	Tis	N0	M0
Stage IA1	T1mi	N0	M0
	T1a	N0	M0
Stage IA2	T1b	N0	M0
Stage IA3	T1c	N0	M0
Stage IB	T2a	N0	M0
Stage IIA	T2b	N0	M0
Stage IIB	T1a	N1	M0
	T1b	N1	M0
	T1c	N1	M0
	T2a	N1	M0
	T2b	N1	M0
	T3	N0	M0
Stage IIIA	T1a	N2	M0
	T1b	N2	M0
	T1c	N2	M0
	T2a	N2	M0
	T2b	N2	M0
	T3	N1	M0
	T4	N0	M0
	T4	N1	M0

Prognostic Stage Groups

	T	N	M
Stage IIIB	T1a	N3	M0
	T1b	N3	M0
	T1c	N3	M0
	T2a	N3	M0
	T2b	N3	M0
	T3	N2	M0
	T4	N2	M0
Stage IIIC	T3	N3	M0
	T4	N3	M0
Stage IV	Any T	Any N	M1
Stage IVA	Any T	Any N	M1a
	Any T	Any N	M1b
Stage IVB	Any T	Any N	M1c

^a Most pleural (pericardial) effusions with lung cancer are a result of the tumor. In a few patients, however, multiple microscopic examinations of pleural (pericardial) fluid are negative for tumor, and the fluid is nonbloody and not an exudate. If these elements and clinical judgment dictate that the effusion is not related to the tumor, the effusion should be excluded as a staging descriptor.

©, 2017, American College of Surgeons, All Rights Reserved. *Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging System.*



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

ABBREVIATIONS

3D-CRT	three dimensional conformal radiation therapy	ICI	immune checkpoint inhibitor	SRS	stereotactic radiosurgery
4D-CT	four-dimensional computed tomography	IGRT	image-guided radiation therapy	SRT	stereotactic radiation therapy
ACTH	adrenocorticotrophic hormone	IMRT	intensity-modulated radiation therapy	SVC	superior vena cava
ADH	antidiuretic hormone secretion	INSM1	insulinoma-associated protein 1	TKI	tyrosine kinase inhibitor
AIS	adenocarcinoma in situ	LCNEC	large-cell neuroendocrine carcinoma	TNM	tumor node metastasis
AUC	area under the curve	LEMS	Lambert-Eaton myasthenic syndrome	TTF-1	thyroid transcription factor-1
BUN	blood urea nitrogen	LFT	liver function test	VMAT	volumetric modulated arc therapy
C/A/P	chest/abdomen/pelvis	LS-SCLC	limited stage small cell lung cancer	WBRT	whole brain radiation therapy
CBC	complete blood count	NSCLC	non-small cell lung cancer		
CNS	central nervous system	PCI	prophylactic cranial irradiation		
CONVERT	concurrent once-daily versus twice-daily radiotherapy	PD-1	programmed cell death protein 1		
CREST	carotid revascularization endarterectomy versus Stenting Trial	PD-L1	programmed death ligand 1		
CRT	conformal radiation therapy	PFT	pulmonary function test		
CTFI	chemotherapy-free interval	PS	performance status		
EBRT	external beam radiation therapy	RBC	red blood cell		
ENI	elective nodal irradiation	SABR	stereotactic ablative radiotherapy		
EORTC	European Organisation for Research and Treatment of Cancer	SBRT	stereotactic body radiation therapy		
ES-SCLC	extensive stage small cell lung cancer	SCIS	squamous cell carcinoma in situ		
FDG	fluorodeoxyglucose	SCLC	small cell lung cancer		
G-CSF	granulocyte colony-stimulating factor	SER	start of any therapy to the end of RT		
GM-CSF	granulocyte-macrophage colony-stimulating factor	SIB	simultaneous integrated boost		
GTV	gross tumor volume	SIADH	syndrome of inappropriate antidiuretic hormone secretion		
H&E	hematoxylin and eosin				
H&P	history and physical				
HA	hippocampal avoidance				

NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

NCCN Evidence Blocks™

Discussion

- The Discussion on the following pages describes the data and clinical information that support the recommendations in the algorithm.
- The presence of a "Discussion update in progress" watermark indicates that the section is being updated. The date of the last update is shown on MS-2.

Guidelines Update Methodology

- For complete details, refer to [Development and Update of Guidelines](#).
- Prior to the update of the NCCN Guidelines, an electronic search of the PubMed database (<https://pubmed.ncbi.nlm.nih.gov>) using relevant search terms was performed to obtain key literature in the field published since the previous Guidelines update. The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature. The search results were narrowed by selecting studies in humans published in English. Results were confined to select article types including, but not limited to: Clinical Trial; Randomized Controlled Trial; Meta-Analysis; Systematic Review; and Validation Study. The data from key PubMed articles as well as articles from additional sources deemed as relevant to these Guidelines as discussed by the Panel during the Guidelines update have been included in the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel's review of lower-level evidence and expert opinion.
- NCCN Guidelines strive to use language that is person-first, non-stigmatizing, and inclusive and representative of all individuals to whom the Guideline recommendations may apply. Therefore, NCCN Guidelines incorporate organ-specific, non-gendered recommendations and respectful terminology that avoids harmful or disparaging stereotypes. NCCN's Guidance on Inclusive Language can be found at <https://www.nccn.org/business-policy/policy-and-advocacy-program/advancing-cancer-equity>.
- For information about systemic therapy regimen naming methods used within the NCCN Guidelines algorithm, refer to the NCCN Chemotherapy Order Templates® [Systemic Therapy Regimen Naming Methods: Appendix I](#).
- For information about standardized use of biomarker terminology within the NCCN Guidelines algorithm, refer to [Tjota MY, et al. J Natl Compr Canc Netw](#).



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Discussion

This discussion corresponds to the NCCN Guidelines for Small Cell Lung Cancer. Last updated: July 1, 2026

Table of Contents

Overview	MS-3
Diagnosis	MS-4
Screening.....	MS-4
Manifestations.....	MS-4
Pathology.....	MS-5
Evaluation	MS-6
Staging Systems	MS-6
Initial Evaluation.....	MS-6
Prognostic Factors	MS-8
Treatment.....	MS-8
Surgical Resection of Stage I–IIA SCLC	MS-8
Systemic Therapy	MS-9
Older Patients	MS-14
Surveillance for Relapse	MS-15
Subsequent Systemic Therapy	MS-15
Radiation Therapy.....	MS-20
Radiation for Extensive-Stage SCLC	MS-23
Summary.....	MS-27
References.....	MS-28



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Overview

Approximately 13% of lung tumors will be classified as small cell lung cancer (SCLC) in 2026.¹ In 2026, an estimated 30,000 new cases of SCLC will be diagnosed in the United States.¹ Although the incidence of SCLC has been decreasing, the frequency in females is increasing and the male-to-female incidence ratio has been close to 1:1 in the last few years.² Nearly all cases of SCLC are attributable to cigarette smoking.³ Patients with SCLC who also continue to smoke tobacco during treatment have increased toxicity and shorter survival.⁴ Therefore, tobacco smoking cessation counseling and intervention should be strongly promoted in patients with SCLC and other high-grade neuroendocrine carcinomas (see the NCCN Clinical Practice Guidelines in Oncology [NCCN Guidelines®] for Smoking Cessation, available at www.NCCN.org).⁵ Patients who previously smoked tobacco should be strongly encouraged to remain abstinent. Programs using behavioral counseling combined with U.S. Food and Drug Administration (FDA)-approved medications that promote smoking cessation can be very useful.

SCLC is characterized by a rapid doubling time, high growth fraction, and early development of widespread metastases. While most patients with SCLC present with hematogenous metastases; approximately one third present with limited disease confined to the chest. Although 95% of small cell carcinomas originate in the lung, they can also arise from extrapulmonary sites, including the nasopharynx, gastrointestinal tract, and genitourinary tract.^{6,7} Both pulmonary and extrapulmonary small cell carcinomas have a similar clinical and biologic behavior with increased potential for widespread metastases. Management of SCLC is described in the NCCN Guidelines® for Small Cell Lung Cancer that includes the algorithm and this supporting Discussion text. Management of other lung neuroendocrine tumors (LNTs) and non-small cell lung cancer (NSCLC) are described in the NCCN Guidelines for Neuroendocrine and Adrenal Tumors and NCCN Guidelines for Non-Small Cell Lung Cancer, respectively, available at www.NCCN.org.

The definitions for limited- and extensive-stage SCLC incorporate tumor, node, metastasis (TNM) staging. The Panel recommends that the workup for SCLC should be expedited and if possible, studies should be performed in parallel. In patients with limited-stage SCLC, the goal of treatment is cure using chemotherapy plus thoracic radiation therapy (RT). However, some patients with resectable tumors (stage I–IIA) are eligible for curative surgery followed by systemic therapy with or without mediastinal RT.^{8,9} The Panel recommends multidisciplinary evaluation before any surgery. In patients with stage I–IIA SCLC, including medically inoperable circumstances or when the decision not to pursue surgical resection is made, stereotactic ablative radiotherapy (SABR) followed by systemic therapy is an option.^{10–15} The benefits of prophylactic cranial irradiation (PCI) are unclear in patients with stage I SCLC (T1–2a, N0, M0) who have received definitive therapy. The Panel recommends that MRI brain surveillance be considered for all patients with limited-stage SCLC who do not receive PCI. In most patients with extensive-stage SCLC, systemic therapy with or without RT can palliate symptoms and prolong survival; however, long-term survival is rare.¹⁶ SCLC is highly sensitive to initial chemotherapy and RT; however, most patients eventually die of recurrent disease.¹⁷ Despite recent advances, the recommended therapy options for SCLC need improvement. Clinical trials generally represent state-of-the-art treatment for patients with SCLC. Thus, participation in clinical trials is strongly encouraged.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Diagnosis

Screening

Ideally, a screening test should detect disease at an early stage while it is still curable. The National Lung Screening Trial reported that screening with annual, low-dose, spiral CT scans detected early-stage NSCLC and decreased lung cancer-specific mortality in individuals who are asymptomatic and at high risk (see the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org).¹⁸ Low-dose CT may be less effective for SCLC screening, because symptomatic disease can develop between annual scans due to the aggressive nature of the disease, thereby limiting the potential for reducing mortality through screening.¹⁸⁻²¹ Nevertheless, the same population of patients is eligible for and should benefit from LDCT screening for mortality reduction from NSCLC.

Manifestations

Patients with SCLC typically present with a large hilar mass and bulky mediastinal lymphadenopathy that causes cough and dyspnea.²² Frequently, patients present with symptoms of widespread metastatic disease, such as weight loss, debility, bone pain, and neurologic compromise. It is uncommon for patients to present with a solitary peripheral nodule without central adenopathy. In this situation, fine-needle aspiration (FNA) may not adequately differentiate small cell carcinoma (which is a high-grade neuroendocrine carcinoma) from low-grade (typical carcinoid), intermediate-grade (atypical carcinoid), or large-cell neuroendocrine carcinoma (LCNEC) (which is also a high-grade neuroendocrine carcinoma) (see *Neuroendocrine Tumors* of the Lung in the NCCN Guidelines for Neuroendocrine and Adrenal Tumors, available at www.NCCN.org).^{23,24}

Many neurologic and endocrine paraneoplastic syndromes, including Lambert-Eaton myasthenic syndrome (LEMS), encephalomyelitis, and sensory neuropathy, are associated with SCLC.²⁵⁻²⁷ Patients with LEMS present with proximal leg weakness that is caused by antibodies directed against the voltage-gated calcium channels.^{28,29} Neurologic workup may include but is not limited to tests for PQ- and N-type voltage-gated calcium channel (VGCC) antibodies. In consultation with neurology, amifampridine or intravenous immunoglobulin (IVIG) can be considered for LEMS treatment. The Panel clarified that both the tests for syndrome and treatment should be performed in consultation with neurology. Paraneoplastic encephalomyelitis may precede a diagnosis of SCLC.³⁰ Given the occurrence of paraneoplastic neurologic syndromes in individuals with SCLC, the Panel recommends considering early subspecialty consultation for most recent management and a comprehensive paraneoplastic antibody panel and/or neurologic consultation if neurologic paraneoplastic syndrome is suspected.

SCLC tumors sometimes produce polypeptide hormones, including vasopressin and adrenocorticotrophic hormone, that cause syndrome of inappropriate antidiuretic hormone secretion (SIADH) and Cushing syndrome, respectively.^{31,32} SIADH occurs more frequently than Cushing syndrome in patients with SCLC. Primary treatment for SIADH includes fluid restriction (which is difficult for patients because of increased thirst) and demeclocycline. Cancer treatment (eg, cisplatin) and/or supportive care (e.g., opiates) may also cause hyponatremia.³³ Hyponatremia usually improves after successful treatment of SCLC. However, vasopressin receptor inhibitors (i.e., conivaptan, tolvaptan) can be used for refractory hyponatremia.³³⁻³⁵



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Pathology

SCLC is a poorly differentiated malignant epithelial tumor that is categorized as a high-grade neuroendocrine carcinoma.^{23,36} Up to 30% of tumors from patients with SCLC, particularly those who received treatment, reveal areas of NSCLC differentiation (mainly large cell carcinoma).³⁷ In cases of diagnostic dilemma, including carcinoids, the Panel strongly recommends getting a second opinion from a pathologist specializing in diagnosis of thoracic malignancies. In good-quality samples, the classic and distinctive histology on hematoxylin and eosin (H&E), including small blue cells with scant cytoplasm, high nuclear/cytoplasmic ratio, fine granular nuclear chromatin, and absent or inconspicuous nucleoli, may be sufficient for identifying SCLC.^{23,38} Compared to atypical and typical carcinoids, the mitotic count is high in SCLC. However, mitotic figures are difficult to count in small biopsy samples with crushed or necrotic cells; immunohistochemistry is useful in such samples.³⁹

Using immunohistochemistry as one of the tools to diagnose and distinguish SCLC from NSCLC or other neuroendocrine tumors is important, especially because these cancer types have different treatment recommendations.^{23,39-43} Ki-67 is useful for distinguishing SCLC from carcinoid tumors.^{39,43-45} Nearly all SCLCs are immunoreactive for cytokeratin (AE1/AE3, CAM5.2) and 85% to 90% of SCLCs are positive for thyroid transcription factor-1 (TTF-1).^{23,46-48} Most SCLCs (~95%) stain positively for markers of neuroendocrine differentiation, including insulinoma-associated protein 1 (INSM1), chromogranin A, NCAM (CD56), and synaptophysin.^{23,49,50} However, approximately 10% of NSCLCs will be immunoreactive for at least one of these neuroendocrine markers and therefore cannot be used alone to distinguish SCLC from NSCLC.⁵¹ For suspicious cases of SCLC morphology without any expression of neuroendocrine markers, POU2F3 IHC staining can be considered.^{52,53} Napsin A (adenocarcinoma marker) and p40 (or p63, squamous cell carcinoma marker) are generally negative in SCLC and useful for distinguishing SCLC from poorly differentiated NSCLC and combined SCLC.⁵⁴ It is important to note that p40 (or p63) can be focally positive in SCLC. Immunohistochemical stains for Rb1 and p53 can also be used to differentiate atypical carcinoid and SCLC. While the carcinoid tumors retain RB1 expression and show a wild-type p53 expression, most high-grade neuroendocrine tumors of the lung show RB1 loss and/or p53 mutation-type expression with either strong and diffuse nuclear staining or null (null) type staining. The Panel clarified that biomarker testing via blood, tissue, or both may be considered in rare cases, particularly for patients with extensive-stage/relapsed SCLC who have never smoked or lightly smoked, remote smoking history, diagnostic/therapeutic dilemma, or at time of relapse if not done previously (as this may change management).⁵⁵⁻⁶⁰

The WHO classification recognizes two types of SCLC: pure and combined SCLC.^{43,56,61-64} Combined SCLC, which consists of both SCLC and NSCLC histology (squamous cell, adenocarcinoma, spindle/pleomorphic, and/or large cell carcinoma), is more frequent in patients with limited-stage SCLC.^{43,56,63,65} Any presence of NSCLC histology (no minimal percentage) results in a classification of combined SCLC. The only exception is combined SCLC and LCNEC where at least 10% of the tumor should show LCNEC morphology.^{37,66} Patients with combined SCLC are treated using regimens for SCLC, because it is the more aggressive cancer.⁶⁶ Patients with NSCLC can also transform to SCLC after treatment with epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors or immune checkpoint inhibitors (ICIs).^{67,68}



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Evaluation

Staging Systems

The Veterans Administration (VA) Lung Study Group's 2-stage classification scheme has historically been used to define the extent of disease in patients with SCLC: 1) limited-stage is disease confined to the ipsilateral hemithorax, which can be safely encompassed within a radiation field; and 2) extensive-stage is disease beyond the ipsilateral hemithorax, including malignant pleural/pericardial effusion or hematogenous metastases.⁶⁹ Contralateral mediastinal and ipsilateral supraclavicular lymphadenopathy are generally classified as limited-stage SCLC, whereas the classification of contralateral hilar and supraclavicular lymphadenopathy is more controversial and treatment is individualized.^{17,70,71} Approximately 66% of patients present with overt hematogenous metastases, which commonly involve the contralateral lung, liver, adrenal glands, brain, bones, and/or bone marrow. Most studies use the VA definitions of limited- or extensive-stage SCLC for clinical decision-making. But the TNM system is particularly useful for selecting patients with T1–2, N0 disease who are eligible for surgery and RT.⁷⁰ The American Joint Committee on Cancer (AJCC) revised the TNM staging system (8th edition) for lung cancer in 2017.^{72,73} Clinical studies that include the TNM system will allow for more precise assessments of prognosis and specific therapy.⁷²

The Panel adopted a combined approach of using the AJCC TNM staging system and the older VA scheme for SCLC staging.^{17,70} *Limited-stage* SCLC is defined as stage I–III (T any, N any, M0) that can be safely treated with definitive RT. This excludes T3–4 due to multiple extensive lung nodules or disease with tumor/nodal volume that is too large to be encompassed in a tolerable radiation plan. *Extensive-stage* SCLC is defined as stage IV (T any, N any, M1a/b/c) or T3–4 as described above.

Initial Evaluation

The workup for SCLC should be expedited with studies done in parallel whenever possible. Staging primarily provides a therapeutic guideline for thoracic RT for patients with limited-stage SCLC. However, staging should not delay the onset of treatment for >1 week because of the aggressive nature of SCLC. Many patients may become more seriously ill in the interval with a significant decline in performance status (PS). The Panel recommends early integration of palliative care so that management of cancer-related symptoms and goals of care are discussed during initial evaluation.

The initial diagnostic evaluation includes a history and physical examination; pathology review; clinical laboratory tests; CT scan with contrast of the chest/abdomen/pelvis (C/A/P); and MRI is preferred over CT scan with contrast for brain imaging.^{71,74} An FDG-PET/CT scan (skull base to mid-thigh), which is superior to PET alone, is recommended, if needed, to clarify the extent of disease.^{17,70,75} For most metastatic sites, FDG-PET/CT is superior to CT imaging; however, for the detection of brain metastases FDG-PET/CT is inferior to MRI (or contrast-enhanced CT as an alternative when MRI is not possible) (see the NCCN Guidelines for Central Nervous System Cancers, available at www.NCCN.org).⁷⁶ FDG-PET scans can also increase staging accuracy because SCLC is a highly metabolically active disease.^{75,77,78} Approximately 19% of patients who undergo FDG-PET are upstaged from limited-stage to extensive-stage SCLC, whereas 8% are downstaged from extensive-stage to limited-stage SCLC.⁷¹ Although FDG-PET/CT seems to improve staging accuracy in SCLC, pathologic confirmation is recommended for isolated or equivocal lesions if their involvement changes clinical



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

management. In select studies, FDG-PET staging improved detection of intrathoracic disease sites in approximately 27% of patients and led to alteration of the planned radiation field.^{71,78,79}

Once extensive-stage SCLC has been established, further staging is optional and dependent on the clinical situation. Further brain imaging with MRI (preferred) or CT with contrast is recommended. Brain imaging can identify central nervous system (CNS) metastases in 10% to 15% of patients at diagnosis, of which approximately 30% are asymptomatic. Early treatment of brain metastases results in less chronic neurologic morbidity, arguing for the usefulness of early diagnosis in asymptomatic patients. Therefore, staging should broaden from sites of symptomatic disease, or those sites suggested by laboratory tests.

Bone imaging with radiographs or MRI can be performed if FDG-PET/CT is equivocal or not available. Bone scans are positive in <30% of patients without bone pain or abnormal alkaline phosphatase levels. However, <5% of patients have bone marrow involvement as the only site of extensive-stage SCLC. Unilateral bone marrow aspirates and biopsies may be indicated in select patients with no other evidence of metastatic disease, with nucleated red blood cells on peripheral blood smear and neutropenia, or thrombocytopenia suggestive of bone marrow infiltration. The Panel recommends performing bone marrow biopsy in select patients with pathologic confirmation for isolated or equivocal lesions, only if it changes clinical management. Any compelling evidence of distant disease consistent with malignancy should change the diagnosis to extensive-stage SCLC.

To confirm FDG-PET/CT scan results and rule out occult nodal disease before surgical resection, pathologic lymph node staging is recommended in patients with clinical limited-stage I–IIA SCLC (T1–2, N0, M0).¹⁷ Pathologic lymph node staging can be considered to help determine RT fields in patients with clinical stage IIB–IIIC SCLC (T1–4, N0, M0; T1–4, N1–3, M0), especially for those with clinical N0 disease. Lymph node staging procedures include endobronchial ultrasound-guided biopsy (preferred, if expertise exists), mediastinoscopy, mediastinotomy, esophageal ultrasound-guided biopsy, and video-assisted thoracoscopy.^{80,81} The Panel modified these recommendations to consider endoscopic procedures first, when expertise exists, before other invasive staging procedures are performed. Staging biopsies of visible nodes in all accessible nodal stations should be performed to inform RT targeting, but additional modalities for mediastinal staging are not necessary. The Panel further clarified that if nodal involvement is established by one modality of mediastinal staging, additional modalities are not necessary, but staging biopsies of visible nodes in all accessible nodal stations is encouraged to inform radiotherapy targeting.

Thoracentesis with cytologic analysis is recommended if pleural effusion is large enough to be safely accessed via ultrasound guidance. If thoracentesis does not show malignant cells, then thoracoscopy can be considered to document pleural involvement, which is suggestive of extensive-stage SCLC. The effusion should be excluded as a staging element if: 1) multiple cytopathologic examinations of the pleural fluid are negative for cancer; 2) the fluid is not bloody and not an exudate; and 3) clinical judgment concludes that the effusion is not directly related to the cancer. Pericardial effusions are classified using the same criteria.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Prognostic Factors

Age <70 years, normal lactate dehydrogenase (LDH), and stage I disease are associated with favorable prognosis in patients with limited-stage SCLC. Younger age, good PS, normal creatinine level, normal LDH, and a single metastatic site are favorable prognostic factors in patients with extensive-stage SCLC.^{82,83} Poor PS (3–4), extensive-stage SCLC, weight loss, and markers (such as LDH) associated with bulky disease are the most important adverse prognostic factors.

Treatment

Surgical Resection of Stage I–IIA SCLC

Fewer than 5% of patients with SCLC have true stage I–IIA disease.⁸⁴ The general recommendation for surgery is restricted to certain patients with stage I–IIA disease (T1–2, N0, M0). Most of the data regarding the role of surgery in SCLC are from retrospective studies.^{85–90} These studies report favorable 5-year survival rates of 40% to 60% in patients with stage I disease. In most series, survival rates decline significantly in patients with advanced disease and lymph node involvement. Analyses of the SEER database suggest that surgery is appropriate for some patients with localized disease.^{15,91} However, retrospective studies and analyses of the SEER database are limited by the lack of information on chemotherapy. In addition, comparison of the survival of surgical patients to those who did not undergo surgery is inherently flawed by selection bias. Ultimately, the role of surgery in SCLC will not be fully delineated until prospective trials compare surgery plus adjuvant chemotherapy with concurrent chemoradiotherapy in patients who are rigorously staged. SABR, discussed later, is another option for stage I–IIA SCLC.

The Lung Cancer Study Group conducted the only prospective randomized trial evaluating the role of surgery in SCLC.⁹² Patients with limited-stage SCLC, excluding those with solitary peripheral nodules, that responded to 5 cycles of chemotherapy with CAV (cyclophosphamide, doxorubicin, and vincristine) were randomly assigned to undergo thoracic RT with or without resection. The overall survival (OS) rates of patients in the two arms were equivalent, suggesting no benefit to surgery in this setting. However, only 19% of enrolled patients had clinical limited-stage I (T1–2, N0, M0) disease. Additionally, data show that patients with SCLC who have nodal disease (i.e., T1–3, N1–3, M0–1) do not benefit from surgery.⁹²

The Panel recommends that small lesions that are presumed to be small cell carcinoma on biopsy in patients who do not smoke tobacco should be resected, because they are likely carcinoids that have been misdiagnosed (see the NCCN Guidelines for Neuroendocrine and Adrenal Tumors, available at www.NCCN.org). The Panel also recommends surgery for certain patients with clinical stage I–IIA (T1–2, N0) SCLC with negative nodes that have been confirmed by pathologic lymph node staging.^{13,85,93} Surgery can include patients with clinical limited-stage IIA SCLC based on the staging criteria that includes tumors <5 cm in diameter (T2b) without lymph node involvement (N0). The Panel added recommendations for patients who did not have a preoperative biopsy but have an intraoperative diagnosis of likely SCLC. If resection is performed, the Panel recommends lobectomy (preferred) with mediastinal lymph node dissection or systematic lymph node sampling (e.g., ≥3 N2 and ≥1 N1 stations). The Panel does not recommend pneumonectomy if nodal metastatic disease needs to be encompassed or under other circumstances. Chemoradiation is the preferred alternative over any resection requiring pneumonectomy.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Following surgery, adjuvant chemotherapy is recommended for patients with limited-stage I–IIA (T1–2, N0) SCLC with negative margins (R0 resection).^{88,94–96} Concurrent or sequential chemotherapy and postoperative mediastinal RT are recommended for patients with N+ disease.⁹⁷ Although Panel members agree that postoperative mediastinal RT is recommended for nodal metastases, it should be based on the extent of nodal sampling/dissection and extent of nodal positivity. For patients with limited-stage I–IIA (T1–2, N0) SCLC with positive margins (R1/R2 resection), the Panel recommends concurrent chemoradiation following surgery.

Systemic Therapy

For all patients with SCLC, systemic therapy is an essential component of primary/adjuvant treatment. Adjuvant chemotherapy is recommended for patients with limited-stage SCLC who have had surgery or SABR. Systemic therapy following concurrent chemoradiation or SABR is also recommended for patients with limited-stage I–IIA (T1–2, N0, M0) SCLC who are medically inoperable or do not want to pursue surgical resection. For patients with limited-stage SCLC who are not eligible for surgery or SABR or have positive lymph node staging, chemotherapy with concurrent thoracic RT (category 1 for patients with PS 0–2) is the recommended primary treatment.^{9,98,99} For patients with extensive-stage SCLC, systemic therapy alone is recommended. RT may be used in select patients with extensive-stage disease for palliation of symptoms (see NCCN Guidelines for Palliative Care, available at www.NCCN.org). The systemic therapy recommendations are based on studies described in the following sections.

Cisplatin Versus Carboplatin

Randomized trials and retrospective analyses of patients with SCLC suggest similar efficacy between cisplatin and carboplatin regimens.^{100–102} In a meta-analysis of 663 patients with limited- (32%) and extensive-stage SCLC (68%), no significant difference was observed in response rate (67% vs. 66%), progression-free survival (PFS) (5.5 vs. 5.3 months), or OS (9.6 vs. 9.4 months) between cisplatin- versus carboplatin-containing regimens.¹⁰³ Carboplatin is frequently substituted for cisplatin to reduce the risk of emesis, neuropathy, and nephropathy.¹⁰⁴ The use of carboplatin carries a greater risk of myelosuppression.¹⁰⁴

Limited-Stage SCLC

Adjuvant chemotherapy alone is recommended for patients who have undergone surgical resection or SABR for limited-stage SCLC. Cisplatin/etoposide is the most commonly used first-line combination chemotherapy regimen for patients with limited-stage SCLC.^{105,106} Cisplatin/etoposide replaced alkylator/anthracycline-based regimens based on its superiority in both efficacy and toxicity.^{107–109} If pathologic lymph node involvement is found at surgery (N+) for patients with stage I–IIA (T1–2, N0, M0), then thoracic RT can be added concurrently or sequentially to cisplatin/etoposide. Treatment with cisplatin/etoposide plus definitive thoracic RT showed response rates of 70% to 90% with a median OS of 25 to 30 months and 5-year OS rates of 31% to 34%.¹⁰⁵ Cisplatin/etoposide in combination with thoracic RT increases the risk of esophagitis, as well as pulmonary and hematologic toxicity.¹¹⁰ Thoracic RT improved local control rates by 25% and is associated with improved survival in patients with limited-stage SCLC.^{98,99} Data suggest that chemoradiotherapy may also be indicated for patients with limited-stage SCLC who have cytologically negative or indeterminate pleural effusions but not for those with pericardial effusions.^{111,112}



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

For patients with limited-stage IIB–IIIC (T3–4, N0, M0; T1–4, N1–3, M0) SCLC, the Panel recommends different cisplatin/etoposide regimens plus concurrent thoracic RT (category 1).^{98,99,113,114} The preferred cisplatin/etoposide regimens for limited-stage SCLC are based on the dosing used in the CONVERT trial.¹⁰⁵ Carboplatin/etoposide is also listed as a preferred option based on the subgroup analysis in the ADRIATIC trial.^{115,116} Other recommended options for limited-stage SCLC include additional cisplatin/etoposide doses. The use of myeloid growth factors is not recommended in patients undergoing concurrent chemoradiation (category 1 for not using granulocyte-macrophage colony-stimulating factor [GM-CSF]).¹¹⁷

Response assessment is an important aspect of the management of SCLC. For patients with limited-stage SCLC, the Panel recommends response assessment using CT with contrast after completion of adjuvant chemotherapy alone or chemotherapy with concurrent RT. Response assessment can be measured using CT with contrast of the C/A/P and brain MRI (preferred) or brain CT with contrast. For systemic therapy alone or sequential systemic therapy followed by RT in patients with limited-stage SCLC, the Panel recommends response assessment using CT with contrast of the C/A/P after every 2 cycles of systemic therapy and at completion of therapy. In the absence of new symptoms, repeating scans to assess response during adjuvant or initial treatment is not recommended.

Durvalumab Consolidation Therapy

The ADRIATIC trial included patients with stage I–III limited-stage SCLC, PS 0–1 and no disease progression on chemoradiation. The median OS was 55.9 months in those patients receiving consolidation therapy ($n = 264$) compared to 33.4 months ($P = .01$) in patients receiving placebo ($n = 266$).¹¹⁵ The study revealed a statistically significant and clinically meaningful OS and PFS (16.6 vs. 9.2 months, $P = .02$) improvement in patients receiving consolidation durvalumab compared to placebo. Based on these data, the Panel recommends durvalumab consolidation therapy in those patients with limited-stage SCLC who did not experience disease progression after systemic therapy + concurrent RT (category 1). Durvalumab may be continued until disease progression or intolerable toxicity, or for a maximum of 24 months. The Panel advises imaging every 8 weeks for the first 18 months and every 12 weeks after. Additionally, the Panel recommends if PCI is being considered in this setting it should be given prior to consolidation durvalumab.

Extensive-Stage SCLC

The Panel recommends certain combination chemotherapy plus immunotherapy regimens as preferred options for patients with extensive-stage SCLC.^{118–120} In patients with extensive-stage SCLC and brain metastases, systemic therapy can be given either before or after brain RT depending on whether the patient has neurologic symptoms.^{16,121} The Panel recommends steroid initiation for patients with spinal cord compression or brain metastasis who have symptomatic neurologic disease. For example, dexamethasone 10 mg loading dose followed by 4 to 6 mg maintenance dose (IV or oral [PO] every 4–6 hours [or as appropriate]) can be initiated.¹²² If systemic therapy is given first, brain RT is administered after completion of systemic therapy. However, if brain metastases progress while on systemic therapy, it is recommended that brain RT is initiated before completion of systemic therapy. Systemic therapy with or without RT to localized symptomatic sites is recommended for patients with extensive disease, which includes the superior vena cava (SVC), lobar obstruction, and bone metastasis. Prophylactic RT can be considered for patients with high fracture risk due to osseous structural impairment.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

During systemic therapy for patients with extensive-stage SCLC, the Panel recommends response assessment using CT with contrast of the C/A/P after every 2 to 3 cycles of systemic therapy and at completion of therapy.

For patients with known brain metastases, brain MRI (preferred) or brain CT with contrast should be obtained every 3–4 months, or at a frequency based on clinical indication. If systemic therapy was initiated prior to brain RT, brain MRI (preferred) or CT with contrast is recommended to be repeated after every 2 cycles of systemic therapy until brain RT is initiated.

The preferred regimens for patients with extensive-stage SCLC include etoposide/platinum regimens in combination with the programmed death ligand 1 (PD-L1)–targeted ICIs, atezolizumab or durvalumab. Contraindications for treatment with programmed cell death protein 1 (PD-1)/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or concurrent use of immunosuppressive agents. The Panel recommends continuation of maintenance immunotherapy until disease progression or intolerable toxicity. Atezolizumab or durvalumab may cause unique immune-mediated adverse events that are not seen with traditional cytotoxic chemotherapy. Therefore, health care providers should be aware of the spectrum and management of potential immune-mediated adverse events and have a discussion with patients about possible side effects (see the NCCN Guidelines for Management of Immune Checkpoint Inhibitor-Related Toxicities, available at www.NCCN.org). High-dose corticosteroids are generally recommended for immune-mediated adverse events based on the severity of the reaction. Atezolizumab or durvalumab should be withheld or discontinued for severe or life-threatening immune-mediated adverse events when indicated (see prescribing information).

Chemotherapy + Atezolizumab

IMpower133, a phase 3 randomized trial, assessed the addition of atezolizumab (treatment and maintenance) to carboplatin/etoposide in 403 patients with previously untreated extensive-stage SCLC and compared outcomes to carboplatin/etoposide alone.¹²⁰ The 1-year OS rate was 51.9% for the atezolizumab regimen versus 39.0% for chemotherapy alone. The median OS was 12.3 months (95% CI, 10.8–15.8) with the addition of atezolizumab versus 10.3 months (95% CI, 9.3–11.3) with chemotherapy alone (hazard ratio [HR], 0.76; 95% CI, 0.6–0.95; $P = .0154$).¹¹⁸ Response rates were similar in both arms (60% with atezolizumab/chemotherapy vs. 64% with chemotherapy alone). The rate of grade 3 or 4 adverse events was similar in both groups (67.7% for the atezolizumab regimen vs. 63.3% for chemotherapy alone). There were 4 deaths (2%) in the atezolizumab group versus 11 deaths (5.6%) in the chemotherapy alone group. Different doses for maintenance atezolizumab have been FDA-approved for patients with extensive-stage SCLC. In light of these data and FDA approval, the Panel recommends carboplatin/etoposide + atezolizumab as a category 1 and preferred first-line systemic therapy option followed by maintenance atezolizumab for patients with extensive-stage SCLC.^{118,120} The Panel recommends either 1200 (every 3 weeks) or 1680 mg (every 4 weeks) of maintenance atezolizumab. The category 1 recommendation is only for 1200 mg of maintenance atezolizumab based on the dose used in the clinical trial.^{118,120}

IMforte, a phase 3 randomized trial, assessed the addition of lurbinectedin to the carboplatin/etoposide + atezolizumab regimen during the maintenance phase with atezolizumab. PFS (stratified HR, 0.54; 95% CI, 0.43–0.67; $P < .0001$) and OS (stratified HR, 0.73; 95% CI, 0.57–0.95; $P = .017$) were longer in the lurbinectedin plus atezolizumab group than in the atezolizumab group. The patients in this trial were randomized after platinum therapy only if they



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

had either a response or stable disease; <10% of patients on the control arm received lurbinectedin after progression. The trial excluded patients with brain metastases at diagnosis. Based on these data, the Panel recommends the addition of lurbinectedin to maintenance atezolizumab only in patients who have achieved at least stable disease following four cycles of induction chemoimmunotherapy, have an Eastern Cooperative Oncology Group (ECOG) PS of 0–1, and have no history of brain metastases. The Panel indicates that atezolizumab and hyaluronidase-tqjs subcutaneous injection may be substituted for IV atezolizumab, and notes that atezolizumab and hyaluronidase-tqjs has different dosing and administration instructions compared to IV atezolizumab infusion.

Chemotherapy + Durvalumab

CASPIAN, a phase 3 randomized trial, assessed adding durvalumab to etoposide and either carboplatin or cisplatin followed by maintenance durvalumab in 537 patients with previously untreated extensive-stage SCLC and compared response to etoposide/platinum regimens.^{119,123} Most patients received carboplatin (78%). A 3-year analysis showed that the median OS was 13.0 months (95% CI, 11.5–14.8) in the chemotherapy + durvalumab group and 10.3 months (95% CI, 9.3–11.2) in the chemotherapy alone group (HR, 0.73; 95% CI, 0.59–0.91; $P = .0047$).¹²⁴ The 1-year OS rate was 52.8% for the durvalumab regimen versus 39.3% for chemotherapy alone. The rate of serious adverse events was similar in both groups (32% vs. 36%). The death rate from adverse events was also similar (2% vs. 1%). In this trial, adding tremelimumab to carboplatin (or cisplatin)/etoposide + durvalumab did not improve OS compared with platinum/etoposide (10.4 vs. 10.5 months; HR, 0.82; 95% CI, 0.68–1.0). Based on the data and FDA approval, the Panel recommends carboplatin or cisplatin/etoposide + durvalumab as a category 1 and preferred first-line systemic therapy option followed by maintenance durvalumab for patients with extensive-stage SCLC, including those with asymptomatic untreated brain metastases.^{119,123,125}

Other Primary Systemic Therapies

Other recommended first-line systemic therapy regimens for extensive-stage SCLC include cisplatin or carboplatin/etoposide combinations. Additional chemotherapy combination regimens evaluated in patients with extensive-stage SCLC show inconsistent evidence of benefit compared with cisplatin/etoposide. For example, a phase 3 trial in Japan reported that patients with extensive-stage SCLC treated with cisplatin/irinotecan had a median survival of 12.8 months compared with 9.4 months for patients treated with cisplatin/etoposide ($P = .002$).¹²⁶ The 2-year survival was 19.5% in the cisplatin/irinotecan group versus 5.2% in the cisplatin/etoposide group.¹²⁶ Two subsequent large phase 3 trials performed in the United States comparing cisplatin/irinotecan versus cisplatin/etoposide showed no significant difference in response rate or OS.^{127,128} A phase 3 randomized trial of 220 patients with extensive-stage SCLC found that median OS slightly improved with carboplatin/irinotecan compared with carboplatin/oral etoposide (8.5 vs. 7.1 months; $P = .04$).¹²⁹ In addition, a meta-analysis suggested an improvement in PFS and OS with irinotecan/platinum regimens compared with etoposide/platinum regimens.¹³⁰ The Panel recommends the irinotecan-based regimens as primary therapy options (useful in certain circumstances) for patients with extensive-stage SCLC. However, the relatively small absolute survival benefit needs to be balanced against the toxicity profile of irinotecan-based regimens.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

The use of maintenance or consolidation chemotherapy beyond the recommended 4 to 6 cycles results in a minor increase in the duration of response without improving survival and carries a greater risk of cumulative toxicity.^{131,132} The inability to destroy residual cells, despite the initial chemosensitivity of SCLC, suggests the existence of cancer stem cells that are relatively resistant to cytotoxic therapy. Alternating or sequential combination therapies have been designed to overcome drug resistance by exposing the tumor to as many active cytotoxic agents as possible during initial treatment.¹³³ However, this approach has not improved PFS or OS in randomized trials.^{134,135} Therefore, the Panel recommends 4 cycles of systemic cytotoxic therapy for patients with limited- (with concurrent radiation) and extensive-stage (with concurrent immunotherapy) SCLC. However, some patients with extensive-stage SCLC may receive up to 6 cycles based on response and tolerability.

Attempts to improve long-term survival rates in patients with SCLC through the addition of more agents or the use of dose-intensive chemotherapy, maintenance therapy, or alternating non-cross-resistant chemotherapy regimens have not shown significant advantages compared to recommended approaches. In two trials, the addition of ifosfamide (or anthracycline/cyclophosphamide) to cisplatin/etoposide showed a modest survival advantage.^{136,137} However, these findings have not been uniformly observed, and the addition of an alkylating agent, with or without an anthracycline, significantly increases hematologic toxicity when compared to cisplatin/etoposide alone.¹³⁸ Two different phase 3 trials assessing the combination of epirubicin/etoposide/ifosfamide versus cisplatin/etoposide, and carboplatin/etoposide with or without palifosfamide confirmed the lack of improvement in survival with three-drug chemotherapy regimens in patients with extensive-stage SCLC.^{139,140} Similarly, the addition of paclitaxel to either cisplatin or carboplatin/etoposide yielded promising results in phase 2 studies, but did not improve survival and was associated with unacceptable toxicity in a phase 3 trial.¹⁴¹

The role of high chemotherapy doses for patients with SCLC remains controversial. Patients receiving high chemotherapy doses compared with those given conventional doses of the same agents had higher complete and partial response rates, and modestly longer median survival times.¹⁴² However, randomized trials comparing conventional chemotherapy doses to incremental increases in dose intensity (up to 2 times the conventional dose) have not consistently shown an increase in response rate or survival.¹⁴³⁻¹⁴⁶ In addition, a meta-analysis of trials that studied dose-intensive variations of the CAV and cisplatin/etoposide regimens found that increased relative dose intensity resulted in only a small, clinically insignificant enhancement of median survival in patients with extensive-stage SCLC.¹⁴⁷ Early phase 2 results designed to increase dose intensity by weekly cyclic multidrug chemotherapy were promising, but favorable patient selection was of some concern.^{148,149} No survival benefit was documented in randomized trials, and excessive treatment-related mortality was noted with weekly cyclic multidrug chemotherapy regimens.¹⁵⁰⁻¹⁵³

Despite the success with chemotherapy + atezolizumab or chemotherapy + durvalumab regimens, other immunotherapy-based strategies have not been as favorable. A phase 3 randomized trial in patients with extensive-stage SCLC reported that the addition of ipilimumab to cisplatin or carboplatin/etoposide as first-line therapy did not improve either OS or PFS compared with chemotherapy alone.¹⁵⁴ Likewise, another phase 3 randomized trial showed no improvement in OS in patients with extensive-stage SCLC treated with first-line etoposide/platinum + pembrolizumab followed by maintenance pembrolizumab compared with chemotherapy alone.⁵⁶



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Antiangiogenic therapy has also been evaluated in SCLC. In patients with limited-stage SCLC, a phase 2 study of irinotecan, carboplatin, and bevacizumab with concurrent RT followed by maintenance bevacizumab was terminated early because of an unacceptable incidence of tracheoesophageal fistulae. In extensive-stage SCLC, phase 2 trials of platinum-based chemotherapy plus bevacizumab showed promising response and survival data.¹⁵⁵⁻¹⁵⁸ However, at least two randomized trials have demonstrated no survival benefit for the addition of bevacizumab to standard chemotherapy.^{159,160} Currently, the Panel does not recommend use of bevacizumab in patients with SCLC.

Cytokines (e.g., GM-CSF, granulocyte colony-stimulating factor [G-CSF]) can ameliorate chemotherapy-induced myelosuppression and reduce the incidence of febrile neutropenia, but cumulative thrombocytopenia remains dose-limiting. Although trials involving patients with SCLC were instrumental in obtaining FDA approval for the clinical use of cytokines,¹⁶¹ maintenance of dose intensity with growth factors does not prolong disease-free survival or OS.^{162,163} The Panel does not recommend GM-CSF (category 1 for not using it) or G-CSF for patients with limited-stage SCLC receiving systemic therapy/RT.^{117,164} Trilaciclib or G-CSF may be used as prophylactic supportive care options to decrease the incidence of chemotherapy-induced myelosuppression when administering certain regimens for patients with extensive-stage SCLC.^{117,164-169} Trilaciclib is a transient cyclin-dependent kinase (CDK) 4/6 inhibitor that has been shown to reduce chemotherapy-induced myelosuppression in patients with extensive stage SCLC. In clinical trials, trilaciclib demonstrated reductions in the incidence and duration of severe neutropenia, as well as reductions in the incidence of severe anemia and thrombocytopenia.^{167,170} It is important to note that trilaciclib or G-CSF are not treatment options.

Transformed SCLC from NSCLC with an Oncogenic Driver

The Panel recognizes that transformed SCLC from NSCLC is a rare population of patients with limited data including treatment options. The Panel recommends considering referral to a center with expertise in managing transformed SCLC with an oncogenic driver, that can be actionable genomic alterations. The Panel recommends systemic cytotoxic chemotherapy using the NCCN Guidelines for Small Cell Lung Cancer due to lack of data for transformed SCLC. The role of immunotherapy in transformed SCLC is unclear and should be avoided if tyrosine kinase inhibitors are being used. The Panel clarified that in patients who develop SCLC as a resistance mechanism to tyrosine kinase inhibitors used for actionable genomic alteration-driven NSCLC, ICIs should be avoided if tyrosine kinase inhibitors are continued. This is due to known toxicity that is primarily driven by the long half-life of ICIs.

Older Patients

The incidence of SCLC increases with age with the median age at diagnosis being >70 years.¹⁷¹ The functional status of an individual patient is more useful than chronologic age in guiding clinical decision-making that includes considering treatment tolerance (see the NCCN Guidelines for Older Adult Oncology, available at www.NCCN.org). Greater attention to the needs and support systems of a patient's functional status is recommended to provide optimal care. Patients who are able to perform activities of daily living (ADLs) should be treated with combination systemic therapy and RT, if indicated.¹⁷²⁻¹⁷⁴ For example, a subgroup analysis of the CONVERT trial suggests that concurrent chemoradiation yields equivalent median survival in patients <70 years and ≥70 years with limited-stage SCLC (29 vs. 30 months; $P = .38$).¹⁷² However, myelosuppression, fatigue, and lower organ reserves



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

are encountered more frequently in patients ≥ 70 years. Therefore, the study recommended closer surveillance for patients ≥ 70 years during treatment to avoid excessive risk.¹⁷²

Randomized trials have indicated that less-intense treatment (e.g., single-agent etoposide) is inferior to combination chemotherapy (e.g., etoposide/platinum) in patients of all ages with good PS (0–2).^{175,176} A retrospective analysis in 8637 patients ≥ 70 years with limited-stage SCLC reported that chemoradiation increased survival compared with chemotherapy alone.¹⁷³ Several other strategies have been evaluated in patients ≥ 65 years with SCLC.^{102,177–179} The use of 4 cycles of carboplatin/etoposide seems to yield favorable results, because the area-under-the-curve (AUC) dosing of carboplatin takes into account the declining renal function of the patient.¹⁷⁹ Targeting carboplatin to an AUC of 5, rather than 6, is more reasonable in the population >70 years.¹⁸⁰ A short, 2-week, full-intensity course of chemotherapy showed acceptable results in patients (median age, 73 and poor PS), but this approach has not been directly compared with 4 to 6 cycles of therapy.¹⁸¹

PCI should be used with caution in older patients. A Dutch analysis of >5000 patients suggests that median survival is lower in patients ≥ 70 years compared with patients <70 years treated with PCI, regardless of stage.¹⁸² Patients aged ≥ 60 years are at increased risk for cognitive decline after PCI; therefore, the risks and benefits of PCI versus close MRI surveillance need to be discussed.^{183–186}

Surveillance for Relapse

Although SCLC responds to initial treatment, disease relapse is observed in most patients.^{187,188} Most NCCN Member Institutions use chest CT ($\pm$ abdomen/pelvis) every 2 to 6 months (more frequently in years 1–2 and less frequently thereafter). The frequency of surveillance decreases during subsequent years because of the declining risk of recurrence.¹⁸⁹ If a new pulmonary nodule develops, it should prompt evaluation for a new primary lung cancer, because second primary tumors post-treatment frequently occur in patients with no evidence of SCLC.^{190,191} It is important to monitor for brain metastases prior to the development of potentially debilitating neurologic symptoms and to allow for early treatment. The Panel recommends brain MRI (preferred) or CT with contrast every 3 to 4 months during year 1 for all patients and then every 6 months in year 2, and as clinically indicated thereafter, regardless of the PCI status. MRI is more sensitive than CT for identifying brain metastases.⁷⁴ The Panel maintains that FDG-PET/CT is not recommended for routine follow-up unless contrast CT C/A/P or MRI is contraindicated. Tobacco smoking cessation intervention is recommended for all patients with SCLC, because second primary tumors occur less commonly in patients who quit smoking (see the NCCN Guidelines for Smoking Cessation, available at www.NCCN.org).^{192–194} Patients who previously smoked tobacco should be encouraged to remain abstinent. The Panel also recommends the NCCN Guidelines for Survivorship for appropriate patients (available at www.NCCN.org).

Subsequent Systemic Therapy

Patients with disease relapse or with primary progressive disease may be treated with subsequent systemic therapy. Consider genomic profiling of relapsed tumors, if not previously performed, to determine clinical trial eligibility. The Panel no longer recommends treatment based on the time frame for chemotherapy free intervals (CTFI). Subsequent therapy options for all patients with SCLC should be chosen based on clinical expertise and trial data



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

regardless of CTFI. The optimal duration of subsequent systemic therapy has not been fully explored. For cytotoxic chemotherapy agents, the duration of treatment is usually short, and the cumulative toxicity is frequently limiting even in patients who experience response. For these reasons, subsequent systemic therapy should be continued until progression of disease or development of unacceptable toxicity. For CNS progression only, the Panel recommends continuing systemic therapy and treating brain metastases with RT. Additional subsequent systemic therapy (e.g., third line) can be considered if patients are still PS 0–2. The Panel recommends response assessment using CT with contrast of the C/A/P after every 2 to 3 cycles. Dose reduction or growth factor support should be considered for patients with a PS of 2.

Tarlatamab-dlle

Initial phase 1 and 2 trials in patients heavily pretreated for relapsed/refractory SCLC showed promising durability and acceptable safety profile with tarlatamab, a bispecific T-cell engager antibody that targets delta-like ligand 3 (DLL3) and CD3.¹⁹⁵ DLL3 is abnormally expressed on the surface of the majority of SCLC cells (85–94%) and inhibits Notch. Because of the favorable benefit to risk ratio in patients who received 10 mg versus 100 mg, the lower dose has been chosen for subsequent studies. In the phase 3 trial, patients were randomly assigned to receive tarlatamab (n = 254) or chemotherapy (n = 255, topotecan or lurbinectedin depending on the site). Tarlatamab-dlle significantly improved median OS compared to the chemotherapy group (13.6 vs 8.3 months).¹⁹⁶ OS in the tarlatamab group was 76% at 6 months and 53% at 12 months as compared with 62% at 6 months and 37% at 12 months in the chemotherapy group. Grade 3 or higher adverse events were lower with tarlatamab compared to chemotherapy (54% vs 80%). The most common treatment-related adverse events in the tarlatamab group included neutropenia, lymphopenia, lymphocyte count decrease, fatigue, neutrophil count decrease, hyponatremia, and anemia. Additionally, cytokine release syndrome were observed in 56% of the patients treated with tarlatamab. To manage unique toxicities with tarlatamab-dlle administration, the Panel refers to the NCCN Guidelines for Management of Car T-Cell and Lymphocyte Engager-Related Toxicities (available at www.NCCN.org). The Panel recommends tarlatamab-dlle (10 mg dosing schedule) as a category 1, preferred regimen for subsequent therapy.

Irinotecan

In a phase 2 study, disease responded to irinotecan in 47% of patients (95% CI, 21.4%–1.9%) with relapsed or refractory SCLC.¹⁹⁷ Myelosuppression, diarrhea, and pulmonary toxicity were reported in patients receiving irinotecan.¹⁹⁷ Another phase 2/3 trial showed that irinotecan and topotecan had comparable activity in patients with relapsed or refractory SCLC and metastasis.¹⁹⁸ The Panel recommends irinotecan as a subsequent therapy option (preferred) for patients with SCLC. The Panel added a consideration for irinotecan for patients with CNS disease.

Lurbinectedin

A phase 2 basket trial assessed lurbinectedin (3.2 mg/m² every 3 weeks) as second-line therapy in 105 patients with SCLC who previously received etoposide/platinum; 57% of patients had not received chemotherapy for ≥3 months.¹⁹⁹ The overall response rate with lurbinectedin was 35% (95% CI, 26.2%–45.2%). The response rate ranged from 22% (95% CI, 11.2%–37.1%) to 45% (95% CI, 32.1%–58.4%). Common grade 3–4 adverse events included anemia, leukopenia, neutropenia, and thrombocytopenia. There were no reported treatment-related deaths. In a subset analysis of this trial, the



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

overall response rate associated with lurbinectedin was 60% (95% CI, 36.1%–86.9%). The median OS was 16.2 months (95% CI, 9.6–upper level not reached).²⁰⁰ After 1 year, 60.9% of patients were alive and after 2 years, 27.1% were alive. Common grade 3–4 adverse events included neutropenia, anemia, thrombocytopenia, fatigue, and increased liver function tests (including gamma-glutamyl transferase [GGT], alanine aminotransferase [ALT] and alkaline phosphatase [ALP]). The FDA granted approval for lurbinectedin at a dose of 3.2 mg/m² every 3 weeks based on clinical trial data.¹⁹⁹ The Panel recommends lurbinectedin as a subsequent therapy option (preferred) and if not previously used for patients with SCLC.^{199,200}

Platinum-Based Therapy

A phase 3 randomized trial assessed carboplatin/etoposide compared with oral topotecan in 162 patients with SCLC relapse after >3 months on first-line etoposide/platinum therapy.²⁰¹ The median PFS was 4.7 months (90% CI, 3.9–5.5) in the carboplatin/etoposide group versus 2.7 months (90% CI, 2.3–3.2) in the oral topotecan group (HR, 0.57; 90% CI, 0.41–0.73; *P* = .004). Grade 3–4 adverse events included thrombocytopenia, neutropenia, anemia, febrile neutropenia, and asthenia. In the topotecan group, two treatment-related deaths occurred; no deaths occurred in the carboplatin/etoposide group. The Panel recommends subsequent therapy with platinum-based regimens based on the clinical trial data.^{201–205} The Panel maintains that rechallenging with the original regimen with or without immunotherapy, or similar platinum-based regimen, is recommended as an option (preferred) if there has been prolonged disease-free time.^{201–205}

Topotecan

Topotecan is also recommended as a subsequent therapy option based on clinical trial data.²⁰¹ A randomized phase 3 trial for subsequent treatment compared single-agent IV topotecan with CAV in patients with SCLC relapse at least 60 days after therapy.²⁰⁶ Both arms had similar response rates (topotecan, 24.3%; CAV, 18.3%) and survival (25.0 vs. 24.7 weeks). Compared to CAV, topotecan caused less grade 4 neutropenia (37.8% vs. 51.4%; *P* < .001) and improved symptoms of dyspnea, anorexia, hoarseness, and fatigue. There are conflicting data regarding the usefulness of weekly topotecan in patients with relapsed SCLC.^{207,208} Many practicing oncologists have noted excessive toxicity when using 1.5 mg/m² of IV topotecan for 5 days, and studies suggest that an attenuated dose may be equally efficacious with lower toxicity.²⁰⁹ In another phase 3 trial, oral topotecan improved OS compared with best supportive care (26 vs. 14 weeks).²¹⁰ The efficacy and toxicity of oral and IV topotecan seem to be similar and therefore either route may be used.^{210,211} The Panel recommends oral or IV topotecan as a subsequent therapy option (preferred) for patients with SCLC.^{201,206,210,212,213}

Other Recommended Subsequent Therapy Options

CAV

A phase 3 trial (JCOG0605) from Japan in patients with sensitive, relapsed SCLC reported that the combination of cisplatin/etoposide/irinotecan improved survival compared with topotecan (median survival, 18.2 vs. 12.5 months; HR, 0.67; 90% CI, 0.51–0.88; *P* = .0079). However, the toxicity of this regimen was significant.²¹⁴ The Panel recommends CAV as a subsequent therapy option based on clinical trial data.²⁰⁶



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Docetaxel and Paclitaxel

Docetaxel and paclitaxel, which belong to the taxane class of drugs, have been assessed in patients with refractory or relapsed SCLC. In a phase 2 study in patients with refractory or relapsed SCLC, 24% of patients' disease responded to paclitaxel.²¹⁵ Grade 3–4 toxicity included neutropenia, infection, rash, neuropathy, and pulmonary toxicity. Another phase 2 study of paclitaxel in patients with refractory SCLC yielded a response rate of 29% (95% CI, 12%–51%).²¹⁶ A retrospective study in 185 patients showed a response rate of 17% during third-line or fourth-line therapy with paclitaxel. Toxicity rates were similar in patients with PS 2 compared with PS 0–1 (63% vs. 62%).²¹⁷ Twenty-five percent of patients' disease responded to docetaxel in a phase 2 trial in patients with previously treated SCLC. Reported toxicities included neutropenia and asthenia.²¹⁸

Gemcitabine

Gemcitabine was assessed in a phase 2 trial in 42 patients with sensitive or refractory SCLC.²¹⁹ The median survival was 7.1 months. The overall objective response rate was 11.9%. One patient with refractory SCLC and four patients with sensitive SCLC had disease respond to therapy. Grade 3–4 toxicities included neutropenia (27%), thrombocytopenia (27%), neurologic toxicity (14%), and pulmonary toxicity (9%). Another phase 2 trial assessed gemcitabine in 38 patients with resistant SCLC that progressed within 3 months of last therapy.²²⁰ The median survival was 17 weeks (range, 4–84). The response rate was 13% (95% CI, 6%–27%); there were five partial responses and no complete responses. Grade 3 toxicities included thrombocytopenia (29%) and leukopenia (18%).

Nivolumab and Pembrolizumab

ICIs have been evaluated in patients with relapsed SCLC.^{221–224} CheckMate 032, a phase 1/2 trial, assessed nivolumab alone ($n = 147$) or various doses of ipilimumab + nivolumab ($n = 96$) for relapsed SCLC.^{221,222} Response rates were 11.6% for nivolumab and 21.9% for ipilimumab + nivolumab.²²¹ The 12- and 24-month OS rates were similar (nivolumab, 30.5% and 17.9%; ipilimumab + nivolumab, 30.2% and 16.9%, respectively). Grade 3–4 adverse events were 12.9% for nivolumab alone and 37.5% for ipilimumab + nivolumab. In patients receiving nivolumab alone, the most common grade 3 or 4 treatment-related adverse events were pneumonitis and increased levels of lipase and aspartate aminotransferase.

CheckMate 331, a randomized phase 3 trial, assessed nivolumab monotherapy versus amrubicin or topotecan in 569 patients with relapsed SCLC.²¹³ Data show that OS was 7.5 months in patients receiving nivolumab versus 8.4 months in those receiving chemotherapy (HR, 0.86; 95% CI, 0.72–1.04; $P = .11$).²¹³ OS was similar regardless of PD-L1 levels. Response rates were 13.7% for nivolumab compared with 16.5% for chemotherapy. Treatment-related deaths occurred in two patients receiving nivolumab and in three patients receiving chemotherapy. Fewer grade 3–4 adverse events occurred in patients receiving nivolumab compared with chemotherapy (14% vs. 73%, respectively). A study reported that third-line therapy with nivolumab was associated with longer survival (5.7 months; 95% CI, 3.5–8.0) compared with other treatments such as paclitaxel or topotecan (3.8 months; 95% CI, 2.8–4.9; HR, 0.63; 95% CI, 0.44–0.90).²²⁵ The 1-year OS rate is 28% with nivolumab versus 4% with the other treatments.

A combined analysis of two trials, phase 1b (KEYNOTE-028) and phase 2 (KEYNOTE-158), evaluated the activity of pembrolizumab in 83 patients with relapsed SCLC who had received ≥ 2 lines of therapy. PD-L1 positivity ($\geq 1\%$) was found in 56.6% of tumors and 84.3% patients did not have brain



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

metastases.²²⁶ This analysis reported an objective response rate of 19.3% (95% CI, 11.4%–29.4%). The median OS was 7.7 months (95% CI, 5.2–10.1); the estimated 12-month OS rate was 34.3%. Both OS and response rate were higher in those who were PD-L1 positive; however, one patient with a complete response had a tumor that was PD-L1 negative. Grade 3 or 5 adverse events occurred in 9.6% (8/83) of patients; two patients died from treatment-related adverse events (pneumonitis and encephalitis).

The FDA has withdrawn the subsequent therapy indications for nivolumab or pembrolizumab for patients with relapsed SCLC, because phase 3 randomized trial data did not show an improvement in OS.^{213,221-223,226,227} The Panel decided that nivolumab or pembrolizumab are just as effective as (sometimes better than) and less toxic than the other subsequent therapy options.^{213,225} In addition, many agents recommended as subsequent therapy options for patients with SCLC do not have an FDA indication in this setting, but data show that they are effective. Patients who have not previously received ICIs may benefit from subsequent therapy with nivolumab or pembrolizumab. Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV nivolumab. Similarly, pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab. The use of nivolumab and pembrolizumab is discouraged in patients whose disease progresses while on maintenance atezolizumab or durvalumab as part of first-line therapy. There are no data to suggest that giving patients subsequent ICIs is effective if their disease previously progressed on other ICIs.

Health care providers should be aware of the spectrum of potential immune-mediated adverse events unique to immunotherapeutic agents, such as nivolumab and pembrolizumab, know how to manage these events, and discuss possible side effects with patients (see the NCCN Guidelines for Management of Immune Checkpoint Inhibitor-Related Toxicities, available at www.NCCN.org).^{228,229} For patients with immune-mediated adverse events, high-dose corticosteroids are generally recommended based on the severity of the reaction. Nivolumab or pembrolizumab should be withheld or discontinued for severe or life-threatening immune-mediated adverse events (see prescribing information).

Etoposide

Oral etoposide was assessed in a phase 2 trial in 22 patients with recurrent SCLC.²³⁰ Ten patients (45%; 95% CI, 27%–65%) had a complete or partial response. Median survival was ≥ 3.5 (range, 1 to 15+). Five patients were hospitalized because of neutropenia and fever. Two patients died from sepsis. Another phase 2 trial assessed oral etoposide in 26 patients with refractory SCLC.²³¹ The overall response rate was 23%; there was one complete response and five partial responses.

Temozolomide

Data suggest that temozolomide may be useful for patients with SCLC, especially those with brain metastases and methylated O⁶-methylguanine-DNA methyltransferase (MGMT).²³²⁻²³⁴ Temozolomide was assessed in a phase 2 trial in patients with relapsed or refractory SCLC. In patients with sensitive SCLC, the overall response rate was 23% (95% CI, 12%–37%). The response rate improved for patients with methylated MGMT compared to those with unmethylated MGMT (38% vs. 7%; $P = .08$).



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

Radiation Therapy

The American Radium Society appropriate use criteria and the American Society for Radiation Oncology (ASTRO) guidelines are useful resources.²³⁵⁻²³⁸ The *Principles of Radiation Therapy* section in the NSCLC algorithm may also be useful (see the NCCN Guidelines for Non-Small Cell Lung Cancer, available at www.NCCN.org).

Thoracic Radiation Therapy for Limited-Stage SCLC

Achieving long-term local control using conventional chemoradiotherapy for patients with limited-stage SCLC remains a challenge. The addition of thoracic RT has improved survival for patients with limited-stage SCLC. Meta-analyses that include >2000 patients show that thoracic radiation for limited-stage SCLC yields a 25% to 30% reduction in local progression, and a corresponding 5% to 7% improvement in 2-year OS compared with chemotherapy alone.^{98,99} A phase 3 trial reported 5-year OS of >30%, which is close to outcomes of locally advanced NSCLC of similar stage.¹⁰⁵

Timing of Radiation with Chemotherapy

Optimal thoracic RT is impacted by several factors, including the timing of chemotherapy and RT (concurrent vs. sequential), timing of RT (early vs. late), RT target volume (original tumor volume vs. shrinking field as the tumor responds), radiation dose, and RT fractionation. Early concurrent chemoradiotherapy is recommended for patients with limited-stage SCLC based on randomized trials. A randomized phase 3 trial by the Japanese Cooperative Oncology Group (9104) assessed sequential versus concurrent thoracic RT combined with cisplatin/etoposide for 231 patients with limited-stage SCLC. OS was 27.2 months for those receiving concurrent chemoradiation versus 19.7 months for those receiving sequential chemoradiation ($P = .097$).¹¹⁰ Patients receiving concurrent chemoradiation had more severe hematologic toxicity. Severe esophagitis occurred in 9% of patients receiving concurrent chemoradiation and 4% receiving sequential chemoradiation.

Several systematic reviews and meta-analyses on the timing of thoracic RT in limited-stage SCLC have reported that early concurrent chemoradiation results in a small, but significant improvement in OS compared with late concurrent or sequential chemoradiation.^{239,240} A randomized phase 3 trial (by the National Cancer Institute of Canada) compared RT beginning with either cycle 2 or cycle 6 of chemotherapy and showed that early RT was associated with improved local and systemic control and longer survival.²⁴¹ Meta-analysis in patients with limited-stage SCLC showed that survival improved with rapid completion of the chemoradiotherapy regimen (start of any chemotherapy until the end of RT).²⁴² Another meta-analysis of individual patient data from 12 trials ($n = 2668$) reported that early concurrent chemoradiation, associated with increase in acute esophagitis, had higher 5-year OS (HR, 0.79; 95% CI, 0.69–0.91) compared with late concurrent chemoradiation.²⁴³

Radiation Fractionation

The ECOG/Radiation Therapy Oncology Group (RTOG) compared accelerated to conventionally fractionated RT with cisplatin/etoposide.²⁴⁴ In this trial, 412 patients with limited-stage SCLC were treated with concurrent chemoradiation using a total dose of 45 Gy delivered either twice daily over 3 weeks (accelerated) or once daily over 5 weeks (conventional). Median OS was 23 versus 19 months ($P = .04$), and 5-year survival rates were 26% versus 16%



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

in the accelerated and conventional RT arms, respectively.²⁴⁴ A higher incidence of grade 3–4 esophagitis was seen with the accelerated regimen compared with the conventional regimen (when including elective mediastinal irradiation as was done during this era).²⁴⁴ A criticism of this trial in retrospect is that the 45 Gy conventional regimen provided suboptimal dose intensity compared to modern conventionally fractionated regimens using higher total doses.

CONVERT, a phase 3 randomized trial, compared accelerated 45 Gy (given twice daily over 3 weeks) with higher dose conventionally fractionated 66 Gy (given once daily over 6.5 weeks) in 547 patients with limited-stage SCLC.¹⁰⁵ Median OS was similar between the 2 groups (30 vs. 25 months, accelerated vs. conventional). However, the CONVERT trial was not powered to show equivalence. Toxicity was generally similar between the arms; however, patients receiving accelerated 45 Gy had more grade 4 neutropenia compared with those receiving conventional 66 Gy (49% vs. 38%; $P = .05$).

CALGB 30610 (Alliance)/RTOG 0538, a randomized phase 3 trial, compared high-dose conventional 70 Gy (once daily over 7 weeks) with accelerated 45 Gy (twice daily over 3 weeks) in 638 patients with limited-stage SCLC.²⁴⁵ Originally, there was a 61.2 Gy concomitant boost group in this trial, but it was removed based on a planned interim toxicity analysis.²⁴⁶ Median OS was 30.5 months in the conventional 70 Gy arm versus 28.5 months in the accelerated 45 Gy arm (HR, 0.94; 95% CI, 0.75–1.17; $P = .591$). There were 5 deaths in the conventional 70 Gy arm and 2 deaths in the accelerated 45 Gy arm. OS and toxicity were similar, although it did not demonstrate superiority of the high-dose conventional arm. The conventional 70 Gy arm had better quality-of-life (QOL) scores at 3 weeks with patients reporting it to be more convenient.²⁴⁵

A randomized phase 2 trial assessed concurrent chemoradiation with two similarly accelerated regimens, 42 Gy given as once-daily fractions over 3 weeks compared with 45 Gy given as twice-daily fractions also over 3 weeks in 157 patients with limited-stage SCLC.²⁴⁷ The OS curves overlapped with median OS of 18.8 months in the once-daily arm and 25.1 months in the twice-daily arm ($P = .61$). A retrospective study assessed concurrent chemoradiation with accelerated 40 Gy in 3 weeks given as once-daily fractionation in 68 patients with limited-stage SCLC.²⁴⁸ The median survival was 28 months, which is comparable to outcomes of similarly accelerated twice-daily fractionation.

Despite multiple trials, high-dose conventional RT (66–70 Gy over 6.5–7 weeks) has not been shown to be superior to standard-dose accelerated RT (45 Gy over 3 weeks), though survival and toxicity appear to be comparable.^{105,245,249} Studies have also evaluated the potential advantage of high-dose accelerated RT. Two randomized phase 2 trials compared high-dose accelerated RT with standard-dose accelerated RT. One trial in 182 patients with limited-stage SCLC compared concurrent chemoradiation with high-dose accelerated 65 Gy given as once-daily fractions over approximately 5 weeks with standard-dose accelerated 45 Gy given as twice-daily fractions over 3 weeks.²⁵⁰ Estimated PFS was 17.2 months in the high-dose group versus 13.4 months in the standard-dose group ($P = .031$). OS was 39.3 months in the high-dose group versus 33.6 months in the standard-dose group ($P = .137$). Grade 3 or higher esophagitis (high-dose, 17.4% vs. standard-dose, 15.3%), grade 3 or higher pneumonitis (high-dose, 3.3% vs. standard-dose, 2.4%), and treatment-related deaths (high-dose, 2.2% vs. standard-dose, 1.2%) were similar in each group. The second trial in 176 patients with limited-stage SCLC compared concurrent chemoradiation using high-dose accelerated RT with 60 Gy given as twice-daily fractions over 4 weeks with accelerated standard-dose 45 Gy given as twice-daily fractions over 3 weeks.²⁵¹ After 2 years, 74.2% of patients were alive in the 60 Gy group (95% CI,



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

63.8%–82.9%) versus 48.1% in the 45 Gy group (95% CI, 36.9%–59.5%). The 5-year survival rates were 39.3% (60 Gy) and 28.4% (45 Gy). Three treatment-related deaths occurred in each group. A randomized phase 3 trial demonstrated a significant survival advantage of high-dose accelerated RT given as 54 Gy simultaneous integrated boost to the gross tumor volume in 30 twice-daily fractions over 3 weeks compared to standard 45 Gy in 30 twice-daily fractions over 3 weeks (60.7 months vs. 39.5 months, $P = .003$) in patients with limited-stage disease.²⁵² Radiotherapy-related toxicities were similar between the two groups. This outcome was similar to conclusions from a retrospective study that investigated a real-world cohort study of patients with limited-stage SCLC and found that median PFS was higher in the high-dose accelerated RT (29.7 months; 95% CI, 15.9–not reached]) compared to standard RT (15.0 months; 95% CI, 11.8–30.4).²⁵³

In general, accelerated RT (whether given once or twice daily) is superior to similar doses of conventionally fractionated RT and comparable to higher dose conventionally fractionated RT. The Panel recommends accelerated 45 Gy given as twice-daily fractions over 3 weeks (category 1) or conventionally fractionated 66 to 70 Gy given as once-daily fractions over 6.5 to 7 weeks as acceptable options depending on individual circumstances^{105,245,249} since the twice-daily thoracic RT can be logistically challenging for patients and RT centers. The Panel maintains that higher doses of 66 to 70 Gy are preferred if using once-daily fractionation.²⁴⁹

Patients with no disease progression after chemoradiation and who receive adjuvant immunotherapy with durvalumab have a significant survival advantage.¹¹⁵ High-dose accelerated RT appears superior to standard-dose accelerated RT when delivered concurrently with chemotherapy.²⁵⁰⁻²⁵² The lack of increased toxicity with these regimens suggests that the historical determination of 45 Gy in 3 weeks as the maximum tolerated dose is no longer applicable with modern conformal radiotherapy techniques and the omission of elective nodal irradiation. The role of high-dose accelerated RT in the context of the new standard of adjuvant immunotherapy remains to be determined.

Technical Considerations

The minimum technical requirement for thoracic irradiation is CT-planned three dimension (3D)-conformal RT. Intensity-modulated RT (IMRT) is preferred over 3D-conformal external-beam RT (EBRT) because of lower toxicity. The normal tissue constraints used for NSCLC are appropriate for SCLC when using similar RT doses (see *Principles of Radiation Therapy* in the algorithm and the NCCN Guidelines for Non-Small Cell Lung Cancer, available at www.NCCN.org).²⁵⁴⁻²⁵⁹ More advanced technologies, such as four dimensional (4D)-CT and proton therapy, may also be appropriate to limit normal tissue toxicity. The radiation target volumes can be defined on the FDG-PET/CT scan obtained at the time of RT planning, as well as any positive biopsies, using definitions in Reports 50 and 62 from the International Commission on Radiation Units & Measurements.²⁵⁵ However, the pre-chemotherapy FDG-PET/CT scan should be reviewed to include the original involved lymph node regions in the treatment fields if chemotherapy begins before RT.^{260,261} When using accelerated schedules (e.g., 3–5 weeks), the spinal cord constraints from the CALGB 30610/RTOG 0538 protocol can be used as a guide.^{249,262-264}



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

SABR

Emerging data suggest that SABR (stereotactic ablative radiotherapy, also known as stereotactic body RT (SBRT)) is effective for patients with clinical limited-stage I–IIA (T1–2, N0) SCLC, especially for medically inoperable circumstances or in those who refuse surgery.^{12,265–270} A meta-analysis of 7 studies (399 patients) summarized outcomes in patients with limited-stage SCLC who received SABR; 94% of the patients in this study were deemed inoperable.²⁷⁰ Forty-four percent of patients received chemotherapy and 13.8% of patients received PCI. OS was 86% (95% CI, 74%–95%) at 1 year and 64% (95% CI, 46%–80%) at 2 years. Nodal recurrence rate was 18% (95% CI, 7.5%–31%) and distal recurrence rate was 27% (95% CI, 7.4%–53%). Grade 3 toxicity was observed in 1.4% of patients (95% CI, 0%–5.3%). A multicenter analysis of 74 patients suggested that the addition of chemotherapy typically after SABR improves survival for patients with clinically limited-stage SCLC.^{14,271} Most of these patients had FDG-PET staging but not pathologic nodal staging. Patients who received chemotherapy after SABR had a median OS of 31.4 months versus 14.3 months for those who received SABR alone ($P = .02$).

An analysis of 2107 patients with histologically confirmed T1–T2, N0, M0 from the National Cancer Database found that 7.1% had upfront SABR followed by adjuvant chemotherapy and 92.9% had concurrent chemoradiation.¹⁰ Compared with patients receiving upfront concurrent chemoradiation, those receiving SABR were often older, had T1 disease, and were treated recently in academic medical settings. Median survival was 29.2 months in those receiving SABR/chemotherapy versus 31.2 months in those receiving chemoradiation ($P = .77$). Both ASTRO and the American Radium Society recommend SABR followed by adjuvant chemotherapy as an option for medically inoperable patients with clinically limited-stage I–IIA SCLC (T1–2, N0).^{236,237}

The Panel recommends SABR followed by systemic therapy as an option for select patients with clinically limited-stage I–IIA (T1–2, N0) SCLC who are medically inoperable or decline surgery.^{12,271} The Panel added a caveat that systemic therapy may be initiated first, if time to initiation of SABR will be prolonged. The NCCN Guidelines for Non-Small Cell Lung Cancer provide detailed recommendations for SABR that may be useful for SCLC (see *Principles of Radiation Therapy* in the NCCN Guidelines for Non-Small Cell Lung Cancer, available at www.NCCN.org).

Radiation for Extensive-Stage SCLC

Sequential Thoracic Radiation for Extensive-Stage SCLC

A randomized trial by Jeremic et al assessed sequential (consolidative) thoracic RT in patients experiencing a complete response at distant metastatic sites after 3 cycles of cisplatin/etoposide.²⁷⁴ Patients were randomized to receive either 1) further cisplatin/etoposide; or 2) accelerated hyperfractionated RT (i.e., 54 Gy in 36 fractions over 18 treatment days) in combination with carboplatin/etoposide.²⁷⁴ The addition of RT resulted in improved median OS (17 vs. 11 months). The Dutch CREST trial, a phase 3 randomized trial in patients with extensive-stage SCLC, reported that the addition of consolidative thoracic RT (30 Gy in 10 fractions) did not improve the primary endpoint of 1-year OS (33% vs. 28%; $P = .066$). However, a secondary analysis found improvement in 2-year OS (13% vs. 3%; $P = .004$) and 6-month PFS compared with patients who did not receive consolidative thoracic RT.²⁷⁵ A trial involving 32 patients who received consolidative thoracic RT reported that only 16% of patients had symptomatic chest recurrences.²⁷⁶ Consolidative



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

thoracic RT appears to mainly benefit patients with residual thoracic disease after chemotherapy (without immunotherapy) and low-bulk extrathoracic metastatic disease that has responded to systemic therapy.²⁷⁷ Based on the limited data, the American Radium Society recommends that consolidative thoracic RT be considered for select patients with extensive-stage SCLC.²³⁵ International experts (International Association for the Study of Lung Cancer [IASLC] and European Society for Radiotherapy and Oncology [ESTRO]) recommend consolidative thoracic RT in select patients with stage IV SCLC (that has responded to first-line chemotherapy) and limited extrathoracic tumor burden.²⁷⁸

The Panel recommends that consolidative thoracic RT be considered in select patients with low-bulk, extrathoracic, metastatic, extensive-stage disease who have a complete or near complete response after initial systemic therapy before maintenance immunotherapy.^{235,274,275} Sequential thoracic RT can be considered for selected patients, during or before maintenance immunotherapy; however, there are limited data on optimal sequencing. The benefit of thoracic RT in the context of chemoimmunotherapy is under evaluation in the RAPTOR/NRG LU007 trial (NCT04402788).

Radiotherapy for brain metastases

Prophylactic Cranial Irradiation (PCI)

Intracranial metastases occur in >50% of patients with SCLC. Randomized studies show that PCI is effective in decreasing the incidence of cerebral metastases, but most individual studies did not have sufficient power to show a meaningful survival advantage.²⁷⁹ Several meta-analyses suggest that PCI after complete resection may benefit patients with pathologic stage IIB or stage III SCLC.²⁸⁰⁻²⁸² A meta-analysis of all randomized PCI trials reported a reduction in the 3-year incidence of brain metastases, from 58.6% in the control group to 33.3% in the PCI-treated group.²⁸¹ Thus, PCI seems to prevent, and not simply delay, the emergence of brain metastases. This meta-analysis also reported an increase in 3-year OS from 15.3% in the control group to 20.7% in the PCI group.²⁸¹ Although the number of patients with extensive-stage SCLC was small, the observed benefit was similar in patients with both limited-stage and extensive-stage SCLC. A retrospective study of patients with limited-stage SCLC also found that PCI increased survival at 2, 5, and 10 years compared with those who did not receive PCI.²⁸³

None of the abovementioned studies in limited-stage SCLC used MRI of the brain with FDG-PET scans for staging. A retrospective study ($n = 184$) with limited-stage SCLC assessed PCI versus no PCI in patients whose disease responded to chemoradiotherapy, and who had no brain metastases on MRI imaging before and after primary treatment.²⁸⁴ In patients receiving PCI, median OS was 26 months (range, 19.4–32.6 months) versus 14 months for those without PCI (range, 11.4–16.6 months; $P < .0001$). A more modern retrospective study included 49 patients with limited-stage SCLC who were staged with brain MRI before treatment.²⁸⁵ The median OS was 55 months in patients with limited-stage SCLC who received PCI versus 24 months in those who did not receive PCI ($P < .05$). At 1 year, the probability of developing symptomatic brain metastases was 4% in patients with limited-stage SCLC who received PCI versus 22% in those who did not receive PCI ($P < .05$). To date, there is a lack of studies on PCI in the context of adjuvant immunotherapy to treat limited-stage SCLC.

For patients with extensive-stage SCLC, but without brain metastases, a large retrospective analysis of 4257 patients showed that PCI improved median OS compared with no PCI (13.9 vs. 11.1 months; $P < .0001$).²⁸⁶ Another analysis of patients with extensive-stage SCLC ($n = 397$) reported that PCI



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

improved OS compared with no PCI (13.5 vs. 8.5 months; HR, 0.55; 95% CI, 0.39–0.77; $P = .0005$); however, these patients did not receive routine brain imaging surveillance.²⁸⁷

The EORTC performed a randomized trial that assessed PCI versus no PCI in 286 patients with extensive-stage SCLC that had responded to initial chemotherapy. PCI decreased symptomatic brain metastases (14.6% vs. 40.4%) and increased the 1-year survival rate (27.1% vs. 13.3%) compared with controls.²⁸⁸ However, the study did not require brain imaging prior to PCI and did not standardize the PCI dose or fractionation. Conflicting data from a randomized phase 3 trial found that median OS is not improved in patients receiving PCI compared with MRI surveillance (11.6 months; 95% CI, 9.5–13 vs. 13.7 months; 95% CI, 10.2–16.4) (HR, 1.27; 95% CI, 0.96–1.68; $P = .094$).²⁸⁹ In this trial, patients were required to have an MRI to confirm that they did not have brain metastases prior to PCI, and the PCI regimen was standardized at 25 Gy in 10 fractions. The study also required close MRI surveillance in patients to allow for early treatment of brain metastases. Based on the limited data, the American Radium Society recommends either PCI or brain MRI surveillance for patients with extensive-stage SCLC and without brain metastases.²³⁵ Early prevention of brain metastases does not directly correlate to greater QOL or neurocognitive function in these trials. A review article summarizing the use of PCI in SCLC notes that prospective trials demonstrating the benefits of PCI over MRI surveillance in the modern era are needed.²⁹⁰ A randomized trial (SWOG S1827/MAVERICK) is currently addressing this exact question and assessing brain MRI surveillance alone compared to brain MRI surveillance plus PCI for patients with limited- and extensive-stage SCLC. Late neurologic sequelae have been attributed to PCI, particularly in studies using fractions >3 Gy and/or administering PCI concurrently with chemotherapy.^{184,291,292} Thus, PCI is not recommended for patients with poor PS (3–4) or impaired neurocognitive function.^{96,293} PCI has also been associated with chronic neurotoxicity in patients who are aged ≥60 years.^{183,185}

The Panel has gradually revised the adjuvant recommendations for patients whose disease showed a complete or partial response after primary treatment based on conflicting clinical trial data and concerns about the neurocognitive toxicity of PCI. Before a decision is made to administer PCI, a balanced discussion is necessary between the patient and physician.^{184,294} The Panel recommends considering PCI for patients with limited-stage SCLC that has achieved a complete or partial response.^{96,281,288} PCI should be administered prior to consolidation durvalumab therapy. The Panel maintains that the benefit of PCI is unclear in patients with very-early-stage SCLC (pathologic limited-stage I [T1–2a, N0, M0]) who have had definitive therapy (ie, surgery, SABR). These patients have a lower risk of developing brain metastases than patients with more advanced limited-stage SCLC and may not benefit from PCI.^{271,280,295} The Panel recommends MRI brain surveillance for all patients with limited-stage SCLC who do not receive PCI, as well as for post-treatment surveillance in those receiving PCI. In patients with extensive-stage SCLC, the Panel recommends MRI brain surveillance with or without consideration of PCI based on the conflicting trial results from Japan and the EORTC.^{288,289} Brain imaging surveillance for metastases is recommended using either MRI (preferred) or CT with contrast in patients who are unable to undergo MRI.²⁸⁹

Technical Considerations

Higher PCI doses (e.g., 36 Gy) increased mortality and toxicity compared with lower doses (25 Gy).^{183,296} Therefore, the preferred dose for PCI is 25 Gy in 10 daily fractions (2.5 Gy/fraction).^{281,288,296} A shorter course of PCI may be appropriate (e.g., 20 Gy in 5 fractions) for selected patients with



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

extensive-stage SCLC.²⁸⁸ PCI should not be given concurrently with chemotherapy, and high total RT dose (>30 Gy) should be avoided because of the increased risk of neurotoxicity.¹⁸³ After the acute toxicities of initial systemic therapy have resolved, PCI can be administered. When given after the completion of chemotherapy and at a low dose per fraction, PCI may cause less neurologic toxicity. Fatigue, headache, and nausea/vomiting are the most common acute toxic effects after PCI.^{293,296}

The Panel recommends that memantine be considered for patients receiving PCI or therapeutic whole-brain irradiation. Memantine is a N-methyl-D-aspartate (NMDA) receptor antagonist that may delay cognitive dysfunction in patients receiving brain RT.²⁹⁷ Patients receiving memantine have a longer time before cognitive decline (HR, 0.78; 95% CI, 0.62–0.99; $P = .01$). NRG Oncology CC001, a phase 3 randomized trial, assessed HA whole-brain IMRT/memantine compared with conventional whole-brain RT/memantine in patients with brain metastases who were not diagnosed with SCLC.²⁹⁸ Cognitive preservation and patient-reported outcomes were improved with HA-IMRT (HR, 0.74; 95% CI, 0.58–0.95; $P = .02$). However, conflicting data have been reported with HA-PCI versus conventional PCI. PREMER, a phase 3 randomized trial, reported improved cognitive preservation with HA-PCI.²⁹⁹ However, another phase 3 randomized trial reported no differences in cognition with HA-PCI.³⁰⁰ A large randomized trial (NRG CC003) assessed HA-PCI versus conventional PCI and found that addition of HA to PCI prevented neurocognitive functioning failure (adjusted HR, 0.78; 95% CI, 0.61–0.99; $P = .039$).³⁰¹ Overall, hippocampal avoidance should be considered when administering whole brain radiotherapy for PCI or treatment of brain metastases.

Brain metastases

Brain metastases have conventionally been treated with whole brain RT (WBRT) in patients with SCLC due to the frequent occurrence of multiple metastases. The recommended dose for WBRT is 30 Gy in 10 daily fractions.³⁰⁵ Also, see the NCCN Guidelines for Non-Small Cell Lung Cancer, available at www.NCCN.org. IMRT, SABR, or stereotactic radiosurgery (SRS) may be appropriate for select patients (e.g., those whose tumors are in close proximity to organs at risk).

A retrospective multicenter cohort study assessed stereotactic radiosurgery (SRS) versus WBRT in 710 patients with SCLC who had a limited number of brain metastases; OS was 6.5 months (95% CI, 5.5–8.0) for SRS and 5.2 months (95% CI, 4.4–6.7) for whole-brain RT ($P = .003$).³⁰⁶ A meta-analysis of nine observational studies (1638 patients) also reported favorable lesion control and survival outcomes with SRS versus whole-brain RT.³⁰⁷ Phase 2 and phase 3 trials in patients with brain metastases, who did not receive any prior brain directed radiation, investigated whether SRS improves symptoms or neurological death compared to WBRT (historical controls or in a randomized controlled trial).^{272,273} Neurologic death in the phase 2 trial was defined as marked, progressive, radiologic brain progression accompanied by corresponding neurologic symptomatology without systemic disease progression or systemic symptoms of a life-threatening nature. This trial studying patients with small cell lung cancer and 1-10 brain metastases showed that SRS resulted in lower neurologic death compared to historical WBRT controls (11% vs 17.5%). The phase 3 trial showed that compared to hippocampal-avoidance (HA) WBRT in patients with mostly non-small cell lung cancer, breast, melanoma, or others with 5 to 20 brain metastases, SRS was associated with less severe symptoms and interference with daily functioning.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

A randomized trial (NRG CC009) is comparing SRS to HA whole-brain IMRT/memantine in this setting. The Panel recommends that SRS may be used for selected patients with limited brain metastases based on available data, pending outcomes of the ongoing trials.³⁰⁶ In patients who develop brain metastases after PCI, SRS (preferred) or repeat WBRT (in carefully selected patients) may be considered.^{308,309} For patients with a better prognosis (e.g., ≥ 4 months), HA whole-brain IMRT/memantine is preferred to conventional WBRT/memantine because it produces less of a decrease in cognitive function. However, patients with metastases within 5 mm of the hippocampi, leptomeningeal metastases, and other high-risk features were not eligible for HA whole-brain IMRT in the phase 3 NRG CC001 trial.²⁹⁸

Palliative Radiation Therapy

For patients with localized symptomatic sites of disease (i.e., painful bony lesions, spinal cord compression, obstructive atelectasis) or with brain metastases, RT can provide excellent palliation (see the algorithm and the NCCN Guidelines for Non-Small Cell Lung Cancer, available at www.NCCN.org).³⁰²⁻³⁰⁴ Orthopedic stabilization may be useful in patients at high risk for fracture because of osseous structural impairment. Because patients with SCLC often have a short life span, surgery is usually not recommended for spinal cord compression. Radiation dose and fractionation for extracranial metastases include 30 Gy in 10 fractions, 20 Gy in 5 fractions, or 8 Gy in 1 fraction based on common dose-fractionation regimens used for other solid tumors, and more definitive doses may be appropriate in selected circumstances.

Summary

In summary, the V.1.2027 NCCN Guidelines for Small Cell Lung Cancer have recommendations for diagnosis, evaluation, therapy, and surveillance for limited- and extensive-stage SCLC based on data from clinical trials and Panel expertise.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

References

1. Siegel RL, Kratzer TB, Wagle NS, et al. Cancer statistics, 2026. CA Cancer J Clin 2026;76:e70043. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41528114>.
2. Wells LE, Cohen S, Brennan B, et al. Epidemiology of SCLC in the United States From 2000 to 2019: A study utilizing the Surveillance, Epidemiology, and End Results Registry. JTO Clin Res Rep 2025;6:100799. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40104389>.
3. Pesch B, Kendzia B, Gustavsson P, et al. Cigarette smoking and lung cancer--relative risk estimates for the major histological types from a pooled analysis of case-control studies. Int J Cancer 2012;131:1210–1219. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22052329>.
4. Videtic GMM, Stitt LW, Dar AR, et al. Continued cigarette smoking by patients receiving concurrent chemoradiotherapy for limited-stage small-cell lung cancer is associated with decreased survival. J Clin Oncol 2003;21:1544–1549. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12697879>.
5. Leone FT, Evers-Casey S, Toll BA, Vachani A. Treatment of tobacco use in lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e61S–e77S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649454>.
6. Brenner B, Tang LH, Klimstra DS, Kelsen DP. Small-cell carcinomas of the gastrointestinal tract: a review. J Clin Oncol 2004;22:2730–2739. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15226341>.
7. Galanis E, Frytak S, Lloyd RV. Extrapulmonary small cell carcinoma. Cancer 1997;79:1729–1736. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9128989>.
8. Kalemkerian GP. Advances in the treatment of small-cell lung cancer. Semin Respir Crit Care Med 2011;32:94–101. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21500128>.
9. Stinchcombe TE, Gore EM. Limited-stage small cell lung cancer: current chemoradiotherapy treatment paradigms. Oncologist 2010;15:187–195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20145192>.
10. Verma V, Hasan S, Wegner RE, et al. Stereotactic ablative radiation therapy versus conventionally fractionated radiation therapy for stage I small cell lung cancer. Radiother Oncol 2019;131:145–149. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30773182>.
11. Haque W, Verma V, Polamraju P, et al. Stereotactic body radiation therapy versus conventionally fractionated radiation therapy for early stage non-small cell lung cancer. Radiother Oncol 2018;129:264–269. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30031630>.
12. Shioyama Y, Onishi H, Takayama K, et al. Clinical outcomes of stereotactic body radiotherapy for patients with stage I small-cell lung cancer: analysis of a subset of the Japanese radiological society multi-institutional SBRT study group database. Technol Cancer Res Treat 2018;17:1533033818783904. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29983096>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

13. Yang CF, Chan DY, Shah SA, et al. Long-term survival after surgery compared with concurrent chemoradiation for node-negative small cell lung cancer. *Ann Surg* 2018;268:1105–1112. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28475559>.
14. Verma V, Simone CB, 2nd, Allen PK, et al. Multi-institutional experience of stereotactic ablative radiation therapy for stage I small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2017;97:362–371. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28011047>.
15. Yu JB, Decker RH, Detterbeck FC, Wilson LD. Surveillance epidemiology and end results evaluation of the role of surgery for stage I small cell lung cancer. *J Thorac Oncol* 2010;5:215–219. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20101146>.
16. Demedts IK, Vermaelen KY, van Meerbeeck JP. Treatment of extensive-stage small cell lung carcinoma: current status and future prospects. *Eur Respir J* 2010;35:202–215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20044461>.
17. Jett JR, Schild SE, Kesler KA, Kalemkerian GP. Treatment of small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e400S–e419S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649448>.
18. National Lung Screening Trial Research Team, Aberle DR, Adams AM, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* 2011;365:395–409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21714641>.
19. Cuffe S, Moua T, Summerfield R, et al. Characteristics and outcomes of small cell lung cancer patients diagnosed during two lung cancer computed tomographic screening programs in heavy smokers. *J Thorac Oncol* 2011;6:818–822. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21623258>.
20. Thomas A, Pattanayak P, Szabo E, Pinsky P. Characteristics and outcomes of small cell lung cancer detected by CT screening. *Chest* 2018;154:1284–1290. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30080997>.
21. Kondo R, Yoshida K, Kawakami S, et al. Different efficacy of CT screening for lung cancer according to histological type: analysis of Japanese-smoker cases detected using a low-dose CT screen. *Lung Cancer* 2011;74:433–440. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21663995>.
22. Rivera MP, Mehta AC, Wahidi MM. Establishing the diagnosis of lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e142S–e165S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649436>.
23. Travis WD. Advances in neuroendocrine lung tumors. *Ann Oncol* 2010;21 Suppl 7:vii65–71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20943645>.
24. Renshaw AA, Haja J, Lozano RL, et al. Distinguishing carcinoid tumor from small cell carcinoma of the lung: correlating cytologic features and performance in the College of American Pathologists Non-Gynecologic Cytology Program. *Arch Pathol Lab Med* 2005;129:614–618. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15859631>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

25. Gandhi L, Johnson BE. Paraneoplastic syndromes associated with small cell lung cancer. J Natl Compr Canc Netw 2006;4:631–638. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16813730>.
26. Kazarian M, Laird-Offringa IA. Small-cell lung cancer-associated autoantibodies: potential applications to cancer diagnosis, early detection, and therapy. Mol Cancer 2011;10:33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21450098>.
27. Marchioli CC, Graziano SL. Paraneoplastic syndromes associated with small cell lung cancer. Chest Surg Clin N Am 1997;7:65–80. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9001756>.
28. Titulaer MJ, Wirtz PW, Willems LN, et al. Screening for small-cell lung cancer: a follow-up study of patients with Lambert-Eaton Myasthenic Syndrome. J Clin Oncol 2008;26:4276–4281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18779614>.
29. Meriney SD, Hulsizer SC, Lennon VA, Grinnell AD. Lambert-Eaton Myasthenic Syndrome immunoglobulins react with multiple types of calcium channels in small-cell lung carcinoma. Ann Neurol 1996;40:739–749. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8957015>.
30. Graus F, Keime-Guibert F, Rene R, et al. Anti-Hu-associated paraneoplastic encephalomyelitis: analysis of 200 patients. Brain 2001;124:1138–1148. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11353730>.
31. Delisle L, Boyer MJ, Warr D, et al. Ectopic corticotropin syndrome and small-cell carcinoma of the lung. Clinical features, outcome, and complications. Arch Intern Med 1993;153:746–752. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8383484>.
32. Johnson BE, Chute JP, Rushin J, et al. A prospective study of patients with lung cancer and hyponatremia of malignancy. Am J Respir Crit Care Med 1997;156:1669–1678. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9372692>.
33. Castillo JJ, Vincent M, Justice E. Diagnosis and management of hyponatremia in cancer patients. Oncologist 2012;17:756–765. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22618570>.
34. Schrier RW, Gross P, Gheorghiade M, et al. Tolvaptan, a selective oral vasopressin V2-receptor antagonist, for hyponatremia. N Engl J Med 2006;355:2099–2112. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17105757>.
35. Verbalis JG, Zeltser D, Smith N, et al. Assessment of the efficacy and safety of intravenous conivaptan in patients with euvolaemic hyponatraemia: subgroup analysis of a randomized, controlled study. Clin Endocrinol (Oxf) 2008;69:159–168. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18034777>.
36. Travis WD. Pathology and diagnosis of neuroendocrine tumors: lung neuroendocrine. Thorac Surg Clin 2014;24:257–266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25065926>.
37. Nicholson SA, Beasley MB, Brambilla E, et al. Small cell lung carcinoma (SCLC): a clinicopathologic study of 100 cases with surgical specimens. Am J Surg Pathol 2002;26:1184–1197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12218575>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

38. Zakowski MF. Pathology of small cell carcinoma of the lung. *Semin Oncol* 2003;30:3–8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12635085>.
39. Pelosi G, Rodriguez J, Viale G, Rosai J. Typical and atypical pulmonary carcinoid tumor overdiagnosed as small-cell carcinoma on biopsy specimens: a major pitfall in the management of lung cancer patients. *Am J Surg Pathol* 2005;29:179–187. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15644774>.
40. Bellizzi AM. Immunohistochemistry in the diagnosis and classification of neuroendocrine neoplasms: what can brown do for you? *Hum Pathol* 2020;96:8–33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31857137>.
41. Thunnissen E, Borczuk AC, Flieder DB, et al. The use of immunohistochemistry improves the diagnosis of small cell lung cancer and its differential diagnosis. An international reproducibility study in a demanding set of cases. *J Thorac Oncol* 2017;12:334–346. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27998793>.
42. Rindi G, Klersy C, Inzani F, et al. Grading the neuroendocrine tumors of the lung: an evidence-based proposal. *Endocr Relat Cancer* 2014;21:1–16. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24344249>.
43. Travis WD, Brambilla E, Burke AP, et al. WHO classification of tumours of the lung, pleura, thymus and heart. Fourth edition. Geneva, Switzerland: World Health Organization; 2015.
44. Rindi G, Klimstra DS, Abedi-Ardekani B, et al. A common classification framework for neuroendocrine neoplasms: an International Agency for Research on Cancer (IARC) and World Health Organization (WHO) expert consensus proposal. *Mod Pathol* 2018;31:1770–1786. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30140036>.
45. Pelosi G, Rindi G, Travis WD, Papotti M. Ki-67 antigen in lung neuroendocrine tumors: unraveling a role in clinical practice. *J Thorac Oncol* 2014;9:273–284. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24518085>.
46. Masai K, Tsuta K, Kawago M, et al. Expression of squamous cell carcinoma markers and adenocarcinoma markers in primary pulmonary neuroendocrine carcinomas. *Appl Immunohistochem Mol Morphol* 2013;21:292–297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23060301>.
47. Ordonez NG. Value of thyroid transcription factor-1 immunostaining in distinguishing small cell lung carcinomas from other small cell carcinomas. *Am J Surg Pathol* 2000;24:1217–1223. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10976695>.
48. Kaufmann O, Dietel M. Expression of thyroid transcription factor-1 in pulmonary and extrapulmonary small cell carcinomas and other neuroendocrine carcinomas of various primary sites. *Histopathology* 2000;36:415–420. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10792482>.
49. Rooper LM, Sharma R, Li QK, et al. INSM1 demonstrates superior performance to the individual and combined use of synaptophysin, chromogranin and CD56 for diagnosing neuroendocrine tumors of the thoracic cavity. *Am J Surg Pathol* 2017;41:1561–1569. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28719469>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

50. Sakakibara R, Kobayashi M, Takahashi N, et al. Insulinoma-associated Protein 1 (INSM1) is a better marker for the diagnosis and prognosis estimation of small cell lung carcinoma than neuroendocrine phenotype markers such as chromogranin a, synaptophysin, and CD56. *Am J Surg Pathol* 2020;44:757–764. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32118626>.
51. Guinee DG, Jr., Fishback NF, Koss MN, et al. The spectrum of immunohistochemical staining of small-cell lung carcinoma in specimens from transbronchial and open-lung biopsies. *Am J Clin Pathol* 1994;102:406–414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7524299>.
52. Baine MK, Febres-Aldana CA, Chang JC, et al. POU2F3 in SCLC: Clinicopathologic and genomic analysis with a focus on its diagnostic utility in neuroendocrine-low sclc. *J Thorac Oncol* 2022;17:1109–1121. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35760287>.
53. Wang Y, Jin Y, Shen X, et al. POU2F3: A sensitive and specific diagnostic marker for neuroendocrine-low/negative small cell lung cancer. *Am J Surg Pathol* 2023;47:1059–1066. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37357936>.
54. Rekhtman N, Pietanza CM, Sabari J, et al. Pulmonary large cell neuroendocrine carcinoma with adenocarcinoma-like features: napsin A expression and genomic alterations. *Mod Pathol* 2018;31:111–121. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28884744>.
55. George J, Lim JS, Jang SJ, et al. Comprehensive genomic profiles of small cell lung cancer. *Nature* 2015;524:47–53. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26168399>.
56. Rudin CM, Brambilla E, Faivre-Finn C, Sage J. Small-cell lung cancer. *Nat Rev Dis Primers* 2021;7:3. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33446664>.
57. Su S, Zou JJ, Zeng YY, et al. Tumor mutational burden and genomic alterations in chinese small cell lung cancer measured by whole-exome sequencing. *Biomed Res Int* 2019;2019:6096350. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31781628>.
58. Wakuda K, Kenmotsu H, Serizawa M, et al. Molecular profiling of small cell lung cancer in a Japanese cohort. *Lung Cancer* 2014;84:139–144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24657128>.
59. Liguori NR, Lee Y, Borges W, et al. Absence of biomarker-driven treatment options in small cell lung cancer, and selected preclinical candidates for next generation combination therapies. *Front Pharmacol* 2021;12:747180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34531756>.
60. Ogino A, Choi J, Lin M, et al. Genomic and pathological heterogeneity in clinically diagnosed small cell lung cancer in never/light smokers identifies therapeutically targetable alterations. *Mol Oncol* 2021;15:27–42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32191822>.
61. Tsao MS, Nicholson AG, Maleszewski JJ, et al. Introduction to 2021 WHO classification of thoracic tumors. *J Thorac Oncol* 2022;17:e1–e4. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34930611>.
62. WHO classification of tumours editorial board. Thoracic tumours. In: Who classification of tumours series. 5th ed. WHO classification of tumours series. Lyon, France: International Agency for Research on Cancer; 2021.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

63. Sanguedolce F, Zanelli M, Palicelli A, et al. The classification of neuroendocrine neoplasms of the lung and digestive system according to WHO, 5th edition: similarities, differences, challenges, and unmet needs. *Panminerva Med* 2022;64:259–264. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35146989>.
64. Travis WD, Brambilla E, Nicholson AG, et al. The 2015 World Health Organization classification of lung tumors: Impact of genetic, clinical and radiologic advances since the 2004 classification. *J Thorac Oncol* 2015;10:1243–1260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26291008>.
65. Travis WD, Brambilla E, Burke AP, et al. Introduction to the 2015 world health organization classification of tumors of the lung, pleura, thymus, and heart. *J Thorac Oncol* 2015;10:1240–1242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26291007>.
66. Qin J, Lu H. Combined small-cell lung carcinoma. *Onco Targets Ther* 2018;11:3505–3511. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29950855>.
67. Marcoux N, Gettinger SN, O'Kane G, et al. EGFR-mutant adenocarcinomas that transform to small-cell lung cancer and other neuroendocrine carcinomas: Clinical outcomes. *J Clin Oncol* 2019;37:278–285. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30550363>.
68. Sehgal K, Varkaris A, Viray H, et al. Small cell transformation of non-small cell lung cancer on immune checkpoint inhibitors: uncommon or under-recognized? *J Immunother Cancer* 2020;8:e000697. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32581048>.
69. Micke P, Faldut A, Metz T, et al. Staging small cell lung cancer: Veterans Administration Lung Study Group versus International Association for the Study of Lung Cancer--what limits limited disease? *Lung Cancer* 2002;37:271–276. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12234695>.
70. Kalemkerian GP, Gadgeel SM. Modern staging of small cell lung cancer. *J Natl Compr Canc Netw* 2013;11:99–104. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23307985>.
71. Kalemkerian GP. Staging and imaging of small cell lung cancer. *Cancer Imaging* 2012;11:253–258. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22245990>.
72. Amin MB, Greene FL, Byrd DR, et al. *AJCC cancer staging manual*, 8th edition: Springer International Publishing; 2016:1-1024.
73. Nicholson AG, Chansky K, Crowley J, et al. The international association for the study of lung cancer lung cancer staging project: Proposals for the revision of the clinical and pathologic staging of small cell lung cancer in the forthcoming eighth edition of the TNM classification for lung cancer. *J Thorac Oncol* 2016;11:300–311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26723244>.
74. Seute T, Leffers P, ten Velde GP, Twijnstra A. Detection of brain metastases from small cell lung cancer: consequences of changing imaging techniques (CT versus MRI). *Cancer* 2008;112:1827–1834. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18311784>.
75. Fischer BM, Mortensen J, Langer SW, et al. A prospective study of PET/CT in initial staging of small-cell lung cancer: comparison with CT, bone scintigraphy and bone marrow analysis. *Ann Oncol* 2007;18:338–345. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17060487>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

76. Brink I, Schumacher T, Mix M, et al. Impact of [18F]FDG-PET on the primary staging of small-cell lung cancer. *Eur J Nucl Med Mol Imaging* 2004;31:1614–1620. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15258700>.
77. Podoloff DA, Ball DW, Ben-Josef E, et al. NCCN task force: clinical utility of PET in a variety of tumor types. *J Natl Compr Canc Netw* 2009;7 Suppl 2:S1–26. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19555588>.
78. Bradley JD, Dehdashti F, Mintun MA, et al. Positron emission tomography in limited-stage small-cell lung cancer: a prospective study. *J Clin Oncol* 2004;22:3248–3254. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15310768>.
79. Kamel EM, Zwahlen D, Wyss MT, et al. Whole-body (18)F-FDG PET improves the management of patients with small cell lung cancer. *J Nucl Med* 2003;44:1911–1917. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14660716>.
80. Rintoul RC, Tournoy KG, El Daly H, et al. EBUS-TBNA for the clarification of PET positive intra-thoracic lymph nodes-an international multi-centre experience. *J Thorac Oncol* 2009;4:44–48. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19096305>.
81. Medford AR, Bennett JA, Free CM, Agrawal S. Mediastinal staging procedures in lung cancer: EBUS, TBNA and mediastinoscopy. *Curr Opin Pulm Med* 2009;15:334–342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19395972>.
82. Foster NR, Mandrekar SJ, Schild SE, et al. Prognostic factors differ by tumor stage for small cell lung cancer: a pooled analysis of North Central Cancer Treatment Group trials. *Cancer* 2009;115:2721–2731. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19402175>.
83. Albain KS, Crowley JJ, LeBlanc M, Livingston RB. Determinants of improved outcome in small-cell lung cancer: an analysis of the 2,580-patient Southwest Oncology Group data base. *J Clin Oncol* 1990;8:1563–1574. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2167954>.
84. Ignatius Ou SH, Zell JA. The applicability of the proposed IASLC staging revisions to small cell lung cancer (SCLC) with comparison to the current UICC 6th TNM Edition. *J Thorac Oncol* 2009;4:300–310. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19156001>.
85. Schneider BJ, Saxena A, Downey RJ. Surgery for early-stage small cell lung cancer. *J Natl Compr Canc Netw* 2011;9:1132–1139. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21975913>.
86. Rostad H, Naalsund A, Jacobsen R, et al. Small cell lung cancer in Norway. Should more patients have been offered surgical therapy? *Eur J Cardiothorac Surg* 2004;26:782–786. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15450573>.
87. Inoue M, Miyoshi S, Yasumitsu T, et al. Surgical results for small cell lung cancer based on the new TNM staging system. Thoracic Surgery Study Group of Osaka University, Osaka, Japan. *Ann Thorac Surg* 2000;70:1615–1619. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11093496>.
88. Brock MV, Hooker CM, Syphard JE, et al. Surgical resection of limited disease small cell lung cancer in the new era of platinum chemotherapy: Its time has come. *J Thorac Cardiovasc Surg* 2005;129:64–72. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15632826>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

89. Lim E, Belcher E, Yap YK, et al. The role of surgery in the treatment of limited disease small cell lung cancer: time to reevaluate. *J Thorac Oncol* 2008;3:1267–1271. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18978561>.
90. Shields TW, Higgins GA, Jr., Matthews MJ, Keehn RJ. Surgical resection in the management of small cell carcinoma of the lung. *J Thorac Cardiovasc Surg* 1982;84:481–488. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6289013>.
91. Schreiber D, Rineer J, Weedon J, et al. Survival outcomes with the use of surgery in limited-stage small cell lung cancer: should its role be re-evaluated? *Cancer* 2010;116:1350–1357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20082453>.
92. Lad T, Piantadosi S, Thomas P, et al. A prospective randomized trial to determine the benefit of surgical resection of residual disease following response of small cell lung cancer to combination chemotherapy. *Chest* 1994;106:320S–323S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7988254>.
93. Inoue M, Nakagawa K, Fujiwara K, et al. Results of preoperative mediastinoscopy for small cell lung cancer. *Ann Thorac Surg* 2000;70:1620–1623. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11093497>.
94. Shepherd FA, Evans WK, Feld R, et al. Adjuvant chemotherapy following surgical resection for small-cell carcinoma of the lung. *J Clin Oncol* 1988;6:832–838. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2835443>.
95. Tsuchiya R, Suzuki K, Ichinose Y, et al. Phase II trial of postoperative adjuvant cisplatin and etoposide in patients with completely resected stage I–IIa small cell lung cancer: The Japan clinical oncology lung cancer study group trial (JCOG9101). *J Thorac Cardiovasc Surg* 2005;129:977–983. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15867769>.
96. Yang CF, Chan DY, Speicher PJ, et al. Role of adjuvant therapy in a population-based cohort of patients with early-stage small-cell lung cancer. *J Clin Oncol* 2016;34:1057–1064. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26786925>.
97. Zhang C, Zhao G, Wu H, et al. Application of postoperative adjuvant radiotherapy in limited-stage small cell lung cancer: A systematic review and meta-analysis. *Radiother Oncol* 2024;193:110123. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38309584>.
98. Pignon JP, Arriagada R, Ihde DC, et al. A meta-analysis of thoracic radiotherapy for small-cell lung cancer. *N Engl J Med* 1992;327:1618–1624. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1331787>.
99. Warde P, Payne D. Does thoracic irradiation improve survival and local control in limited-stage small-cell carcinoma of the lung? A meta-analysis. *J Clin Oncol* 1992;10:890–895. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1316951>.
100. Hatfield LA, Huskamp HA, Lamont EB. Survival and toxicity after cisplatin plus etoposide versus carboplatin plus etoposide for extensive-stage small-cell lung cancer in elderly patients. *J Oncol Pract* 2016;12:666–673. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27352949>.
101. Skarlos DV, Samantas E, Kosmidis P, et al. Randomized comparison of etoposide-cisplatin vs. etoposide-carboplatin and irradiation in small-cell lung cancer. A Hellenic co-operative oncology group study. *Ann Oncol* 1994;5:601–607. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7993835>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

102. Okamoto H, Watanabe K, Kunikane H, et al. Randomised phase III trial of carboplatin plus etoposide vs split doses of cisplatin plus etoposide in elderly or poor-risk patients with extensive disease small-cell lung cancer: JCOG 9702. *Br J Cancer* 2007;97:162–169. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17579629>.
103. Rossi A, Di Maio M, Chiodini P, et al. Carboplatin- or cisplatin-based chemotherapy in first-line treatment of small-cell lung cancer: the COCIS meta-analysis of individual patient data. *J Clin Oncol* 2012;30:1692–1698. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22473169>.
104. Griesinger F, Korol EE, Kayaniyl S, et al. Efficacy and safety of first-line carboplatin-versus cisplatin-based chemotherapy for non-small cell lung cancer: A meta-analysis. *Lung Cancer* 2019;135:196–204. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31446995>.
105. Faivre-Finn C, Snee M, Ashcroft L, et al. Concurrent once-daily versus twice-daily chemoradiotherapy in patients with limited-stage small-cell lung cancer (CONVERT): an open-label, phase 3, randomised, superiority trial. *Lancet Oncol* 2017;18:1116–1125. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28642008>.
106. Evans WK, Shepherd FA, Feld R, et al. VP-16 and cisplatin as first-line therapy for small-cell lung cancer. *J Clin Oncol* 1985;3:1471–1477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2997406>.
107. Pujol JL, Carestia L, Daures JP. Is there a case for cisplatin in the treatment of small-cell lung cancer? A meta-analysis of randomized trials of a cisplatin-containing regimen versus a regimen without this alkylating agent. *Br J Cancer* 2000;83:8–15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10883661>.
108. Mascaux C, Paesmans M, Berghmans T, et al. A systematic review of the role of etoposide and cisplatin in the chemotherapy of small cell lung cancer with methodology assessment and meta-analysis. *Lung Cancer* 2000;30:23–36. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11008007>.
109. Sundstrom S, Bremnes RM, Kaasa S, et al. Cisplatin and etoposide regimen is superior to cyclophosphamide, epirubicin, and vincristine regimen in small-cell lung cancer: results from a randomized phase III trial with 5 years' follow-up. *J Clin Oncol* 2002;20:4665–4672. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12488411>.
110. Takada M, Fukuoka M, Kawahara M, et al. Phase III study of concurrent versus sequential thoracic radiotherapy in combination with cisplatin and etoposide for limited-stage small-cell lung cancer: results of the Japan clinical oncology group Study 9104. *J Clin Oncol* 2002;20:3054–3060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12118018>.
111. Niho S, Kubota K, Yoh K, et al. Clinical outcome of chemoradiation therapy in patients with limited-disease small cell lung cancer with ipsilateral pleural effusion. *J Thorac Oncol* 2008;3:723–727. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18594317>.
112. Niho S, Kubota K, Yoh K, et al. Clinical outcome of small cell lung cancer with pericardial effusion but without distant metastasis. *J Thorac Oncol* 2011;6:796–800. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21258253>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

113. Kubota K, Hida T, Ishikura S, et al. Etoposide and cisplatin versus irinotecan and cisplatin in patients with limited-stage small-cell lung cancer treated with etoposide and cisplatin plus concurrent accelerated hyperfractionated thoracic radiotherapy (JCOG0202): a randomised phase 3 study. *Lancet Oncol* 2014;15:106–113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24309370>.
114. Saito H, Takada Y, Ichinose Y, et al. Phase II study of etoposide and cisplatin with concurrent twice-daily thoracic radiotherapy followed by irinotecan and cisplatin in patients with limited-disease small-cell lung cancer: West Japan thoracic oncology group 9902. *J Clin Oncol* 2006;24:5247–5252. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17114657>.
115. Cheng Y, Spigel DR, Cho BC, et al. Durvalumab after chemoradiotherapy in limited-stage small-cell lung cancer. *N Engl J Med* 2024;391:1313–1327. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39268857>.
116. Pan B, Shen B. The ADRIATIC study: revolutionizing the standard treatment paradigm for concurrent chemoradiotherapy in limited-stage small cell lung cancer. *J Natl Cancer Cent* 2025;5:1–2. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40040876>.
117. Bunn PA, Jr., Crowley J, Kelly K, et al. Chemoradiotherapy with or without granulocyte-macrophage colony-stimulating factor in the treatment of limited-stage small-cell lung cancer: a prospective phase III randomized study of the Southwest Oncology Group. *J Clin Oncol* 1995;13:1632–1641. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7602352>.
118. Liu SV, Reck M, Mansfield AS, et al. Updated overall survival and PD-L1 subgroup analysis of patients with extensive-stage small-cell lung cancer treated with atezolizumab, carboplatin, and etoposide (IMpower133). *J Clin Oncol* 2021;39:619–630. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33439693>.
119. Paz-Ares L, Dvorkin M, Chen Y, et al. Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *Lancet* 2019;394:1929–1939. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31590988>.
120. Horn L, Mansfield AS, Szczesna A, et al. First-line atezolizumab plus chemotherapy in extensive-stage small-cell lung cancer. *N Engl J Med* 2018;379:2220–2229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30280641>.
121. Postmus PE, Haaxma-Reiche H, Gregor A, et al. Brain-only metastases of small cell lung cancer; efficacy of whole brain radiotherapy. An EORTC phase II study. *Radiother Oncol* 1998;46:29–32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9488124>.
122. Kumar A, Weber MH, Gokaslan Z, et al. Metastatic spinal cord compression and steroid treatment: A systematic review. *Clin Spine Surg* 2017;30:156–163. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28437329>.
123. Goldman JW, Dvorkin M, Chen Y, et al. Durvalumab, with or without tremelimumab, plus platinum-etoposide versus platinum-etoposide alone in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): updated results from a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2021;22:51–65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33285097>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

124. Paz-Ares L, Chen Y, Reinmuth N, et al. Durvalumab, with or without tremelimumab, plus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer: 3-year overall survival update from CASPIAN. ESMO Open 2022;7:100408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35279527>.
125. Mathieu L, Shah S, Pai-Scherf L, et al. FDA approval summary: Atezolizumab and durvalumab in combination with platinum-based chemotherapy in extensive stage small cell lung cancer. Oncologist 2021;26:433–438. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33687763>.
126. Noda K, Nishiwaki Y, Kawahara M, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. N Engl J Med 2002;346:85–91. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11784874>.
127. Lara PN, Jr., Natale R, Crowley J, et al. Phase III trial of irinotecan/cisplatin compared with etoposide/cisplatin in extensive-stage small-cell lung cancer: clinical and pharmacogenomic results from SWOG S0124. J Clin Oncol 2009;27:2530–2535. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19349543>.
128. Hanna N, Bunn PA, Jr, Langer C, et al. Randomized phase III trial comparing irinotecan/cisplatin with etoposide/cisplatin in patients with previously untreated extensive-stage disease small-cell lung cancer 10.1200/JCO.2005.04.8595. J Clin Oncol 2006;24:2038–2043. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16648503>.
129. Hermes A, Bergman B, Bremnes R, et al. Irinotecan plus carboplatin versus oral etoposide plus carboplatin in extensive small-cell lung cancer: a randomized phase III trial. J Clin Oncol 2008;26:4261–4267. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18779613>.
130. Lima JP, dos Santos LV, Sasse EC, et al. Camptothecins compared with etoposide in combination with platinum analog in extensive stage small cell lung cancer: systematic review with meta-analysis. J Thorac Oncol 2010;5:1986–1993. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20978445>.
131. Schiller JH, Adak S, Cella D, et al. Topotecan versus observation after cisplatin plus etoposide in extensive-stage small-cell lung cancer: E7593--a phase III trial of the Eastern Cooperative Oncology Group. J Clin Oncol 2001;19:2114–2122. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11304763>.
132. Zhou H, Zeng C, Wei Y, et al. Duration of chemotherapy for small cell lung cancer: a meta-analysis. PLoS One 2013;8:e73805. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24023692>.
133. Goldie JH, Coldman AJ. A mathematic model for relating the drug sensitivity of tumors to their spontaneous mutation rate. Cancer Treat Rep 1979;63:1727–1733. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/526911>.
134. Fukuoka M, Furuse K, Saijo N, et al. Randomized trial of cyclophosphamide, doxorubicin, and vincristine versus cisplatin and etoposide versus alternation of these regimens in small-cell lung cancer. J Natl Cancer Inst 1991;83:855–861. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1648142>.
135. Roth BJ, Johnson DH, Einhorn LH, et al. Randomized study of cyclophosphamide, doxorubicin, and vincristine versus etoposide and cisplatin versus alternation of these two regimens in extensive small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. J Clin Oncol 1992;10:282–291. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1310103>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

136. Loehrer PJ, Sr., Ansari R, Gonin R, et al. Cisplatin plus etoposide with and without ifosfamide in extensive small-cell lung cancer: a Hoosier Oncology Group study. *J Clin Oncol* 1995;13:2594–2599. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7595712>.
137. Pujol JL, Daures JP, Riviere A, et al. Etoposide plus cisplatin with or without the combination of 4'-epidoxorubicin plus cyclophosphamide in treatment of extensive small-cell lung cancer: a French Federation of Cancer Institutes multicenter phase III randomized study. *J Natl Cancer Inst* 2001;93:300–308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11181777>.
138. Miyamoto H, Nakabayashi T, Isobe H, et al. A phase III comparison of etoposide/cisplatin with or without added ifosfamide in small-cell lung cancer. *Oncology* 1992;49:431–435. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1334539>.
139. Berghmans T, Scherpereel A, Meert AP, et al. A phase III randomized study comparing a chemotherapy with cisplatin and etoposide to a etoposide regimen without cisplatin for patients with extensive small-cell lung cancer. *Front Oncol* 2017;7:217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28975084>.
140. Jalal SI, Lavin P, Lo G, et al. Carboplatin and etoposide with or without palifosfamide in untreated extensive-stage small-cell lung cancer: A multicenter, adaptive, randomized phase III study (MATISSE). *J Clin Oncol* 2017;35:2619–2623. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28605291>.
141. Niell HB, Herndon JE, 2nd, Miller AA, et al. Randomized phase III intergroup trial of etoposide and cisplatin with or without paclitaxel and granulocyte colony-stimulating factor in patients with extensive-stage small-cell lung cancer: Cancer and Leukemia Group B Trial 9732. *J Clin Oncol* 2005;23:3752–3759. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15923572>.
142. Cohen MH, Creaven PJ, Fossieck BE, Jr., et al. Intensive chemotherapy of small cell bronchogenic carcinoma. *Cancer Treat Rep* 1977;61:349–354. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/194691>.
143. Johnson DH, Einhorn LH, Birch R, et al. A randomized comparison of high-dose versus conventional-dose cyclophosphamide, doxorubicin, and vincristine for extensive-stage small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. *J Clin Oncol* 1987;5:1731–1738. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2824707>.
144. Ihde DC, Mulshine JL, Kramer BS, et al. Prospective randomized comparison of high-dose and standard-dose etoposide and cisplatin chemotherapy in patients with extensive-stage small-cell lung cancer. *J Clin Oncol* 1994;12:2022–2034. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7931470>.
145. Arriagada R, Le Chevalier T, Pignon JP, et al. Initial chemotherapeutic doses and survival in patients with limited small-cell lung cancer. *N Engl J Med* 1993;329:1848–1852. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8247036>.
146. Thatcher N, Girling DJ, Hopwood P, et al. Improving survival without reducing quality of life in small-cell lung cancer patients by increasing the dose-intensity of chemotherapy with granulocyte colony-stimulating factor support: Results of a british medical research council multicenter randomized trial. Medical research council lung cancer working party. *J Clin Oncol* 2000;18:395–404. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10637255>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

147. Klasa RJ, Murray N, Coldman AJ. Dose-intensity meta-analysis of chemotherapy regimens in small-cell carcinoma of the lung. *J Clin Oncol* 1991;9:499–508. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1847968>.
148. Miles DW, Earl HM, Souhami RL, et al. Intensive weekly chemotherapy for good-prognosis patients with small-cell lung cancer. *J Clin Oncol* 1991;9:280–285. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1846406>.
149. Murray N, Gelmon K, Shah A, et al. Potential for long-term survival in extensive stage small-cell lung cancer (ESCLC) with code chemotherapy and radiotherapy. *Lung Cancer* 1994;11:99. Available at: <https://www.sciencedirect.com/science/article/pii/0169500294941505>.
150. Sculier JP, Paesmans M, Bureau G, et al. Multiple-drug weekly chemotherapy versus standard combination regimen in small-cell lung cancer: a phase III randomized study conducted by the European Lung Cancer Working Party. *J Clin Oncol* 1993;11:1858–1865. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8410110>.
151. Souhami RL, Rudd R, Ruiz de Elvira MC, et al. Randomized trial comparing weekly versus 3-week chemotherapy in small-cell lung cancer: A cancer research campaign trial. *J Clin Oncol* 1994;12:1806–1813. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8083704>.
152. Fukuoka M, Masuda N, Negoro S, et al. CODE chemotherapy with and without granulocyte colony-stimulating factor in small-cell lung cancer. *Br J Cancer* 1997;75:306–309. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9010043>.
153. Murray N, Livingston RB, Shepherd FA, et al. Randomized study of CODE versus alternating CAV/EP for extensive-stage small-cell lung cancer: an Intergroup Study of the National Cancer Institute of Canada Clinical Trials Group and the Southwest Oncology Group. *J Clin Oncol* 1999;17:2300–2308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10561291>.
154. Reck M, Luft A, Szczesna A, et al. Phase III randomized trial of ipilimumab plus etoposide and platinum versus placebo plus etoposide and platinum in extensive-stage small-cell lung cancer. *J Clin Oncol* 2016;34:3740–3748. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27458307>.
155. Petrioli R, Roviello G, Laera L, et al. Cisplatin, etoposide, and bevacizumab regimen followed by oral etoposide and bevacizumab maintenance treatment in patients with extensive-stage small cell lung cancer: A single-institution experience. *Clin Lung Cancer* 2015;16:e229–234. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26072097>.
156. Spigel DR, Townley PM, Waterhouse DM, et al. Randomized phase II study of bevacizumab in combination with chemotherapy in previously untreated extensive-stage small-cell lung cancer: results from the SALUTE trial. *J Clin Oncol* 2011;29:2215–2222. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21502556>.
157. Spigel DR, Greco FA, Zubkus JD, et al. Phase II trial of irinotecan, carboplatin, and bevacizumab in the treatment of patients with extensive-stage small-cell lung cancer. *J Thorac Oncol* 2009;4:1555–1560. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19875975>.
158. Horn L, Dahlberg SE, Sandler AB, et al. Phase II study of cisplatin plus etoposide and bevacizumab for previously untreated, extensive-stage small-cell lung cancer: Eastern Cooperative Oncology Group Study E3501. *J Clin Oncol* 2009;27:6006–6011. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19826110>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

159. Tiseo M, Boni L, Ambrosio F, et al. Italian, multicenter, phase III, randomized study of cisplatin plus etoposide with or without bevacizumab as first-line treatment in extensive-disease small-cell lung cancer: The GOIRC-AIFA FARM6PMFJM trial. *J Clin Oncol* 2017;35:1281–1287. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28135143>.
160. Pujol JL, Lavole A, Quoix E, et al. Randomized phase II-III study of bevacizumab in combination with chemotherapy in previously untreated extensive small-cell lung cancer: results from the IFCT-0802 trial†. *Ann Oncol* 2015;26:908–914. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25688059>.
161. Crawford J, Ozer H, Stoller R, et al. Reduction by granulocyte colony-stimulating factor of fever and neutropenia induced by chemotherapy in patients with small-cell lung cancer. *N Engl J Med* 1991;325:164–170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1711156>.
162. Berghmans T, Paesmans M, Lafitte JJ, et al. Role of granulocyte and granulocyte-macrophage colony-stimulating factors in the treatment of small-cell lung cancer: a systematic review of the literature with methodological assessment and meta-analysis. *Lung Cancer* 2002;37:115–123. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12140132>.
163. Sculier JP, Paesmans M, Lecomte J, et al. A three-arm phase III randomised trial assessing, in patients with extensive-disease small-cell lung cancer, accelerated chemotherapy with support of haematological growth factor or oral antibiotics. *Br J Cancer* 2001;85:1444–1451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11720426>.
164. Wang C, Zhu S, Miao C, et al. Safety and efficacy of pegylated recombinant human granulocyte colony-stimulating factor during concurrent chemoradiotherapy for small-cell lung cancer: a retrospective, cohort-controlled trial. *BMC Cancer* 2022;22:542. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35562713>.
165. Ferrarotto R, Anderson I, Medgyasszay B, et al. Trilaciclib prior to chemotherapy reduces the usage of supportive care interventions for chemotherapy-induced myelosuppression in patients with small cell lung cancer: Pooled analysis of three randomized phase 2 trials. *Cancer Med* 2021;10:5748–5756. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34405547>.
166. Hussein M, Maglakelidze M, Richards DA, et al. Myeloprotective effects of trilaciclib among patients with small cell lung cancer at increased risk of chemotherapy-induced myelosuppression: Pooled results from three phase 2, randomized, double-blind, placebo-controlled studies. *Cancer Manag Res* 2021;13:6207–6218. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34408488>.
167. Weiss J, Goldschmidt J, Andric Z, et al. Effects of trilaciclib on chemotherapy-induced myelosuppression and patient-reported outcomes in patients with extensive-stage small cell lung cancer: Pooled results from three phase II randomized, double-blind, placebo-controlled studies. *Clin Lung Cancer* 2021;22:449–460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33895103>.
168. Hart LL, Ferrarotto R, Andric ZG, et al. Myelopreservation with trilaciclib in patients receiving topotecan for small cell lung cancer: Results from a randomized, double-blind, placebo-controlled phase ii study. *Adv Ther* 2021;38:350–365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33123968>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

169. Cheng Y, Wu L, Huang D, et al. Myeloprotection with trilaciclib in Chinese patients with extensive-stage small cell lung cancer receiving chemotherapy: Results from a randomized, double-blind, placebo-controlled phase III study (TRACES). *Lung Cancer* 2024;188:107455. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38224653>.
170. Liu Y, Wu L, Huang D, et al. Effect of trilaciclib administered before chemotherapy in patients with extensive-stage small-cell lung cancer: A pooled analysis of four randomized studies. *Cancer Treat Res Commun* 2024;42:100869. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39823755>.
171. Hurria A, Kris MG. Management of lung cancer in older adults. *CA Cancer J Clin* 2003;53:325–341. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15224973>.
172. Christodoulou M, Blackhall F, Mistry H, et al. Compliance and outcome of elderly patients treated in the concurrent once-daily versus twice-daily radiotherapy (CONVERT) trial. *J Thorac Oncol* 2019;14:63–71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30391573>.
173. Corso CD, Rutter CE, Park HS, et al. Role of chemoradiotherapy in elderly patients with limited-stage small-cell lung cancer. *J Clin Oncol* 2015;33:4240–4246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26481366>.
174. Gridelli C, Casaluce F, Sgambato A, et al. Treatment of limited-stage small cell lung cancer in the elderly, chemotherapy vs. sequential chemoradiotherapy vs. concurrent chemoradiotherapy: that's the question. *Transl Lung Cancer Res* 2016;5:150–154. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27186510>.
175. Girling DJ. Comparison of oral etoposide and standard intravenous multidrug chemotherapy for small-cell lung cancer: a stopped multicentre randomised trial. Medical Research Council Lung Cancer Working Party. *Lancet* 1996;348:563–566. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8774567>.
176. Souhami RL, Spiro SG, Rudd RM, et al. Five-day oral etoposide treatment for advanced small-cell lung cancer: randomized comparison with intravenous chemotherapy. *J Natl Cancer Inst* 1997;89:577–580. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9106647>.
177. Neubauer M, Schwartz J, Caracandas J, et al. Results of a phase II study of weekly paclitaxel plus carboplatin in patients with extensive small-cell lung cancer with Eastern Cooperative Oncology Group Performance Status of 2, or age > or = 70 years. *J Clin Oncol* 2004;22:1872–1877. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15143079>.
178. Westeel V, Murray N, Gelmon K, et al. New combination of the old drugs for elderly patients with small-cell lung cancer: a phase II study of the PAVE regimen. *J Clin Oncol* 1998;16:1940–1947. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9586913>.
179. Okamoto H, Watanabe K, Nishiwaki Y, et al. Phase II study of area under the plasma-concentration-versus-time curve-based carboplatin plus standard-dose intravenous etoposide in elderly patients with small-cell lung cancer. *J Clin Oncol* 1999;17:3540–3545. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10550152>.
180. Matsui K, Masuda N, Yana T, et al. Carboplatin calculated with Chatelut's formula plus etoposide for elderly patients with small-cell lung cancer. *Intern Med* 2001;40:603–606. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11506300>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

181. Murray N, Grafton C, Shah A, et al. Abbreviated treatment for elderly, infirm, or noncompliant patients with limited-stage small-cell lung cancer. *J Clin Oncol* 1998;16:3323–3328. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9779708>.
182. Damhuis RAM, Senan S, Belderbos JS. Usage of prophylactic cranial irradiation in elderly patients with small-cell lung cancer. *Clin Lung Cancer* 2018;19:e263–e267. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29208355>.
183. Wolfson AH, Bae K, Komaki R, et al. Primary analysis of a phase II randomized trial radiation therapy oncology group (RTOG) 0212: impact of different total doses and schedules of prophylactic cranial irradiation on chronic neurotoxicity and quality of life for patients with limited-disease small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2011;81:77–84. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20800380>.
184. Le Pechoux C, Laplanche A, Faivre-Finn C, et al. Clinical neurological outcome and quality of life among patients with limited small-cell cancer treated with two different doses of prophylactic cranial irradiation in the intergroup phase III trial (PCI99-01, EORTC 22003-08004, RTOG 0212 and IFCT 99-01). *Ann Oncol* 2011;22:1154–1163. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21139020>.
185. Farooqi AS, Holliday EB, Allen PK, et al. Prophylactic cranial irradiation after definitive chemoradiotherapy for limited-stage small cell lung cancer: Do all patients benefit? *Radiother Oncol* 2017;122:307–312. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28073578>.
186. Lok BH, Ma J, Foster A, et al. Factors influencing the utilization of prophylactic cranial irradiation in patients with limited-stage small cell lung cancer. *Adv Radiat Oncol* 2017;2:548–554. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29204521>.
187. Hurwitz JL, McCoy F, Scullin P, Fennell DA. New advances in the second-line treatment of small cell lung cancer. *Oncologist* 2009;14:986–994. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19819917>.
188. Schneider BJ. Management of recurrent small cell lung cancer. *J Natl Compr Canc Netw* 2008;6:323–331. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18377850>.
189. Manapov F, Klocking S, Niyazi M, et al. Timing of failure in limited disease (stage I-III) small-cell lung cancer patients treated with chemoradiotherapy: a retrospective analysis. *Tumori* 2013;99:656–660. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24503787>.
190. Johnson BE, Linnoila RI, Williams JP, et al. Risk of second aerodigestive cancers increases in patients who survive free of small-cell lung cancer for more than 2 years. *J Clin Oncol* 1995;13:101–111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7799009>.
191. Johnson BE. Second lung cancers in patients after treatment for an initial lung cancer. *J Natl Cancer Inst* 1998;90:1335–1345. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9747865>.
192. Richardson GE, Tucker MA, Venzon DJ, et al. Smoking cessation after successful treatment of small-cell lung cancer is associated with fewer smoking-related second primary cancers. *Ann Intern Med* 1993;119:383–390. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8393311>.
193. Kawahara M, Ushijima S, Kamimori T, et al. Second primary tumours in more than 2-year disease-free survivors of small-cell lung cancer in Japan: the role of smoking cessation. *Br J Cancer* 1998;78:409–412. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9703291>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

194. Parsons A, Daley A, Begh R, Aveyard P. Influence of smoking cessation after diagnosis of early stage lung cancer on prognosis: systematic review of observational studies with meta-analysis. *BMJ* 2010;340:b5569. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20093278>.
195. Paz-Ares L, Champiat S, Lai WV, et al. Tarlatamab, a first-in-class DLL3-targeted bispecific T-cell engager, in recurrent small-cell lung cancer: An open-label, phase I study. *J Clin Oncol* 2023;41:2893–2903. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36689692>.
196. Mountzios G, Sun L, Cho BC, et al. Tarlatamab in small-cell lung cancer after platinum-based chemotherapy. *N Engl J Med* 2025;393:349–361. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40454646>.
197. Masuda N, Fukuoka M, Kusunoki Y, et al. CPT-11: a new derivative of camptothecin for the treatment of refractory or relapsed small-cell lung cancer. *J Clin Oncol* 1992;10:1225–1229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1321891>.
198. Edelman MJ, Dvorkin M, Laktionov K, et al. Randomized phase 3 study of the anti-disialoganglioside antibody dinutuximab and irinotecan vs irinotecan or topotecan for second-line treatment of small cell lung cancer. *Lung Cancer* 2022;166:135–142. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35278766>.
199. Trigo J, Subbiah V, Besse B, et al. Lurbinectedin as second-line treatment for patients with small-cell lung cancer: a single-arm, open-label, phase 2 basket trial. *Lancet Oncol* 2020;21:645–654. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32224306>.
200. Subbiah V, Paz-Ares L, Besse B, et al. Antitumor activity of lurbinectedin in second-line small cell lung cancer patients who are candidates for re-challenge with the first-line treatment. *Lung Cancer* 2020;150:90–96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33096421>.
201. Baize N, Monnet I, Greillier L, et al. Carboplatin plus etoposide versus topotecan as second-line treatment for patients with sensitive relapsed small-cell lung cancer: an open-label, multicentre, randomised, phase 3 trial. *Lancet Oncol* 2020;21:1224–1233. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32888454>.
202. Naito Y, Yamada K, Imamura Y, et al. Rechallenge treatment with a platinum-based regimen in patients with sensitive relapsed small-cell lung cancer. *Med Oncol* 2018;35:61. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29610997>.
203. Genestreti G, Tiseo M, Kenmotsu H, et al. Outcomes of platinum-sensitive small-cell lung cancer patients treated with platinum/etoposide rechallenge: A multi-institutional retrospective analysis. *Clin Lung Cancer* 2015;16:e223–228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25983005>.
204. Giaccone G, Ferrati P, Donadio M, et al. Reinduction chemotherapy in small cell lung cancer. *Eur J Cancer Clin Oncol* 1987;23:1697–1699. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2828074>.
205. Postmus PE, Berendsen HH, van Zandwijk N, et al. Retreatment with the induction regimen in small cell lung cancer relapsing after an initial response to short term chemotherapy. *Eur J Cancer Clin Oncol* 1987;23:1409–1411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2824211>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

206. von Pawel J, Schiller JH, Shepherd FA, et al. Topotecan versus cyclophosphamide, doxorubicin, and vincristine for the treatment of recurrent small-cell lung cancer. *J Clin Oncol* 1999;17:658–667. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10080612>.
207. Shah C, Ready N, Perry M, et al. A multi-center phase II study of weekly topotecan as second-line therapy for small cell lung cancer. *Lung Cancer* 2007;57:84–88. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17399850>.
208. Shipley DL, Hainsworth JD, Spigel DR, et al. Topotecan: Weekly intravenous (IV) schedule similar to standard 5-day IV schedule as second-line therapy for relapsed small cell lung cancer (SCLC)--A Minnie Pearl Cancer Research Network phase II trial [abstract]. *J Clin Oncol* 2006;24 (Suppl 18):Abstract 7083. Available at: https://ascopubs.org/doi/abs/10.1200/jco.2006.24.18_suppl.7083.
209. Huber RM, Reck M, Gosse H, et al. Efficacy of a toxicity-adjusted topotecan therapy in recurrent small cell lung cancer. *Eur Respir J* 2006;27:1183–1189. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16481389>.
210. O'Brien ME, Ciuleanu TE, Tsekov H, et al. Phase III trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J Clin Oncol* 2006;24:5441–5447. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17135646>.
211. Eckardt JR, von Pawel J, Pujol JL, et al. Phase III study of oral compared with intravenous topotecan as second-line therapy in small-cell lung cancer. *J Clin Oncol* 2007;25:2086–2092. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17513814>.
212. Aix SP, Ciuleanu TE, Navarro A, et al. Combination lurbinectedin and doxorubicin versus physician's choice of chemotherapy in patients with relapsed small-cell lung cancer (ATLANTIS): a multicentre, randomised, open-label, phase 3 trial. *Lancet Respir Med* 2023;11:74–86. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36252599>.
213. Spigel DR, Vicente D, Ciuleanu TE, et al. Second-line nivolumab in relapsed small-cell lung cancer: CheckMate 331(☆). *Ann Oncol* 2021;32:631–641. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33539946>.
214. Goto K, Ohe Y, Shibata T, et al. Combined chemotherapy with cisplatin, etoposide, and irinotecan versus topotecan alone as second-line treatment for patients with sensitive relapsed small-cell lung cancer (JCOG0605): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2016;17:1147–1157. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27312053>.
215. Yamamoto N, Tsurutani J, Yoshimura N, et al. Phase II study of weekly paclitaxel for relapsed and refractory small cell lung cancer. *Anticancer Res* 2006;26:777–781. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16739353>.
216. Smit EF, Fokkema E, Biesma B, et al. A phase II study of paclitaxel in heavily pretreated patients with small-cell lung cancer. *Br J Cancer* 1998;77:347–351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9461009>.
217. von Eidf D, Bozorgmehr F, Chung I, et al. Paclitaxel for treatment of advanced small cell lung cancer (SCLC): a retrospective study of 185 patients. *J Thorac Dis* 2020;12:782–793. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32274145>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

218. Smyth JF, Smith IE, Sessa C, et al. Activity of docetaxel in small cell lung cancer. The early clinical trials group of the EORTC. Eur J Cancer 1994;30A:1058–1060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7654428>.
219. Masters GA, Declerck L, Blanke C, et al. Phase II trial of gemcitabine in refractory or relapsed small-cell lung cancer: Eastern Cooperative Oncology Group Trial 1597. J Clin Oncol 2003;21:1550–1555. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12697880>.
220. van der Lee I, Smit EF, van Putten JW, et al. Single-agent gemcitabine in patients with resistant small-cell lung cancer. Ann Oncol 2001;12:557–561. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11398892>.
221. Ready NE, Ott PA, Hellmann MD, et al. Nivolumab monotherapy and nivolumab plus ipilimumab in recurrent small cell lung cancer: Results from the CheckMate 032 randomized cohort. J Thorac Oncol 2020;15:426–435. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31629915>.
222. Antonia SJ, Lopez-Martin JA, Bendell J, et al. Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. Lancet Oncol 2016;17:883–895. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27269741>.
223. Ott PA, Elez E, Hirt S, et al. Pembrolizumab in patients with extensive-stage small-cell lung cancer: Results from the phase Ib KEYNOTE-028 study. J Clin Oncol 2017;35:3823–3829. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28813164>.
224. Horn L, Reck M, Spigel DR. The future of immunotherapy in the treatment of small cell lung cancer. Oncologist 2016;21:910–921. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27354668>.
225. Keeping ST, Cope S, Chan K, et al. Comparative effectiveness of nivolumab versus standard of care for third-line patients with small-cell lung cancer. J Comp Eff Res 2020;9:1275–1284. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33140652>.
226. Chung HC, Piha-Paul SA, Lopez-Martin J, et al. Pembrolizumab after two or more lines of previous therapy in patients with recurrent or metastatic SCLC: Results from the KEYNOTE-028 and KEYNOTE-158 studies. J Thorac Oncol 2020;15:618–627. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31870883>.
227. Rea F, Rizzardi G, Zuin A, et al. Outcome and surgical strategy in bronchial carcinoid tumors: single institution experience with 252 patients. Eur J Cardiothorac Surg 2007;31:186–191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17140801>.
228. Davies M, Duffield EA. Safety of checkpoint inhibitors for cancer treatment: strategies for patient monitoring and management of immune-mediated adverse events. Immunotargets Ther 2017;6:51–71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28894725>.
229. Khunger M, Rakshit S, Pasupuleti V, et al. Incidence of pneumonitis with use of programmed death 1 and programmed death-ligand 1 inhibitors in non-small cell lung cancer: A systematic review and meta-analysis of trials. Chest 2017;152:271–281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28499515>.
230. Johnson DH, Greco FA, Strupp J, et al. Prolonged administration of oral etoposide in patients with relapsed or refractory small-cell lung cancer: a phase II trial. J Clin Oncol 1990;8:1613–1617. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2170589>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

231. Einhorn LH, Pennington K, McClean J. Phase II trial of daily oral VP-16 in refractory small cell lung cancer: a Hoosier Oncology Group study. *Semin Oncol* 1990;17:32–35. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2154857>.
232. Pietanza MC, Waqar SN, Krug LM, et al. Randomized, double-blind, phase II study of temozolomide in combination with either veliparib or placebo in patients with relapsed-sensitive or refractory small-cell lung cancer. *J Clin Oncol* 2018;36:2386–2394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29906251>.
233. Zauderer MG, Drilon A, Kadota K, et al. Trial of a 5-day dosing regimen of temozolomide in patients with relapsed small cell lung cancers with assessment of methylguanine-DNA methyltransferase. *Lung Cancer* 2014;86:237–240. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25194640>.
234. Pietanza MC, Kadota K, Huberman K, et al. Phase II trial of temozolomide in patients with relapsed sensitive or refractory small cell lung cancer, with assessment of methylguanine-DNA methyltransferase as a potential biomarker. *Clin Cancer Res* 2012;18:1138–1145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22228633>.
235. Higgins KA, Simone CB, 2nd, Amini A, et al. American radium society appropriate use criteria on radiation therapy for extensive-stage sclc. *J Thorac Oncol* 2021;16:54–65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33011389>.
236. Chun SG, Simone CB, 2nd, Amini A, et al. American radium society appropriate use criteria: Radiation therapy for limited-stage SCLC 2020. *J Thorac Oncol* 2021;16:66–75. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33166720>.
237. Simone CB, 2nd, Bogart JA, Cabrera AR, et al. Radiation therapy for small cell lung cancer: An ASTRO clinical practice guideline. *Pract Radiat Oncol* 2020;10:158–173. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32222430>.
238. Kong FM, Lally BE, Chang JY, et al. ACR appropriateness criteria® radiation therapy for small-cell lung cancer. *Am J Clin Oncol* 2013;36:206–213. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23511336>.
239. Fried DB, Morris DE, Poole C, et al. Systematic review evaluating the timing of thoracic radiation therapy in combined modality therapy for limited-stage small-cell lung cancer. *J Clin Oncol* 2004;22:4837–4845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15570087>.
240. Pijls-Johannesma M, De Ruyscher D, Vansteenkiste J, et al. Timing of chest radiotherapy in patients with limited stage small cell lung cancer: a systematic review and meta-analysis of randomised controlled trials. *Cancer Treat Rev* 2007;33:461–473. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17513057>.
241. Murray N, Coy P, Pater JL, et al. Importance of timing for thoracic irradiation in the combined modality treatment of limited-stage small-cell lung cancer. The National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* 1993;11:336–344. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8381164>.
242. De Ruyscher D, Bremer RH, Koppe F, et al. Omission of elective node irradiation on basis of CT-scans in patients with limited disease small cell lung cancer: a phase II trial. *Radiother Oncol* 2006;80:307–312. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16949169>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

243. De Ruysscher D, Lueza B, Le Pechoux C, et al. Impact of thoracic radiotherapy timing in limited-stage small-cell lung cancer: usefulness of the individual patient data meta-analysis. *Ann Oncol* 2016;27:1818–1828. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27436850>.
244. Turrisi AT, 3rd, Kim K, Blum R, et al. Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with cisplatin and etoposide. *N Engl J Med* 1999;340:265–271. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9920950>.
245. Ganti AK, Fruth B, Rimner A, et al. Quality of life outcomes in patients participating in the CALGB 30610 trial (CALGB 70702): Alliance. *Cancer* 2025;131:e35663. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39580636>.
246. Bogart JA, Wang X, Masters GA, et al. Short communication: Interim toxicity analysis for patients with limited stage small cell lung cancer (LSCLC) treated on CALGB 30610 (alliance) / RTOG 0538. *Lung Cancer* 2021;156:68–71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33894496>.
247. Gronberg BH, Halvorsen TO, Flotten O, et al. Randomized phase II trial comparing twice daily hyperfractionated with once daily hypofractionated thoracic radiotherapy in limited disease small cell lung cancer. *Acta Oncol* 2016;55:591–597. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26494411>.
248. Turgeon GA, Souhami L, Kopek N, et al. Thoracic irradiation in 3weeks for limited-stage small cell lung cancer: Is twice a day fractionation really needed? *Cancer Radiother* 2017;21:89–98. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28325618>.
249. Bogart J, Wang X, Masters G, et al. High-dose once-daily thoracic radiotherapy in limited-stage small-cell lung cancer: CALGB 30610 (alliance)/RTOG 0538. *J Clin Oncol* 2023;41:2394–2402. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36623230>.
250. Qiu B, Li Q, Liu J, et al. Moderately hypofractionated once-daily compared with twice-daily thoracic radiation therapy concurrently with etoposide and cisplatin in limited-stage small cell lung cancer: A multicenter, phase II, randomized trial. *Int J Radiat Oncol Biol Phys* 2021;111:424–435. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33992717>.
251. Gronberg BH, Killingberg KT, Flotten O, et al. High-dose versus standard-dose twice-daily thoracic radiotherapy in limited-stage SCLC: Final survival data, long-term toxicity, and relapse patterns in a randomized, open-label, phase ii trial. *J Thorac Oncol* 2025;20:1108–1119. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40258573>.
252. Yu J, Jiang L, Zhao L, et al. High-dose hyperfractionated simultaneous integrated boost radiotherapy versus standard-dose radiotherapy for limited-stage small-cell lung cancer in China: a multicentre, open-label, randomised, phase 3 trial. *Lancet Respir Med* 2024;12:799–809. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39146944>.
253. Yu J, Zhang J, Chang X, et al. Real-world clinical outcomes in patients with limited-stage small cell lung cancer who received high-dose hyperfractionated simultaneous integrated boost radiation therapy versus standard-dose radiation therapy. *Pract Radiat Oncol* 2026. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/42034286>.
254. Shirvani SM, Juloori A, Allen PK, et al. Comparison of 2 common radiation therapy techniques for definitive treatment of small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2013;87:139–147. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23920393>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

255. ICRU report 83: Prescribing, recording, and reporting intensity-modulated photon-beam therapy. Bethesda, MD: International Commission on Radiation Units and Measurements (ICRU); 2010. Available at: <https://www.icru.org/report/prescribing-recording-and-reporting-intensity-modulated-photon-beam-therapy-imrticru-report-83>.
256. Gregoire V, Mackie TR. State of the art on dose prescription, reporting and recording in Intensity-Modulated Radiation Therapy (ICRU report No. 83). Cancer Radiother 2011;15:555–559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21802333>.
257. Hartford AC, Palisca MG, Eichler TJ, et al. American society for therapeutic radiology and oncology (ASTRO) and american college of radiology (ACR) practice guidelines for intensity-modulated radiation therapy (IMRT). Int J Radiat Oncol Biol Phys 2009;73:9–14. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19100920>.
258. Shirvani SM, Komaki R, Heymach JV, et al. Positron emission tomography/computed tomography-guided intensity-modulated radiotherapy for limited-stage small-cell lung cancer. Int J Radiat Oncol Biol Phys 2012;82:e91–97. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21489716>.
259. Chun SG, Hu C, Choy H, et al. Impact of intensity-modulated radiation therapy technique for locally advanced non-small-cell lung cancer: A secondary analysis of the NRG oncology RTOG 0617 randomized clinical trial. J Clin Oncol 2017;35:56–62. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28034064>.
260. Bogart JA, Herndon JE, 2nd, Lyss AP, et al. 70 Gy thoracic radiotherapy is feasible concurrent with chemotherapy for limited-stage small-cell lung cancer: analysis of Cancer and Leukemia Group B study 39808. Int J Radiat Oncol Biol Phys 2004;59:460–468. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15145163>.
261. Liengswangwong V, Bonner JA, Shaw EG, et al. Limited-stage small-cell lung cancer: patterns of intrathoracic recurrence and the implications for thoracic radiotherapy. J Clin Oncol 1994;12:496–502. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8120547>.
262. Kim TH, Cho KH, Pyo HR, et al. Dose-volumetric parameters for predicting severe radiation pneumonitis after three-dimensional conformal radiation therapy for lung cancer. Radiology 2005;235:208–215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15703313>.
263. Rose J, Rodrigues G, Yaremko B, et al. Systematic review of dose-volume parameters in the prediction of esophagitis in thoracic radiotherapy. Radiother Oncol 2009;91:282–287. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18950881>.
264. Kong FM, Ritter T, Quint DJ, et al. Consideration of dose limits for organs at risk of thoracic radiotherapy: atlas for lung, proximal bronchial tree, esophagus, spinal cord, ribs, and brachial plexus. Int J Radiat Oncol Biol Phys 2011;81:1442–1457. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20934273>.
265. Videtic GM, Stephans KL, Woody NM, et al. Stereotactic body radiation therapy-based treatment model for stage I medically inoperable small cell lung cancer. Pract Radiat Oncol 2013;3:301–306. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24674402>.
266. Alongi F, Arcangeli S, De Bari B, et al. Stage-I small cell lung cancer: A new potential option for stereotactic ablative radiation therapy? A review of literature. Crit Rev Oncol Hematol 2017;112:67–71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28325266>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

267. Rathod S, Koul R, Bashir B, et al. Role of stereotactic body radiation therapy in early stage small cell lung cancer in the era of lung cancer screening: A systematic review. *Am J Clin Oncol* 2019;42:123–130. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30418179>.
268. Shioyama Y, Nakamura K, Sasaki T, et al. Clinical results of stereotactic body radiotherapy for stage I small-cell lung cancer: a single institutional experience. *J Radiat Res* 2013;54:108–112. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22923748>.
269. Li C, Xiong Y, Zhou Z, et al. Stereotactic body radiotherapy with concurrent chemotherapy extends survival of patients with limited stage small cell lung cancer: a single-center prospective phase II study. *Med Oncol* 2014;31:369. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25416052>.
270. Safavi AH, Mak DY, Boldt RG, et al. Stereotactic ablative radiotherapy in T1-2N0M0 small cell lung cancer: A systematic review and meta-analysis. *Lung Cancer* 2021;160:179–186. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34330566>.
271. Verma V, Simone CB, 2nd, Allen PK, Lin SH. Outcomes of stereotactic body radiotherapy for T1-T2N0 small cell carcinoma according to addition of chemotherapy and prophylactic cranial irradiation: a multicenter analysis. *Clin Lung Cancer* 2017;18:675–681 e671. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28408183>.
272. Aizer AA, Shin KY, Catalano PJ, et al. Treatment for brain metastases with stereotactic radiation vs hippocampal-avoidance whole brain radiation: A randomized clinical trial. *JAMA* 2026;335:1127–1136. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41712219>.
273. Aizer AA, Tanguturi SK, Shi DD, et al. Stereotactic radiosurgery in patients with small cell lung cancer and 1-10 brain metastases: A multi-institutional, phase II, prospective clinical trial. *J Clin Oncol* 2025;43:2986–2997. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40644657>.
274. Jeremic B, Shibamoto Y, Nikolic N, et al. Role of radiation therapy in the combined-modality treatment of patients with extensive disease small-cell lung cancer: A randomized study. *J Clin Oncol* 1999;17:2092–2099. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10561263>.
275. Slotman BJ, van Tinteren H, Praag JO, et al. Use of thoracic radiotherapy for extensive stage small-cell lung cancer: a phase 3 randomised controlled trial. *Lancet* 2015;385:36–42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25230595>.
276. Yee D, Butts C, Reiman A, et al. Clinical trial of post-chemotherapy consolidation thoracic radiotherapy for extensive-stage small cell lung cancer. *Radiother Oncol* 2012;102:234–238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21930323>.
277. Slotman BJ, van Tinteren H, Praag JO, et al. Radiotherapy for extensive stage small-cell lung cancer - Authors' reply. *Lancet* 2015;385:1292–1293. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25890910>.
278. Putora PM, Glatzer M, De Ruyscher D, et al. Consolidative thoracic radiotherapy in stage IV small cell lung cancer: Selection of patients amongst European IASLC and ESTRO experts. *Radiother Oncol* 2019;135:74–77. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31015173>.
279. Arriagada R, Le Chevalier T, Borie F, et al. Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission. *J Natl Cancer Inst* 1995;87:183–190. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7707405>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

280. Yang Y, Zhang D, Zhou X, et al. Prophylactic cranial irradiation in resected small cell lung cancer: A systematic review with meta-analysis. *J Cancer* 2018;9:433–439. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29344290>.
281. Auperin A, Arriagada R, Pignon JP, et al. Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission. Prophylactic Cranial Irradiation Overview Collaborative Group. *N Engl J Med* 1999;341:476–484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10441603>.
282. Tomassen ML, Pomp J, van der Stap J, et al. The overall survival impact of prophylactic cranial irradiation in limited-stage small-cell lung cancer: A systematic review and meta-analysis. *Clin Transl Radiat Oncol* 2022;33:145–152. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35243025>.
283. Patel S, Macdonald OK, Suntharalingam M. Evaluation of the use of prophylactic cranial irradiation in small cell lung cancer. *Cancer* 2009;115:842–850. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19117355>.
284. Eze C, Roengvoraphoj O, Niyazi M, et al. Treatment response and prophylactic cranial irradiation are prognostic factors in a real-life limited-disease small-cell lung cancer patient cohort comprehensively staged with cranial magnetic resonance imaging. *Clin Lung Cancer* 2017;18:e243–e249. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28065620>.
285. Held MK, Hansen O, Schytte T, et al. Outcomes of prophylactic cranial irradiation in patients with small cell lung cancer in the modern era of baseline magnetic resonance imaging of the brain. *Acta Oncol* 2022;61:185–192. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34583620>.
286. Sharma S, McMillan MT, Doucette A, et al. Effect of prophylactic cranial irradiation on overall survival in metastatic small-cell lung cancer: A propensity score-matched analysis. *Clin Lung Cancer* 2018;19:260–269.e263. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29358031>.
287. Bang A, Kendal WS, Laurie SA, et al. Prophylactic cranial irradiation in extensive stage small cell lung cancer: Outcomes at a comprehensive cancer centre. *Int J Radiat Oncol Biol Phys* 2018;101:1133–1140. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29908788>.
288. Slotman B, Faivre-Finn C, Kramer G, et al. Prophylactic cranial irradiation in extensive small-cell lung cancer. *N Engl J Med* 2007;357:664–672. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17699816>.
289. Takahashi T, Yamanaka T, Seto T, et al. Prophylactic cranial irradiation versus observation in patients with extensive-disease small-cell lung cancer: a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2017;18:663–671. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28343976>.
290. Edelman MJ. Prophylactic cranial irradiation for small-cell lung cancer: Time for a reassessment. *Am Soc Clin Oncol Educ Book* 2020;40:24–28. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32421453>.
291. Lee JS, Umsawasdi T, Lee YY, et al. Neurotoxicity in long-term survivors of small cell lung cancer. *Int J Radiat Oncol Biol Phys* 1986;12:313–321. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3007407>.
292. Slotman BJ, Senan S. Radiotherapy in small-cell lung cancer: lessons learned and future directions. *Int J Radiat Oncol Biol Phys* 2011;79:998–1003. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21353159>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

293. Slotman BJ, Mauer ME, Bottomley A, et al. Prophylactic cranial irradiation in extensive disease small-cell lung cancer: short-term health-related quality of life and patient reported symptoms: results of an international phase III randomized controlled trial by the EORTC radiation oncology and lung cancer groups. *J Clin Oncol* 2009;27:78–84. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19047288>.
294. Pechoux CL, Sun A, Slotman BJ, et al. Prophylactic cranial irradiation for patients with lung cancer. *Lancet Oncol* 2016;17:e277–e293. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27396646>.
295. Eze C, Roengvoraphoj O, Manapov F. Prophylactic cranial irradiation in resected early-stage small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2017;98:612–614. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28581402>.
296. Le Pechoux C, Dunant A, Senan S, et al. Standard-dose versus higher-dose prophylactic cranial irradiation (PCI) in patients with limited-stage small-cell lung cancer in complete remission after chemotherapy and thoracic radiotherapy (PCI 99-01, EORTC 22003-08004, RTOG 0212, and IFCT 99-01): a randomised clinical trial. *Lancet Oncol* 2009;10:467–474. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19386548>.
297. Brown PD, Pugh S, Laack NN, et al. Memantine for the prevention of cognitive dysfunction in patients receiving whole-brain radiotherapy: a randomized, double-blind, placebo-controlled trial. *Neuro Oncol* 2013;15:1429–1437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23956241>.
298. Brown PD, Gondi V, Pugh S, et al. Hippocampal avoidance during whole-brain radiotherapy plus memantine for patients with brain metastases: Phase III trial NRG oncology CC001. *J Clin Oncol* 2020;38:1019–1029. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32058845>.
299. Rodriguez de Dios N, Counago F, Murcia-Mejia M, et al. Randomized phase III trial of prophylactic cranial irradiation with or without hippocampal avoidance for small-cell lung cancer (PREMER): A GICOR-GOECF-SEOR study. *J Clin Oncol* 2021;39:3118–3127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34379442>.
300. Belderbos JSA, De Ruyscher DKM, De Jaeger K, et al. Phase 3 randomized trial of prophylactic cranial irradiation with or without hippocampus avoidance in SCLC (NCT01780675). *J Thorac Oncol* 2021;16:840–849. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33545387>.
301. Gondi V, Pugh SL, Mehta MP, et al. Hippocampal avoidance during prophylactic cranial irradiation for patients with small cell lung cancer: Randomized phase II/III trial NRG-CC003. *J Clin Oncol* 2025;43:3516–3525. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40789106>.
302. Maranzano E, Trippa F, Casale M, et al. 8Gy single-dose radiotherapy is effective in metastatic spinal cord compression: results of a phase III randomized multicentre Italian trial. *Radiother Oncol* 2009;93:174–179. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19520448>.
303. Lutz S, Berk L, Chang E, et al. Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. *Int J Radiat Oncol Biol Phys* 2011;79:965–976. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21277118>.
304. Ferrell B, Koczywas M, Grannis F, Harrington A. Palliative care in lung cancer. *Surg Clin North Am* 2011;91:403–417, ix. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21419260>.



NCCN Guidelines Version 1.2027

Small Cell Lung Cancer

305. Tsao MN, Lloyd N, Wong RK, et al. Whole brain radiotherapy for the treatment of newly diagnosed multiple brain metastases. Cochrane Database Syst Rev 2012;4:CD003869. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22513917>.
306. Rusthoven CG, Yamamoto M, Bernhardt D, et al. Evaluation of first-line radiosurgery vs whole-brain radiotherapy for small cell lung cancer brain metastases: The FIRE-SCLC cohort study. JAMA Oncol 2020;6:1028–1037. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32496550>.
307. Viani GA, Gouveia AG, Louie AV, Moraes FY. Stereotactic radiosurgery for brain metastases from small cell lung cancer without prior whole-brain radiotherapy: A meta-analysis. Radiother Oncol 2021;162:45–51. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34171453>.
308. Bernhardt D, Bozorgmehr F, Adeberg S, et al. Outcome in patients with small cell lung cancer re-irradiated for brain metastases after prior prophylactic cranial irradiation. Lung Cancer 2016;101:76–81. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27794411>.
309. Wegner RE, Olson AC, Kondziolka D, et al. Stereotactic radiosurgery for patients with brain metastases from small cell lung cancer. Int J Radiat Oncol Biol Phys 2011;81:e21–27. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21345622>.