



National Comprehensive
Cancer Network®


NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Kidney Cancer

NCCN Evidence Blocks™

Version 2.2026 — April 14, 2026

**NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.**





NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

***Robert J. Motzer, MD/Chair † ♢**
Memorial Sloan Kettering Cancer Center

***Eric Jonasch, MD/Vice-chair †**
The University of Texas
MD Anderson Cancer Center

Neeraj Agarwal, MD ‡ †
Huntsman Cancer Institute
at the University of Utah

Hilary Bagshaw, MD §
Stanford Cancer Institute

Michael Baine, MD, PhD §
Fred & Pamela Buffet Cancer Center

Maria I. Carlo, MD †
Memorial Sloan Kettering Cancer Center

Toni K. Choueiri, MD † ♢
Dana-Farber/Brigham and Women's
Cancer Center | Mass General Cancer Center

Brian A. Costello, MD, MS †
Mayo Clinic Comprehensive Cancer Center

Ithaar H. Derweesh, MD ω
UC San Diego Moores Cancer Center

Arpita Desai, MD † ♢
UCSF Helen Diller Family
Comprehensive Cancer Center

Yasser Ged, MBBS †
Johns Hopkins Kimmel Cancer Center

Saby George, MD †
Roswell Park Comprehensive Cancer Center

Nikhil Gopal, MD ω
St. Jude Children's Research Hospital/The
University of Tennessee Health Science
Center

John L. Gore, MD, MS ω
Fred Hutchinson Cancer Center

Andrew Gunn, MD ∩
O'Neal Comprehensive Cancer Center at UAB

Naomi Haas, MD †
Abramson Cancer Center at the
University of Pennsylvania

Angela Y. Jia, MD, PhD §
Case Comprehensive Cancer Center/University
Hospitals Seidman Cancer Center and
Cleveland Clinic Taussig Cancer Institute

Samuel Kaffenberger, MD ω
University of Michigan Rogel Cancer Center

Payal Kapur, MD ≠
UT Southwestern Simmons
Comprehensive Cancer Center

Jennifer King, MD ♢
Indiana University Melvin and Bren Simon
Comprehensive Cancer Center

Christos Kyriakopoulos, MD ‡
University of Wisconsin Carbone Cancer Center

Elaine T. Lam, MD †
University of Colorado Cancer Center

Primo N. Lara Jr., MD †
UC Davis Comprehensive Cancer Center

Clayton Lau, MD ω
City of Hope National Medical Center

Bryan Lewis, JD ≠
KidneyCAN

David C. Madoff, MD ∩
Yale Cancer Center/Smilow Cancer Hospital

***Brandon Manley, MD ω**
Moffitt Cancer Center

M. Dror Michaelson, MD, PhD †
Dana-Farber/Brigham and Women's
Cancer Center | Mass General Cancer Center

Amir Mortazavi, MD †
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Kent Perry, Jr, MD ω
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Brian Rini, MD †
Vanderbilt-Ingram Cancer Center

Brian Shuch, MD ω
UCLA Jonsson Comprehensive
Cancer Center

Randy F. Sweis, MD †
The UChicago Medicine
Comprehensive Cancer Center

Lewis Thomas, MD ω
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

Matthew Zibelman, MD †
Fox Chase Cancer Center

NCCN
Ryan Schonfeld, BA
MaryElizabeth Stein, PhD

‡ Hematology/Hematology oncology
♢ Internal medicine
∩ Interventional radiology
† Medical oncology
≠ Patient advocacy
≠ Pathology
§ Radiotherapy/Radiation oncology
ω Urology
*Discussion writing committee member

[NCCN Guidelines Panel Disclosures](#)



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

[NCCN Kidney Cancer Panel Members](#)
[NCCN Evidence Blocks Definitions \(EB-DEF\)](#)

Kidney Cancer

[Initial Workup \(KID-1\)](#)

[Primary Treatment and Follow-Up for Stage I \(KID-1\)](#)

[Primary Treatment and Follow-Up for Stages II and III \(KID-2\)](#)

[Primary Treatment for Stage IV \(KID-3\)](#)

[Treatment for Stage IV or Relapsed Disease \(KID-4\)](#)

[General Principles of Management for Renal Cell Carcinoma \(KID-A\)](#)

[Principles of Radiation Therapy \(KID-B\)](#)

[Follow-up \(KID-C\)](#)

[Principles of Systemic Therapy for Stage IV or Relapsed Disease \(KID-D\)](#)

[Risk Models to Direct Treatment \(KID-E\)](#)

Hereditary Renal Cell Carcinoma

[Criteria for Further Genetic Risk Evaluation for Hereditary RCC Syndromes \(HERED-RCC-1\)](#)

[Hereditary RCC Syndromes Overview \(HERED-RCC-2\)](#)

[Genetic Testing \(GENE-1\)](#)

[Kidney-Specific Screening Recommendations for Patients With Confirmed Hereditary RCC](#)

[Who Do Not Yet Have a Radiographic or Pathologic Diagnosis of RCC \(HERED-RCC-B\)](#)

[Kidney-Specific Surgical Recommendations for Patients With Confirmed Hereditary RCC \(HERED-RCC-C\)](#)

[Kidney-Specific Systemic Therapy for Patients With Confirmed Hereditary RCC \(HERED-RCC-D\)](#)

Find an NCCN Member Institution:

<https://www.nccn.org/home/member-institutions>.

NCCN Categories of

Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference:

All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

NCCN Guidelines for Patients®
available at www.nccn.org/patients

[Evidence Blocks](#)
[Staging \(ST-1\)](#)
[Abbreviations \(ABBR-1\)](#)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS

5					
4					
3					
2					
1					

E S Q C A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

Example Evidence Block

5					
4					
3					
2					
1					

E S Q C A

E = 4
S = 4
Q = 3
C = 4
A = 3

Efficacy of Regimen/Agent

5	Highly effective: Cure likely and often provides long-term survival advantage
4	Very effective: Cure unlikely but sometimes provides long-term survival advantage
3	Moderately effective: Modest impact on survival, but often provides control of disease
2	Minimally effective: No, or unknown impact on survival, but sometimes provides control of disease
1	Palliative: Provides symptomatic benefit only

Safety of Regimen/Agent

5	Usually no meaningful toxicity: Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
4	Occasionally toxic: Rare significant toxicities or low-grade toxicities only; little interference with ADLs
3	Mildly toxic: Mild toxicity that interferes with ADLs
2	Moderately toxic: Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
1	Highly toxic: Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

Note: For significant chronic or long-term toxicities, score decreased by 1

Quality of Evidence


5	High quality: Multiple well-designed randomized trials and/or meta-analyses
4	Good quality: One or more well-designed randomized trials
3	Average quality: Low quality randomized trial(s) or well-designed non-randomized trial(s)
2	Low quality: Case reports or extensive clinical experience
1	Poor quality: Little or no evidence

Consistency of Evidence

5	Highly consistent: Multiple trials with similar outcomes
4	Mainly consistent: Multiple trials with some variability in outcome
3	May be consistent: Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
2	Inconsistent: Meaningful differences in direction of outcome between quality trials
1	Anecdotal evidence only: Evidence in humans based upon anecdotal experience

Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)

5	Very inexpensive
4	Inexpensive
3	Moderately expensive
2	Expensive
1	Very expensive



^a Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health. See [NCCN Guidelines for Distress Management](#) (DIS-A).

^b Imaging with and without contrast is strongly preferred, such as a renal protocol for abdomen.

^c Biopsy of small lesions may be considered to obtain or confirm a diagnosis of malignancy and guide decisions on surveillance, percutaneous ablation, and/or surgery.

d General Principles of Management for Renal Cell Carcinoma (KID-A).

^e SBRT is an option for patients considered non-optimal for either surgery or percutaneous ablation for clinical stage T1a kidney cancer (category 2A). SBRT is an option for non-optimal surgical candidates for clinical stage T1b kidney cancer (category 2A). SBRT may be considered for non-optimal surgical candidates with stage II (category 2B) or III (category 3) kidney cancer. See [Principles of Radiation Therapy \(KID-B\)](#).

f Follow-up (KID-C).

⁹ No single follow-up plan is appropriate for all patients. Follow-up should be individualized based on patient requirements.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

STAGE	PRIMARY TREATMENT ^{d,e}	ADJUVANT TREATMENT ^f	FOLLOW-UP ^g (CATEGORY 2B)
Stage II →	Partial nephrectomy or Radical nephrectomy or SBRT (in select patients) (category 2B)	Clear cell histology: Surveillance or Adjuvant Pembrolizumab ⁱ (category 1) (Grade 4 tumors with clear cell histology ± sarcomatoid features) Non-clear cell histology: Surveillance	Follow-up → (KID-C) Relapse or progression (KID-4)
Stage III ^h →	Radical nephrectomy or Partial nephrectomy, if clinically indicated or SBRT (in select patients) (category 3)	Clear cell histology: Adjuvant Pembrolizumab ⁱ (category 1) or Surveillance Non-clear cell histology: Surveillance or clinical trial	

[See Evidence Blocks on EB-1](#)

^d [General Principles of Management for Renal Cell Carcinoma \(KID-A\)](#).

^e SBRT is an option for patients considered non-optimal for either surgery or percutaneous ablation for clinical stage T1a kidney cancer (category 2A). SBRT is an option for non-optimal surgical candidates for clinical stage T1b kidney cancer (category 2A). SBRT may be considered for non-optimal surgical candidates with stage II (category 2B) or III (category 3) kidney cancer. See [Principles of Radiation Therapy \(KID-B\)](#).

^f [Follow-up \(KID-C\)](#).

^g No single follow-up plan is appropriate for all patients. Follow-up should be individualized based on patient requirements.

^h For patients with clinical T3 or T4 tumors without evidence of metastatic disease but in whom complete resection is not feasible up front, or which have imperative indication for nephron preservation (eg, solitary kidney, chronic kidney disease, bilateral tumors) and in which a partial nephrectomy is not feasible, utilization of presurgical neoadjuvant therapy may be considered following confirmation of clear cell histology to achieve cytoreduction, which may enable complete resection or facilitate a partial nephrectomy, respectively (Hakimi K, et al. BJU Int 2024;133:425-431).

ⁱ Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™



^d [General Principles of Management for Renal Cell Carcinoma \(KID-A\)](#).

^f [Follow-up \(KID-C\)](#).

^g No single follow-up plan is appropriate for all patients. Follow-up should be individualized based on patient requirements.

^h For patients with clinical T3 or T4 tumors without evidence of metastatic disease but in whom complete resection is not feasible up front, or which have imperative indication for nephron preservation (eg, solitary kidney, chronic kidney disease, bilateral tumors) and in which a partial nephrectomy is not feasible, utilization of presurgical neoadjuvant therapy may be considered following confirmation of clear cell histology to achieve cytoreduction, which may enable complete resection or facilitate a partial nephrectomy, respectively (Hakimi K, et al. BJU Int 2024;133:425-431).

ⁱ Pembrolizumab and berahyaluronidase alfa-pmpH subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmpH has different dosing and administration instructions compared to IV pembrolizumab.

^j Individualize treatment based on symptoms and extent of metastatic disease.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**




NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

STAGE IV OR RELAPSED DISEASE



[See Evidence Blocks on EB-3](#)

ⁱ Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

^k [Principles of Radiation Therapy \(KID-B\)](#).

^l Best supportive care can include radiation therapy (RT) where SBRT is the preferred approach, bisphosphonates, or receptor activator of nuclear factor kappa-B (RANK) ligand inhibitors for bony metastases. An FDA-approved biosimilar is an appropriate substitute for denosumab.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)
All recommendations are category 2A unless otherwise indicated.



GENERAL PRINCIPLES OF MANAGEMENT FOR RENAL CELL CARCINOMA

- Nephron-sparing surgery (partial nephrectomy) is recommended in select patients, such as:
 - ▶ Patients with unilateral stage I–III tumors, where technically feasible
 - ▶ Patients with uninephric state, renal insufficiency, bilateral renal masses, and familial renal cell cancer
 - ▶ Patients at relative risk for developing progressive chronic kidney disease due to young age or medical risk factors (ie, hypertension, diabetes, nephrolithiasis)
- Open, laparoscopic, or robotic surgical techniques may be used to perform radical and partial nephrectomies.
- Regional lymph node dissection is optional but should be considered for patients with resectable adenopathy on preoperative imaging or palpable/visible adenopathy at time of surgery.
- Lymph node dissection: Right-sided tumors¹
 - ▶ Renal hilar, paracaval, precaval nodes (crus of the diaphragm to aortic bifurcation)
 - ▶ Extended lymph node dissection: Inter-aortocaval/retrocaval and right common iliac nodes
- Lymph node dissection: Left-sided tumors¹
 - ▶ Renal hilar, preaortic, and paraaortic nodes (crus of the diaphragm to aortic bifurcation)
 - ▶ Extended lymph node dissection: Inter-aortocaval/retroaortic and left common iliac nodes
- If adrenal gland is uninvolved, adrenalectomy may be omitted.
- Special teams or referral to high-volume centers may be required for extensive inferior vena cava involvement.
- Percutaneous ablation (eg, cryosurgery, radiofrequency ablation, microwave ablation) is an option for the management of clinical stage T1 renal lesions.
 - ▶ Percutaneous ablation is suitable for renal masses ≤3 cm.
 - ▶ Percutaneous ablation is an option for clinical T1b masses in select patients not eligible for surgery.
 - ▶ Biopsy of lesions is recommended to be done prior to or at time of ablation.
 - ▶ Percutaneous ablation may require retreatment to achieve the same local oncologic outcomes as conventional surgery.^{2,3}
- SBRT is an option for patients considered non-optimal for either surgery or percutaneous ablation for clinical stage T1a kidney cancer (category 2A). SBRT is an option for non-optimal surgical candidates for clinical stage T1b kidney cancer (category 2A). SBRT may be considered for non-optimal surgical candidates with stage II (category 2B) or III (category 3) kidney cancer ([KID-1](#), [KID-2](#)).
- Active surveillance is an option for the initial management of clinical stage T1 renal lesions, for example:
 - ▶ It is an option for renal masses <2 cm given the high rates of benign tumors and low metastatic potential of these masses.
 - ▶ Active surveillance of patients with T1a tumors (≤4 cm) that have a predominantly cystic component is recommended.
 - ▶ It is an option for patients with clinical stage T1 masses and significant competing risks of death or morbidity from intervention.
 - ▶ Active surveillance entails serial abdomen imaging with timely intervention should the mass demonstrate changes (eg, increasing tumor size, growth rate, infiltrative pattern) indicative of increasing metastatic potential.
 - ▶ Active surveillance should include periodic metastatic survey including blood work and chest imaging, particularly if the mass demonstrates growth.
- Generally, patients who would be candidates for cytoreductive nephrectomy prior to systemic therapy have:
 - ▶ Excellent performance status (ECOG PS <2)
 - ▶ No brain metastasis
- Patients either with large-volume distant metastases or tumors with large sarcomatoid burdens should receive systemic therapy prior to cytoreductive nephrectomy.

¹ Campi R, Sessa F, Di Maida F, et al. Templates of lymph node dissection for renal cell carcinoma: A systematic review of the literature. *Front Surg* 2018;5:76.

² Campbell S, Uzzo R, Allaf M, et al. Renal mass and localized renal cancer: AUA Guideline. *J Urol* 2017;198:520-529.

³ Pierorazio P, Johnson M, Patel H, et al. Management of renal masses and localized renal cancer: Systematic review and meta-analysis. *J Urol* 2016;196:989-999.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY

General Treatment Information:

- Consider radiation therapy (RT) in the following situations:

- Modalities: SBRT should be considered as the primary radiation modality in all situations unless precluded by anatomic site, proximity to organs at risk (OAR), or past treatments.

Primary Disease:

- Definitive therapy^{1,2}

- Definitive radiation using SBRT may be considered as a treatment option for non-optimal surgical candidates. Consider referral to a high-volume center with specialized expertise.
- Patient selection: SBRT can be considered for patients with T1 and T2a tumors (≤ 10 cm in diameter). There are insufficient data to pursue SBRT in tumors > 7 cm. When delivering SBRT to tumors abutting bowel, consideration should be given to the use of a 5-fraction schedule, with strict adherence (eg, priority 1) to bowel constraints, with target coverage designated as priority 2. Volumetric image guidance with motion management is required. When available, strong consideration should be given to on-table adaptive technology—such as CT-guided or MR-guided adaptive techniques. In the absence of online adaptation, planning volume (PRV) constraints must be applied. To further mitigate risk of gastrointestinal toxicity, an every-other-day schedule may be considered.
- Dosing regimens: SBRT should be delivered using over 1–5 fractions. Conventional fractionation is discouraged. Dose and fractionation options should attempt to keep the biologically effective dose (BED) to ≥ 80 Gy assuming an alpha/beta ratio of 10 due to association with improved local control. Established dosing regimens include:
 - ◊ 25–26 Gy in 1 fraction^a
 - ◊ 39–48 Gy in 3 fractions
 - ◊ 48 Gy in 4 fractions
 - ◊ 40–50 Gy in 5 fractions^b
- For multi-fraction dosing, treatment can be delivered on consecutive or non-consecutive days.
- Treatment technique:
 - ◊ For simulation and treatment planning four-dimensional (4D)-CT and fusion with a renal protocol CT and/or renal MRI is strongly encouraged.
 - ◊ Motion management is essential, with method dependent upon treatment platform and institutional practice. An internal target volume (ITV) should be utilized pending method of motion management.
 - ◊ When available, on-table adaptive technology such as CT-guided or MR-guided adaptive techniques should be utilized in the planning of tumors abutting bowel.
 - ◊ Use of daily pretreatment imaging including MRI if available or cone-beam CT is recommended. Volumetric image guidance is required when treating tumors abutting bowel.
 - ◊ OAR should be contoured to include a 3-mm PRV and to account for motion on 4D-CT.^c Bowel PRV of 5 mm should be used when planning tumor abutting bowel.
 - ◊ OAR constraints for SBRT 1- and 3-fraction regimens per FASTRACK II protocol.³ OAR constraints for SBRT 5-fraction regimen per UK Consensus guidelines.⁴

^a The use of single-fraction regimens has been largely limited to small (≤ 4 cm) tumors.

^b The regimen of 40 Gy in 5 fractions has a lower BED than others listed; its use is favored only in settings where dose-volume criteria for OAR cannot be met with higher BED regimens (eg, larger tumors or those with unfavorable anatomic relationships).

^c For specific representative OAR dose constraints, see TROG 15.03 FASTRACK II Protocol: Siva S, et al. Lancet Oncol 2024;25:308-316.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

KID-B
1 OF 5



PRINCIPLES OF RADIATION THERAPY

Distant Metastatic Disease:

- **Ablative treatment for intact extracranial metastases:**⁵⁻⁷
 - SBRT may offer more durable local control.⁸ SBRT should be considered for patients with oligometastasis unless metastasectomy is planned or SBRT cannot be delivered due to anatomic site, proximity to OAR, or past treatments.⁸⁻¹³ Strict adherence to normal tissue constraints is recommended. For multi-fraction dosing, treatment can be delivered on consecutive or non-consecutive days.
- **Spine SBRT regimens include but are not limited to:**
 - 16–24 Gy in 1 fraction¹⁴
 - 20–24 Gy in 2 fractions¹⁵
 - 24–27 Gy in 3 fractions¹⁶
 - 25–40 Gy in 5 fractions
- **SBRT regimens for other body sites include but are not limited to:**
 - 16–24 Gy in 1 fraction¹⁴
 - 48–60 Gy in 3 fractions^{8,17}
 - 40–60 Gy in 4–5 fractions^{8,18}
- **Palliative treatment of symptomatic extracranial metastases:**
 - A variety of treatment regimens are acceptable depending on location and/or clinical indication. Higher doses and/or hypofractionated regimens may be associated with more robust and durable palliation.^{14,19,20} SBRT dosing and fractionation should be pursued when possible. Strict adherence to normal tissue constraints is recommended. For multi-fraction dosing, treatment can be delivered on consecutive or non-consecutive days.
- **Preferred regimens include:**
 - 20–24 Gy in 2 fractions (spine only)¹⁵
 - 24–27 Gy in 3 fractions^{21,22}
 - 32–48 Gy in 4 fractions²³
 - 30–50 Gy in 5 fractions
 - 36 Gy in 6 fractions
- **Other potential regimens include:**
 - 8 Gy in 1 fraction²⁴
 - 20 Gy in 5 fractions²⁴
 - 30 Gy in 10 fractions²⁴

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

KID-B
2 OF 5



PRINCIPLES OF RADIATION THERAPY

• Brain metastases:

- ▶ **Stereotactic radiosurgery (SRS) and fractionated stereotactic RT (SRT)** are techniques for delivering a high dose of radiation to a specific target while delivering a minimal dose to surrounding tissues, generally in the brain and spine and in 1 to 5 sessions. Image-guided RT (IGRT) should be used to improve accuracy of radiotherapy delivery, where clinically appropriate.
- ▶ **SRS or SRT as primary treatment**
 - ◊ Smaller tumors may be treated with maximal doses of 15–24 Gy in 1 fraction according to volume guidelines based on maximum tolerated dose results from the RTOG 90-05 dose escalation study (shown below).²⁵ Caution is recommended for lesions >3 cm, and single-fraction radiosurgery is not typically recommended for lesions >4 cm.
 - Lesions with maximum diameter ≤20 mm receive up to 24 Gy
 - Lesions with maximum diameter 21–30 mm receive up to 18 Gy
 - Lesions with maximum diameter 31–40 mm receive up to 15 Gy
 - ◊ Larger tumors, however, may be treated with fractionated SRT. Potential regimens include, but are not limited to:^{26,27}
 - 24–27 Gy in 3 fractions
 - 25–35 Gy in 5 fractions
- ▶ **SRS/SRT as adjuvant treatment**
 - ◊ Smaller cavities may be treated with single-fraction SRS maximal doses ranging from 12–20 Gy depending on cavity volume per the NCCTG N107C trial protocol.²⁸
 - Lesions <4.2 cc receive 20 Gy
 - Lesions ≥4.2 cc to <8.0 cc receive 18 Gy
 - Lesions ≥8.0 cc to <14.4 cc receive 17 Gy
 - Lesions ≥14.4 cc to <20 cc receive 15 Gy
 - Lesions ≥20 cc to <30 cc receive 14 Gy
 - Lesions ≥30 cc to <5 cm receive 12 Gy
 - ◊ In general, single-fraction adjuvant SRS is not recommended for cavities >5 cm.
 - ◊ Larger cavities, however, may be treated with fractionated SRT. Potential regimens include, but are not limited to:
 - 24–27 Gy in 3 fractions
 - 25–35 Gy in 5 fractions

▶ Palliative whole brain RT (WBRT)

- ◊ Only consider for palliative purposes when SRS/SRT is not feasible in patients with good PS for whom disease has progressed.
- ◊ The pros and cons of WBRT should be considered carefully in the context of individual patient preferences/goals of care.²⁹
- ◊ WBRT can be considered if radiographic, clinical, or pathologic signs of leptomeningeal carcinomatosis are present (see LEPT-1 in the [NCCN Guidelines for Central Nervous System Cancers](#)).
- ◊ Common WBRT regimens include:
 - Standard doses include 30 Gy in 10 fractions and 20 Gy in 5 fractions. WBRT can be done with or without hippocampal avoidance (HA) + memantine. HA-WBRT (plus memantine) 30 Gy in 10 fractions is preferred for patients with a better prognosis (≥4 months) and no metastases within 5 mm of the hippocampi.³⁰
 - For patients with poor predicted prognosis and with symptomatic brain metastases, standard WBRT of 20 Gy in 5 fractions is a reasonable option.³¹ If WBRT is given, for patients with a better prognosis, consider memantine during and after WBRT for a total of 6 months.³²
- ▶ **Adjuvant WBRT**
 - ◊ Adjuvant WBRT after resection or SRS/SRT is not recommended for patients with kidney cancer.³³
 - Recent data from a randomized trial suggest that adjuvant WBRT is associated with worse cognitive decline when compared to adjuvant SRS/SRT alone.²⁸ Although local control appears superior with adjuvant WBRT, there were no differences in overall survival (OS).
 - ◊ For dosing, see Palliative WBRT section above.
- ▶ Also see [NCCN Guidelines for Central Nervous System Cancers](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

References

KID-B
3 OF 5



PRINCIPLES OF RADIATION THERAPY REFERENCES

- ¹ Siva S, Louie AV, Kotecha R, et al. Stereotactic body radiotherapy for primary renal cell carcinoma: a systematic review and practice guideline from the International Society of Stereotactic Radiosurgery (ISRS). *Lancet Oncol* 2024;25:e18-e28.
- ² Siva S, Bressel M, Sidhom M, et al. Stereotactic ablative body radiotherapy for primary kidney cancer (TROG 15.03 FASTER II): a non-randomised phase 2 trial. *Lancet Oncol* 2024;25:308-316.
- ³ Siva S, Chesson B, Bressel M, et al. TROG 15.03 phase II clinical trial of focal ablative stereotactic radiosurgery for cancers of the kidney - FASTER II. *BMC Cancer* 2018;18:1030.
- ⁴ Hanna GG, Murray L, Patel R, et al. UK consensus on normal tissue dose constraints for stereotactic radiotherapy. *Clin Oncol (R Coll Radiol)* 2018;30:5-14.
- ⁵ Masini C, Iotti C, De Giorgi U, et al. Nivolumab in combination with stereotactic body radiotherapy in pretreated patients with metastatic renal cell carcinoma. Results of the phase II NIVES study. *Eur Urol* 2022;81:274-282.
- ⁶ Cheung P, Patel S, North SA, et al. Stereotactic radiotherapy for oligoprogression in metastatic renal cell cancer patients receiving tyrosine kinase inhibitor therapy: A phase 2 prospective multicenter study. *Eur Urol* 2021;80:693-700.
- ⁷ Correa RJM, Louie AV, Zaorsky NG, et al. The emerging role of stereotactic ablative radiotherapy for primary renal cell carcinoma: A systematic review and meta-analysis. *Eur Urol Focus* 2019;5:958-969.
- ⁸ Stinauer MA, Kavanagh BD, Schefter TE, et al. Stereotactic body radiation therapy for melanoma and renal cell carcinoma: impact of single fraction equivalent dose on local control. *Radiat Oncol* 2011;6:34.
- ⁹ Zaorsky NG, Lehrer EJ, Kothari G, et al. Stereotactic ablative radiation therapy for oligometastatic renal cell carcinoma (SABR ORCA): a meta-analysis of 28 studies. *Eur Urol Oncol* 2019;2:515-523.
- ¹⁰ Hannan R, Christensen M, Hammers H, et al. Phase II trial of stereotactic ablative radiation for oligoprogressive metastatic kidney cancer. *Eur Urol Oncol* 2022;5:216-224.
- ¹¹ Tang C, Msaouel P, Hara K, et al. Definitive radiotherapy in lieu of systemic therapy for oligometastatic renal cell carcinoma: a single-arm, single-centre, feasibility, phase 2 trial. *Lancet Oncol* 2021;22:1732-1739.
- ¹² Ali M, Mooi J, Lawrentschuk N, et al. The role of stereotactic ablative body radiotherapy in renal cell carcinoma. *Eur Urol* 2022;82:613-622.
- ¹³ Palma DA, Olson R, Harrow S, et al. Stereotactic ablative radiotherapy versus standard of care palliative treatment in patients with oligometastatic cancers (SABR-COMET): A randomised, phase 2, open-label trial. *Lancet* 2019;393:2051-2058.
- ¹⁴ Gerszten PC, Burton SA, Quinn AE, et al. Radiosurgery for the treatment of spinal melanoma metastases. *Stereotact Funct Neurosurg* 2005;83:213-221.
- ¹⁵ Sahgal A, Roberge D, Schellenberg D, et al. The Canadian Association of Radiation Oncology scope of practice guidelines for lung, liver and spine stereotactic body radiotherapy. *Clin Oncol (R Coll Radiol)* 2012;24:629-639.
- ¹⁶ Wang XS, Rhines LD, Shiu AS, et al. Stereotactic body radiation therapy for management of spinal metastases in patients without spinal cord compression: a phase 1-2 trial. *Lancet Oncol* 2012;13:395-402.
- ¹⁷ Seung SK, Curti BD, Crittenden M, et al. Phase 1 study of stereotactic body radiotherapy and interleukin-2--tumor and immunological responses. *Sci Transl Med* 2012;4:137ra74.
- ¹⁸ Singh D, Chen Y, Hare MZ, et al. Local control rates with five-fraction stereotactic body radiotherapy for oligometastatic cancer to the lung. *J Thorac Dis* 2014;6:369-374.
- ¹⁹ Olivier KR, Schild SE, Morris CG, et al. A higher radiotherapy dose is associated with more durable palliation and longer survival in patients with metastatic melanoma. *Cancer* 2007;110:1791-1795.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY REFERENCES

- ²⁰ Sahgal A, Myrehaug SD, Siva S, et al. Stereotactic body radiotherapy versus conventional external beam radiotherapy in patients with painful spinal metastases: an open-label, multicentre, randomised, controlled, phase 2/3 trial. *Lancet Oncol* 2021;22:1023-1033.
- ²¹ Overgaard J, Gonzalez Gonzalez D, Hulshof MC, et al. Randomised trial of hyperthermia as adjuvant to radiotherapy for recurrent or metastatic malignant melanoma. *European Society for Hyperthermic Oncology. Lancet* 1995;345:540-543.
- ²² Overgaard J, von der Maase H, Overgaard M. A randomized study comparing two high-dose per fraction radiation schedules in recurrent or metastatic malignant melanoma. *Int J Radiat Oncol Biol Phys* 1985;11:1837-1839.
- ²³ Sause WT, Cooper JS, Rush S, et al. Fraction size in external beam radiation therapy in the treatment of melanoma. *Int J Radiat Oncol Biol Phys* 1991;20:429-432.
- ²⁴ Huguenin PU, Kieser S, Glanzmann C, et al. Radiotherapy for metastatic carcinomas of the kidney or melanomas: an analysis using palliative end points. *Int J Radiat Oncol Biol Phys* 1998;41:401-405.
- ²⁵ Shaw E, Scott C, Souhami L, et al. Single dose radiosurgical treatment of recurrent previously irradiated primary brain tumors and brain metastases: final report of RTOG protocol 90-05. *Int J Radiat Oncol Biol Phys* 2000;47:291-298.
- ²⁶ Minniti G, D'Angelillo RM, Scaringi C, et al. Fractionated stereotactic radiosurgery for patients with brain metastases. *J Neurooncol* 2014;117:295-301.
- ²⁷ Rajakesari S, Arvold ND, Jimenez RB, et al. Local control after fractionated stereotactic radiation therapy for brain metastases. *J Neurooncol* 2014;120:339-346.
- ²⁸ Brown PD, Ballman KV, Cerhan JH, et al. Postoperative stereotactic radiosurgery compared with whole brain radiotherapy for resected metastatic brain disease (NCCTG N107C/CEC.3): a multicentre, randomised, controlled, phase 3 trial. *Lancet Oncol* 2017;18:1049-1060.
- ²⁹ Mulvenna P, Nankivell M, Barton R, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): results from a phase 3, non-inferiority, randomised trial. *Lancet* 2016;388:2004-2014.
- ³⁰ Brown PD, Gondi V, Pugh S, et al. Hippocampal avoidance during whole-brain radiotherapy plus memantine for patients with brain metastases: Phase III trial NRG Oncology CC001. *J Clin Oncol* 2020;38:1019-1029.
- ³¹ Andrews DW, Scott CB, Sperduto PW, et al. Whole brain radiation therapy with or without stereotactic radiosurgery boost for patients with one to three brain metastases: phase III results of the RTOG 9508 randomized trial. *Lancet* 2004;363:1665-1672.
- ³² Brown PD, Pugh S, Laack NN, et al. Memantine for the prevention of cognitive dysfunction in patients receiving whole-brain radiotherapy: a randomized, doubleblind, placebo-controlled trial. *Neuro Oncol* 2013;15:1429-1437.
- ³³ Hong AM, Fogarty GB, Dolven-Jacobsen K, et al. Adjuvant whole-brain radiation therapy compared with observation after local treatment of melanoma brain metastases: a multicenter, randomized phase III trial. *J Clin Oncol* 2019;37:3132-3141.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

FOLLOW-UP^a (category 2B)

Stage I

Follow-up During Active Surveillance¹⁻⁶

- H&P annually
- Laboratory tests annually, as clinically indicated
- Abdomen imaging:
 - Abdomen CT or MRI both with and without IV contrast (unless otherwise contraindicated) within 6 months of surveillance initiation, then CT, MRI, or ultrasound (US) at least annually
- Chest imaging:
 - Chest x-ray or CT at baseline and annually as clinically indicated to assess for pulmonary metastases
 - Consider repeat chest imaging if intervention is being contemplated
- Consider renal mass biopsy at initiation of active surveillance or at follow-up, as clinically indicated
- Follow-up may be individualized based on surgical status, treatment schedules, side effects, comorbidities, and symptoms

Follow-up After Percutaneous Ablation^{1,7-10}

- H&P annually
- Laboratory tests annually, as clinically indicated
- Abdomen imaging:
 - Abdomen CT or MRI both with and without IV contrast (unless otherwise contraindicated), or contrast-enhanced US at 1–3 months, 6 months, and 12 months after ablation, then annually thereafter. If patient is unable to receive IV contrast, MRI or contrast-enhanced US are the preferred imaging modalities
 - If there is imaging or clinical concern for residual or recurrent disease, then renal mass biopsy or further treatment may be indicated
- Chest imaging:
 - Chest CT annually for 5 years for patients who have biopsy-proven low-risk pathologic features (no sarcomatoid, low-grade [grade 1/2] renal cell carcinoma [RCC]), nondiagnostic biopsies, or no prior biopsy

Follow-up After SBRT

- Abdomen and chest imaging:
 - Abdomen/chest CT every 3 months for year 1, every 6 months for year 2, every 9 months for years 3–4, and annually for year 5
- Renal function:
 - Evaluate every 3 months for year 1, every 6 months for year 2, every 9 months for years 3–4, and annually for year 5

^a No single follow-up plan is appropriate for all patients. Follow-up frequency and duration should be individualized based on patient requirements, and may be extended beyond 5 years ([KID-C, 5 of 5](#)). Further study is required to define optimal follow-up duration.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

FOLLOW-UP^a (category 2B)

Stage I

Follow-up After a Partial or Radical Nephrectomy¹

- H&P annually
- Laboratory tests annually, as clinically indicated
- Abdomen imaging:
 - Baseline abdomen CT or MRI (preferred) both with and without IV contrast (unless otherwise contraindicated) within 3–12 months of surgery, then annually for up to 5 years or longer as clinically indicated
 - A more rigorous imaging schedule can be considered if there are positive margins or adverse pathologic features (such as sarcomatoid, high-grade [grade 3/4])
- Chest imaging:
 - Chest CT annually for at least 5 years, then as clinically indicated
 - A more rigorous imaging schedule can be considered if there are positive margins or adverse pathologic features

Stage II

Follow-up After a Partial or Radical Nephrectomy¹

- H&P annually
- Laboratory tests annually, as clinically indicated
- Abdomen and pelvis imaging:
 - Baseline abdomen/pelvis CT or MRI (preferred) both with and without IV contrast (unless otherwise contraindicated), every 6 months for 2 years, then annually for up to 5 years or longer as clinically indicated
 - A more rigorous imaging schedule can be considered if there are positive margins or adverse pathologic features (such as sarcomatoid, high-grade [grade 3/4])
- Chest imaging:
 - Chest CT annually for at least 5 years, then as clinically indicated
 - A more rigorous imaging schedule can be considered if there are positive margins or adverse pathologic features
- Patients who are being considered candidates for adjuvant therapy should undergo restaging prior to initiating treatment

^a No single follow-up plan is appropriate for all patients. Follow-up frequency and duration should be individualized based on patient requirements, and may be extended beyond 5 years ([KID-C, 5 of 5](#)). Further study is required to define optimal follow-up duration.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

FOLLOW-UP^a (category 2B)

Follow-up for Stage III, or T4 NXM0 Resected^{1,11}

- H&P every 3–6 months for 3 years, then annually for up to 5 years, and as clinically indicated thereafter
- Comprehensive metabolic panel and other tests as indicated every 3–6 months for 3 years, then annually for up to 5 years, and as clinically indicated thereafter
- Abdomen and pelvis imaging:
 - ▶ Baseline abdomen/pelvis CT or MRI both with and without IV contrast (unless otherwise contraindicated) within 3–6 months, then CT or MRI (preferred), or US (US is category 2B for stage III), every 3–6 months for at least 3 years and then annually for up to 5 years
 - ▶ Imaging beyond 5 years: as clinically indicated
- Chest imaging:
 - ▶ Baseline chest CT within 3–6 months with continued imaging (CT preferred) every 3–6 months for at least 3 years and then annually for up to 5 years
 - ▶ Imaging beyond 5 years: as clinically indicated based on individual patient characteristics and tumor risk factors
- Additional imaging (ie, bone scan, brain imaging):
 - ▶ As symptoms warrant
- Fluorodeoxyglucose (FDG)-PET is useful in certain circumstances (fumarate hydratase [FH]-deficient RCC or succinate dehydrogenase complex subunit B [SDHB]-deficient RCC)^{12,13}
- Patients who are being considered candidates for adjuvant therapy should undergo restaging prior to initiating treatment

^a No single follow-up plan is appropriate for all patients. Follow-up frequency and duration should be individualized based on patient requirements, and may be extended beyond 5 years ([KID-C, 5 of 5](#)). Further study is required to define optimal follow-up duration.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

FOLLOW-UP (category 2B)

Follow-up Before, During, and After Adjuvant Therapy

- Patients who received adjuvant therapy should receive clinical follow-up as for stage III disease

Follow-up for Relapsed or Stage IV and Surgically Unresectable Disease^b

- H&P every 6–16 weeks for patients receiving systemic therapy, or more frequently as clinically indicated and adjusted for type of systemic therapy patient is receiving
- Laboratory evaluation as per requirements for therapeutic agent being used
- Chest, abdomen, and pelvis imaging:
 - ▶ CT or MRI both with and without IV contrast (unless otherwise contraindicated) to assess baseline pretreatment or prior to observation
 - ▶ Follow-up imaging every 6–16 weeks as per physician discretion, patient clinical status, and therapeutic schedule. Imaging interval to be adjusted shorter or longer according to rate of disease change and sites of active disease
- Consider MRI (preferred) or CT of head at baseline and as clinically indicated. Annual surveillance scans at physician discretion
- MRI of spine as clinically indicated
- Bone scan as clinically indicated
- FDG-PET is useful in certain circumstances (bone-predominant disease, assessment prior to metastasectomy, FH-deficient RCC or SDHB-deficient RCC)^{12,13}

^b No single follow-up plan is appropriate for all patients. Follow-up should be individualized based on treatment schedules, side effects, comorbidities, and symptoms.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



FOLLOW-UP (category 2B)

Long-Term Follow-Up (>5 years)^{2,14,15}

- Follow-up should be considered based on assessment of competing sources of mortality, personal risk factors for RCC, patient PS, and patient preference.
- Follow-up may be performed by a primary care physician if appropriate.
- H&P should be performed annually.
- Laboratory tests should be performed annually in surgical patients to evaluate renal function and determine glomerular filtration rate.
- Imaging:
 - ▶ Abdomen imaging may continue beyond recommended follow-up with increasing intervals given low but significant risk of metachronous tumors and/or late recurrences.
 - ▶ Consider chest imaging for higher stage disease and increasing intervals given low but significant risk of late recurrence.

¹ Donat SM, Diaz M, Bishoff JT, et al. Follow-up for clinically localized renal neoplasms: AUA Guideline. J Urol 2013;190:407-416.

² McIntosh AG, Ristau BT, Ruth K, et al. Active surveillance for localized renal masses: Tumor growth, delayed intervention rates, and >5-yr clinical outcomes. Eur Urol 2018;74:157-164.

³ Gupta M, Alam R, Patel HD, et al. Use of delayed intervention for small renal masses initially managed with active surveillance. Urol Oncol 2019;37:18-25.

⁴ Kassiri B, Cheaib JG, Pierorazio PM. Patients with small renal masses undergoing active surveillance: Is yearly chest imaging necessary? J Urol 2019;201:1061-1063.

⁵ Chandrasekar T, Ahmad AE, Fadaak K, et al. Natural history of complex renal cysts: Clinical evidence supporting active surveillance. J Urol 2018;199:633-640.

⁶ Jewett MA, Mattar K, Basiuk J, et al. Active surveillance of small renal masses: progression patterns of early stage kidney cancer. Eur Urol 2011;60:39-44.

⁷ Lay AH, Faddegon S, Olweny EO, et al. Oncologic efficacy of radio frequency ablation for small renal masses: Clear cell vs papillary subtype. J Urol 2015;194:653-657.

⁸ Beksac AT, Rivera-Sanfeliz G, Dufour CA, et al. Impact of tumor histology and grade on treatment success of percutaneous renal cryoablation. World J Urol 2017;35:633-640.

⁹ Haddad MM, Schmit GD, Kurup AN, et al. Percutaneous cryoablation of solitary, sporadic renal cell carcinoma: Outcome analysis based on clear-cell versus papillary subtypes. J Vasc Interv Radiol 2018;29:1122-1126.

¹⁰ Pierorazio PM, Johnson MH, Patel HD, et al. Management of renal masses and localized renal cancer: Systematic review and meta-analysis. J Urol 2016;196:989-99.

¹¹ Gershman B, Moreira DM, Thompson RH, et al. Renal cell carcinoma with isolated lymph node involvement: Long-term natural history and predictors of oncologic outcomes following surgical resection. Eur Urol 2017;72:300-306.

¹² Wu HC, Yen RF, Shen YY, et al. Comparing whole body 18F-2-deoxyglucose positron emission tomography and technetium-99m methylene diphosphate bone scan to detect bone metastases in patients with renal cell carcinomas - a preliminary report. J Cancer Res Clin Oncol 2002;128:503-506.

¹³ Nikolovski I, Carlo MI, Chen YB, et al. Imaging features of fumarate hydratase-deficient renal cell carcinomas: a retrospective study. Cancer Imaging 2021;21:24.

¹⁴ Narayan V, Puligandla M, Haas NB, et al. Patterns of relapse and implications for post-nephrectomy surveillance for patients with high-risk non-clear cell renal cell carcinoma: Subgroup analysis of the phase 3 ECOG-ACRIN E2805 trial. J Urol 2019;201:62-68.

¹⁵ Dabestani S, Beisland C, Stewart GD, et al. Long-term outcomes of follow-up for initially localised clear cell renal cell carcinoma: RECUR database analysis. Eur Urol Focus 2019;5:857-866.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY FOR STAGE IV (M1 OR UNRESECTABLE T4, M0) OR RELAPSED DISEASE

[See Evidence Blocks on EB-4](#)

FIRST-LINE THERAPY FOR CLEAR CELL HISTOLOGY (IN ALPHABETICAL ORDER BY CATEGORY)			
Risk	Preferred	Other Recommended	Useful in Certain Circumstances
Favorable^a	• Axitinib + Pembrolizumab^{b,c} (category 1) • Cabozantinib + Nivolumab^{b,d} (category 1) • Ipilimumab + Nivolumab^{b,e} (category 1) • Lenvatinib + Pembrolizumab^{b,c} (category 1)	• Axitinib + Avelumab^b • Pazopanib • Sunitinib • Cabozantinib (category 2B)	• Active surveillance^{1,2,3} • Axitinib (category 2B)
Poor/ intermediate^a	• Axitinib + Pembrolizumab^{b,c} (category 1) • Cabozantinib + Nivolumab^{b,d} (category 1) • Ipilimumab + Nivolumab^{b,e} (category 1) • Lenvatinib + Pembrolizumab^{b,c} (category 1) • Cabozantinib	• Axitinib + Avelumab^b • Pazopanib • Sunitinib	• Axitinib (category 2B)

Footnotes:

^a [Risk Models to Direct Treatment \(IMDC Criteria or MSKCC Prognostic Model\) \(KID-E\)](#).

^b [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).

^c Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

^d Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

^e Nivolumab and hyaluronidase-nvhy is not approved for concurrent use with IV Ipilimumab; however, for Nivolumab monotherapy, Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

References:

¹ Rini BI, Dorff TB, Elson P, et al. Active surveillance in metastatic renal-cell carcinoma: a prospective, phase 2 trial. *Lancet Oncol* 2016;17:1317-1324.

² Harrison MR, Costello BA, Bhavsar NA, et al. Active surveillance of metastatic renal cell carcinoma: Results from a prospective observational study (MaRCC). *Cancer* 2021;127:2204-2212.

³ Bex A. Increasing the evidence for surveillance of metastatic renal cancer. *Cancer* 2021;127:2184-2186.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY FOR STAGE IV OR RELAPSED DISEASE

[See Evidence Blocks on EB-5](#)

SUBSEQUENT THERAPY FOR CLEAR CELL HISTOLOGY (IN ALPHABETICAL ORDER BY CATEGORY)			
Immuno-oncology (IO) Therapy History Status	Preferred	Other Recommended	Useful in Certain Circumstances
IO Therapy Naïve	• None	• Axitinib + Pembrolizumab^{b,c} • Cabozantinib • Cabozantinib + Nivolumab^{b,d} • Everolimus/Lenvatinib • Ipilimumab + Nivolumab^{b,e} • Lenvatinib + Pembrolizumab^{b,c} • Nivolumab^{b,d}	• Axitinib • Everolimus • Lenvatinib • Pazopanib • Sunitinib • Tivozanib • Belzutifan (category 2B) • Bevacizumab^g (category 2B)
Prior IO Therapy	• None	• Axitinib • Belzutifan^f • Cabozantinib • Everolimus/Lenvatinib • Tivozanib	• Everolimus • Ipilimumab + Nivolumab^{b,e} • Lenvatinib • Pazopanib • Sunitinib • Axitinib + Pembrolizumab^{b,c} (category 2B) • Bevacizumab^g (category 2B) • Cabozantinib + Nivolumab^{b,d} (category 2B) • Lenvatinib + Pembrolizumab^{b,c} (category 2B)

^b [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).

^c Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

^d Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

^e Nivolumab and hyaluronidase-nvhy is not approved for concurrent use with IV Ipilimumab; however, for Nivolumab monotherapy, Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

^f This regimen is for patients who have received a programmed cell death protein 1 (PD-1) or programmed death ligand 1 (PD-L1) inhibitor and a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI).

^g An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY FOR STAGE IV (M1 OR UNRESECTABLE T4, M0)^h OR RELAPSED DISEASE

SYSTEMIC THERAPY FOR NON-CLEAR CELL HISTOLOGY ⁱ (IN ALPHABETICAL ORDER BY CATEGORY)		
Preferred	Other Recommended	Useful in Certain Circumstances
• Clinical trial^j • Cabozantinib • Cabozantinib + Nivolumab^{b,d} • Lenvatinib + Pembrolizumab^{b,c}	• Erlotinib + Bevacizumab^g for selected patients with advanced papillary RCC including hereditary leiomyomatosis and renal cell cancer (HLRCC)-associated RCC (HERED-RCC-D) • Everolimus/Lenvatinib • Nivolumab^{b,d} • Pembrolizumab^{b,c} • Sunitinib	• Axitinib • Everolimus • Everolimus + Bevacizumab^g • Ipilimumab + Nivolumab^{b,e} (category 2B)

[See Evidence Blocks on EB-6](#)

^b [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).

^c Pembrolizumab and bevacizumab subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and bevacizumab has different dosing and administration instructions compared to IV pembrolizumab.

^d Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

^e Nivolumab and hyaluronidase-nvhy is not approved for concurrent use with IV Ipilimumab; however, for Nivolumab monotherapy, Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

^g An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.

^h For first-line only.

ⁱ For collecting duct or medullary subtypes, partial responses have been observed with cytotoxic chemotherapy (Gemcitabine/Carboplatin, Paclitaxel/Carboplatin, or Gemcitabine/Cisplatin) and other platinum-based chemotherapies currently used for urothelial carcinomas. Doxorubicin/Gemcitabine can also produce responses in renal medullary carcinoma (RMC) (Wilson NR, et al. Clin Genitourin Cancer 2021;19:e401-e408). Oral targeted therapies generally do not produce responses in patients with RMC; Erlotinib + Bevacizumab can produce responses even in heavily pretreated patients with RMC. Outside of clinical trials, platinum-based chemotherapy regimens should be the preferred first-line therapy for RMC.

^j Consider tumor genomic testing for clinical trial eligibility.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



RISK MODELS TO DIRECT TREATMENT

Memorial Sloan Kettering Cancer Center (MSKCC) Prognostic Model¹

Prognostic Factors

- Interval from diagnosis to treatment of <1 year
- Karnofsky PS <80%
- Serum lactate dehydrogenase (LDH) >1.5 times the upper limit of normal (ULN)
- Corrected serum calcium greater than the ULN
- Serum hemoglobin less than the lower limit of normal (LLN)

Prognostic Risk Groups

- Low-risk group: no prognostic factors
- Intermediate-risk group: 1 or 2 prognostic factors
- Poor-risk group: ≥3 prognostic factors

International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) Criteria²

Prognostic Factors

- <1 year from time of diagnosis to systemic therapy
- PS <80% (Karnofsky)
- Hemoglobin less than the LLN (Normal: 120 g/L or 12 g/dL)
- Calcium greater than the ULN (Normal: 8.5–10.2 mg/dL)
- Neutrophil greater than the ULN (Normal: 2.0–7.0×10⁹/L)
- Platelets greater than the ULN (Normal: 150,000–400,000)

Prognostic Risk Groups

- Favorable-risk group: no prognostic factors
- Intermediate-risk group: 1 or 2 prognostic factors
- Poor-risk group: 3–6 prognostic factors

¹ Motzer RJ, Bacik J, Murphy BA, et al. Interferon-alfa as a comparative treatment for clinical trials of new therapies against advanced renal cell carcinoma. J Clin Oncol 2002;20:289-296.

² Heng DY, Xie W, Regan MM, et al. Prognostic factors for overall survival in patients with metastatic renal cell carcinoma treated with vascular endothelial growth factor-targeted agents: Results from a large, multicenter study. J Clin Oncol 2009;27:5794-5799.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

CRITERIA FOR FURTHER GENETIC RISK EVALUATION FOR HEREDITARY RCC SYNDROMES^a

1. An individual with a close blood relative ^b with a known pathogenic/likely pathogenic variant in a cancer susceptibility gene
2. An individual with RCC with any of the following criteria: ▶ Diagnosed at age ≤46 y^c ▶ Bilateral or multifocal tumors ▶ ≥1 first- or second-degree relative^b with RCC ▶ Personal or family history of mesothelioma or uveal melanoma
3. An individual whose tumors have the following histologic characteristics: ▶ Multifocal papillary histology ▶ HLRCC-associated RCC, RCC with FH deficiency or other histologic features associated with HLRCC ▶ Birt-Hogg-Dubé syndrome (BHDS)-related histology (multiple chromophobe, oncocytoma, or oncocytic hybrid) ▶ Angiomyolipomas of the kidney and one additional tuberous sclerosis complex (TSC) criterion in the same person (Table 1) ▶ Succinate dehydrogenase (SDH)-deficient RCC histology^d
4. An unaffected individual ^{e,f} with any of the following criteria: ▶ ≥2 first- or second-degree relatives^b with RCC (on the same side of the family) ▶ Any first-degree relative who meets the criteria in boxes 2 or 3 who is unable or unwilling to genetically test

→ [GENE-1](#)

→ Consider referral to cancer genetics professional and Refer to specific syndromes - See [Hereditary RCC Syndromes Overview \(HERED-RCC-2\)](#) and [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate: Principles of Cancer Risk Assessment and Counseling \(EVAL-A\) and Pedigree \(EVAL-B\)](#)

→ [GENE-1](#)

^a Table adapted from ACMG Practice Guidelines. Hampel H, Bennett RL, Buchanan A, et al. A practice guideline from the American College of Medical Genetics and Genomics and the National Society of Genetic Counselors: Referral indications for cancer predisposition assessment. *Genet Med* 2015;17:70-87. Schuch B, Vourganti S, Ricketts CJ, et al. Defining early-onset kidney cancer: Implications for germline and somatic mutation testing and clinical management. *J Clin Oncol* 2014;32:431-437.

^b Close blood relatives include the patient's first-degree (ie, parents, siblings, children) and second-degree (ie, half-siblings, aunts, uncles, nieces, nephews, grandparents, grandchildren) relatives.

^c Using age as a sole criterion for genetic risk evaluation is generally not a sensitive method.

^d Tumors that show loss of staining for SDHB have been termed SDH-deficient. Morphology of these tumors may include: solid or focally cystic growth, uniform cytology with eosinophilic flocculent cytoplasm, intracytoplasmic vacuolations and inclusions, and round to oval low-grade nuclei. (Ricketts CJ, Shuch B, Vocke CD, et al. Succinate dehydrogenase kidney cancer: an aggressive example of the Warburg effect in cancer. *J Urol* 2012;188:2063-2071; Gill AJ, Hes O, Papathomas T, et al. Succinate dehydrogenase [SDH]-deficient renal carcinoma: a morphologically distinct entity: a clinicopathologic series of 36 tumors from 27 patients. *Am J Surg Pathol* 2014;38:1588-1602; Gill AJ. Succinate dehydrogenase [SDH] and mitochondrial driven neoplasia. *Pathology* 2012;44:285-292.)

^e If unaffected, when possible, test family member with highest likelihood of a pathogenic/likely pathogenic variant before testing an unaffected individual.

^f Unnecessary in translocational RCC or medullary RCC.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

HEREDITARY RCC SYNDROMES OVERVIEW

Syndrome/Gene	Common Histologies	Inheritance Pattern Major Clinical Manifestations	Other Specialists Involved in Screening
von Hippel-Lindau (VHL)/ <i>VHL</i> gene	Clear cell	- Autosomal dominant Table 2	- Neurosurgery - Ophthalmology - Audiology - Endocrinology - Endocrine surgery
Hereditary papillary renal carcinoma (HPRC)/ <i>MET</i> gene	Papillary	- Autosomal dominant - Multifocal, bilateral renal cell tumors	Nephrology
Birt-Hogg-Dubé syndrome (BHDS)/ <i>FLCN</i> gene ^{1,2}	Chromophobe, hybrid oncocyctic tumors, clear cell, oncocytomas, angiomyolipomas, papillary RCC	- Autosomal dominant - Cutaneous fibrofolliculoma or trichodiscoma, pulmonary cysts, and spontaneous pneumothorax	- Pulmonology - Dermatology
Tuberous sclerosis complex (TSC)/ <i>TSC1</i> , <i>TSC2</i> genes	Angiomyolipoma (and other PEComas), renal cysts, eosinophilic solid and cystic RCC, RCC with fibromyxomatous stroma, eosinophilic vacuolated tumor, low-grade oncocyctic tumor, clear cell	- Autosomal dominant Table 1	- Neurology - Dermatology
Hereditary leiomyomatosis and renal cell cancer (HLRCC)/ <i>FH</i> gene	HLRCC-associated RCC or FH-deficient RCC	- Autosomal dominant - Leiomyomas of skin and uterus, unilateral, solitary, and aggressive renal cell tumors. PET-positive adrenal adenomas	- Gynecology - Dermatology
<i>BAP1</i> tumor predisposition syndrome (TPDS)/ <i>BAP1</i> gene ^{3,4}	Clear cell	- Autosomal dominant - Melanoma (uveal and cutaneous), kidney cancer, mesothelioma	- Dermatology - Ophthalmology - Thoracic oncology
Hereditary paraganglioma/pheochromocytoma (PGL/PCC) syndrome/ <i>SDHA</i> / <i>B/C/D</i> genes	SDH-deficient RCC	- Autosomal dominant - Head and neck PGL and adrenal or extra-adrenal PCCs, gastrointestinal stromal tumors (GIST)	- Endocrine - Endocrine surgery

- ¹ Schmidt LS, Nickerson ML, Warren MB, et al. Germline BHD-mutation spectrum and phenotype analysis of a large cohort of families with Birt-Hogg-Dubé syndrome. *Am J Hum Genet* 2005;76:1023-1033.
- ² Sattler EC, Steinlein OK. Birt-Hogg-Dubé Syndrome. 2006 Feb 27 [Updated 2020 Jan 30]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. *GeneReviews*® [Internet]. Seattle (WA): University of Washington, Seattle;1993-2020.
- ³ Peña-Llopis S, Vega-Ruweather bín-de-Celis S, Liao A. BAP1 loss defines a new class of renal cell carcinoma. *Nat Genet* 2012;44:751-759.
- ⁴ Hakimi AA, Ostrovnaya I, Reva B. Adverse outcomes in clear cell renal cell carcinoma with mutations of 3p21 epigenetic regulators BAP1 and SETD2: a report by MSKCC and the KIRC TCGA Research Network. *Clin Cancer Res* 2013;19:3259-3267.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

See [GENE-1](#)


HERED-RCC-2



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™



^a In individuals who meet diagnostic criteria, but in whom no germline mutations are identified, consider workup for mosaicism.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

Table 1: Features of Tuberous Sclerosis (TSC)

Major Features	Minor Features
• Renal angiomyolipoma^{a,b} • Cardiac rhabdomyoma • Cortical dysplasias, including tubers and cerebral white matter migration lines • Angiofibromas (≥3) or fibrous cephalic plaque • Hypomelanotic macules (3 to >5 mm in diameter) • Lymphangiomaleiomyomatosis (LAM)^a • Multiple retinal nodular hamartomas • Shagreen patch • Subependymal giant cell astrocytoma (SEGA) • Subependymal nodules (SENs) • Ungual fibromas (≥2)	• Multiple renal cysts • "Confetti" skin lesions (numerous 1- to 3-mm hypopigmented macules scattered over regions of the body such as the arms and legs) • Dental enamel pits (>3) • Intraoral fibromas (≥2) • Nonrenal hamartomas • Retinal achromic patch

Table 2: Features of Von Hippel-Lindau (VHL) Disease

Major Features	Minor Features
• Hemangioblastomas of the retina, spine, or brain • Clear cell RCC • PCC • PGL of abdomen, thorax, or neck • Retinal angiomas	• Endolymphatic sac tumors • Papillary cystadenomas of the epididymis or broad ligament • Pancreatic serous cystadenoma (>1) • Pancreatic neuroendocrine tumor (pNET) or multiple pancreatic cysts (>1)

^a The combination of angiomyolipoma and LAM does not meet criteria for definite diagnosis.

^b Multiple angiomyolipoma are a major feature.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

KIDNEY-SPECIFIC SCREENING RECOMMENDATIONS FOR PATIENTS WITH CONFIRMED HEREDITARY RCC WHO DO NOT YET HAVE A RADIOGRAPHIC OR PATHOLOGIC DIAGNOSIS OF RCC

General

- Follow-up should be individualized based on treatment schedules, side effects, comorbidities, and symptoms.
- Whenever possible, screening should be coordinated with another specialist involved in the patient's care.
- Patients of childbearing age who are planning conception should consider renal imaging prior to pregnancy.
- If there is a family member with an early diagnosis, screening should begin 10 years before earliest age of diagnosis in family member.
- CT of the abdomen can be used for surgical planning but should be limited if possible for surveillance due to lifetime radiation exposure for hereditary syndromic patients.
- Imaging frequency would be increased once lesions are detected based on growth rate and size of lesion(s).
- For surgical recommendations for each syndrome, see [HERED-RCC-C](#); for systemic therapy, see [HERED-RCC-D](#).

Syndrome	Screening Recommendations
BAP1-TPDS	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, every 2 y starting at age 30 y ¹
BHDS	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, every 3 y starting at age 20 y ²
HLRCC	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, annually starting at age 8–10 y ³
HPRC	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, every 1–2 y starting at age 30 y ^{4,5}
PGL/PCC^a	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, every 2 years concurrently with PGL/PCC screening recommendations starting at age 12 y ^{5,6,7}
TSC	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, every 1–3 y starting at age 12 y ⁸
VHL	• Abdomen MRI (preferred) or CT, both exams obtained with and without IV contrast, to assess kidneys, pancreas, and adrenals every 2 y starting at age 15 y ^{5,9}

^a See [NCCN Guidelines for Neuroendocrine and Adrenal Tumors](#) for full screening recommendations.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

HERED-RCC-B
1 OF 2



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

KIDNEY-SPECIFIC SCREENING RECOMMENDATIONS FOR PATIENTS WITH CONFIRMED HEREDITARY RCC WHO DO NOT YET HAVE A RADIOGRAPHIC OR PATHOLOGIC DIAGNOSIS OF RCC

REFERENCES

- ¹ Star P, Goodwin A, Kapoor R, et al. Germline BAP1-positive patients: the dilemmas of cancer surveillance and a proposed interdisciplinary consensus monitoring strategy. *Eur J Cancer* 2018;92:48-53.
- ² Menko F, van Steensel M, Giraud S, et al. Birt-Hogg-Dubé syndrome: Diagnosis and management. *Lancet Oncol* 2009;10:1199-1206.
- ³ Menko F, Maher E, Schmidt L, et al. Hereditary leiomyomatosis and renal cell cancer (HLRCC): Renal cancer risk, surveillance and treatment. *Fam Cancer* 2014;13:637-644.
- ⁴ Ornstein DK, Lubensky IA, Venzon D, et al. Prevalence of microscopic tumors in normal appearing renal parenchyma of patients with hereditary papillary renal cancer. *J Urol* 2000;163:431-433.
- ⁵ Rednam SP, Erez A, Druker H, et al. Von Hippel-Lindau and hereditary pheochromocytoma/paraganglioma syndromes: Clinical features, genetics, and surveillance recommendations in childhood. *Clin Cancer Res* 2017;23:e68-e75.
- ⁶ Tufton N, Sahdev A, Akker SA. Radiological surveillance screening in asymptomatic succinate dehydrogenase mutation carriers. *J Endocr Soc* 2017;1:897-907.
- ⁷ Eijkelenkamp K, Osinga TE, de Jong MM, et al. Calculating the optimal surveillance for head and neck paraganglioma in SDHB-mutation carriers. *Fam Cancer* 2017;16:123-130.
- ⁸ Northrup H, Aronow ME, Bebin EM, et al. Updated international tuberous sclerosis complex diagnostic criteria and surveillance and management recommendations. *Pediatr Neurol* 2021;123:50-66.
- ⁹ Binderup MLM, Smerdel M, Borgwadt L, et al. von Hippel-Lindau disease: Updated guideline for diagnosis and surveillance. *Eur J Med Genet* 2022;65:104538.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



KIDNEY-SPECIFIC SURGICAL RECOMMENDATIONS FOR PATIENTS WITH CONFIRMED HEREDITARY RCC

- Preoperative alert: Patients with a suspected or known diagnosis of PGL/PCC or VHL are at increased risk of PCCs and should have blood and/or urine screening for this prior to any surgical procedure.

BAP1-TPDS

- There are no specific guidelines in surgical management for this syndrome ([KID-A](#)).

BHDS

- Nephron-sparing surgery is the treatment of choice for renal tumors whenever possible, with consideration that an individual may have multiple tumors during their lifetime.¹
- Percutaneous ablation options may be considered for those with significant medical or surgical risk to undergo an operation.

HLRCC/FH-Deficient RCC

- As these tumors can be aggressive, surveillance of renal tumors is not recommended, and total radical nephrectomy should be considered.²
- FDG-PET is useful for staging pre-nephrectomy and/or for surveillance.^{3,4}

HPRC

- Nephron-sparing surgery is the treatment of choice for renal tumors whenever possible, with consideration that an individual may have multiple tumors during their lifetime.
- Percutaneous ablation options may be considered for those with significant medical or surgical risk to undergo an operation.

PGL/PCC/SDH-Deficient RCC

- Malignant tumors absent aggressive histology and early stage should undergo surgical resection; partial nephrectomy can be considered.
- For larger tumors and those with aggressive histology (eg, high grade, sarcomatoid), radical nephrectomy should be considered.⁵
- FDG-PET is useful for staging pre-nephrectomy and/or for surveillance.^{3,4}

TSC

- Angiomyolipoma is a benign lesion associated with TSC and managed separately.^{6,7,8}
- Nephron-sparing surgery is the treatment of choice for malignant renal tumors whenever possible, with consideration that an individual may have multiple tumors during their lifetime.⁹
- Percutaneous ablation options may be considered for those with significant medical or surgical risk to undergo an operation.

VHL

- Management of localized renal masses in patients with VHL is typically guided under the “3 cm rule.”⁹
- The idea is to intervene at a time point of maximal benefit to the patient to limit the chance of development of metastatic disease but also to consider the recurrent and multiple resections many of these patients will have over the course of their lifetime with subsequent development of chronic and progressive renal failure.^{9,10}
- Patient should undergo partial nephrectomy if at all possible and consider referral to centers with surgical expertise in complex partial nephrectomies and comprehensive care of patients with VHL.¹⁰
- Percutaneous ablation options may be considered for those with significant medical or surgical risk to undergo an operation.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

KIDNEY-SPECIFIC SURGICAL RECOMMENDATIONS FOR PATIENTS WITH CONFIRMED HEREDITARY RCC

REFERENCES

- ¹ Pavlovich CP, Grubb RL 3rd, Hurley K, et al. Evaluation and management of renal tumors in the Birt-Hogg-Dubé syndrome. J Urol 2005;173:1482-1486.
- ² Menko FH, Maher ER, Schmidt LS, et al. Hereditary leiomyomatosis and renal cell cancer (HLRCC): renal cancer risk, surveillance and treatment. Fam Cancer 2014;13:637-644.
- ³ Wu HC, Yen RF, Shen YY, et al. Comparing whole body 18F-2-deoxyglucose positron emission tomography and technetium-99m methylene diphosphate bone scan to detect bone metastases in patients with renal cell carcinomas - a preliminary report. J Cancer Res Clin Oncol 2002;128:503-506.
- ⁴ Nikolovski I, Carlo MI, Chen YB, et al. Imaging features of fumarate hydratase-deficient renal cell carcinomas: a retrospective study. Cancer Imaging 2021;21:24.
- ⁵ Gill AJ, Hes O, Papathomas T, et al. Succinate dehydrogenase (SDH)-deficient renal carcinoma: a morphologically distinct entity: a clinicopathologic series of 36 tumors from 27 patients. Am J Surg Pathol 2014;38:1588-1602.
- ⁶ Krueger DA, Northrup H; International Tuberous Sclerosis Complex Consensus Group. Tuberous sclerosis complex surveillance and management: recommendations of the 2012 International Tuberous Sclerosis Complex Consensus Conference. Pediatr Neurol 2013;49:255-265.
- ⁷ Muller A, Rouviere O. Renal artery embolization—indications, technical approaches and outcomes Nat Rev Nephrol 2015;11:288-301.
- ⁸ Nelson CP, Sanda MG. Contemporary diagnosis and management of renal angiomyolipoma. J Urol 2002;168:1315-1325.
- ⁹ Shuch B, Singer EA, Bratslavsky G. The surgical approach to multifocal renal cancers. Urol Clin North Am 2012;39:133-148.
- ¹⁰ Singer EA, Vourganti S, Lin KY, et al. Outcomes of patients with surgically treated bilateral renal masses and a minimum of 10 years of followup. J Urol 2012;188:2084-2088.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Hereditary Renal Cell Carcinoma

NCCN Evidence Blocks™

KIDNEY-SPECIFIC SYSTEMIC THERAPY FOR PATIENTS WITH CONFIRMED HEREDITARY RCC

Syndrome	Kidney-Specific Systemic Therapy
HLRCC	Other Recommended • Erlotinib + Bevacizumab^{a,b,1} Useful in Certain Circumstances • Cabozantinib + Nivolumab^{c,d}
TSC	Preferred • Everolimus^{e,2} Other Recommended • Sirolimus³
VHL	Preferred • Belzutifan^{f,4} Useful in Certain Circumstances • Pazopanib^{g,5}

[See Evidence Blocks on EB-7](#)

Footnotes:

^a An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.

^b There are no specific FDA-approved therapies for HLRCC. Treatment with Erlotinib plus Bevacizumab demonstrated benefit in patients with metastatic RCC from HLRCC.

^c [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).

^d Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

^e Everolimus is an FDA-approved therapy for asymptomatic, growing angiomyolipoma measuring >3 cm in diameter.

^f Belzutifan is FDA-approved for the treatment of VHL-associated-RCC, central nervous system (CNS) hemangioblastomas, or pNET, not requiring immediate surgery.

^g Pazopanib was associated with a >50% objective response rate in renal lesions in a 31-patient phase II study.

References:

¹ Srinivasan R, Gurram S, Al Harthy M, et al. Results from a phase II study of bevacizumab and erlotinib in subjects with advanced hereditary leiomyomatosis and renal cell cancer (HLRCC) or sporadic papillary renal cell cancer [abstract]. J Clin Oncol 2020;38(Suppl):Abstract 5004.

² Bissler JJ, Kingswood JC, Radzikowska E, et al. Everolimus for angiomyolipoma associated with tuberous sclerosis complex or sporadic lymphangioleiomyomatosis (EXIST-2): A multicentre, randomised, double-blind, placebo-controlled trial. Lancet 2013;381:817-824.

³ Bissler JJ, McCormack FX, Young LR, et al. Sirolimus for angiomyolipoma in tuberous sclerosis complex or lymphangioleiomyomatosis. N Engl J Med 2008;358:140-151.

⁴ Jonasch E, Donskov F, Iliopoulos O, et al. Belzutifan for renal cell carcinoma in von Hippel–Lindau disease. N Engl J Med 2021;385:2036-2046.

⁵ Jonasch E, McCutcheon IE, Gombos DS, et al. Pazopanib in patients with von Hippel-Lindau disease: a single-arm, single-centre, phase 2 trial. Lancet Oncol 2018;19:1351-1359.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

EVIDENCE BLOCKS FOR SYSTEMIC ADJUVANT TREATMENT FOLLOWING NEPHRECTOMY FOR CLEAR CELL HISTOLOGY AND HIGH-RISK RENAL CELL CARCINOMA

Stage II	
Pembrolizumab	

Stage III	
Pembrolizumab	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

4					
3					
2					
1					
	E	S	Q	C	A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

EVIDENCE BLOCK FOR ADJUVANT TREATMENT FOLLOWING NEPHRECTOMY FOR STAGE IV (RESECTABLE T4, M0) RCC WITH CLEAR CELL HISTOLOGY

Pembrolizumab	*
---------------	---

*Evidence Block development in progress

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

4					
3					
2					
1					
	E	S	Q	C	A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

EVIDENCE BLOCK FOR ADJUVANT TREATMENT FOLLOWING METASTASECTOMY FOR STAGE IV RCC WITH CLEAR CELL HISTOLOGY

Pembrolizumab



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



EVIDENCE BLOCKS FOR FIRST-LINE SYSTEMIC THERAPIES FOR CLEAR CELL HISTOLOGY

FAVORABLE-RISK PATIENTS	
Preferred Regimens	
Axitinib + Pembrolizumab	
Cabozantinib + Nivolumab	
Ipilimumab + Nivolumab	
Lenvatinib + Pembrolizumab	
Other Recommended Regimens	
Axitinib + Avelumab	
Pazopanib	
Sunitinib	
Cabozantinib	
Useful in Certain Circumstances	
Axitinib	

POOR/INTERMEDIATE-RISK PATIENTS	
Preferred Regimens	
Axitinib + Pembrolizumab	
Cabozantinib + Nivolumab	
Ipilimumab + Nivolumab	
Lenvatinib + Pembrolizumab	
Cabozantinib	
Other Recommended Regimens	
Axitinib + Avelumab	
Pazopanib	
Sunitinib	
Useful in Certain Circumstances	
Axitinib	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



4					
3					
2					
1					
	E	S	Q	C	A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

EVIDENCE BLOCKS FOR SUBSEQUENT SYSTEMIC THERAPIES FOR CLEAR CELL HISTOLOGY

IMMUNO-ONCOLOGY (IO) THERAPY HISTORY STATUS	
IO THERAPY NAÏVE	
Other Recommended Regimens	
Axitinib + Pembrolizumab	
Cabozantinib	
Cabozantinib + Nivolumab	
Everolimus/Lenvatinib	
Ipilimumab + Nivolumab	
Lenvatinib + Pembrolizumab	
Nivolumab	
Useful in Certain Circumstances	
Axitinib	
Everolimus	
Lenvatinib	*
Pazopanib	
Sunitinib	
Tivozanib	
Belzutifan	
Bevacizumab	

IMMUNO-ONCOLOGY (IO) THERAPY HISTORY STATUS	
PRIOR IO THERAPY	
Other Recommended Regimens	
Axitinib	
Belzutifan	
Cabozantinib	
Everolimus/Lenvatinib	
Tivozanib	
Useful in Certain Circumstances	
Everolimus	
Ipilimumab + Nivolumab	
Lenvatinib	*
Pazopanib	
Sunitinib	
Axitinib + Pembrolizumab	
Bevacizumab	
Cabozantinib + Nivolumab	
Lenvatinib + Pembrolizumab	

*Evidence Block development in progress

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



EVIDENCE BLOCKS FOR SYSTEMIC THERAPIES FOR NON-CLEAR CELL HISTOLOGY

SYSTEMIC THERAPY FOR NON-CLEAR CELL HISTOLOGY	
Preferred Regimen	
Cabozantinib	
Cabozantinib + Nivolumab	
Lenvatinib + Pembrolizumab	*
Other Recommended Regimens	
Erlotinib + Bevacizumab for selected patients with advanced papillary RCC including hereditary leiomyomatosis and renal cell carcinoma (HLRCC)-associated RCC	
Everolimus/Lenvatinib	
Nivolumab	
Pembrolizumab	
Sunitinib	
Useful in Certain Circumstances	
Axitinib	
Everolimus	
Everolimus + Bevacizumab	
Ipilimumab + Nivolumab	

*Evidence Block development in progress

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



4					
3					
2					
1					
	E	S	Q	C	A

EVIDENCE BLOCKS FOR KIDNEY-SPECIFIC SYSTEMIC THERAPY FOR PATIENTS WITH CONFIRMED HEREDITARY RCC

HLRCC	
Other Recommended Regimen	
Erlotinib + Bevacizumab	
Useful in Certain Circumstances	
Cabozantinib + Nivolumab	*

TSC	
Preferred Regimen	
Everolimus	
Other Recommended Regimen	
Sirolimus	*

VHL	
Preferred Regimen	
Belzutifan	
Useful in Certain Circumstances	
Pazopanib	

*Evidence Block development in progress

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



**Table 1. American Joint Committee on Cancer (AJCC)
TNM Staging System for Kidney Cancer (8th ed., 2017)**

T	Primary Tumor
TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
T1	Tumor ≤7 cm in greatest dimension, limited to the kidney
T1a	Tumor ≤4 cm in greatest dimension, limited to the kidney
T1b	Tumor >4 cm but ≤7 cm in greatest dimension, limited to the kidney
T2	Tumor >7 cm in greatest dimension, limited to the kidney
T2a	Tumor >7 cm but ≤10 cm in greatest dimension, limited to the kidney
T2b	Tumor >10 cm, limited to the kidney
T3	Tumor extends into major veins or perinephric tissues, but not into the ipsilateral adrenal gland and not beyond Gerota's fascia
T3a	Tumor extends into the renal vein or its segmental branches, or invades the pelvicalyceal system, or invades perirenal and/or renal sinus fat but not beyond Gerota's fascia
T3b	Tumor extends into the vena cava below the diaphragm
T3c	Tumor extends into the vena cava above the diaphragm or invades the wall of the vena cava
T4	Tumor invades beyond Gerota's fascia (including contiguous extension into the ipsilateral adrenal gland)
N	Regional Lymph Nodes
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in regional lymph node(s)
M	Distant Metastasis
M0	No distant metastasis
M1	Distant metastasis

Table 2. AJCC Prognostic Groups

	T	N	M
Stage I	T1	N0	M0
Stage II	T2	N0	M0
Stage III	T1-T2	N1	M0
	T3	NX,N0-N1	M0
Stage IV	T4	Any N	M0
	Any T	Any N	M1

Table 3. Histologic Grade (G)

GX	Grade cannot be assessed
G1	Nucleoli absent or inconspicuous and basophilic at 400x magnification
G2	Nucleoli conspicuous and eosinophilic at 400x magnification, visible but not prominent at 100x magnification
G3	Nucleoli conspicuous and eosinophilic at 100x magnification
G4	Marked nuclear pleomorphism and/or multinucleate giant cells and/or rhabdoid and/or sarcomatoid differentiation

Used with the permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.



ABBREVIATIONS

4D-CT	four-dimensional computed tomography	IMDC	International Metastatic Renal Cell Carcinoma Database Consortium	SBRT	stereotactic body radiation therapy
BED	biologically effective dose	IO	immuno-oncology	SDH	succinate dehydrogenase
BHDS	Birt-Hogg-Dubé syndrome	ITV	internal target volume	SDHB	succinate dehydrogenase complex subunit B
CBC	complete blood count	LAM	lymphangioleiomyomatosis	SEGA	subependymal giant cell astrocytoma
CNS	central nervous system	LDH	lactate dehydrogenase	SENs	subependymal nodules
ECOG	Eastern Cooperative Oncology Group	LLN	lower limit of normal	SRS	stereotactic radiosurgery
		OAR	organ at risk	SRT	stereotactic radiation therapy
		OS	overall survival	TPDS	tumor predisposition syndrome
FDG	fluorodeoxyglucose			TSC	tuberous sclerosis complex
FH	fumarate hydratase	PCC	pheochromocytoma		
FNA	fine-needle aspiration	PD-1	programmed cell death protein 1	ULN	upper limit of normal
		PD-L1	programmed death ligand 1		
GIST	gastrointestinal stromal tumor	PGL	paraganglioma	VEGF-TKI	vascular endothelial growth factor tyrosine kinase inhibitor
		pNET	pancreatic neuroendocrine tumor		
HA	hippocampal avoidance	PRV	planning organ at risk volume	VHL	von Hippel-Lindau
H&P	history and physical	PS	performance status		
HLRCC	hereditary leiomyomatosis and renal cell cancer			WBRT	whole brain radiation therapy
HPRC	hereditary papillary renal carcinoma	RCC	renal cell carcinoma		
		RMC	renal medullary carcinoma		
IGRT	image-guided radiation therapy				



NCCN Guidelines Version 2.2026

Kidney Cancer

NCCN Evidence Blocks™

NCCN Categories of Evidence and Consensus

Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference

Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 2.2026

Kidney Cancer

Discussion

This discussion corresponds to the NCCN Guidelines for Kidney Cancer. Last updated: April 14, 2026

Table of Contents

Overview	MS-2
Guidelines Update Methodology	MS-2
Literature Search Criteria	MS-2
Sensitive/Inclusive Language Usage	MS-3
Initial Evaluation and Staging	MS-3
Treatment of Localized Disease	MS-4
Management of Stage I (T1a or T1b) Disease	MS-6
Management of Stages II, III, and IV (Resectable T4,M0) Disease	MS-7
Follow-up After Treatment of Localized Disease	MS-9
Follow-up During Active Surveillance for Stage I	MS-9
Follow-up After Percutaneous Ablation for Stage I	MS-9
Follow-up After SBRT for Stage I	MS-10
Follow-up After Partial or Radical Nephrectomy for Stages I–II	MS-10
Follow-up for Patients with Stage III or Resected T4,NXM0 RCC	MS-10
Prognostic Tools for Localized Disease	MS-11
Management of Relapsed or Stage IV (M1 or Unresectable T4,M0) Disease	MS-11
Prognostic Models for Metastatic Disease	MS-11
Treatment Options for Patients with Relapsed or Stage IV (M1 or Unresectable T4,M0) Disease	MS-12
Systemic Therapy Options for Patients with Relapsed or Stage IV Disease	MS-13
First-Line Systemic Therapy Options for Patients with Stage IV (M1 or Unresectable T4,M0) or Relapsed Clear Cell RCC	MS-14

Subsequent Systemic Therapy Options for Patients with Clear Cell RCC ..MS-18

Systemic Therapy for Patients with Stage IV (M1 or Unresectable T4,M0) or Relapsed Non-Clear Cell RCCMS-23

Follow-up Recommendations for Relapsed or Stage IV Disease and Surgically Unresectable DiseaseMS-26

Supportive Care.....MS-27

Hereditary RCC Syndromes.....MS-28

Genetic Testing and Surveillance Recommendations for Individuals with a Personal or Family History of an RCC SyndromeMS-28

Genetic Testing and Screening Recommendations for Patients with a Clinical Diagnosis of RCC Who Have Characteristics Consistent with Inherited RCCMS-29

Kidney-Specific Surgical Recommendations for Patients with a Confirmed Hereditary RCC SyndromeMS-29

Kidney-Specific Systemic Therapy for Patients with Confirmed Hereditary RCCMS-30

Data Summary.....MS-31

Table 1: Key Studies on First-Line Therapy for Patients with Clear Cell RCC (ccRCC).....MS-32

Table 2: Key Studies on Subsequent Therapy for Patients with Clear Cell RCC (ccRCC).....MS-35

Table 3: Key Studies on Systemic Therapy for Patients with Non-Clear Cell RCC (nccRCC).....MS-37

References.....MS-39



NCCN Guidelines Version 2.2026

Kidney Cancer

Overview

An estimated 80,450 Americans will be diagnosed with cancers of the kidney and renal pelvis and 15,160 will die of the disease in the United States in 2026.¹ These cancers comprise approximately 4% of all new cancers, with a median age at diagnosis of 65 years.² Global data analysis shows that in the United States, there has been a 0.9% increase in kidney cancer cases and a 1.6% decrease in mortality from kidney cancer.³ About 66% of cases are diagnosed as localized disease, and the 5-year survival rate between 2015 and 2021 is 79%.¹ The most important prognostic determinants of 5-year survival are the tumor stage, grade, local extent of the tumor, presence of regional nodal metastases, and evidence of metastatic disease at presentation.⁴⁻¹³ RCC primarily metastasizes to the lung, bone, liver, lymph nodes, adrenal gland, and brain.¹⁴⁻¹⁶

Approximately 85% of kidney tumors are renal cell carcinoma (RCC), and approximately 80% to 90% of these have a clear cell histology (clear cell RCC, ccRCC).¹⁷⁻²⁰ Other less common cell types include papillary, oncocytic and chromophobe, molecularly defined RCC such as *TFE3*-rearranged and *TFEB*-altered (translocation) RCC, and collecting duct carcinoma.²¹ *SMARCB1*-deficient medullary renal carcinoma is a rare and aggressive RCC variant that almost exclusively arises in patients who have sickle-cell trait or hemoglobin sickle cell disease, or rarely sickle cell disease.²² An update in the pathologic classification system from the Genitourinary Pathology Society has close to 20 types of RCC with several additional emerging entities.²³ The histologic diagnosis of RCC is established after surgical removal of renal tumors or after biopsy.

Smoking, obesity, and hypertension are established risk factors for RCC development. Several hereditary types of RCC also exist, with von Hippel-Lindau (VHL) disease being the most common. VHL disease is caused by an autosomal-dominant constitutional mutation in the *VHL* gene that predisposes individuals to benign and malignant cysts/tumors.^{15,24-26}

Other hereditary types include fumarate hydratase (FH)-deficient and succinate dehydrogenase (SDH)-deficient RCC, associated with germline genetic alterations (also see *Hereditary RCC Syndromes* in this Discussion).

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Kidney Cancer provide multidisciplinary recommendations for the clinical management of ccRCC and non-clear cell RCC (nccRCC). These NCCN Guidelines® are intended to assist with clinical decision-making, but they cannot incorporate all possible clinical variations and are not intended to replace good clinical judgment or individualization of treatments. Medical practitioners should note that unusual patient scenarios (presenting in <5% of patients) are not specifically discussed in these Guidelines.

Guidelines Update Methodology

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Literature Search Criteria

Prior to the update of this version of the NCCN Guidelines for Kidney Cancer, an electronic search of the PubMed database was performed to obtain key literature on Kidney Cancer published since the previous Guidelines update, using the following search terms: Renal Cell Carcinoma, RCC, renal carcinoma, or Kidney Cancer. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV;



NCCN Guidelines Version 2.2026

Kidney Cancer

Guideline; Practice Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles as well as articles from additional sources deemed as relevant to these Guidelines as discussed by the Panel during the Guidelines update have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel's review of lower-level evidence and expert opinion.

Sensitive/Inclusive Language Usage

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation. NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-weight biased; and inclusive of individuals of all sexual orientations and gender identities.²⁷ NCCN Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will continue to use the terms men, women, female, and male when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Initial Evaluation and Staging

Patients with RCC typically present with a suspicious mass involving the kidney that has been visualized using a radiographic study, often a CT scan. As the use of imaging methods (eg, abdomen CT with or without pelvis CT, MRI) has become more widespread, the frequency of incidental detection of RCC has increased,^{28,29} and fewer patients present with the typical triad symptoms (hematuria, flank mass, and flank pain).

Less frequently, patients present with signs or symptoms resulting from metastatic disease, including bone pain, adenopathy, and pulmonary symptoms attributable to lung parenchyma or mediastinal metastases. Other presentations include fever, weight loss, anemia, or a varicocele.

A thorough physical examination should be performed along with obtaining a complete medical history of the patient. Laboratory evaluation includes a complete blood count (CBC) and comprehensive metabolic panel. The metabolic panel may include serum corrected calcium, serum creatinine, liver function studies, and urinalysis.

CT of the abdomen with or without pelvis CT or MRI, and CT chest (preferred) or chest x-ray are essential studies in the initial workup.^{30 31,32} Abdomen MRI is used to evaluate the inferior vena cava if tumor involvement is suspected, or it can be used instead of CT for detecting renal masses and for staging when contrast material cannot be administered because of allergy or moderate renal insufficiency.^{33,34} All imaging studies should be performed with and without contrast, such as renal protocol for the abdomen.

A central renal mass may suggest the presence of urothelial carcinoma; if so, urine cytology, ureteroscopy, or percutaneous mass biopsy should be considered.



NCCN Guidelines Version 2.2026

Kidney Cancer

Most bone and brain metastases are symptomatic at diagnosis. Therefore, a bone scan is not routinely performed unless the patient has an elevated serum alkaline phosphatase (ALP) or complains of bone pain.³⁵ MRI of the brain can be performed if clinical signs, presentation, and symptoms suggest brain metastases.

The recommended abdomen imaging studies provide high diagnostic accuracy. Therefore, a core needle biopsy is not always necessary before surgery, especially in patients whose imaging studies are consistent with RCC. In selected individuals, core needle biopsy may be considered to establish the diagnosis of RCC and to guide decisions on active surveillance strategies.³⁶ Biopsy of small lesions may also be considered to guide decisions on percutaneous ablation and/or surgery. Biopsy should also be considered if a central lesion or a homogeneous infiltration of renal parenchyma is observed on scans to rule out urothelial carcinoma or lymphoma, respectively.

The value of PET in RCC remains to be determined. Currently, PET or PET/CT is not an imaging tool that is routinely recommended to diagnose kidney cancer or to follow for evidence of relapse after nephrectomy.³⁷ However, in certain patients, FDG-PET may be useful for follow-up and extent of disease evaluation.^{38,39} See *Follow-up* sections for stage III and IV disease below for additional information.

RCC in younger patients (≤ 46 years) may indicate an inheritable disorder,⁴⁰ and these patients should be referred to a hereditary cancer clinic for further evaluation. If patients present with multiple renal masses, are ≤ 46 years of age at diagnosis, or have a family history of RCC, they should consider genetic evaluation (see *Hereditary RCC Syndromes* in this Discussion).

Treatment of Localized Disease

Surgical resection remains an effective therapy for clinically localized RCC, with options including radical nephrectomy and nephron-sparing surgery—each detailed below. Each of these modalities is associated with its benefits and risks, the balance of which should optimize long-term renal function and expected cancer-free survival.

Nephron-Sparing Surgery and Radical Nephrectomy

A radical nephrectomy includes a perifascial resection of the kidney, perirenal fat, regional lymph nodes, and in some instances, the ipsilateral adrenal gland. Radical nephrectomy is the preferred treatment if the tumor extends into the inferior vena cava, involves $>50\%$ of the kidney, or partial nephrectomy would result in a poorly functioning kidney. Open, laparoscopic, or robotic surgical techniques may be used to perform radical nephrectomy. Long-term outcomes data indicate that all techniques for laparoscopic and open radical nephrectomies have equivalent cancer-free survival rates.⁴¹⁻⁴⁸

Originally, partial nephrectomy (nephron-sparing surgery) was indicated only in clinical settings in which a radical nephrectomy would render the patient functionally anephric, necessitating dialysis. These settings include RCC in a solitary kidney, RCC in one kidney with inadequate contralateral renal function, and bilateral synchronous RCC.

Partial nephrectomy has well-established oncologic outcomes data comparable to radical nephrectomy.⁴⁹⁻⁵⁴ Radical nephrectomy can lead to an increased risk for chronic kidney disease^{55,56} and is associated with increased risks of cardiovascular morbidity and mortality according to population-based studies.⁵⁷ When compared with radical nephrectomy, partial nephrectomy can achieve preserved renal function, decreased overall mortality, and reduced frequency of cardiovascular events.⁵⁷⁻⁶¹ Patients with a hereditary form of RCC, such as VHL disease, should also



NCCN Guidelines Version 2.2026

Kidney Cancer

be considered for nephron-sparing surgery (see *Hereditary RCC Syndromes* in this Discussion). Nephron-sparing surgery has been used increasingly in patients with T1a and T1b renal tumors (ie, ≤ 7 cm in greatest dimension) and a normal contralateral kidney, with equivalent outcomes to radical nephrectomy.^{52,62-64} Radical nephrectomy should not be employed when nephron sparing can be achieved. One study showed that among Medicare beneficiaries with early-stage kidney cancer, treatment with partial rather than radical nephrectomy was associated with improved survival.⁶⁵

Studies with limited follow-up data show that the oncologic outcome for laparoscopic versus open nephron-sparing surgery appears to be similar.^{66,67} A study of oncologic outcomes at 7 years after surgery found metastasis-free survival to be 97.5% and 97.3% ($P = .47$) after laparoscopic and open nephron-sparing surgery, respectively.⁶⁸

The goals of nephron-sparing surgery should be obtaining optimal locoregional tumor control while minimizing ischemia time to ideally less than 30 minutes.⁶⁹ However, in some patients with localized RCC, nephron-sparing surgery may not be suitable because of locally advanced tumor growth or because the tumor is in an unfavorable location. Laparoscopic, robotic, and open partial nephrectomy all offer comparable outcomes in the hands of skilled surgeons. Patients in satisfactory medical condition should undergo surgical excision of stage I through III tumors.

Lymph Node Dissection

Lymph node dissection has not been consistently shown to provide therapeutic benefit. The EORTC phase III trial compared radical nephrectomy with a complete lymph node dissection to radical nephrectomy alone. The results showed no significant differences in overall survival (OS), time to progression of the disease, or progression-free survival (PFS) between the two study groups.⁷⁰ However, primary tumor pathologic features such as nuclear grade, sarcomatoid component,

tumor size, stage, and presence of tumor necrosis were all factors that influenced the likelihood of regional lymph node involvement at the time of radical nephrectomy.⁷¹ Assessment of lymph node status is based on enlargement of imaging (CT/MRI) and on assessment by direct palpation at the time of surgery. CT/MRI may not detect small metastases in normal lymph nodes.⁷² A systematic review and meta-analysis reported that nephrectomy with routine lymph node dissection did not show any OS and PFS benefit for non-metastatic RCC patients and had negative effects on cancer-specific survival.⁷³

The NCCN Kidney Cancer Panel indicates regional lymph node dissection should be considered for patients with palpable or enlarged lymph nodes detected on preoperative imaging tests. See *General Principles of Management for Renal Cell Carcinoma* in the algorithm for dissection templates for left- and right-sided renal tumors.⁷⁴

Adrenalectomy

Ipsilateral adrenal gland resection should be considered for patients with large upper pole tumors or abnormal-appearing adrenal glands on CT.⁷⁵⁻⁷⁷ Adrenalectomy is not indicated when imaging shows a normal adrenal gland or if the tumor is not high risk, based on size and location.⁷⁸

Active Surveillance, Percutaneous Ablation, and Stereotactic Body Radiation Therapy (SBRT)

Active surveillance^{79,80} is defined as the initial monitoring of tumors using abdominal imaging techniques with delayed intervention when indicated. The Panel recommends active surveillance as an option for certain patients with small renal masses (< 2 cm), T1a tumors (≤ 4 cm), and competing comorbidities. Patients who are older and those with small renal masses and other comorbidities often have low RCC-specific mortality.⁸¹ Active surveillance and percutaneous ablation such as cryosurgery, microwave ablation, or radiofrequency ablation are alternative strategies for selected patients, particularly for those who are



NCCN Guidelines Version 2.2026

Kidney Cancer

older, those with competing health risks, and those with T1b masses not eligible for surgery. Randomized phase III comparison of percutaneous ablation with surgical resection (ie, radical or partial nephrectomy by open or laparoscopic techniques) has not been performed. Stereotactic body radiation therapy (SBRT), a noninvasive approach, may also be considered for nonoptimal surgical candidates.

The NCCN Kidney Cancer Panel has addressed the utility of each of the above-mentioned treatment modalities for localized disease in the context of tumor stages: stage I (T1a and T1b), stage II, stage III, and resectable stage IV (T4,M0). See below for specific recommendations and more information regarding these primary treatment options.

Management of Stage I (T1a or T1b) Disease

Nephrectomy

The Panel prefers surgical excision by partial nephrectomy for the management of clinical stage I (T1a) renal masses. Adequate expertise and careful patient selection are important. Partial nephrectomy is most appropriate in patients with small unilateral stage I–III tumors or whenever preservation of renal function is a primary issue, such as in patients having one kidney or those with renal insufficiency, bilateral renal masses, or familial RCC. Partial nephrectomy is also appropriate for patients at relative risk of developing progressive chronic kidney disease due to young age or medical risk factors (eg, hypertension, diabetes, nephrolithiasis). Open, laparoscopic, or robotic approaches to partial nephrectomy can be considered, depending on tumor size, location, and the surgeon's expertise.

Some localized renal tumors may not be amenable to partial nephrectomy, in which case radical nephrectomy is recommended. The NCCN Guidelines also list radical nephrectomy as an alternative for patients with stage I (T1a) RCC if a partial nephrectomy is not technically feasible as

determined by the urologic surgeon. Partial nephrectomy for localized RCC has an oncologic outcome similar to that of radical surgery for T1b tumors.^{82,83} Surgery by partial nephrectomy, whenever feasible, or by radical nephrectomy is the recommendation for clinical T1b tumors according to the NCCN Kidney Cancer Panel.

Active Surveillance

Active surveillance is an option for the management of localized renal masses and should be a primary consideration for patients with decreased life expectancy or extensive comorbidities that would place them at excessive risk for more invasive intervention. Short- and intermediate-term oncologic outcomes indicate that an appropriate strategy is to initially monitor small renal masses (<2 cm), and, if required, treat for progression.⁷⁹ Active surveillance is an option for patients with stage T1a disease and for select patients with T1b disease.

Percutaneous Ablation

Although distant recurrence-free survival rates of percutaneous ablation and conventional surgery are comparable, percutaneous ablation may require retreatment to achieve the same local oncologic outcomes as conventional surgery.⁸⁴⁻⁸⁶ A meta-analysis of 32 observational studies and 1 randomized controlled trial (RCT) concluded that ablative therapy in T1a patients resulted in worse OS (hazard ratio [HR], 1.64; 95% confidence interval [CI], 1.39–1.95) as compared to partial nephrectomy but resulted in similar local recurrence-free survival (HR, 1.54; 95% CI, 0.88–2.71) and a smaller decline in estimated glomerular filtration rate postoperatively (mean differences [MD]: -7.42; 95% CI, -13.1 to -1.70). Oncologic outcomes in T1b patients showed some potential benefit, although more clinical evidence in this regard is lacking.⁸⁷ Judicious patient selection and counseling remain of paramount importance for these less invasive technologies. The NCCN Guidelines recommend percutaneous ablation



NCCN Guidelines Version 2.2026

Kidney Cancer

only in patients with stage I RCC (T1a [category 2A] and in select patients with T1b tumors who are not surgical candidates [category 2B]).

Stereotactic Body Radiation Therapy (SBRT)

SBRT may be considered as an option for non-optimal surgical candidates with T1 and T2a tumors (≤ 10 cm). There are insufficient data supporting use of SBRT with tumors > 7 cm. The phase II TROG 15.03 FASTRACK II trial investigated the use of SBRT in patients with localized RCC who were not surgical candidates or who declined surgery ($n = 70$).⁸⁸ Most patients enrolled had T1a ($n = 24$) or T1b ($n = 39$) disease, and 70% of patients had ccRCC. SBRT was done in single fraction (26 Gy) for tumors ≤ 4 cm or multiple fractions (42 Gy, 3 fractions) for tumors 4 to 10 cm.⁸⁸ At 12 months, local control was 100% ($P < .0001$), as was cancer-specific survival. Additionally, OS rates at 12 and 36 months were 99% and 82%, respectively. Most patients had grade 1–2 treatment-related adverse events (AEs), and 10% of patients had grade 3 (eg, nausea or vomiting; abdomen, flank, or tumor pain).⁸⁸

A meta-analysis from IROCK included median 5-year follow-up data from 190 patients with non-metastatic RCC, most with clear cell histology, who were treated with SBRT.⁸⁹ Patients received SBRT in single or multiple fractions (> 5 Gy) and median tumor size was 4 cm. Local failure incidence was 5.5% (95% CI, 2.8–9.5) at 3 and 5 years and 8.4% (95% CI, 4.2–14.3) at 7 years.⁸⁹ Less local failure was observed for patients who received single fractions versus multiple fractions. The 5-year cancer-specific survival rate was 92% and median PFS was 6.7 years. Grade 1–2 AEs were observed, and no grade 3 AEs or deaths related to treatment were reported.⁸⁹

The Panel recommends SBRT as a treatment option for non-optimal candidates for surgery or percutaneous ablation with stage T1a disease. It is also an option for selected patients with stage T1b disease who are not optimal surgical candidates. SBRT should be considered as the primary

radiation modality in all situations unless precluded by anatomic site, proximity to organs at risk, or past treatments. Referral to a high-volume center with specialized expertise should be considered. Detailed recommendations on SBRT technique, dosing, fractionation, and constraints for organs at risk for primary disease can be found in the *Principles of Radiation* section of these Guidelines.

Management of Stages II, III, and IV (Resectable T4,M0) Disease

Primary Treatment

The curative therapy for patients with stages II and III disease remains radical nephrectomy.⁴⁷ Radical nephrectomy is the preferred treatment for tumors that extend into the inferior vena cava. Resection of a caval or atrial thrombus often requires the assistance of cardiovascular surgeons because treatment-related mortality may reach 10%, depending on the local extent of the primary tumor and the level of vena caval extension. Partial nephrectomy is generally not suitable for patients with locally advanced tumors; however, it may be performed in patients with locally advanced tumors if technically feasible and clinically indicated. For example, partial nephrectomy may be considered for those with small, polar, unilateral tumors. The Panel lists radical nephrectomy or partial nephrectomy, if feasible or indicated, as options for stage II and III tumors. Nephrectomy is also recommended as primary treatment for resectable stage IV (T4,M0) disease. For patients who are non-optimal surgical candidates, SBRT may be considered for stage II RCC (category 2B) and stage III RCC (category 3).

Neoadjuvant Treatment

Robust data are lacking to support neoadjuvant systemic therapy for localized RCC, with currently only a few non-randomized clinical trials and retrospective analyses that have small sample sizes.⁹⁰⁻⁹³ However, the Panel felt that neoadjuvant therapy may be a treatment consideration in practice for certain patients with localized T3 or T4 ccRCC.



NCCN Guidelines Version 2.2026

Kidney Cancer

Therefore, the Panel indicates neoadjuvant therapy may be considered following confirmation of clear cell histology to achieve cytoreduction specifically for patients with clinical T3 or T4 tumors with no metastatic disease but in whom complete resection is not feasible up front, or if the patient has imperative indication for nephron preservation (eg, solitary kidney, chronic kidney disease, bilateral tumors), and in which a partial nephrectomy is not feasible. This may enable complete resection or facilitate a partial nephrectomy.⁹⁰ The Panel does not recommend any specific regimens in this setting.

Adjuvant Treatment for Clear Cell, High-Risk, Localized RCC

Individuals who have undergone resection of clear-cell, localized RCC with intermediate or high risk of recurrence may benefit from adjuvant pembrolizumab. Adjuvant radiation therapy after nephrectomy has not shown benefit, even in patients with nodal involvement or incomplete tumor resection.

Over the years, several vascular endothelial growth factor receptor (VEGFR) targeted tyrosine kinase inhibitors (TKIs) have been evaluated in the adjuvant setting with contrasting results. The ASSURE, PROTECT, and ATLAS trials investigating adjuvant sorafenib or sunitinib, pazopanib, and axitinib, respectively, did not show survival benefit and reported high toxicities.⁹⁴⁻⁹⁶ The phase III S-TRAC trial was the first to show benefit in disease-free survival (DFS) with adjuvant sunitinib following nephrectomy in patients with ccRCC, but grade 3 or higher AEs occurred in ~63% of patients receiving sunitinib compared to ~22% with placebo.^{97,98} Two recent meta-analyses of five RCTs evaluating adjuvant TKI monotherapies also concluded that they offer no benefit in OS or DFS and have significantly higher AE risks.^{99,100} Due to concerns about toxicity, lack of a demonstrated OS benefit, and conflicting results between the S-TRAC trial and the ASSURE/ATLAS/PROTECT trials, the use of sunitinib or other TKI therapy in the adjuvant setting is not recommended.

Immune checkpoint inhibitors (ICIs) that target programmed cell death protein 1 (PD-1) and programmed death ligand 1 (PD-L1) on T cells have also been investigated in the adjuvant setting. The phase III, multicenter, randomized, double-blind, placebo-controlled KEYNOTE-564 trial investigated the use of pembrolizumab versus placebo in 994 patients with locoregional RCC with a clear cell histology and an intermediate-to-high or high risk of recurrence (ie, tumor stage II with nuclear grade 4 or sarcomatoid differentiation, tumor stage III or higher, regional lymph node metastasis) after nephrectomy, or stage M1 with NED (no evidence of disease) status after nephrectomy and resection of metastatic lesions.¹⁰¹ DFS was noted in 77.3% of patients treated with pembrolizumab as compared to 68.1% of patients given placebo at 24 months (HR for recurrence or death, 0.68; 95% CI, 0.53–0.87; $P = .002$). The 30-month and 57-month follow-up analyses were consistent with these data, demonstrating a clinical benefit of pembrolizumab in this setting.^{102,103} At median follow-up of 57.2 months, OS favored pembrolizumab in the intention-to-treat population (HR for death, 0.62; 95% CI, 0.44–0.87; $P = .005$).¹⁰³ Additionally, time to subsequent therapy or any-cause death was prolonged in those treated with pembrolizumab compared to placebo. Grade 3 or higher AEs occurred in 32.4% of pembrolizumab-treated patients versus 17.7% of those who received placebo.^{101,102}

Based on the KEYNOTE-564 trial results, the Panel recommends pembrolizumab as a category 1 adjuvant treatment for both patients with stage II RCC (grade 4 tumors with clear cell histology ± sarcomatoid features), and for patients with stage III ccRCC, after a discussion with the patient about the potential benefits as well as the risks of adjuvant therapy. Adjuvant pembrolizumab is also an option for patients with resectable T4,M0 ccRCC. Surveillance is another option for patients in this setting.

The Panel also recommends considering adjuvant pembrolizumab for treatment of stage IV (M1 or unresectable T4,M0) or relapsed ccRCC after



NCCN Guidelines Version 2.2026

Kidney Cancer

metastasectomy with complete resection of disease, within a year of nephrectomy. Other recommendations for stage IV disease are covered later in this Discussion.

Due to the lack of evidence on the role of adjuvant pembrolizumab therapy for patients with RCC with non-clear cell histology (nccRCC), the Panel does not recommend including it as a treatment option for non-clear cell histology. Instead, surveillance is recommended following primary treatment for stage II, III, and resectable T4,M0 nccRCC. For patients with stage III or resectable T4,M0 nccRCC, clinical trial is also an option after primary treatment.

Follow-up After Treatment of Localized Disease

After surgical excision, 20% to 30% of patients with localized tumors experience relapse. Lung metastasis is the most common site of distant recurrence, occurring in 50% to 60% of patients.¹⁰⁴ The median time to relapse after surgery is 1 to 2 years, with most relapses occurring within 3 years.¹⁰⁵

The Panel has provided a framework for follow-up of patients undergoing surveillance of a small renal mass and for patients who underwent surgery or ablative therapy for primary RCC. The Panel has reiterated in a footnote that no single follow-up plan is appropriate for everyone, and follow-up should be modified for the individual patient using clinical judgment. Since uniform consensus among the Panel members regarding the most appropriate follow-up plan is lacking, these recommendations are listed as category 2B. Also, the guidance for follow-up has been provided for the first 5 years after nephrectomy, with follow-up evaluation to be extended beyond 5 years at the discretion of the physician. Results from a retrospective analysis indicate that in a subset of patients, relapses occur more than 5 years after surgery for their primary RCC.¹⁰⁶ The analysis suggests that continued follow-up/surveillance after 5 years may be of

potential value in some patients. Another retrospective analysis suggests that patients with lower risk are more likely to relapse later.¹⁰⁷ Identification of subsets of patients with higher risk who require longer follow-up has not been defined, and further research is required to refine follow-up strategies for patients with RCC.

The NCCN Guidelines incorporate a risk-stratified use of imaging that may target those patients most in need of intensive surveillance and/or imaging tests during follow-up.

Follow-up During Active Surveillance for Stage I

For follow-up during active surveillance, the Panel recommends an annual history and physical (H&P) examination and annual laboratory tests as clinically indicated. In order to study the growth rate of the tumor, the Panel recommends abdomen imaging (CT or MRI both with and without IV contrast, unless otherwise contraindicated) within 6 months from initiation of active surveillance; subsequent imaging (with CT, MRI, or ultrasound [US]) may be performed annually thereafter. All three modalities (US, CT, and MRI) have been found to accurately predict pathologic tumor size in a retrospective analysis.¹⁰⁸ Therefore, best clinical judgment should be used in choosing the imaging modality. The Panel recommends chest x-ray or chest CT at baseline and annually as clinically indicated to assess pulmonary metastases. Repeat chest imaging can be considered if intervention is being contemplated. The Panel notes that follow-up may be individualized based on surgical status, treatment schedules, side effects, comorbidities, and symptoms.

Follow-up After Percutaneous Ablation for Stage I

Most follow-up tests after percutaneous ablation included by the Panel are similar to those recommended during active surveillance. For imaging, the Panel recommends abdomen CT and MRI both with and without IV contrast (unless otherwise contraindicated), or contrast-enhanced US at 1



NCCN Guidelines Version 2.2026

Kidney Cancer

to 3 months, 6 months, and 12 months following ablation. Subsequent imaging is recommended annually. If the patient cannot receive IV contrast, MRI or contrast-enhanced US are preferred. If imaging results or clinical findings suggest residual disease or recurrence, then biopsy or further treatment may be indicated.

For those who have biopsy-proven low-risk pathologic features (no sarcomatoid, low-grade [grade 1/2] RCC), non-diagnostic biopsies, or no prior biopsy, the Panel also recommends annual chest CT for 5 years to assess for pulmonary metastases.

Follow-up After SBRT for Stage I

The Panel also includes follow-up recommendations for imaging and assessing renal function for patients who have stage I RCC and received SBRT. An abdomen/chest CT is recommended every 3 months in the first year, every 6 months in the second year, every 9 months in the third and fourth year, and annually in the fifth year. Renal function should be assessed using the same schedule as imaging.

Follow-up After Partial or Radical Nephrectomy for Stages I–II

For patients with stage I or II RCC who underwent a partial or radical nephrectomy, the Panel recommends an annual H&P examination and annual laboratory tests as clinically indicated. For patients with stage I RCC, the Panel recommends a baseline abdomen CT or MRI (preferred) both with and without contrast (unless contraindicated) within 3 to 12 months following renal surgery, then annually for up to 5 years or longer as clinically indicated. For patients with stage II RCC, the Panel recommends an increase in abdomen imaging frequency and inclusion of pelvis imaging, with baseline abdomen/pelvis CT or MRI (preferred), both with and without contrast (unless contraindicated) every 6 months for 2 years, then annually for up to 5 years or longer as clinically indicated. A more rigorous imaging schedule can be considered if the patient has

positive margins or adverse pathologic features (eg, sarcomatoid, grade 3/4 RCC). The rates of local recurrence for smaller tumors after partial nephrectomy are 1.4% to 2% versus 10% for larger tumors.^{66,109,110} The Panel also recommends yearly chest CT for at least 5 years and as clinically indicated thereafter. As mentioned above, a more rigorous imaging schedule (CT preferred) can be considered if the patient has positive margins or adverse pathologic features. Additionally, for any patients with stage II RCC who are being considered for adjuvant therapy, comprehensive restaging including chest CT should be performed prior to initiating treatment.

Follow-up for Patients with Stage III or Resected T4,NXM0 RCC

For patients with stage III RCC, larger tumors have a substantially higher risk of both local and metastatic recurrence, which warrants an increased follow-up frequency compared with patients with stage I or II RCC. Therefore, for these patients, the Panel recommends an H&P examination every 3 to 6 months for 3 years, then annually for up to 5 years. The follow-up evaluation may be extended beyond 5 years at the discretion of the physician as clinically indicated. A comprehensive metabolic panel and other tests are recommended as indicated every 3 to 6 months for 3 years, then annually up to 5 years, and as clinically indicated thereafter.

The Panel recommends baseline abdomen and pelvis CT or MRI (both with and without contrast, unless contraindicated) within 3 to 6 months following surgery, followed by CT, MRI (preferred), or US every 3 to 6 months for at least 3 years, and annually thereafter for up to 5 years. There is disagreement among the Panel members regarding the usefulness of US in patients with stage III disease. Therefore, it is listed as a category 2B option specifically for patients with stage III disease.



NCCN Guidelines Version 2.2026

Kidney Cancer

The Panel also recommends baseline chest CT within 3 to 6 months following surgery, followed by continued imaging (CT preferred) every 3 to 6 months for at least 3 years, and annually thereafter for up to 5 years.

While the use of US imaging for follow-up is an option for patients with low-risk RCC, CT or MRI is the preferred modality for those with a high risk of recurrence. The Panel notes that imaging beyond 5 years may be performed as clinically indicated, and additional site-specific imaging (eg, bone scan, brain imaging) may be performed as symptoms warrant.

FDG-PET is also recommended in certain circumstances. One retrospective study that identified imaging features for FH-deficient RCC in 16 patients showed elevated metabolic activity in the tumors imaged using FDG-PET/CT (n = 3).³⁸ The Panel recommends FDG-PET be used during follow-up if a patient has FH-deficient RCC, and also recommends this imaging modality for SDHB-deficient RCC.

For patients who are being considered for adjuvant therapy, restaging should be done prior to initiating treatment. Notably, the Panel recommends that all patients who receive adjuvant therapy also be given follow-up before, during, and after therapy based on the recommendations for stage III disease.

Prognostic Tools for Localized Disease

Alternate surveillance programs have been proposed, such as risk-based calculators and nomograms that can give 5-year survival estimates. The Panel does not explicitly recommend a risk-based calculator for identifying high-risk localized disease; however, two examples from the literature for identifying patients with higher risk of recurrence include the surveillance protocol based on the University of California Los Angeles (UCLA) Integrated Staging System (UISS) and the ASSURE RCC prognostic nomogram from Fox Chase Cancer Center.^{111,112} The UISS is an evidence-based system in which patients are stratified based on the 1997

TNM (tumor, node, metastasis) stage, grade, and ECOG performance status into low-, intermediate-, or high-risk groups for developing recurrence or metastases for post-surgical treatment of localized or locally advanced RCC.¹¹¹ The ASSURE RCC prognostic nomogram was created for prediction of recurrence in intermediate- and high-risk localized disease derived from characteristics of the patient population in the phase III ASSURE trial mentioned previously. This model includes histology, tumor size and grade, necrosis, and nodal involvement as characteristics, and is independent of AJCC TNM staging. It was applied to DFS, OS, and early disease progression measures for validation.¹¹²

Management of Relapsed or Stage IV (M1 or Unresectable T4,M0) Disease

Prognostic Models for Metastatic Disease

Prognostic scoring systems have been developed to define risk groups of patients by combining independent prognostic factors for survival in patients with metastatic RCC.^{113,114}

The first prognostic factor model to be widely applied was from Memorial Sloan Kettering Cancer Center (MSKCC). The model was derived from examining prognostic factors in patients (n = 463) with metastatic RCC enrolled in clinical trials and treated with interferon (IFN).¹¹³ Prognostic factors for multivariable analysis included five variables: interval from diagnosis to treatment of <1 year; Karnofsky Performance Status (KPS) <80%; serum LDH >1.5 times the upper limit of normal (ULN); corrected serum calcium greater than the ULN; and serum hemoglobin less than the lower limit of normal (LLN). Patients with none of these factors are in the favorable risk category with good prognosis, those with one or two factors present are considered intermediate risk, and patients with three or more of the factors are considered poor risk. The MSKCC criteria have been additionally validated by an independent group at the Cleveland Clinic.¹¹⁵



NCCN Guidelines Version 2.2026

Kidney Cancer

A prognostic model derived from a population of patients with metastatic RCC treated with vascular endothelial growth factor (VEGF)-targeted therapy followed the International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) model.¹¹⁴ This model was derived from a retrospective study of 645 patients with metastatic RCC treated with sunitinib, sorafenib, or bevacizumab plus IFN. Patients who received prior immunotherapy (ie, received their targeted therapy as second-line treatment) also were included in the analysis. The analysis identified six clinical parameters to stratify patients into favorable, intermediate, and poor prognosis groups. Four of the five adverse prognostic factors are those previously identified by MSKCC as independent predictors of short survival: hemoglobin less than the LLN, serum-corrected calcium greater than the ULN, KPS <80%, and time from the initial diagnosis to initiation of therapy of <1 year. Additional, independent, adverse prognostic factors validated in this model are absolute neutrophil count (ANC) greater than the ULN and platelets greater than the ULN.¹¹⁴

Patients with none of the identified six adverse factors were in the favorable-risk category (n = 133; 22.7%) in which a median OS was not reached and a 2-year OS was 75% (95% CI, 65%–82%). Patients with one or two adverse factors were in the intermediate-risk category (n = 301; 51.4%) in which a median OS was 27 months and a 2-year OS was 53% (95% CI, 46%–59%). Finally, those patients with three to six adverse factors were in the poor-risk category (n = 152; 25.9%) in which a median OS was 8.8 months and a 2-year OS was 7% (95% CI, 2%–16%).¹¹⁴ This model was validated in an independent dataset.¹¹⁶

Treatment Options for Patients with Relapsed or Stage IV (M1 or Unresectable T4,M0) Disease

Certain patients with stage IV disease also may benefit from surgery. For example, lymph nodes suspicious of metastatic disease on CT may be

hyperplastic and not involved with the tumor; thus, the presence of minimal regional adenopathy does not preclude surgery.

Cytoreductive nephrectomy before systemic therapy may be considered in select patients with stage IV, M1 disease and a potentially surgically resectable primary tumor mass. Generally, candidates for upfront cytoreductive nephrectomy have excellent performance status and no brain metastasis. A retrospective analysis conducted in the cytokine era indicated that patients most likely to benefit from cytoreductive nephrectomy before systemic therapy were those with lung-only metastases, good prognostic features, and good performance status.¹¹⁷ Retrospective data from the IMDC suggested that cytoreductive nephrectomy continues to play a role in patients treated with VEGF-targeted agents.¹¹⁸ The efficacy of newer systemic therapies is challenging the standard in some patients with metastatic disease. Results from the CARMENA phase III trial of patients with metastatic RCC who were eligible for cytoreductive nephrectomy found that sunitinib alone was non-inferior to sunitinib after nephrectomy.¹¹⁹ The median OS was 18.4 months in the sunitinib-alone group and 13.9 months in the sunitinib after nephrectomy group (HR, 0.89; 95% CI, 0.71–1.10), which did not exceed the fixed non-inferiority limit (1.20). However, many of the patients in this trial had poor-risk features, underscoring the importance of patient selection to obtain the greatest benefit from nephrectomy or targeted therapy.^{119,120} A post-hoc analysis of the CARMENA trial reported that for patients with only one IMDC risk factor, OS was longer following nephrectomy (31.4 months vs. 25.2 months).¹²¹ At this point, there are no prospective data defining the role of cytoreductive nephrectomy in patients who subsequently receive checkpoint antibody therapy. However, the ongoing open-label, randomized, prospective SEVURO-CN trial is investigating initial cytoreductive nephrectomy prior to IO-based therapy, as well as deferred cytoreductive nephrectomy, compared to IO therapy only in patients with synchronous metastatic RCC (Clinical Trial ID:



NCCN Guidelines Version 2.2026

Kidney Cancer

[NCT05753839](#)). Further study will better define the role and timing of cytoreductive nephrectomy in the rapidly evolving treatment landscape for RCC. There are two ongoing randomized phase III trials investigating deferred cytoreductive nephrectomy following immuno-oncology (IO)-based therapy in patients with synchronous metastatic RCC: the PROBE (SWOG 1931) trial (Clinical Trial ID: [NCT04510597](#)) and the NORDIC-SUN trial (Clinical Trial ID: [NCT03977571](#)).

Besides cytoreductive nephrectomy, the Panel also recommends upfront systemic therapy for patients with stage IV, M1 potentially resectable primary disease and large-volume distant metastases or large sarcomatoid tumors over initial cytoreductive nephrectomy. For these patients, clinical trial is also an option (category 2B). In patients whose tumors are surgically unresectable (stage IV T4, M0 or M1), the Panel recommends performing tissue sampling to confirm diagnosis of RCC to determine histology and guide subsequent management. Systemic therapy is generally recommended after recurrence, after cytoreductive nephrectomy in patients with multiple metastatic sites, or for patients with surgically unresectable tumors. See *Systemic Therapy Options for Patients with Relapsed or Stage IV Disease* below for detailed recommendations.

Patients with metastatic disease who present with hematuria or other symptoms related to the primary tumor should be offered palliative nephrectomy if they are surgical candidates. In addition, the small subset of patients with potentially surgically resectable primary RCC and oligometastatic sites are candidates for nephrectomy and management of metastases by surgical metastasectomy; alternatively, SBRT or percutaneous ablation are available for selected patients as well. Candidates include patients who: 1) initially present with primary RCC and oligometastatic sites; or 2) develop oligometastases after a prolonged disease-free interval from nephrectomy. Oligometastatic sites that are

amenable to this approach include the lung, bone, and brain. The primary tumor and the metastases may be resected during the same operation or at different times. Most patients who undergo targeted treatment of oligometastases experience recurrence, but long-term relapse-free survival has been reported in these patients. Prospective phase II studies showed some patients may benefit from SBRT treatment for oligometastases, postponing time to systemic therapy.¹²²⁻¹²⁴ SBRT may offer durable local control for metastases.¹²⁵ It should be considered for a patient with oligometastases unless metastasectomy is planned or SBRT cannot be delivered due to anatomic site, proximity to organs at risk, or previous treatment.¹²²⁻¹²⁷ For detailed recommendations on normal tissue constraints, as well as SBRT dosing and fractionation for distant metastatic disease, see *Principles of Radiation Therapy* in the Guidelines.

Patients who have undergone a nephrectomy and years later develop an oligometastatic recurrence also have the option of metastasectomy, SBRT,¹²⁸⁻¹³⁰ or percutaneous ablation, in addition to the first-line therapy options below. Clinical trial is an option for all patients, regardless of histology. Best supportive care is also recommended for all patients in this setting (see the *Supportive Care* section later in this Discussion).

Systemic Therapy Options for Patients with Relapsed or Stage IV Disease

Targeted therapy utilizing TKIs, and/or anti-VEGF antibodies, has been widely used in first- and second-line treatments. Agents targeting the mammalian target of rapamycin (mTOR) are also used in highly selected settings. A number of targeted agents have been approved by the FDA for the treatment of advanced RCC in the first and/or subsequent lines of therapy. ICIs provided a revolution in treatment options. Checkpoint antibodies alter the interaction between immune cells and antigen-presenting cells, including tumor cells. These agents can augment an anti-



NCCN Guidelines Version 2.2026

Kidney Cancer

tumor immune response and have shown promise in a number of tumor indications.

Tumor histology and risk stratification of patients are important in therapy selection. The NCCN Guidelines for Kidney Cancer stratify treatment recommendations by histology. Recommendations for first-line treatment of ccRCC are also stratified by risk group based on the MSKCC Prognostic Model and the IMDC prognostic criteria.

NCCN Categories of Preference

To further guide management of advanced RCC, the NCCN Kidney Cancer Panel has categorized all systemic kidney cancer therapy regimens as “Preferred,” “Other Recommended,” or “Useful in Certain Circumstances.” This categorization provides guidance on treatment selection by considering the efficacy, safety, evidence, and other factors that play a role in treatment selection. These factors include pre-existing comorbidities, nature of the disease, and in some cases consideration of access to agents.

Data Tables According to Line of Treatment and RCC Histology (Key Studies)

Due to the increasing number of NCCN-recommended systemic therapy options for metastatic RCC, the Panel has organized efficacy data from key studies into tables according to RCC histology and line of treatment (when applicable) for category 1 and 2A, preferred, and other recommended regimens; see *Table 1*, *Table 2*, and *Table 3* at the end of this Discussion.

Information about drug mechanism of action, FDA approval, summaries of study conclusions and safety data, and Categories of Evidence and Consensus and Categories of Preference for NCCN-recommended regimens remains below, and is stratified by RCC histology, line of

treatment (when applicable), prior IO therapy status (when applicable), and Categories of Preference.

First-Line Systemic Therapy Options for Patients with Stage IV (M1 or Unresectable T4,M0) or Relapsed Clear Cell RCC

Preferred

Axitinib + Pembrolizumab (All Risk Groups)

Axitinib is a selective, second-generation TKI of VEGFRs, while pembrolizumab is a monoclonal antibody that selectively binds to PD-1 (expressed on activated T cells) and blocks the interaction between PD-1 and its ligands PD-L1 and PD-L2 (both expressed on tumor cells and antigen-presenting cells). In April 2019, the FDA approved axitinib in combination with pembrolizumab for first-line treatment of patients with advanced RCC. Data from the randomized phase III KEYNOTE-426 trial, which included patients with favorable-, intermediate-, or poor-risk RCC, supported the combination therapy’s approval for this indication. Patients received either axitinib + pembrolizumab or sunitinib; those receiving the combination regimen had a significantly higher overall response rate (ORR) and longer PFS than those receiving sunitinib.¹³¹ Subsequent analyses at 31-, 43-, and 67-month median follow-ups showed agreement with these data (see *Table 1* for most recent efficacy data).¹³²⁻¹³⁴ Median OS in the most recent follow-up (median 67.2 months) was longer in the axitinib + pembrolizumab arm compared to the sunitinib arm.¹³⁴ Frequent AEs of grade 3 or higher in the axitinib + pembrolizumab arm included hypertension, diarrhea, and elevated alanine aminotransferase.¹³² Based on these data, the Panel recommends first-line axitinib + pembrolizumab as a category 1, preferred option for patients with ccRCC across all risk groups.

Cabozantinib + Nivolumab (All Risk Groups)

Cabozantinib is a multitargeted TKI of VEGFRs, MET, and AXL, while nivolumab is an anti-PD-1 antibody. In January 2021, the FDA approved



NCCN Guidelines Version 2.2026

Kidney Cancer

cabozantinib in combination with nivolumab for first-line treatment of patients with advanced RCC. Data from the randomized phase III CheckMate 9ER trial, which included patients with favorable-, intermediate-, or poor-risk RCC, supported the combination therapy's approval for this indication. Patients received either cabozantinib + nivolumab or sunitinib monotherapy. Those receiving cabozantinib + nivolumab had significantly higher ORRs and longer PFS than those receiving sunitinib. Additionally, OS was significantly longer with cabozantinib + nivolumab than sunitinib, and the HR favored cabozantinib + nivolumab (HR for death, 0.60; 98.89% CI, 0.40–0.89; $P = .001$).¹³⁵ In a subgroup analysis, the cabozantinib + nivolumab arm showed improved PFS, OS, and ORR in patients with advanced RCC with sarcomatoid features (an aggressive histologic subtype associated with poor prognosis) when compared to sunitinib.¹³⁶ Patients receiving combination treatment also reported delayed time to deterioration of patient-reported outcome scores compared to sunitinib.¹³⁷ Frequent AEs of grade 3 or 4 included hypertension, palmar-plantar erythrodysesthesia, and diarrhea.¹³⁶ In the final analysis at median follow-up of 67.6 months, the cabozantinib + nivolumab combination continued to show clinically meaningful benefit over sunitinib, with longer OS and PFS and higher ORR (see *Table 1* for recent efficacy data).¹³⁸ Based on these data, the Panel recommends first-line cabozantinib + nivolumab as a category 1, preferred option for patients with ccRCC across all risk groups.

Ipilimumab + Nivolumab (All Risk Groups)

Ipilimumab is a monoclonal antibody that selectively blocks the interaction between the negative regulator cytotoxic T-lymphocyte antigen 4 (CTLA-4; expressed early on activated T cells) and its ligands CD80/CD86 (expressed on antigen-presenting cells). Nivolumab's mechanism of action was described previously. In April 2018, the FDA approved ipilimumab in combination with nivolumab for first-line treatment of patients with poor-/intermediate-risk advanced RCC. Data from the randomized phase III

CheckMate 214 trial, which supported the FDA approval, compared combination ipilimumab + nivolumab followed by nivolumab monotherapy with sunitinib monotherapy in patients with advanced RCC.¹³⁹ The study's coprimary endpoints were ORR, OS, and PFS in patients with intermediate- and poor-risk RCC only; exploratory analyses of data in patients with favorable-risk RCC were reported separately. In patients with intermediate-/poor-risk RCC, combination ipilimumab + nivolumab led to a higher ORR and CR rate versus sunitinib monotherapy. Updated analyses continued to show a combined OS and ORR for intermediate-/poor-risk RCC that favored ipilimumab + nivolumab over sunitinib. In the primary analysis for patients with intermediate-/poor-risk RCC, longer OS was reported with ipilimumab + nivolumab versus sunitinib; however, PFS was not statistically significant per the predetermined threshold (HR for disease progression or death, 0.82; 99.1% CI, 0.64–1.05; $P = 0.03$).¹³⁹ Longer follow-up suggested improved PFS, with an HR of 0.73 (95% CI, 0.62–0.85).^{140,141}

In initial analyses, OS benefit for patients with favorable-risk RCC with ipilimumab + nivolumab was not observed (HR, 1.45 at 25-month follow-up);¹³⁹ however, a trend towards improved OS in the combination was observed in an exploratory analysis of this patient population at the 8-year follow-up (HR, 0.82; 95% CI, 0.60–1.13).¹⁴¹ Updated efficacy data for all risk groups are found in *Table 1*. Treatment-related AEs occurred in 93% of patients in the ipilimumab + nivolumab group and 97% of patients in the sunitinib group; grade 3 or 4 events occurred in 46% and 63% of patients, respectively. AEs led to treatment discontinuation in 22% and 12% of patients receiving ipilimumab + nivolumab and sunitinib, respectively. Treatment-related deaths occurred in 8 patients receiving combination therapy and 4 patients receiving sunitinib. Thirty-five percent of patients who developed immune-mediated AEs after ipilimumab + nivolumab treatment received high-dose steroids.¹³⁹ Based on these data, the Panel



NCCN Guidelines Version 2.2026

Kidney Cancer

recommends first-line ipilimumab + nivolumab as a category 1, preferred treatment option for patients with ccRCC across risk groups.

Lenvatinib + Pembrolizumab (All Risk Groups)

Lenvatinib is a multitargeted TKI of VEGFR-1, -2, and -3; fibroblast growth factor receptor (FGFR)-1, -2, -3, and 4; platelet-derived growth factor receptor- α (PDGFR- α); c-KIT; and RET. Pembrolizumab's mechanism of action was described previously. In August 2021, the FDA approved lenvatinib in combination with pembrolizumab for first-line treatment of patients with advanced RCC. Data from the randomized phase III CLEAR trial, which included patients with favorable-, intermediate-, or poor-risk RCC, supported the combination therapy's approval for this indication. Patients received either lenvatinib + pembrolizumab, everolimus/lenvatinib, or sunitinib. Those receiving lenvatinib + pembrolizumab had significantly longer PFS and a higher ORR than those receiving sunitinib, which were maintained in follow-up analyses.¹⁴²⁻¹⁴⁴ At median 49-month follow-up, OS was favorable for lenvatinib + pembrolizumab compared to sunitinib (see *Table 1* for most recent efficacy data).¹⁴⁴ In contrast, OS was not significantly different between the everolimus/lenvatinib and sunitinib groups.¹⁴² The final analysis noted that ~85% of patients treated with lenvatinib + pembrolizumab had treatment-emergent AEs (grade 3 or higher), which included hypertension and diarrhea, versus ~75% of patients who received sunitinib, consistent with prior safety profile analyses.¹⁴⁴ Based on these data, the Panel recommends first-line lenvatinib + pembrolizumab as a category 1, preferred treatment option for patients with ccRCC across all risk groups.

Cabozantinib (Poor-/Intermediate-Risk Groups)

In the open-label, randomized, phase II CABOSUN trial, patients with intermediate- or poor-risk advanced RCC received either cabozantinib or sunitinib.¹⁴⁵ See *Table 1* for efficacy data. Those treated with cabozantinib showed a significantly increased PFS and higher ORR

compared to those treated with sunitinib. The percentage of patients who experienced AEs of any grade or grade 3 or 4 was similar between cabozantinib and sunitinib. The most frequently reported AEs with cabozantinib included diarrhea, hypertension, fatigue, and palmar-plantar erythrodysesthesia.¹⁴⁵ Cabozantinib also increased quality-adjusted time without symptoms of disease or toxicity (Q-TWiST) versus sunitinib (317 days vs. 180 days, respectfully, with no disease progression and no grade 3 or 4 AEs).¹⁴⁶

Based on these results, the Panel recommends first-line cabozantinib as a category 2A, preferred treatment option for patients with poor- and intermediate-risk ccRCC.

Other Recommended

Axitinib + Avelumab (All Risk Groups)

Avelumab is a monoclonal antibody that selectively binds to PD-L1; axitinib's mechanism of action was described previously. In May 2019, the FDA approved axitinib + avelumab for first-line treatment of patients with advanced RCC. Data from the randomized phase III JAVELIN Renal 101 trial, which included patients with favorable-, intermediate-, or poor-risk RCC, supported the combination therapy's approval for this indication.^{147,148} For both the overall population and PD-L1–positive patients, those receiving axitinib + avelumab had significantly longer PFS than those receiving sunitinib. This benefit was observed across all risk groups. For OS, data were immature for all groups in both the primary¹⁴⁷ and 13-month interim¹⁴⁸ analyses. In the final analysis at an approximate median of 74 months, median OS was ~45 months with axitinib + avelumab versus ~39 months with sunitinib in the overall population but was not statistically significant (see *Table 1*).¹⁴⁹ PFS was significantly longer and ORR was higher with the combination versus sunitinib.^{149,150} Incidence of AEs of any grade, or grade 3 or higher, was similar between treatment arms,¹⁴⁷ and no new safety signals were reported in the



NCCN Guidelines Version 2.2026

Kidney Cancer

extended follow-up.^{149,150} Based on these results, the Panel recommends first-line axitinib + avelumab as a category 2A, other recommended treatment option for patients with ccRCC across all risk groups.

The post-hoc analysis of 108 patients with sarcomatoid histology in the phase III JAVELIN Renal 101 trial showed that patients in the avelumab + axitinib treatment arm had improved PFS (stratified HR, 0.57; 95% CI, 0.325–1.003) and a higher ORR (46.8% vs. 21.3%; complete response [CR] in 4.3% vs. 0%) versus those in the sunitinib arm.¹⁵¹

Pazopanib (All Risk Groups)

Pazopanib is an oral multitargeted TKI/angiogenesis inhibitor of VEGFRs, PDGFR- α and - β , and c-KIT, ITK, LCK, and c-FMS. The drug's safety and efficacy were evaluated in an open-label phase III study. Patients with advanced ccRCC who received 0–1 prior treatment received either pazopanib or placebo (see *Table 1* for efficacy data). PFS was significantly longer and ORR was significantly higher with pazopanib versus placebo in the treatment-naïve sub-population,¹⁵² but there was no difference in OS between the two groups.¹⁵³ Notable grade 3 toxicity was hepatotoxicity, indicated by elevated levels of alanine (30%) and aspartate (21%) transaminases.¹⁵² Therefore, it is critical to monitor liver function before and during treatment with the drug.

Additionally, the COMPARZ non-inferiority study of sunitinib versus pazopanib showed that these two drugs have similar safety and efficacy (see *Table 1* for efficacy data).^{154,155} Based on these data, the Panel has listed first-line pazopanib as a category 2A, other recommended regimen for patients with ccRCC across all risk groups.

Sunitinib (All Risk Groups)

Sunitinib is a multikinase inhibitor targeting several receptor tyrosine kinases, including PDGFR- α and - β ; VEGFR-1, -2, and -3; c-KIT; FLT3; CSF-1R; and RET.¹⁵⁶⁻¹⁵⁹ The efficacy of first-line sunitinib was studied in a

randomized phase III trial in which patients with metastatic RCC received either sunitinib or IFN- α .¹⁵⁶ See *Table 1* for efficacy data. PFS was longer in those receiving sunitinib across all risk groups. Updated results demonstrated a strong trend towards OS advantage of sunitinib over IFN- α in the first-line setting.¹⁶⁰ Based on these data, the Panel includes first-line sunitinib as a category 2A, other recommended regimen for patients with ccRCC across all risk groups.

Cabozantinib (Favorable-Risk Group)

Extrapolating on the CABOSUN data for patients with poor-/intermediate-risk (see above), the Panel added first-line cabozantinib as a category 2B, other recommended regimen for patients with favorable-risk ccRCC.

Useful in Certain Circumstances

Active Surveillance for Select, Asymptomatic Patients with ccRCC

A subset of patients with advanced ccRCC show indolent progression of disease and could benefit from initial active surveillance because of the toxicity of systemic therapies. A phase II trial of patients with treatment-naïve, asymptomatic, metastatic RCC followed patients on active surveillance through radiographic assessment at defined intervals until a decision was made to initiate systemic therapy.¹⁶¹ Of the 48 patients included in the analysis, the median time of surveillance from registration to initiation of systemic therapy was 14.9 months. This study demonstrated that a subset of patients with advanced ccRCC can safely undergo active surveillance before starting systemic therapy. In a prospective observational study of 504 patients with metastatic RCC, the median OS was not reached (95% CI, 122 months to not estimable [NE]) in patients who received active surveillance versus 30 months for those treated with systemic therapy.^{162,163} Therefore, the Panel included active surveillance as a category 2A, useful in certain circumstances option for select, asymptomatic patients with favorable-risk ccRCC.



NCCN Guidelines Version 2.2026

Kidney Cancer

Axitinib (All Risk Groups)

As a second-line therapy for patients with ccRCC, axitinib treatment led to higher ORR and longer median PFS compared with sorafenib.¹⁶⁴ In a randomized phase III trial, treatment-naïve patients received either axitinib or sorafenib; PFS was not significantly longer in patients receiving axitinib versus sorafenib but had an acceptable toxicity profile.¹⁶⁵ Based on these data, the Panel has included first-line axitinib as a category 2B, useful in certain circumstances option for patients with ccRCC across all risk groups.

Subsequent Systemic Therapy Options for Patients with Clear Cell RCC

The NCCN Kidney Cancer Panel stratified the subsequent therapies for ccRCC based on whether the patients have received any prior IO therapy. The recommended options are categorized into “IO therapy naïve” and “prior IO therapy.” Certain regimens (ie, axitinib, cabozantinib, nivolumab, tivozanib) are category 2A options in this setting because of the Panel’s observation that randomized registrational trials for these monotherapies began prior to the approval of IO combination therapy, and very few patients enrolled on these trials received upfront IO combination therapy. Therefore, the data do not support category 1 level evidence for subsequent monotherapy after frontline TKIs in the era of IO combination therapy, despite the limited number of phase III trials for IO combinations in this setting.

Additionally, IO/TKI combinations for those who received prior IO therapy are designated as category 2B recommendations in this setting. Data to support the use of IO therapy following progression on IO therapy are lacking. Two randomized phase III trials investigating the use of IO/TKI combinations suggest IO therapy not be considered for patients who previously received it. In the CONTACT-03 trial, patients previously treated with IO with locally advanced or metastatic RCC were randomized

to either atezolizumab (PD-L1 inhibitor) + cabozantinib or cabozantinib monotherapy.¹⁶⁶ PFS was not statistically significant between treatment arms (HR, 1.03; $P = .78$), and HR for death was 0.94 ($P = .69$). Serious AEs were reported for 48% of patients treated with the combination compared to 33% treated with cabozantinib. Discontinuation of treatment due to AEs occurred in 16% of patients with the combination, compared to 4% with monotherapy.¹⁶⁶ In the phase III TiNivo2 trial, patients with metastatic RCC who previously received IO therapy were randomized to either tivozanib + nivolumab or tivozanib alone.¹⁶⁷ Likewise, PFS was 5.7 months versus 7.4 months in the combination versus monotherapy arms, respectively (HR, 1.10; $P = .49$). The authors reported that this result was independent of where in the treatment sequence IO therapy was used.¹⁶⁷

Cabozantinib

In the randomized phase III METEOR trial, patients with disease progression after previous TKI therapy received cabozantinib or everolimus. See *Table 2* for efficacy data. Median PFS was significantly longer and ORR significantly higher in patients receiving cabozantinib versus everolimus.¹⁶⁸ The final analysis of the METEOR trial showed a statistically significant increase in OS in the cabozantinib arm versus the everolimus arm.^{169,170} Common grade 3 or 4 AEs that occurred more frequently with cabozantinib than with everolimus included hypertension, diarrhea, fatigue, hypomagnesaemia, and palmar-plantar erythrodysesthesia syndrome.¹⁶⁸ Additionally, a network meta-analysis comparing the relative effectiveness of subsequent treatment options for advanced and metastatic ccRCC found cabozantinib offered greater PFS, OS, and ORR benefit over everolimus, lenvatinib monotherapy, nivolumab, pazopanib, axitinib, sorafenib, temsirolimus, and tivozanib, but not over everolimus/lenvatinib combination. The odds of severe AEs were improved with cabozantinib over everolimus/lenvatinib and lenvatinib monotherapy.¹⁷¹ Cabozantinib has also been used as the control arm for two studies that included patients who received prior IO therapy



NCCN Guidelines Version 2.2026

Kidney Cancer

(CONTACT-03, described above, and the randomized CANTATA trial). CANTATA was a randomized trial investigating the addition of glutaminase inhibitor telaglenastat to cabozantinib.¹⁷² In both studies, there was no statistically significant survival difference between the experimental and control arms.^{166,172}

Based on these data, the Panel has included cabozantinib as a category 2A, other recommended subsequent therapy option for patients with ccRCC regardless of their prior IO therapy status.

Everolimus/Lenvatinib

In May 2016, the FDA approved lenvatinib, a multitargeted kinase inhibitor, in combination with everolimus, an mTOR inhibitor, for treating advanced RCC following one prior anti-angiogenic therapy. In a randomized phase II trial, patients with metastatic or unresectable, locally advanced ccRCC who had received prior antiangiogenic therapy received either combination everolimus/lenvatinib, single-agent lenvatinib, or single-agent everolimus. See *Table 2* for efficacy data. PFS and OS were significantly longer in patients receiving everolimus/lenvatinib versus everolimus monotherapy.^{173,174} Diarrhea was the most common grade 3 or 4 AE for those receiving everolimus/lenvatinib.¹⁷³ A prospective study of 55 patients with metastatic ccRCC, heavily pretreated with prior ICIs and VEGFR-TKIs, showed a median PFS of 6.2 months and median OS of 12.2 months with everolimus/lenvatinib.¹⁷⁵ Based on the phase II trial data, the Panel considers everolimus/lenvatinib a category 2A, other recommended subsequent therapy option for patients with ccRCC regardless of their prior IO therapy status.

Nivolumab

In the randomized phase III CheckMate 025 trial, patients with advanced ccRCC who were previously treated with one or more lines of anti-angiogenic therapy (excluding mTOR inhibitors) received either nivolumab

or everolimus. See *Table 2* for efficacy data. Patients receiving nivolumab had significantly longer OS and significantly higher ORR than those receiving everolimus.¹⁷⁶ The most prevalent grade 3 or 4 AEs with nivolumab were fatigue, anemia, and elevated levels of alanine and aspartate aminotransferases.¹⁷⁷ Based on these data, the Panel has included nivolumab as a category 2A, subsequent therapy option for patients with ccRCC who have not received any prior IO therapy.

Axitinib

The randomized phase III AXIS study compared second-line axitinib versus sorafenib. See *Table 2* for efficacy data. PFS was significantly longer and ORR significantly higher in patients receiving axitinib versus sorafenib.¹⁶⁴ The most prevalent AEs of grade 3 or higher for those treated with axitinib included hypertension, diarrhea, and fatigue.¹⁷⁸ Based on these data, the Panel included axitinib as a category 2A other recommended subsequent therapy option for patients with prior IO therapy and useful in certain circumstances for patients naïve for any prior IO therapy.

Axitinib + Pembrolizumab

Upon axitinib + pembrolizumab's FDA approval in a first-line setting, the Panel discussed whether the combination therapy might be used in clinical practice as an off-label subsequent treatment option in patients with relapsed or stage IV ccRCC. While they conceded that there were no robust, published data to support the use of axitinib + pembrolizumab in a second-line setting, they thought that clinicians were likely to consider the combination as a treatment option in patients with advanced ccRCC whose disease progressed after first-line sunitinib therapy. A retrospective study on 38 patients with ccRCC who had disease progression and previously received either ICI or VEGFR-TKI therapy were administered axitinib + pembrolizumab.¹⁷⁹ Median PFS was 9.7 months (95% CI, 4.1–15.3) at a median follow-up of 17.1 months. The ORR was 25% (all partial



NCCN Guidelines Version 2.2026

Kidney Cancer

responses [PRs]). Among 17 patients who had previously received ipilimumab + nivolumab and then second-line axitinib + pembrolizumab, the median PFS was 11.1 months and ORR was 31.4%. About 87% of patients in this study experienced AEs associated with the combination regimen, and hypertension, fatigue, and diarrhea were the most prevalent.¹⁷⁹ The Panel added axitinib + pembrolizumab as a category 2A, other recommended option for patients who are IO therapy naïve. This combination is useful in certain circumstances for patients with prior IO therapy (category 2B).

Cabozantinib + Nivolumab

Apolo et al 2020¹⁸⁰ published data from an ongoing phase I dose escalation trial (ie, NCT02496208) in which patients with metastatic urothelial carcinoma or other genitourinary tumors (including three patients with ccRCC) received combination cabozantinib + nivolumab with or without ipilimumab; data from patients with ccRCC were not reported separately. A pooled analysis of the phase I dose-finding cohort and seven subsequent expansion cohorts, which included 16 patients with metastatic ccRCC, was reported.¹⁸¹ See *Table 2* for efficacy data. In these patients, median OS was 43.6 months (95% CI, 19.4–NE). Although there are no prospective or retrospective published data showing the benefit of cabozantinib + nivolumab in later lines of therapy in the treatment of advanced RCC, the Panel's decision was based on available data for combined ICI/TKI combinations of similar class such as lenvatinib + pembrolizumab and axitinib + pembrolizumab. The Panel added cabozantinib + nivolumab as a category 2A, other recommended option for patients who are IO therapy naïve. This combination is useful in certain circumstances for patients with prior IO therapy (category 2B).

Ipilimumab + Nivolumab

The phase I CheckMate 016 trial included treatment-naïve patients and those who had received one to four or more prior treatment regimens.

Only the ORR results were stratified by treatment status: ORRs in the N3I1 and N1I3 were approximately 46% and 39%, respectively. OS and PFS data were not stratified by treatment line, but were similar.¹⁸² In a single-arm, phase II trial (TITAN-RCC), 32% of the 98 patients who had disease progression after second-line nivolumab had an objective response (OR) with ipilimumab/nivolumab. AEs of any grade occurred in 83% of patients, with grade 3 or 4 AEs occurring in 40% of patients.¹⁸³ In the randomized phase II FRACTION-RCC trial, patients whose disease progressed on or after IO therapy had an ORR of 17.4% on ipilimumab/nivolumab and median duration of response (DOR) of 16.4 months (see *Table 2*). Similarly to TITAN-RCC, AEs of any grade occurred in 78% of patients, with AEs of grade 3 or 4 occurring in 28%.¹⁸⁴ Two of the most common AEs of grade 3 or 4 reported in both trials were elevated lipase and diarrhea.^{183,184} Based on these data, the Panel considers ipilimumab + nivolumab as a category 2A, other recommended option for patients who are IO therapy naïve and useful in certain circumstances option for patients with prior IO therapy.

Belzutifan

Belzutifan inhibits the transcription factor hypoxia-inducible factor 2 α (HIF-2 α) and blocks the heterodimerization of HIF-2 α with HIF-1 β , thereby inducing tumor regression. Follow-up from an expansion cohort of patients, with ccRCC in a phase I/II trial of belzutifan, who had received 1 or more prior therapies showed a disease control rate of 80% among 55 patients. Median PFS was 14.5 months with 51% reporting PFS of 12 months. The most common AEs reported were anemia, fatigue, hypoxia, and dyspnea, among others.¹⁸⁵ Based on these results, belzutifan was considered well tolerated with a favorable safety profile as a single agent. Belzutifan was approved by the FDA for patients with advanced RCC who were previously treated with a PD-1 or PD-L1 inhibitor and VEGF-TKI. The open-label, randomized, phase III Litespark-005 trial compared belzutifan to everolimus in 746 patients with advanced ccRCC whose disease had



NCCN Guidelines Version 2.2026

Kidney Cancer

progressed after first-line therapies.¹⁸⁶ At a median follow-up of 18.4 months, median PFS and ORR for belzutifan were significantly favorable compared to everolimus (see *Table 2*). CR was observed in the interim analyses in the belzutifan arm but not in the everolimus arm. No statistically significant difference in OS was reported at a median follow-up of 25.7 months (see *Table 2*). The safety profile was similar to previous studies and incidence of grade 3–5 treatment-related AEs were similar between the two treatment arms.¹⁸⁶

Based on these data, the Panel included belzutifan as a category 2A, other recommended regimen for patients who had prior IO therapy, particularly those who received PD-1/L1 and VEGF inhibitors. For patients who are IO therapy naïve, belzutifan is a category 2B, useful in certain circumstances subsequent therapy option for patients with ccRCC.

Lenvatinib + Pembrolizumab

The phase II KEYNOTE-146 trial included three groups of patients: treatment-naïve; those who had previously received at least one line of treatment that did not include anti-PD-1 or anti-PD-L1 ICIs; and those who had previously received at least one anti-PD-1 or anti-PD-L1 ICI. See *Table 2* for efficacy data. Treatment-naïve patients had the highest ORR and the longest PFS; ORR and PFS were comparable in the ICI-naïve and ICI treatment-experienced groups. OS was longer only in the ICI-naïve group.¹⁸⁷ In a longer follow-up, the ORR for treatment-naïve, previously treated but ICI-naïve, and ICI-pretreated populations were consistent with previous observations (see *Table 2*). The median DOR was 24.2 months, 9 months, and 14.1 months, respectively. The longer follow-up data showed OS and PFS benefit for lenvatinib + pembrolizumab in all three treatment groups with the treatment-naïve population showing the highest benefit, followed by ICI-pretreated groups, and then the pretreated ICI-naïve group.¹⁸⁸ Hypertension, fatigue, and diarrhea were the most frequent treatment-related AEs and

hypothyroidism was the most frequent immune-related AE.¹⁸⁷ Based on these data, the Panel considers lenvatinib + pembrolizumab a category 2A, other recommended option for patients who are IO therapy naïve and a category 2B, useful in certain circumstances option for patients with prior IO therapy.

Tivozanib

In March 2021, the FDA approved tivozanib, a multitargeted TKI of VEGFR-1, -2, and 3; c-KIT; and PDGFR-β for patients with relapsed or refractory advanced RCC who previously received two or more systemic therapies. Data from the randomized phase III TIVO-3 trial, which enrolled treatment-experienced patients with relapsed or refractory advanced ccRCC, supported the drug's approval. See *Table 2* for efficacy data. Patients receiving tivozanib had significantly longer PFS than those receiving sorafenib; OS was similar between the two groups.^{189,190} In the extended follow-up, a statistically meaningful, higher 12-month landmark, PFS-conditioned OS was observed in those treated with tivozanib versus sorafenib, suggesting that some patients may have an OS benefit with tivozanib.^{190,191} Tivozanib also increased Q-TWiST as compared to sorafenib (15.04 months vs. 12.78 months, respectively).¹⁹² Overall, incidence of AEs grade 3 or higher was lower in those treated with tivozanib compared to sorafenib. Common AEs grade 3 or higher were hypertension and asthenia.^{189,190} Additionally, tivozanib monotherapy was used as the control arm in the TiNivo-2 trial, as described previously, where there was no statistically significant PFS difference between the experimental and control arms in individuals who had received prior IO therapy.¹⁶⁷

Based on these data, the Panel considers tivozanib as a category 2A, other recommended subsequent therapy option for patients who have received prior IO therapy and a useful in certain circumstances option for those who are IO therapy naïve.



NCCN Guidelines Version 2.2026

Kidney Cancer

Lenvatinib

The Panel also includes lenvatinib as a subsequent therapy option useful in certain circumstances (category 2A) regardless of prior IO therapy status based on the phase II trial investigating everolimus/lenvatinib, lenvatinib monotherapy, and everolimus monotherapy described above.^{173,174} PFS favored lenvatinib monotherapy over everolimus monotherapy (see *Table 2* for efficacy data). There was no statistically significant difference in OS between lenvatinib and everolimus monotherapy. No significant difference in PFS or OS was observed between lenvatinib monotherapy and the everolimus/lenvatinib combination.^{173,174} Common grade 3 treatment-emergent AEs in those who received lenvatinib monotherapy included proteinuria, hypertension, and diarrhea. Hypothyroidism was reported more frequently in those receiving single-agent lenvatinib or the combination compared to everolimus.¹⁷³

Pazopanib

A phase III trial comparing pazopanib with placebo, detailed earlier under the *Other Recommended Regimens* for first-line ccRCC, also included patients who had received prior cytokine therapy. PFS was significantly longer with pazopanib (7.4 months) versus placebo (4.2 months) in the treatment-experienced sub-population (HR, 0.54; $P < .001$).¹⁵² However, OS was similar between the two groups (23 vs. 19 months, respectively; HR, 0.82).¹⁵³ Additionally, a prospective phase II trial evaluated second-line pazopanib in patients with advanced metastatic RCC previously treated with a targeted agent (ie, bevacizumab, sunitinib). Twenty-seven percent of patients had an OR to pazopanib; 49% had stable disease (SD). Median PFS was 7.5 months, regardless of prior treatment regimen. Estimated OS rate at 24 months was 43%. AEs were mostly grade 1 or 2, and common grade 3 and 4 AEs included diarrhea and hypertension, similar to the phase III study.¹⁹³ Based on these data, the Panel considers pazopanib a category 2A, useful in certain circumstances subsequent

therapy option for patients with ccRCC regardless of their prior IO therapy status.

Sunitinib

Sunitinib also has demonstrated substantial anti-tumor activity as a second-line therapy in patients with metastatic RCC who progressed on cytokine therapy.^{157,194} Sunitinib is considered a category 2A, useful in certain circumstances subsequent therapy option for patients with ccRCC regardless of their prior IO therapy status.

Everolimus

Everolimus (RAD001) is an orally administered mTOR inhibitor. In the randomized phase III RECORD-1 trial, everolimus was compared with placebo for the treatment of metastatic RCC in patients whose disease had progressed on treatment with sunitinib or sorafenib. The median PFS was significantly longer for everolimus versus placebo, but OS was similar between the two groups.^{195,196} Common AEs included stomatitis, fatigue, and rash, which generally were not severe. Pneumonitis of any grade was detected in <15% of patients treated with everolimus.^{195,196} Everolimus is listed as a category 2A, useful in certain circumstances subsequent therapy option for patients with ccRCC regardless of their prior IO therapy status.

Bevacizumab

Phase II trials have shown benefit of bevacizumab monotherapy after prior treatment with a cytokine. In a phase II study investigating low- and high-dose bevacizumab versus placebo, bevacizumab extended time to progression of disease. Some patients receiving bevacizumab experienced hypertension or proteinuria, with no grade 4 or 5 AEs reported.¹⁹⁷ Bevacizumab is a category 2B, useful in certain circumstances subsequent therapy option for patients with ccRCC regardless of their prior IO therapy status.



NCCN Guidelines Version 2.2026

Kidney Cancer

Systemic Therapy for Patients with Stage IV (M1 or Unresectable T4,M0) or Relapsed Non-Clear Cell RCC

Clinical trials of targeted agents have predominantly focused on patients with ccRCC due to the high prevalence of ccRCC.¹⁹⁸ Data from systematic reviews, meta-analyses, and phase II studies with targeted agents also show some activity in patients with nccRCC. Compared with responses in ccRCC, however, the response rates with these agents are significantly lower for nccRCC. Therefore, according to the Panel, enrollment in clinical trials is a preferred strategy for nccRCC and tumor genomic testing should be considered for trial eligibility.

Non-Clear Cell RCC: Preferred

Cabozantinib

The randomized phase II SWOG 1500 trial compared the MET-targeted TKIs cabozantinib, crizotinib, and savolitinib with standard-of-care sunitinib in patients with advanced papillary RCC who had previously received up to 1 previous systemic therapy, excluding VEGF- and MET-targeted TKIs. Assignment to the crizotinib and savolitinib arms was halted due to results of a prespecified futility analysis. See *Table 3* for efficacy data. Patients receiving cabozantinib had significantly longer PFS and a higher ORR than those receiving sunitinib.¹⁹⁹ Final OS analysis did not show any significant difference in survival.²⁰⁰ The incidence of AEs grade 3 or 4 was highest for cabozantinib (74%), and the most common grade 3 or 4 AEs for cabozantinib were hypertension, fatigue, and hand-foot syndrome.¹⁹⁹ Based on these data, the Panel included cabozantinib as a category 2A, preferred option for patients with nccRCC.

Cabozantinib + Nivolumab

Two separate patient cohorts defined by nccRCC histology in a phase II open-label trial received cabozantinib + nivolumab combination.²⁰¹ ORR for patients with papillary, unclassified, or translocation RCC was 48% with a median follow-up time of 34 months. Median PFS was 13 months (95%

CI, 7–16) and median OS was 28 months (95% CI, 23–43).²⁰² The most prevalent AEs of any grade were fatigue, diarrhea, and palmar-plantar erythrodysesthesia syndrome, and the most common grade 3 or 4 event was hypertension.²⁰¹ Study of patients with chromophobe RCC closed early due to the lack of efficacy. Based on these results, the Panel added cabozantinib + nivolumab as a category 2A, preferred option for first or subsequent-line treatment of stage IV (M1 or unresectable T4,M0) or relapsed nccRCC.

Lenvatinib + Pembrolizumab

The phase II KEYNOTE-B61 trial investigated the combination of lenvatinib + pembrolizumab in untreated, advanced nccRCC.^{203,204} The primary endpoint was ORR, which was 49% in the initial analysis.²⁰³ A 22.8-month follow-up showed similar results (51% ORR).²⁰⁴ See *Table 3* for all efficacy data. The most frequent treatment-related AEs of grade 3 or 4 included hypertension, stomatitis, and proteinuria.²⁰³ The Panel included lenvatinib + pembrolizumab as a category 2A, preferred option for patients in this setting.

Non-Clear Cell RCC: Other Recommended

Erlotinib + Bevacizumab for Advanced Papillary RCC, Including Hereditary Leiomyomatosis and Renal Cell Carcinoma Associated RCC

Hereditary leiomyomatosis and renal cell carcinoma (HLRCC) is a hereditary condition in which affected patients are at risk for development of skin and uterine leiomyomas, as well as an aggressive form of papillary kidney cancer.²⁰⁵

In a phase II trial of patients with advanced papillary RCC (HLRCC-associated RCC; n = 43 or sporadic papillary RCC; n = 40), patients were treated with bevacizumab plus erlotinib.²⁰⁶ Some patients who were enrolled with either subtype received previous therapy (21% with HLRCC; 45% with sporadic papillary RCC), including a VEGFR TKI. The majority of



NCCN Guidelines Version 2.2026

Kidney Cancer

patients had intermediate-risk disease. The ORR was 72% for those with HLRCC compared to 35% with sporadic papillary RCC. Median PFS was 21.1 months in the HLRCC group compared to 8.9 months in the sporadic papillary RCC group. The most frequent AEs of grade 1–2 were acneiform rash, diarrhea, proteinuria, and dry skin, and of grade 3 or higher were hypertension and proteinuria.²⁰⁶ Based on these data, the Panel recommends erlotinib + bevacizumab as a category 2A, other recommended option for select patients with nccRCC and papillary histology, including HLRCC.

Everolimus/Lenvatinib

Extrapolating on data from the phase III everolimus/lenvatinib trial in patients with ccRCC¹⁷³ (see *Subsequent Systemic Therapy Options for Patients with ccRCC*), the Panel added the combination therapy as a category 2A, other recommended regimen for patients with nccRCC.

They also reviewed data²⁰⁷ from a single-arm phase II trial (ie, NCT02915783) that enrolled patients with unresectable advanced or metastatic nccRCC who had not previously received prior systemic therapy. All patients in the trial received combination everolimus/lenvatinib. See *Table 3* for efficacy data. Authors reported that ORR was 26% (95% CI, 12–45). Eight patients in the trial achieved a PR (papillary, n = 3; chromophobe, n = 4; unclassified, n = 1); no patients had a CR. The median DOR was NE. Eighteen patients (58.1%) had SD, and the clinical benefit rate (CR + PR + durable SD [duration ≥23 weeks]) was 61% (95% CI, 42–78). The median PFS was 9.2 months (95% CI, 5.5–NE) and OS was 15.6 months (95% CI, 9.2–NE). AEs of any grade that occurred in >50% of patients included fatigue, diarrhea, nausea, vomiting, and decreased appetite.²⁰⁷ While the Panel conceded that the number of enrolled patients was small, they generally felt that everolimus/lenvatinib treatment led to improved patient outcomes across all nccRCC subtypes.

Nivolumab

A retrospective analysis evaluated the response to at least one dose of nivolumab in patients with metastatic nccRCC.²⁰⁸ See *Table 3* for efficacy data. This study evaluated 35 patients for response and found that 20% had a PR and 29% had SD, with a median follow-up of 8.5 months and median PFS of 3.5 months. Fatigue, fever, rash, and hypothyroidism were the most common AEs observed. A separate retrospective analysis found modest responses with PD-1/PD-L1 inhibitors in 43 patients also with metastatic nccRCC.²⁰⁹ An OR was achieved in eight patients (19%), including four patients (13%) who received PD-1/PD-L1 monotherapy. Based on these data, the Panel considers nivolumab a category 2A, other recommended regimen for patients with nccRCC.

Pembrolizumab

Cohort B of the phase II KEYNOTE-427 study assessed the efficacy and safety of pembrolizumab monotherapy in 165 patients with systemic therapy-naïve, newly diagnosed or recurrent stage IV nccRCC.²¹⁰ See *Table 3* for efficacy data. The majority (about 72%) of patients had confirmed papillary RCC, about 13% had chromophobe RCC, and about 16% had unclassified RCC histology. ORR across all subtypes was approximately 27% (ORR by histology was 29% for papillary, 10% for chromophobe, and 31% for unclassified). Overall PFS and OS were 4.2 months and 28.9 months, respectively. Treatment-related AEs of any grade experienced by patients were most commonly pruritus, fatigue, hypothyroidism, and diarrhea.²¹⁰ Based on these data, the Panel added pembrolizumab as a category 2A, other recommended regimen for patients with nccRCC.

Sunitinib

Two randomized phase II studies compared first-line sunitinib with first-line everolimus in patients with nccRCC. See *Table 3* for efficacy data. While data from the ASPEN trial²¹¹ suggested that patients receiving sunitinib



NCCN Guidelines Version 2.2026

Kidney Cancer

had significantly longer PFS than those receiving everolimus, data from the ESPN trial²¹² suggested that both OS and PFS were similar between the two groups. Common AEs of grade 3 or 4 observed in both trials included hypertension, diarrhea, and fatigue. Other AEs of grade 3 or 4 in the ESPN trial that were of lower incidence in the ASPEN trial included neutropenia and anemia.^{211,212}

Additionally, a meta-analysis of randomized clinical trials for patients with nccRCC found that TKI treatment reduced the risk of progression compared with mTOR inhibitors.²¹³ The study found that sunitinib significantly reduced the risk of progression compared to everolimus in the first-line setting. However, no significant differences between TKIs and mTOR inhibitor treatment were found for OS and ORR.

Based on these data and the phase II SWOG 1500 trial results,¹⁹⁹ the Panel decided sunitinib is a category 2A, other recommended option for patients with nccRCC.

Non-Clear Cell RCC: Useful in Certain Circumstances

Axitinib

A phase II trial of axitinib in 40 patients with recurrent or metastatic nccRCC that progressed after treatment with temsirolimus found a median PFS of 7.4 months, median OS of 12.1 months, and ORR of 37.5%. The most common AEs were hypertension, anorexia, cough, and plantar erythrodysesthesia, which were mostly low grade.²¹⁴ The Panel considers axitinib a category 2A, useful in certain circumstances option for patients with nccRCC.

Everolimus

The efficacy and safety of everolimus in patients with metastatic nccRCC were evaluated in a subgroup of 75 patients enrolled in the REACT trial. ORR and rate of SD were similar between patients with ccRCC and

nccRCC.²¹⁵ In a phase II study of treatment-experienced patients with nccRCC,²¹⁶ median OS was 14 months, ORR was 10.2%, and median PFS was 5.2 months. About 47% of patients experienced AEs of grade 3 or higher, which included anemia and hyperglycemia.²¹⁶ According to data from the phase II RAPTOR trial,²¹⁷ median OS ranged from 24 to 28 months and median PFS ranged from 5 to 8 months, depending on type of papillary nccRCC; patients with type 1 nccRCC had better responses than those with type 2 histology. The most common AEs of any grade in >30% of patients were rash, cough, asthenia, mucosal inflammation, diarrhea, and decreased appetite.²¹⁷ Based on these data, the Panel included everolimus as a category 2A, useful in certain circumstances option for patients with nccRCC.

Everolimus + Bevacizumab

A phase II trial of 34 treatment-naïve patients with metastatic nccRCC studied the efficacy and safety of treatment with bevacizumab plus everolimus.²¹⁸ Median PFS, OS, and ORR were 11.0 months, 18.5 months, and 29%, respectively. Patients with tumors that contained appreciable papillary or chromophobe elements showed significantly higher PFS and ORR than other histologies. Most AEs were grade 1 or 2, and AEs of grade 3 or 4 included hypertension, proteinuria, lymphopenia, hyperglycemia, and hypertriglyceridemia.²¹⁸ Based on these data, the Panel recommends everolimus + bevacizumab as a category 2A, useful in certain circumstances option for patients with nccRCC.

Ipilimumab + Nivolumab

A cohort of 52 patients with advanced nccRCC of the phase 3/4 Checkmate 920 trial received four doses of ipilimumab/nivolumab combination followed by nivolumab for ≤2 years or until disease progression. With 24.1 months of minimum study follow-up, the ORR was 19.6% with a median PFS of 3.7 months and median OS of 21.2 months (95% CI, 16.6–NE).²¹⁹ The most common AEs of any grade were rash,



NCCN Guidelines Version 2.2026

Kidney Cancer

diarrhea/colitis, and hypothyroidism/thyroiditis. Reported AEs of grade 3 or 4 included diarrhea/colitis, rash, nephritis/renal dysfunction, adrenal insufficiency, hepatitis, and hypophysitis.

The randomized phase II SUNNIFORECAST trial investigated ipilimumab/nivolumab in patients with untreated, advanced nccRCC.²²⁰ Patients were randomized to either ipilimumab/nivolumab (n = 157) or investigator's choice of a standard approved regimen (n = 152). OS rate at 12 months was 78% (95% CI, 71–84) with ipilimumab/nivolumab versus 68% (95% CI, 60–75) with standard of care ($P = .026$).²²⁰ Median OS, PFS, and ORR with ipilimumab/nivolumab were 33.2 months, 5.4 months, and 33%, respectively. The same measures with standard-of-care therapy were 25.2 months, 5.7 months, and 20%, respectively. Results were similar in subgroup analyses by histology (ie, papillary and non-papillary RCC). Incidence of AEs (any grade or serious AEs) were similar between treatment arms.²²⁰

Based on these data, the Panel includes ipilimumab + nivolumab as a category 2B option, useful in certain circumstances for advanced nccRCC.

Additional Treatment Options for Rare Types of nccRCC

Among the nccRCC histologies, renal medullary carcinoma (RMC) is extremely rare, comprising approximately 2% of all primary renal tumors in young people.^{221,222} Metastatic disease is seen at presentation in 67% to 95% of patients.^{221–223} Chemotherapy remains the focus of treatment for this subtype, although the prognosis remains dismal.

Collecting-duct carcinoma is also a very rare type of nccRCC, often presenting at an advanced stage of disease. Up to 40% of patients have metastatic spread at initial presentation, and most patients die within 1 to 3 years from the time of primary diagnosis.^{224–227} Collecting duct carcinoma shares biologic features with urothelial carcinoma. In a multicenter prospective study, 23 patients with no prior therapy were treated with a

combination of gemcitabine and either cisplatin or carboplatin.²²⁸ The results showed a response rate of 26% and an OS of 10.5 months.²²⁸

The Panel notes that in patients with other nccRCC subtypes such as collecting duct or medullary subtypes, PRs to cytotoxic chemotherapy have been observed (gemcitabine in combination with carboplatin or cisplatin; or paclitaxel with carboplatin) as well as for other platinum-based chemotherapies currently used for urothelial carcinomas. Gemcitabine in combination with doxorubicin can also produce responses in patients with RMC.^{223,229,230} Oral targeted therapies generally do not produce responses in patients with RMC. Erlotinib in combination with bevacizumab can produce responses even in heavily pretreated patients with RMC. Outside of clinical trials, platinum-based chemotherapy regimens should be the preferred first-line therapy for RMC.

Follow-up Recommendations for Relapsed or Stage IV Disease and Surgically Unresectable Disease

The Panel recommends an H&P examination of patients every 6 to 16 weeks for patients receiving systemic therapy, or more frequently as clinically indicated. Other laboratory evaluations may be carried out as per the requirements for the therapeutic agent being used.

Imaging tests such as CT or MRI (both with and without IV contrast unless contraindicated) should be performed prior to initiating systemic treatment/observation; subsequent imaging may be performed every 6 to 16 weeks as per the physician's discretion, patient's clinical status, and therapeutic schedule. Imaging interval frequency should be altered according to rate of disease change and sites of active disease. MRI (preferred) or CT of head at baseline can be considered, as clinically indicated. Annual surveillance scans can be performed at the physician's discretion. The Panel recommends additional imaging such as MRI of spine and bone scan as clinically indicated. As is recommended with follow-up for stage III and resectable stage IV disease, the Panel



NCCN Guidelines Version 2.2026

Kidney Cancer

recommends FDG-PET be used during follow-up if a patient has FH-deficient or SDHB-deficient RCC.³⁸ Another study investigated the use of FDG-PET for detection of bone metastases in 18 patients with RCC and found that FDG-PET was more accurate and sensitive in detecting bone metastases in these patients than bone scan.³⁹ FDG-PET is also recommended if a patient has bone-predominant disease or if evaluation takes place prior to metastasectomy.

Supportive Care

Supportive care remains a mainstay of therapy for all patients with metastatic RCC (see NCCN Guidelines for Palliative Care available at www.NCCN.org). The Panel recommends radiation therapy as a supportive care option, where SBRT is the preferred approach. SBRT can be used for palliative treatment of symptomatic extracranial metastases.²³¹⁻²³³ See *Principles of Radiation Therapy* in these Guidelines for specific recommendations.

Brain metastasis occurs in ~2% to 8% of patients with advanced RCC, depending on histologic type of RCC.²³⁴ Surgery may be an option for certain patients with oligometastatic disease in the brain whose disease is well-controlled extracranially. Whole brain radiation should only be considered for palliative care in patients with good performance status and disease progression where stereotactic radiosurgery and fractionated stereotactic radiation are not feasible. Pros and cons should be weighed based on patient goals.²³⁵ For recommendations and dosing regimens for brain metastases, as well as whole brain radiation regimens, see *Principles of Radiation Therapy* in the Guidelines.

Surgery also may be appropriate for selected patients with malignant spinal cord compression or impending or actual fractures in weight-bearing bones. The frequency of clinic visits or radiographic and laboratory assessments depends on the individual needs of the patient.

Bone metastasis occurs in ~30% to 40% of patients with advanced RCC.^{14,234,236} Bone lesions in patients with RCC are typically osteolytic and cause considerable morbidity, leading to skeletal-related events (SREs), including bone pain with need for surgery or radiotherapy, hypercalcemia, pathologic fractures, and spinal cord compression. Two studies of patients with bone metastases showed an improvement in bone pain using different radiotherapy modalities.^{237,238}

The role of bone-modifying agents such as bisphosphonates (eg, zoledronic acid) has been established in patients with various malignancies.^{239,240} The bone-modifying agent approved for use in patients with RCC that has metastasized to the bone is the RANK-L inhibitor, denosumab. A phase III randomized trial directly compared the development of SREs on either denosumab or zoledronic acid in patients with multiple myeloma or bone metastases with a solid tumor (excluding breast or prostate cancer).²⁴¹ The study enrolled 1597 patients with solid tumors, including patients with RCC (n = 155). Time to first on-study SRE was 21.4 months with denosumab in patients with solid tumors compared to 15.4 months with zoledronic acid (HR, 0.81; 95% CI, 0.68–0.96; *P* = .017).²⁴²

The Panel accepts the use of a bisphosphonate or a RANK ligand inhibitor in select cases for a patient with bony metastases and creatinine clearance ≥ 30 mL/min. Calcium and vitamin D daily supplements are highly recommended with use of these agents.²⁴¹ An FDA-approved biosimilar is an appropriate substitute for denosumab. Treatment for the palliation of symptoms, especially in patients with marginal performance status and evidence of metastatic disease, includes optimal pain management (see NCCN Guidelines for Adult Cancer Pain, available at www.NCCN.org).



NCCN Guidelines Version 2.2026

Kidney Cancer

Hereditary RCC Syndromes

While hereditary RCC is relatively rare (around 3% of all RCC cases),²⁴³ the Panel felt that it was important to provide recommendations for patients with a suspected or confirmed hereditary RCC syndrome. Accordingly, the Guidelines now describe seven of the most common hereditary RCC syndromes that may predispose patients to RCC: *BAP1* tumor predisposition syndrome (*BAP1*-TPDS), Birt-Hogg-Dubé syndrome (BHDS), HLRCC (FH-deficient), hereditary papillary renal carcinoma (HPRC), hereditary paraganglioma/pheochromocytoma (PGL/PCC) syndrome (SDH-deficient), tuberous sclerosis complex (TSC), and VHL disease. The Guidelines describe kidney-specific clinical features and manifestations of each of these syndromes and known associated genes/inheritance patterns. They also provide genetic testing, surveillance, and treatment recommendations for individuals who are suspected or confirmed to have a hereditary RCC syndrome. While published data informed the majority of these recommendations, the Panel also relied on the real-world experience and expertise of the hereditary subcommittee members to develop recommendations in instances of limited data.

The subcommittee notes that there are some syndromes associated with RCC that overlap with other cancers (eg, Cowden syndrome, Lynch syndrome). For Cowden and Lynch syndromes, the Panel refers readers to the information provided in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate (available at www.NCCN.org). Future versions of the Guidelines may be expanded to include other hereditary syndromes such as microphthalmia-associated transcription factor (MITF)-associated cancer syndrome, which predisposes patients to melanoma and/or RCC.²⁵

The subcommittee also notes that patients with hereditary RCC syndromes often experience non-renal manifestations but felt that input

from clinicians from other specialties (eg, dermatology, endocrinology, neurology, ophthalmology, urology) would be necessary to provide consensus-based recommendations for all potential manifestations. Accordingly, the scope is currently limited to kidney-specific clinical features and manifestations, but the subcommittee identified specialists who may be helpful in managing non-renal manifestations in patients with a hereditary RCC syndrome. Recommendations for genetic testing, surveillance, and treatment vary according to the individual's personal and/or family history of a hereditary RCC syndrome or clinical diagnosis of RCC. Below is a summary of recommendations by patient population.

Genetic Testing and Surveillance Recommendations for Individuals with a Personal or Family History of an RCC Syndrome

The Panel recommends that individuals with a personal or family history of an RCC syndrome or individuals with syndrome features should undergo genetic evaluation. For criteria to be met for further genetic risk evaluation for hereditary RCC syndromes and their histologies, inheritance patterns, and clinical manifestations, see *HERED-RCC-1* and *HERED-RCC-2* in the algorithm. If patients harbor a familial pathogenic or likely pathogenic genetic mutation associated with an RCC syndrome, they should undergo screening for the development of RCC.

For kidney-specific screening in patients who are confirmed to have a hereditary RCC syndrome, but who do not yet have a radiographic or pathologic diagnosis of RCC, the Panel recommends use of abdomen MRI (preferred). CT may also be used for surgical planning purposes, but the Panel warns that use of abdomen CT should be limited due to the potential of increased lifetime radiation exposure. The Panel also includes recommendations on testing intervals and the age at which patients should begin regular screening, as both vary widely by the hereditary RCC syndrome in question. While patients with HLRCC should undergo



NCCN Guidelines Version 2.2026

Kidney Cancer

imaging annually,²⁰⁵ those with less aggressive syndromes such as TSC may benefit from testing at longer intervals.²⁴⁴⁻²⁴⁷

The age at which patients should begin screening also varies by hereditary RCC syndrome. The Panel recommends that patients with confirmed HLRCC, PGL/PCC, TSC, and VHL disease should begin screening in childhood.^{205,244-248} In contrast, those with *BAP1*-TPDS, BHDS, or HPRC should begin screening in adulthood (ie, age 20 years for BHDS, age 30 years for *BAP1*-TPDS and HPRC).^{244,249,250,251} However, the Panel notes that if a patient has a known family member with an early diagnosis of hereditary RCC, screening should begin 10 years before the age that the family member was diagnosed, regardless of the syndrome in question.

Genetic Testing and Screening Recommendations for Patients with a Clinical Diagnosis of RCC Who Have Characteristics Consistent with Inherited RCC

The Panel includes recommendations for patients who already have a clinical or pathologic diagnosis of RCC and have characteristics potentially associated with a hereditary syndrome. This includes RCC diagnosis at ≤46 years of age (though not as sensitive when used as a single criterion), presence of bilateral or multifocal tumors, ≥1 known first- or second-degree relative with RCC, and/or a personal or family history of mesothelioma or uveal melanoma. These patients should also undergo genetic risk assessment, and if indicated, genetic testing. The Panel also recommends genetic risk evaluation for hereditary RCC syndromes for unaffected individuals who have ≥2 first- or second-degree relatives with RCC (on the same side of the family) and/or any first-degree relative with clinical or pathologic diagnosis of a hereditary RCC syndrome who is unable or unwilling to genetically test. If inherited RCC is confirmed, patients should undergo screening as described above, in addition to disease stage-appropriate surveillance.

Kidney-Specific Surgical Recommendations for Patients with a Confirmed Hereditary RCC Syndrome

The Panel also provides surgical recommendations for the majority of the included hereditary RCC syndromes, which are based on published data and/or the subcommittee's real-world experience in treating patients with these syndromes. In order to develop these recommendations, they carefully weighed the potential morbidity and mortality of surgical treatment against the potential aggressiveness of each of the syndromes. They agreed that patients with BHDS, HPRC, and TSC may benefit from more conservative treatment, such as nephron-sparing surgery or percutaneous ablation,^{252,253} while patients with HLRCC/FH-deficient RCC should undergo total radical nephrectomy.²⁰⁵ The Panel's recommendations for surgical treatment of PGL/PCC/SDH-deficient RCC vary by tumor size and histology: those with smaller, less aggressive tumors may be eligible for partial nephrectomy, while those with larger, more aggressive tumors (eg, high-grade, sarcomatoid) should undergo radical nephrectomy.²⁵⁴ For both HLRCC/FH-deficient RCC and PGL/PCC/SDH-deficient RCC, the Panel indicates that FDG-PET is useful for staging pre-nephrectomy, as well as for surveillance.^{38,39} Tumor size also factored into the Panel's surgical recommendations for patients with VHL disease; they noted that these patients are likely to undergo multiple surgical resections during their lifetime that may contribute to chronic and progressive renal failure. Thus, the timing of surgical intervention must be carefully determined to limit the development of metastases and morbidity associated with surgical intervention. They agreed that only patients with VHL disease with tumors approaching 3 cm in diameter should undergo partial nephrectomy (or percutaneous ablation if nephrectomy is contraindicated).^{253,255}



NCCN Guidelines Version 2.2026

Kidney Cancer

Kidney-Specific Systemic Therapy for Patients with Confirmed Hereditary RCC

The Guidelines include a limited number of kidney-specific systemic therapy recommendations for patients with hereditary RCC.

Tuberous Sclerosis Complex (TSC) Angiomyolipoma

Everolimus was approved in April 2012 for treating TSC-associated benign renal angiomyolipomas not requiring immediate surgery based on the randomized phase III EXIST-2 trial.²⁵⁶ In the 4-year update, results were reported for both patients who were originally randomized to everolimus and those who later received everolimus in an unblinded extension phase (n = 112). Confirmed angiomyolipoma response was reported in 58% of patients, and ≥75% had a persistent angiomyolipoma volume decrease of at least 30%.²⁵⁷ The Panel included everolimus as a category 2A, preferred recommendation for patients with TSC-associated angiomyolipoma.

The Panel also included sirolimus as an option in this setting (category 2A, other recommended). Sirolimus, another mTOR inhibitor, was investigated for the treatment of patients with TSC or sporadic lymphangiomyomatosis in a phase I/II trial.²⁵⁸ At the 12-month mark, angiomyolipoma volume (primary endpoint) was reduced by ≥30% for 16 out of 20 patients assessed. Angiomyolipoma volume was also evaluated a year after ending sirolimus treatment, and 5 out of 18 evaluable patients continued to show a volume reduction (≥30%). Common AEs included gastrointestinal events and upper respiratory infections.²⁵⁸

Hereditary Leiomyomatosis and Renal Cell Cancer (HLRCC)

The Panel included erlotinib + bevacizumab for patients with HLRCC/FH-deficient-associated metastatic RCC. While this regimen is not FDA-approved for use in this patient population, its inclusion is supported by clinical trial data showing improved patient outcomes. A phase II study investigating erlotinib + bevacizumab treatment in patients with HLRCC (n

= 43) or sporadic papillary RCC (n = 40) was described above (see *Non-Clear Cell RCC: Other Recommended* and *Table 3* for detailed information). As described above, erlotinib + bevacizumab treatment led to a 72% ORR in patients with HLRCC-associated RCC, median PFS of 21.1 months, and median DOR of 19.3 months.²⁰⁶ Based on these data, the Panel considers erlotinib + bevacizumab a category 2A, other recommended option for patients with HLRCC-associated RCC.

Though data are limited for cabozantinib + nivolumab in HLRCC-associated RCC, the Panel felt that for certain patients this combination can be an option (useful in certain circumstances, category 2A). In the phase II trial investigating cabozantinib + nivolumab in nccRCC discussed earlier, a small subset of patients had FH-deficient RCC (n = 5). Though the sample size was small, the authors noted that an OR was observed in these five patients.²⁰¹

von Hippel-Lindau (VHL) Disease

In August 2021, the FDA approved belzutifan for the treatment of patients with VHL disease-associated RCC who require therapy for RCC but do not require immediate surgery. LITESPARK-004, an open-label, phase II clinical trial, enrolled 61 patients with VHL-associated RCC; 97% had previously undergone a tumor reduction procedure.²⁵⁹ The major efficacy endpoint was ORR, which was 49% (95% CI, 36–62) after a median follow-up of 21.8 months, with 30 patients confirming PRs. Updated results from a 50-month follow-up reported an ORR of 67%, with a CR observed in seven patients and a PR in 34 patients.²⁶⁰ SD was identified in another 30 patients (49%). Median time to response was 8.2 months and median DOR was not reached.²⁵⁹ The Panel considers belzutifan a category 2A, preferred option for patients with VHL-associated-RCC.

The Panel also considers pazopanib a category 2A, useful in certain circumstances option for patients with VHL disease-associated nonmetastatic lesions. In a phase II trial, pazopanib led to a 42% ORR and



NCCN Guidelines Version 2.2026

Kidney Cancer

a 52% renal tumor-specific response rate in 31 patients with VHL disease.²⁶¹

Data Summary

The following tables summarize the key data supporting the inclusion of systemic therapy category 1 and 2A, preferred, and other recommended treatment options for ccRCC and nccRCC. Table 1 includes data on recommended first-line systemic therapies for patients with ccRCC. Table 2 includes data on recommended subsequent systemic therapies for patients with ccRCC. Table 3 includes data on recommended systemic therapies for patients with nccRCC. For all tables, the most recent data are reported unless otherwise indicated.

NCCN Guidelines Version 2.2026

Kidney Cancer

Table 1: Key Studies on First-Line Therapy for Patients with Clear Cell RCC (ccRCC)

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	Median PFS (months)	Median OS (months)
Combination Therapy							
JAVELIN Renal 101 Choueiri et al 2020 ¹⁴⁸ Motzer et al 2019 ¹⁴⁷ Haanen et al 2023 ¹⁵⁰ Choueiri et al 2025 ¹⁴⁹	Axitinib + Avelumab	442	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve, advanced ccRCC; ECOG PS 0–1 270 patients in the axitinib/avelumab arm and 290 patients in the sunitinib arm were PD-L1+.	~74	<u>ORR: Overall pop.</u> Axi/Ave: 59.7 (95% CI, 55.0–64.3) Sunitinib: 32.0 (95% CI, 27.7–36.5) <u>CR (%): Overall pop.</u> Axi/Ave: 5.7 Sunitinib: 3.6 <u>PD-L1+</u> Axi/Ave: 64.8 (95% CI, 58.8–70.5) Sunitinib: 31.4 (95% CI, 26.1–37.1) <u>CR (%): PD-L1+</u> Axi/Ave: 7.8 Sunitinib: 4.5	<u>Overall population</u> Axi/Ave: 13.9 (95% CI, 11.1–16.6) Sunitinib: 8.5 (95% CI, 8.2–9.7) HR, 0.66 (95% CI, 0.566–0.769) P < .0001 <u>PD-L1+</u> Axi/Ave: 13.9 (95% CI, 11.0–17.8) Sunitinib: 8.2 (95% CI, 6.9–9.1) HR, 0.57 (95% CI, 0.469–0.697) P < .0001	<u>Overall population</u> Axi/Ave: 44.8 (95% CI, 39.7–51.1) Sunitinib: 38.9 (95% CI, 31.4–45.2) HR, 0.88 (95% CI, 0.749–1.039) P = .0669 <u>PD-L1+</u> Axi/Ave: 43.2 (95% CI, 36.5–51.7) Sunitinib: 36.2 (95% CI, 29.8–44.2) HR, 0.86 (95% CI, 0.701–1.057) P = .0755
	Sunitinib	444	In the extended follow-up, data for PD-L1+ patients were reported for PFS and OS.				
KEYNOTE-426 Rini et al 2019 ¹³¹ Powles et al 2020 ¹³² Plimack et al 2023 ¹³³ Rini et al 2025 ¹³⁴	Axitinib + Pembrolizumab	432	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve, advanced ccRCC; Karnofsky PS ≥70%	67.2	Axi/Pem: 60.6 (95% CI, 55.9–65.3) Sunitinib: 39.6 (95% CI, 35–44.4)	Axi/Pem: 15.7 (95% CI, 13.6–20.2) Sunitinib: 11.1 (95% CI, 8.9–12.5)	Axi/Pem: 47.2 (95% CI, 43.6–54.8) Sunitinib: 40.8 (95% CI, 34.3–47.5)
	Sunitinib	429	Noted in the recent analysis: >60% of patients in each arm received subsequent therapy after study treatment discontinuation		<u>CR (%)</u> Axi/Pem: 11.6 Sunitinib: 4	HR, 0.69 (95% CI, 0.59–0.81)	HR, 0.84 (95% CI, 0.71–0.99)
CheckMate 9ER Choueiri et al 2021 ¹³⁵ Motzer et al 2022 ¹³⁶ Motzer et al 2026 ¹³⁸	Cabozantinib + Nivolumab	323	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve, advanced ccRCC; Karnofsky PS ≥70%	67.6	Cabo/Nivo: 55.7 (95% CI, 50.1–61.2) Sunitinib: 27.4 (95% CI, 23–32.9)	Cabo/Nivo: 16.4 (95% CI, 12.5–19.3) Sunitinib: 8.3 (95% CI, 7.0–9.7)	Cabo/Nivo: 46.5 (95% CI, 40.6–53.8) Sunitinib: 35.5 (95% CI, 29.2–42.8)
	Sunitinib	328	Data presented here are for the ITT population and not based on risk group.		<u>CR (%)</u> Cabo/Nivo: 13.9 Sunitinib: 4.6	HR, 0.58 (95% CI, 0.49–0.70)	HR, 0.79 (95% CI, 0.65–0.96)



NCCN Guidelines Version 2.2026

Kidney Cancer

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	Median PFS (months)	Median OS (months)	
CheckMate 214 Motzer et al 2018 ¹³⁹ Motzer et al 2022 ¹⁴⁰ Tannir et al 2024 ¹⁴¹	Ipilimumab + Nivolumab	550	The study enrolled 425 intermediate-risk, 422 poor-risk, and 249 favorable-risk patients with systemic therapy-naïve, advanced ccRCC; Karnofsky PS ≥70% Note: The study's coprimary endpoints were ORR, OS, and PFS in intermediate- and poor-risk patients. Exploratory analyses of data in favorable-risk patients were reported separately. Combined data for all risk groups are not shown.	99.1	Ipi/Nivo: 39.5 Sunitinib: 33 <u>Intermediate-/poor-risk</u> Ipi/Nivo: 42.4 (95% CI, 37.6–47.2) Sunitinib: 27.5 (95% CI, 23.3–32) <u>CR (%)</u> Ipi/Nivo: 11.8 Sunitinib: 2.6 <u>Favorable-risk</u> Ipi/Nivo: 29.6 (95% CI, 21.8–38.4) Sunitinib: 51.6 (95% CI, 42.5%–60.7%) <u>CR (%)</u> Ipi/Nivo: 12.8 Sunitinib: 6.5	Ipi/Nivo: 12.4 Sunitinib: 12.3 <u>HR, 0.88</u> <u>(95% CI, 0.75–1.03)</u> <u>Intermediate-/poor-risk</u> Ipi/Nivo: 12.4 (95% CI, 8.7–16.8) Sunitinib: 8.5 (95% CI, 7.0–11.1) <u>HR, 0.73</u> <u>(95% CI, 0.61–0.87)</u> <u>Favorable-risk</u> Ipi/Nivo: 12.4 (95% CI, 10.3–18) Sunitinib: 28.9 (95% CI, 23.2–42.8) <u>HR, 1.76</u> <u>(95% CI, 1.25–2.48)</u>	Ipi/Nivo: 52.7 Sunitinib: 37.8 <u>HR, 0.72</u> <u>(95% CI, 0.62–0.83)</u> <u>Intermediate-/poor-risk</u> Ipi/Nivo: 46.7 (95% CI, 35.0–55.7) Sunitinib: 26 (95% CI, 21.8–32.6) <u>HR, 0.69</u> <u>(95% CI, 0.59–0.81)</u> <u>Favorable-risk</u> Ipi/Nivo: 77.9 (95% CI, 64.6–91.6) Sunitinib: 66.7 (95% CI, 56.0–79.9) <u>HR, 0.82</u> <u>(95% CI, 0.60–1.13)</u>	
	Sunitinib	546						
CLEAR Motzer et al 2021 ¹⁴² Choueiri et al 2023 ¹⁴³ Motzer et al 2024 ¹⁴⁴ Note: For Len/Pem the most recent data are shown	Lenvatinib + Pembrolizumab	355	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve, advanced ccRCC; Karnofsky PS ≥70%	49	Len/Pem: 71.3 (95% CI, 66.6–76) Len/Ev: 54 Sunitinib: 36.7 (95% CI, 31.7–41.7) <u>Len/Pem vs. Sunitinib</u> RR: 1.94 (95% CI, 1.67–2.26) <i>P</i> < .0001 <u>Len/Ev vs. Sunitinib</u> RR: 1.48 (95% CI, 1.26–1.74) <u>CR</u> Len/Pem: 16 Lev/Ev: 10 Sunitinib: 4	Len/Pem: 23.9 (95% CI, 20.8–27.7) Len/Ev: 14.7 (95% CI, 11.1–16.7) Sunitinib: 9.2 (95% CI, 6.0–11.0) <u>Len/Pem vs. Sunitinib</u> HR, 0.39 (95% CI, 0.32–0.49) <i>P</i> < .0001 <u>Len/Ev vs. Sunitinib</u> HR, 0.65 (95% CI, 0.53–0.80) <i>P</i> < .001	Len/Pem: 53.7 (95% CI, 48.7–NE) Len/Ev: NR Sunitinib: 54.3 (95% CI, 40.9–NE) <u>Len/Pem vs. Sunitinib</u> HR, 0.79 (95% CI, 0.63–0.99) <i>P</i> = .0424 <u>Len/Ev vs. Sunitinib</u> HR, 1.15 (95% CI, 0.88–1.50) <i>P</i> = .30	
	Everolimus/ Lenvatinib	357						
	Sunitinib	357						



NCCN Guidelines Version 2.2026

Kidney Cancer

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	Median PFS (months)	Median OS (months)
Monotherapy							
VEG105192 Sternberg et al 2013 ¹⁵³ (OS data) Sternberg et al 2010 ¹⁵² (PFS and ORR data)	Pazopanib	290	Favorable-, intermediate-, or poor-risk, locally advanced or metastatic ccRCC; ECOG PS 0–1 Note: Of 435 enrolled patients, 202 received prior cytokine treatment and 233 were systemic therapy-naïve. Data were reported separately. See Table 2 for data for patients who received prior treatment.	Median NR; Up to 24 months for primary outcome	Pazopanib: 32 (95% CI, 24–39) Placebo: 4 (95% CI, 0–8)	Pazopanib: 11.1 Placebo: 2.8 HR, 0.40 (95% CI, 0.27–0.60) <i>P</i> < .0001	Pazopanib: 23 Placebo: 24 HR, 1.01 (95% CI, 0.72–1.42) <i>P</i> value NR
	Placebo	145					
COMPARZ Motzer et al 2013 ¹⁵⁴ Note: In 2014, updated OS data were reported in a correspondence letter to the publishing journal. ¹⁵⁵ Only the most recent OS data are shown.	Pazopanib	557	Favorable- or intermediate-risk, systemic therapy-naïve, advanced or metastatic ccRCC; Karnofsky PS ≥70%	Median NR; Up to 48 months for primary outcome	Pazopanib: 31 Sunitinib: 25 <i>P</i> = .03	Pazopanib: 8.4 (95% CI, 8.3–10.9) Sunitinib: 9.5 (95% CI, 8.3–11.1) HR, 1.05 (95% CI, 0.90–1.22) noninferior	Pazopanib: 28 (95% CI, 26–36) Sunitinib: 29 (95% CI, 25–33) HR, 0.92 (95% CI, 0.79–1.06) <i>P</i> = .24
	Sunitinib	553					
Phase III trial Motzer et al 2007 ¹⁵⁶	Sunitinib	375	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve metastatic ccRCC; ECOG PS 0–1	NR	Sunitinib: 31 (95% CI, 26–36) Interferon: 6 (95% CI, 4–9) <i>P</i> < .001	Sunitinib: 11 (95% CI, 10–12) Interferon: 5 (95% CI, 4–6) HR, 0.42 (95% CI, 0.32–0.54) <i>P</i> < .001	Sunitinib: NR Interferon: NR HR, 0.65 (95% CI, 0.45–0.94) <i>P</i> = .02
	Interferon alfa	375					
CABOSUN Choueiri et al 2017 ¹⁴⁵	Cabozantinib	79	Intermediate- or poor-risk, systemic therapy-naïve, advanced or metastatic ccRCC, ECOG PS 0–2	21.4	Cabo: 33 (95% CI, 23–44) Sunitinib: 12 (95% CI, 5.4–21)	Cabo: 8.2 (95% CI, 6.2–8.8) Sunitinib: 5.6 (95% CI, 3.4–8.1)	Cabo: 30.3 (95% CI, 14.6–35) Sunitinib: 21.8 (95% CI, 16.3–27) HR, 0.80 (95% CI, 0.50–1.26)
	Sunitinib	78					



NCCN Guidelines Version 2.2026

Kidney Cancer

Table 2: Key Studies on Subsequent Therapy for Patients with Clear Cell RCC (ccRCC)

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	PFS (months)	OS (months)
Combination Therapy							
Phase I/II study Apolo et al 2024 ¹⁸¹ Note: Only data for ccRCC are shown	Cabozantinib + Nivolumab +/- Ipilimumab	16	Metastatic ccRCC; 3 patients with sarcomatoid features (>50%); received at least one line of therapy; Karnofsky PS ≥70%	49.2	62.5 (95% CI, 35.4–84.8)	16.3 (95% CI, 6.4–21.8)	43.6 (95% CI, 19.4–NE)
FRACTION-RCC Choueiri et al 2022 ¹⁸⁴ Note: Data for Track 2 patients who had prior IO therapy	Ipilimumab + Nivolumab	46	Favorable-, intermediate-, or poor-risk advanced RCC; enrolled in 1 of 2 tracks; Karnofsky PS ≥70%. Patients in 2 tracks: Track 1 (IO therapy naïve, stratified according to previous TKI); Track 2 (prior IO), received 1 of 5 treatments containing nivolumab; Track 1 patients who progressed were then enrolled in Track 2	33.8	17.4 (95% CI, 7.8–31.4)	3.7 (95% CI, 2–7.3)	23.8 (95% CI, 13.2–NE)
Phase II study Motzer et al 2015 ¹⁷³ Motzer et al 2016 ¹⁷⁴ Note: Data from primary publication shown	Everolimus/Lenvatinib	51	Favorable-, intermediate-, or poor-risk advanced or metastatic ccRCC; received at least one VEGFR-targeted TKI with progression within 9 months of treatment; ECOG PS 0–1	17–19; varied by group	Ev/Len: 43 Ev: 6 Len: 27 <u>Ev/Len vs. Len</u> P < .0001 <u>Len vs. Ev</u> P = .0067	Ev/Len: 14.6 (95% CI, 5.9–20.1) Ev: 5.5 (95% CI, 3.5–7.1) Len: 7.4 (95% CI, 5.6–10.2)	Ev/Len: 25.5 (95% CI, 16.4–NE) Ev: 15.4 (95% CI, 11.8–19.6) Len: 19.1 (95% CI, 13.6–26.2)
	Everolimus	50				<u>Ev/Len vs. Ev</u> HR, 0.40 (95% CI, 0.24–0.68) P = .0005	<u>Ev/Len vs. Ev</u> HR, 0.51 (95% CI, 0.30–0.88) P = .024
	Lenvatinib	52				<u>Ev/Len vs. Len</u> HR, 0.66 (95% CI, 0.39–1.10) P = .12 <u>Len vs. Ev</u> HR, 0.61 (95% CI, 0.39–0.98) P = .048	<u>Len vs. Ev/Len</u> HR, 0.75 (95% CI, 0.43–1.30) P = .32 <u>Len vs. Ev</u> HR, 0.68 (95% CI, 0.41–1.14) P = .12



NCCN Guidelines Version 2.2026

Kidney Cancer

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	PFS (months)	OS (months)
KEYNOTE-146 Lee et al 2021 ¹⁸⁷ Lee et al 2023 ¹⁸⁸	Lenvatinib + Pembrolizumab, previously treated but ICI-naïve (2+L ICI-naïve)	17	Favorable-, intermediate-, or poor-risk metastatic ccRCC; ECOG PS 0–1	6–51 months; varied by outcome	2+L, ICI-naïve: 52.9 (95% CI, 27.8–77.0)	2+L, ICI-naïve: 11.8 (95% CI, 5.5–18.6)	2+L, ICI-naïve: 30.3 (95% CI, 28.7–NE)
	Lenvatinib + Pembrolizumab, ICI treatment-experienced (2+L ICI-TE)	104			2+L, ICI-TE: 58.7 (95% CI, 48.6–68.2)	2+L, ICI-TE: 11.6 (95% CI, 7.6–14.1)	2+L, ICI-TE: 32.1 (95% CI, 26.4–NE)
	Lenvatinib + Pembrolizumab, treatment-naïve (TN)	22			TN: 77.3 (95% CI, 54.6–92.2)	TN: 22.1 (95% CI, 11.6–31.7)	TN: 55.8 (95% CI, 31.4–NE)
Monotherapy							
AXIS Rini et al 2011 ¹⁶⁴ Motzer et al 2013 ¹⁷⁸	Axitinib	361	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve metastatic ccRCC; ECOG PS 0–1	Up to 36 months	Axi: 19 Sor: 9 <i>P</i> = .0001	Axi: 8.3 (95% CI, 6.7–9.2) Sor: 5.7 (95% CI, 4.7–6.5)	Axi: 20.1 (95% CI, 16.7–23.4) Sor: 19.2 (95% CI, 17.5–22.3)
	Sorafenib	362				HR, 0.67 (95% CI, 0.55–0.78) <i>P</i> < .0001	HR, 0.97 (95% CI, 0.80–1.17) <i>P</i> = .37
METEOR Choueiri et al 2016 ¹⁶⁹ Choueiri et al 2015 ¹⁶⁸ Motzer et al 2018 ¹⁷⁰	Cabozantinib	330	Favorable-, intermediate-, or poor-risk advanced or metastatic ccRCC; received at least one VEGFR-targeted TKI with progression within 6 months of treatment; Karnofsky PS ≥70%	OS: 22 ¹⁷⁰ ORR, PFS: 19 ¹⁶⁹	Cabo: 17 Ev: 3 <i>P</i> < .0001	Cabo: 7.4 (95% CI, 6.6–9.1) Ev: 3.9 (95% CI, 3.7–5.1)	Cabo: 21.4 Ev: 17.1
	Everolimus	328				HR, 0.51 (95% CI, 0.41–0.62) <i>P</i> < .0001	HR, 0.70 (95% CI, 0.58–0.85) <i>P</i> = .0002
CheckMate 025 Motzer et al 2015 ¹⁷⁶ Motzer et al 2020 ¹⁷⁷	Nivolumab	406	Favorable-, intermediate-, or poor-risk advanced or metastatic ccRCC; received 1–2 prior antiangiogenic therapies (except mTOR inhibitors); Karnofsky PS ≥70%	72	Nivo: 22.9 (95% CI, 18.9–27.3) Ev: 4.1 (95% CI, 2.4–6.5) OR, 6.86 (95% CI, 4–11.7) <i>P</i> < .0001	Nivo: 4.2 (95% CI, 3.7–5.4) Ev: 4.5 (95% CI, 3.7–5.5)	Nivo: 25.8 (95% CI, 22.2–29.8) Ev: 19.7 (95% CI, 17.6–22.1)
	Everolimus	397				HR, 0.84 (95% CI, 0.72–0.99) <i>P</i> = .0331	HR, 0.73 (95% CI, 0.62–0.85) <i>P</i> < .0001



NCCN Guidelines Version 2.2026

Kidney Cancer

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	PFS (months)	OS (months)
TIVO-3 Rini et al 2020 ¹⁸⁹ Beckermann et al 2024 ¹⁹⁰ (extended follow-up [mean 22.8 mo.] for PFS and OS)	Tivozanib	175	Favorable-, intermediate-, or poor-risk metastatic ccRCC; received 2–3 prior systemic therapies including at least 1 VEGFR-targeted TKI other than sorafenib or tivozanib; ECOG PS 0–1; ITT population shown	19	Tivo: 18 Sor: 8	Tivo: 5.6 (95% CI, 5.3–7.3) Sor: 3.9 (95% CI, 3.7–5.6)	Tivo: 16.4 (95% CI, 13.4–21.9) Sor: 19.1 (95% CI, 14.9–24.2)
	Sorafenib	175				HR, 0.73 (95% CI, 0.56–0.94) <i>P</i> = .016 Extended follow-up: HR, 0.624 (95% CI, 0.49–0.79) <i>P</i> < .0001	HR, 0.89 (95% CI, 0.70–1.14) <i>P</i> = .35 12-mo landmark PFS-conditioned OS: Tivo: 48.3 Sor: 32.8 HR, 0.45 (95% CI, 0.22–0.99) <i>P</i> = .0221
LITESPARK-005 Choueiri et al 2024 ¹⁸⁶	Belzutifan	374	Unresectable, locally advanced, or metastatic ccRCC; 1–3 prior therapies, progressed PD-1 or PD-L1 inhibitor and VEGF-targeted TKI; Karnofsky PS ≥70%	18.4 (PFS, ORR)	Bel: 21.9 (95% CI, 17.8–26.5) Ev: 3.5 (95% CI, 1.9–5.9)	Bel: 5.6 (95% CI, 3.9–7.0) Ev: 5.6 (95% CI, 4.8–5.8)	Bel: 21.4 (95% CI, 18.2–24.3) Ev: 18.1 (95% CI, 15.8–21.8)
	Everolimus	372		25.7 (OS)		HR, 0.75 (95% CI, 0.63–0.90) <i>P</i> < .001	HR, 0.88 (95% CI, 0.73–1.07) <i>P</i> = .20

Table 3: Key Studies on Systemic Therapy for Patients with Non-Clear Cell RCC (nccRCC)

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	PFS (months)	OS (months)
Combination Therapy							
Phase II, cohort study Lee et al 2022 ²⁰¹ Fitzgerald et al 2024 ²⁰²	Cabozantinib + Nivolumab	47	Advanced nccRCC, underwent 0–1 prior systemic therapies	34	48 (95% CI, 32–64)	13 (95% CI, 7–16)	28 (95% CI, 23–43)
KEYNOTE-B61 Albiges et al 2023 ²⁰³ Voss et al 2025 ²⁰⁴ (ORR analysis)	Lenvatinib + Pembrolizumab	158	Untreated, stage IV nccRCC	22.8	51 (95% CI, 43–59)	17.9 (95% CI, 15–21.1)	41.5 (95% CI, 32.8–NR)
Phase II trial Srinivasan et al 2025 ²⁰⁶	Erlotinib + Bevacizumab	HLRCC: 43 Sporadic papillary: 40	Metastatic HLRCC or sporadic papillary RCC; majority of patients had intermediate risk per IMDC; 0–3 lines of therapy, including VEGFR TKI	71.9	HLRCC: 72 (95% CI, 57–83) Sporadic papillary: 35 (95% CI, 22–51)	HLRCC: 21.1 (95% CI, 15.6–26.6) Sporadic papillary: 8.9 (95% CI, 5.5–8.3)	HLRCC: 44.6 (95% CI, 32.7–NE) Sporadic papillary: 18.2 (95% CI, 12.6–29.3)



NCCN Guidelines Version 2.2026

Kidney Cancer

Trial/Author	Regimen	No. of Patients	Patient Characteristics	Median Follow-up (months)	ORR (%)	PFS (months)	OS (months)
Phase II trial Hutson et al 2021 ²⁰⁷	Everolimus/Lenvatinib	31	Unresectable advanced or metastatic nccRCC	NR	PR: 26 SD: 58	9.2 (95% CI, 5.5–NE)	15.6 (95% CI, 9.2–NE)
Monotherapy							
Phase II SWOG 1500 Pal et al 2021 ¹⁹⁹ Barata et al 2024 ²⁰⁰ (final OS analysis) Note: The trial also included savolitinib and crizotinib groups; assignment was halted after a futility analysis.	Cabozantinib	46	Favorable-, intermediate-, or poor-risk metastatic papillary RCC; previously received 0–1 therapies, excluding VEGFR and MET TKIs	NR; up to 36 months follow-up specified in trial 17.5 (for OS)	Cabo: 23 Sun: 4 <i>P</i> = .010	Cabo: 9.0 (95% CI, 6–12) Sun: 5.6 (95% CI, 3–7)	Cabo: 21.5 (95% CI, 12–28.1) Sun: 17.3 (95% CI, 12.8–21.8)
	Sunitinib	44				HR, 0.60 (95% CI, 0.37–0.97) <i>P</i> = .019	HR, 0.83 (95% CI, 0.51–1.36) <i>P</i> = .46
Retrospective study Koshkin et al 2018 ²⁰⁸	Nivolumab	35	Metastatic nccRCC	9	PR: 20 SD: 29	3.5	NR
Phase II KEYNOTE-427 (cohort B) McDermott et al 2021 ²¹⁰	Pembrolizumab	165	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve, newly diagnosed, or recurrent stage IV nccRCC; Karnofsky PS ≥70%	32	27	4.2 (95% CI, 2.9–5.6)	28.9 (95% CI, 24.3–NE)
Phase II ASPEN trial Armstrong et al 2016 ²¹¹	Sunitinib	51	Favorable-, intermediate-, or poor-risk, systemic therapy-naïve metastatic nccRCC (papillary, chromophobe, or unclassified); Karnofsky PS ≥60%	12–13	Sun: 18 Evero: 9	Sun: 8.3 (80% CI, 5.8–11.4) Evero: 5.6 (80% CI, 5.5–6.0)	Sun: 31.5 (95% CI, 14.8–NE) Evero: 13.2 (95% CI, 9.7–37.9)
	Everolimus	57				HR, 1.41 (80% CI, 1.03–1.92) <i>P</i> = .16	HR, 1.12 (95% CI, 0.7–2.1) <i>P</i> = .60
Phase II ESPN trial Tannir et al 2016 ²¹²	Sunitinib	33	Good-, intermediate-, or poor-risk, systemic therapy-naïve metastatic nccRCC (papillary, chromophobe, collecting duct, Xp11.2 translocation, unclassified) or ccRCC with >20% sarcomatoid features; ECOG PS 0–1	24	6	Sun: 6.1 (95% CI, 4.2–9.4) Evero: 4.1 (95% CI, 2.7–10.5) <i>P</i> = .60	Sun: 16.2 (95% CI, 14.2–NE) Evero: 14.9 (95% CI, 8.0–23.4) <i>P</i> = .18
	Everolimus	35					



NCCN Guidelines Version 2.2026

Kidney Cancer

References

1. Siegel RL, Kratzer TB, Wagle NS, et al. Cancer statistics, 2026. CA Cancer J Clin 2026;76:e70043. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41528114>.
2. SEER cancer stat facts: Kidney and renal pelvis cancer. Bethesda, MD: National Cancer Institute; Available at: <https://seer.cancer.gov/statfacts/html/kidrp.html>. Accessed September 2025.
3. Barchuk A, Vignat J, Tikkinen KAO, et al. Global patterns and trends in kidney cancer incidence and mortality. Int J Cancer 2026. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41609424>.
4. Ficarra V, Schips L, Guille F, et al. Multiinstitutional European validation of the 2002 TNM staging system in conventional and papillary localized renal cell carcinoma. Cancer 2005;104:968–974. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16007683>.
5. Frank I, Blute ML, Leibovich BC, et al. Independent validation of the 2002 American Joint Committee on cancer primary tumor classification for renal cell carcinoma using a large, single institution cohort. J Urol 2005;173:1889–1892. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15879769>.
6. Zisman A, Pantuck AJ, Chao D, et al. Reevaluation of the 1997 TNM classification for renal cell carcinoma: T1 and T2 cutoff point at 4.5 rather than 7 cm. better correlates with clinical outcome. J Urol 2001;166:54–58. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11435822>.
7. Klatte T, Patard JJ, Goel RH, et al. Prognostic impact of tumor size on pT2 renal cell carcinoma: an international multicenter experience. J Urol 2007;178:35–40; discussion 40. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17521678>.
8. Lam JS, Klatte T, Patard JJ, et al. Prognostic relevance of tumour size in T3a renal cell carcinoma: a multicentre experience. Eur Urol 2007;52:155–162. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17316970>.
9. Minervini A, Lilas L, Minervini R, Selli C. Prognostic value of nuclear grading in patients with intracapsular (pT1-pT2) renal cell carcinoma. Long-term analysis in 213 patients. Cancer 2002;94:2590–2595. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12173325>.
10. Dall'Oglio MF, Antunes AA, Sarkis AS, et al. Microvascular tumour invasion in renal cell carcinoma: the most important prognostic factor. BJU Int 2007;100:552–555. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17555475>.
11. Dall'Oglio MF, Ribeiro-Filho LA, Antunes AA, et al. Microvascular tumor invasion, tumor size and Fuhrman grade: a pathological triad for prognostic evaluation of renal cell carcinoma. J Urol 2007;178:425–428; discussion 428. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17561167>.
12. Lam JS, Shvarts O, Said JW, et al. Clinicopathologic and molecular correlations of necrosis in the primary tumor of patients with renal cell carcinoma. Cancer 2005;103:2517–2525. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15880379>.
13. Sengupta S, Lohse CM, Leibovich BC, et al. Histologic coagulative tumor necrosis as a prognostic indicator of renal cell carcinoma aggressiveness. Cancer 2005;104:511–520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15973740>.
14. Bianchi M, Sun M, Jeldres C, et al. Distribution of metastatic sites in renal cell carcinoma: a population-based analysis. Ann Oncol 2012;23:973–980. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21890909>.
15. DeVita VT Jr HS, Rosenberg SA. Cancer principles and practice of oncology. (ed 8th). Philadelphia, PA: Lippincott Williams & Wilkins; 2008.
16. Meyer CP, Sun M, Karam JA, et al. Complications After Metastasectomy for Renal Cell Carcinoma-A Population-based



NCCN Guidelines Version 2.2026

Kidney Cancer

Assessment. Eur Urol 2017;72:171–174. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28359734>.

17. Moch H, Gasser T, Amin MB, et al. Prognostic utility of the recently recommended histologic classification and revised TNM staging system of renal cell carcinoma: a Swiss experience with 588 tumors. Cancer 2000;89:604–614. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/10931460>.

18. Leibovich BC, Lohse CM, Crispen PL, et al. Histological subtype is an independent predictor of outcome for patients with renal cell carcinoma. J Urol 2010;183:1309–1315. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20171681>.

19. Mousavi SE, Najafi M, Aslani A, et al. A population-based study on incidence trends of kidney and renal pelvis cancers in the United States over 2000-2020. Sci Rep 2024;14:11294. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/38760399>.

20. Oza B, Frangou E, Eisen T, et al. Determining the impact of histology on the incidence, pattern, and timing of recurrences in patients with renal cell carcinoma: A pooled analysis from the SORCE and ASSURE trials. Eur Urol Open Sci 2025;79:19–26. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/40756159>.

21. Moch H, Amin MB, Berney DM, et al. The 2022 World Health Organization classification of tumours of the urinary system and male genital organs-part a: Renal, penile, and testicular tumours. Eur Urol 2022;82:458–468. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/35853783>.

22. Msaouel P, Hong AL, Mullen EA, et al. Updated recommendations on the diagnosis, management, and clinical trial eligibility criteria for patients with renal medullary carcinoma. Clin Genitourin Cancer 2019;17:1–6. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30287223>.

23. Trpkov K, Hes O, Williamson SR, et al. New developments in existing WHO entities and evolving molecular concepts: The Genitourinary Pathology Society (GUPS) update on renal neoplasia. Mod Pathol

2021;34:1392–1424. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/33664427>.

24. Choyke PL, Glenn GM, Walther MM, et al. Hereditary renal cancers. Radiology 2003;226:33–46. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/12511666>.

25. Schmidt LS, Linehan WM. Genetic predisposition to kidney cancer. Semin Oncol 2016;43:566–574. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27899189>.

26. DeVita VT Jr LT, Rosenberg SA. DeVita, Hellman, and Rosenberg's Cancer Principles and Practice of Oncology. (ed 10th). Philadelphia, PA: Wolters Kluwer Health; 2015.

27. Freedman-Cass DA, Fischer T, Alpert AB, et al. The value and process of inclusion: Using sensitive, respectful, and inclusive language and images in NCCN content. J Natl Compr Canc Netw 2023;21:434–441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37156485>.

28. Jayson M, Sanders H. Increased incidence of serendipitously discovered renal cell carcinoma. Urology 1998;51:203–205. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/9495698>.

29. Luciani LG, Cestari R, Tallarigo C. Incidental renal cell carcinoma-age and stage characterization and clinical implications: study of 1092 patients (1982-1997). Urology 2000;56:58–62. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/10869624>.

30. Israel GM, Bosniak MA. How I do it: evaluating renal masses. Radiology 2005;236:441–450. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16040900>.

31. Lim DJ, Carter MF. Computerized tomography in the preoperative staging for pulmonary metastases in patients with renal cell carcinoma. J Urol 1993;150:1112–1114. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/8371366>.



NCCN Guidelines Version 2.2026 Kidney Cancer

32. Sheth S, Scatarige JC, Horton KM, et al. Current concepts in the diagnosis and management of renal cell carcinoma: role of multidetector ct and three-dimensional CT. Radiographics 2001;21 Spec No:S237–254. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11598260>.

33. Hricak H, Demas BE, Williams RD, et al. Magnetic resonance imaging in the diagnosis and staging of renal and perirenal neoplasms. Radiology 1985;154:709–715. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3969475>.

34. Janus CL, Mendelson DS. Comparison of MRI and CT for study of renal and perirenal masses. Crit Rev Diagn Imaging 1991;32:69–118. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1863349>.

35. Seaman E, Goluboff ET, Ross S, Sawczuk IS. Association of radionuclide bone scan and serum alkaline phosphatase in patients with metastatic renal cell carcinoma. Urology 1996;48:692–695. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8911510>.

36. Shannon BA, Cohen RJ, de Bruto H, Davies RJ. The value of preoperative needle core biopsy for diagnosing benign lesions among small, incidentally detected renal masses. J Urol 2008;180:1257–1261; discussion 1261. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18707712>.

37. Park JW, Jo MK, Lee HM. Significance of 18F-fluorodeoxyglucose positron-emission tomography/computed tomography for the postoperative surveillance of advanced renal cell carcinoma. BJU Int 2009;103:615–619. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19007371>.

38. Nikolovski I, Carlo MI, Chen YB, Vargas HA. Imaging features of fumarate hydratase-deficient renal cell carcinomas: a retrospective study. Cancer Imaging 2021;21:24. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33608050>.

39. Wu HC, Yen RF, Shen YY, et al. Comparing whole body 18F-2-deoxyglucose positron emission tomography and technetium-99m methylene diphosphate bone scan to detect bone metastases in patients with renal cell carcinomas - a preliminary report. J Cancer Res Clin Oncol

2002;128:503–506. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12242515>.

40. Shuch B, Vourganti S, Ricketts CJ, et al. Defining early-onset kidney cancer: implications for germline and somatic mutation testing and clinical management. J Clin Oncol 2014;32:431–437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24378414>.

41. Berger A, Brandina R, Atalla MA, et al. Laparoscopic radical nephrectomy for renal cell carcinoma: oncological outcomes at 10 years or more. J Urol 2009;182:2172–2176. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19758651>.

42. Burgess NA, Koo BC, Calvert RC, et al. Randomized trial of laparoscopic v open nephrectomy. J Endourol 2007;21:610–613. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17638555>.

43. Chung SD, Huang KH, Lai MK, et al. Long-term follow-up of hand-assisted laparoscopic radical nephrectomy for organ-confined renal cell carcinoma. Urology 2007;69:652–655. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17445645>.

44. Gabr AH, Gdor Y, Strobe SA, et al. Patient and pathologic correlates with perioperative and long-term outcomes of laparoscopic radical nephrectomy. Urology 2009;74:635–640. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19616826>.

45. Hemal AK, Kumar A. A prospective comparison of laparoscopic and robotic radical nephrectomy for T1-2N0M0 renal cell carcinoma. World J Urol 2009;27:89–94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18704439>.

46. Hemal AK, Kumar A, Kumar R, et al. Laparoscopic versus open radical nephrectomy for large renal tumors: a long-term prospective comparison. J Urol 2007;177:862–866. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17296361>.

47. Luo JH, Zhou FJ, Xie D, et al. Analysis of long-term survival in patients with localized renal cell carcinoma: laparoscopic versus open radical



NCCN Guidelines Version 2.2026 Kidney Cancer

nephrectomy. World J Urol 2010;28:289–293. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19916010>.

48. Nambirajan T, Jeschke S, Al-Zahrani H, et al. Prospective, randomized controlled study: transperitoneal laparoscopic versus retroperitoneoscopic radical nephrectomy. Urology 2004;64:919–924. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/15533478>.

49. Dash A, Vickers AJ, Schachter LR, et al. Comparison of outcomes in elective partial vs radical nephrectomy for clear cell renal cell carcinoma of 4–7 cm. BJU Int 2006;97:939–945. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16643474>.

50. Lau WK, Blute ML, Weaver AL, et al. Matched comparison of radical nephrectomy vs nephron-sparing surgery in patients with unilateral renal cell carcinoma and a normal contralateral kidney. Mayo Clin Proc 2000;75:1236–1242. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/11126830>.

51. Lee CT, Katz J, Shi W, et al. Surgical management of renal tumors 4 cm. or less in a contemporary cohort. J Urol 2000;163:730–736. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10687966>.

52. Leibovich BC, Blute M, Cheville JC, et al. Nephron sparing surgery for appropriately selected renal cell carcinoma between 4 and 7 cm results in outcome similar to radical nephrectomy. J Urol 2004;171:1066–1070. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14767272>.

53. Zini L, Perrotte P, Capitanio U, et al. Radical versus partial nephrectomy: effect on overall and noncancer mortality. Cancer 2009;115:1465–1471. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19195042>.

54. Lee HJ, Liss MA, Derweesh IH. Outcomes of partial nephrectomy for clinical T1b and T2 renal tumors. Curr Opin Urol 2014;24:448–452. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24921904>.

55. Huang WC, Levey AS, Serio AM, et al. Chronic kidney disease after nephrectomy in patients with renal cortical tumours: a retrospective cohort

study. Lancet Oncol 2006;7:735–740. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16945768>.

56. Go AS, Chertow GM, Fan D, et al. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. N Engl J Med 2004;351:1296–1305. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/15385656>.

57. Thompson RH, Boorjian SA, Lohse CM, et al. Radical nephrectomy for pT1a renal masses may be associated with decreased overall survival compared with partial nephrectomy. J Urol 2008;179:468–471; discussion 472–463. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18076931>.

58. Weight CJ, Lieser G, Larson BT, et al. Partial nephrectomy is associated with improved overall survival compared to radical nephrectomy in patients with unanticipated benign renal tumours. Eur Urol 2010;58:293–298. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20546991>.

59. Weight CJ, Larson BT, Gao T, et al. Elective partial nephrectomy in patients with clinical T1b renal tumors is associated with improved overall survival. Urology 2010;76:631–637. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20451967>.

60. Kim SP, Thompson RH, Boorjian SA, et al. Comparative effectiveness for survival and renal function of partial and radical nephrectomy for localized renal tumors: a systematic review and meta-analysis. J Urol 2012;188:51–57. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/22591957>.

61. Thompson RH, Siddiqui S, Lohse CM, et al. Partial versus radical nephrectomy for 4 to 7 cm renal cortical tumors. J Urol 2009;182:2601–2606. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19836797>.

62. Hollingsworth JM, Miller DC, Dunn RL, et al. Surgical management of low-stage renal cell carcinoma: Technology does not supersede biology. Urology 2006;67:1175–1180. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16765177>.



NCCN Guidelines Version 2.2026 Kidney Cancer

63. Shuch B, Lam JS, Belldegrün AS. Open partial nephrectomy for the treatment of renal cell carcinoma. *Curr Urol Rep* 2006;7:31–38. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16480666>.
64. Chen DY, Uzzo RG. Optimal management of localized renal cell carcinoma: surgery, ablation, or active surveillance. *J Natl Compr Canc Netw* 2009;7:635–642; quiz 643. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19555585>.
65. Tan HJ, Norton EC, Ye Z, et al. Long-term survival following partial vs radical nephrectomy among older patients with early-stage kidney cancer. *JAMA* 2012;307:1629–1635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22511691>.
66. Gill IS, Kavoussi LR, Lane BR, et al. Comparison of 1,800 laparoscopic and open partial nephrectomies for single renal tumors. *J Urol* 2007;178:41–46. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17574056>.
67. Gong EM, Orvieto MA, Zorn KC, et al. Comparison of laparoscopic and open partial nephrectomy in clinical T1a renal tumors. *J Endourol* 2008;22:953–957. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18363510>.
68. Lane BR, Gill IS. 7-year oncological outcomes after laparoscopic and open partial nephrectomy. *J Urol* 2010;183:473–479. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20006866>.
69. Funahashi Y, Hattori R, Yamamoto T, et al. Ischemic renal damage after nephron-sparing surgery in patients with normal contralateral kidney. *Eur Urol* 2009;55:209–215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18706758>.
70. Blom JH, van Poppel H, Marechal JM, et al. Radical nephrectomy with and without lymph-node dissection: final results of European Organization for Research and Treatment of Cancer (EORTC) randomized phase 3 trial 30881. *Eur Urol* 2009;55:28–34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18848382>.
71. Blute ML, Leibovich BC, Cheville JC, et al. A protocol for performing extended lymph node dissection using primary tumor pathological features for patients treated with radical nephrectomy for clear cell renal cell carcinoma. *J Urol* 2004;172:465–469. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15247704>.
72. Capitanio U, Becker F, Blute ML, et al. Lymph node dissection in renal cell carcinoma. *Eur Urol* 2011;60:1212–1220. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21940096>.
73. Shi X, Feng D, Li D, et al. The role of lymph node dissection for non-metastatic renal cell carcinoma: An updated systematic review and meta-analysis. *Front Oncol* 2021;11:790381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35096589>.
74. Campi R, Sessa F, Di Maida F, et al. Templates of lymph node dissection for renal cell carcinoma: A systematic review of the literature. *Front Surg* 2018;5:76. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30619877>.
75. Kuczyk M, Munch T, Machtens S, et al. The need for routine adrenalectomy during surgical treatment for renal cell cancer: the Hannover experience. *BJU Int* 2002;89:517–522. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11942955>.
76. Kuczyk M, Wegener G, Jonas U. The therapeutic value of adrenalectomy in case of solitary metastatic spread originating from primary renal cell cancer. *Eur Urol* 2005;48:252–257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15936136>.
77. O'Malley RL, Godoy G, Kanofsky JA, Taneja SS. The necessity of adrenalectomy at the time of radical nephrectomy: a systematic review. *J Urol* 2009;181:2009–2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19286216>.
78. Lane BR, Tiong HY, Campbell SC, et al. Management of the adrenal gland during partial nephrectomy. *J Urol* 2009;181:2430–2436; discussion 2436–2437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19371896>.



NCCN Guidelines Version 2.2026

Kidney Cancer

79. Rais-Bahrami S, Guzzo TJ, Jarrett TW, et al. Incidentally discovered renal masses: oncological and perioperative outcomes in patients with delayed surgical intervention. *BJU Int* 2009;103:1355–1358. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19239459>.

80. Abouassaly R, Lane BR, Novick AC. Active surveillance of renal masses in elderly patients. *J Urol* 2008;180:505–508; discussion 508–509. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18550113>.

81. Lane BR, Abouassaly R, Gao T, et al. Active treatment of localized renal tumors may not impact overall survival in patients aged 75 years or older. *Cancer* 2010;116:3119–3126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20564627>.

82. Simmons MN, Weight CJ, Gill IS. Laparoscopic radical versus partial nephrectomy for tumors >4 cm: intermediate-term oncologic and functional outcomes. *Urology* 2009;73:1077–1082. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19394509>.

83. Peycelon M, Hupertan V, Comperat E, et al. Long-term outcomes after nephron sparing surgery for renal cell carcinoma larger than 4 cm. *J Urol* 2009;181:35–41. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19012929>.

84. Campbell S, Uzzo RG, Allaf ME, et al. Renal Mass and Localized Renal Cancer: AUA Guideline. *J Urol* 2017;198:520–529. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28479239>.

85. Pierorazio PM, Johnson MH, Patel HD, et al. Management of Renal Masses and Localized Renal Cancer: Systematic Review and Meta-Analysis. *J Urol* 2016;196:989–999. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27157369>.

86. Okhunov Z, Chamberlin J, Moreira DM, et al. Salvage percutaneous cryoablation for locally recurrent renal-cell carcinoma after primary cryoablation. *J Endourol* 2016;30:632–637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27009377>.

87. Chan VW, Abul A, Osman FH, et al. Ablative therapies versus partial nephrectomy for small renal masses - A systematic review and meta-analysis. *Int J Surg* 2022;97:106194. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34958968>.

88. Siva S, Bressel M, Sidhom M, et al. Stereotactic ablative body radiotherapy for primary kidney cancer (TROG 15.03 FASTRACK II): a non-randomised phase 2 trial. *Lancet Oncol* 2024;25:308–316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38423047>.

89. Siva S, Ali M, Correa RJM, et al. 5-year outcomes after stereotactic ablative body radiotherapy for primary renal cell carcinoma: an individual patient data meta-analysis from IROCK (the International Radiosurgery Consortium of the Kidney). *Lancet Oncol* 2022;23:1508–1516. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36400098>.

90. Hakimi K, Campbell SC, Nguyen MV, et al. PADRES: a phase 2 clinical trial of neoadjuvant axitinib for complex partial nephrectomy. *BJU Int* 2024;133:425–431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37916303>.

91. Bilen MA, Vo BT, Liu Y, et al. Neoadjuvant cabozantinib for locally advanced nonmetastatic clear cell renal cell carcinoma: a phase 2 trial. *Nat Cancer* 2025;6:432–444. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40016487>.

92. Huang J, Wang Y, Xu F, et al. Neoadjuvant toripalimab combined with axitinib in patients with locally advanced clear cell renal cell carcinoma: a single-arm, phase II trial. *J Immunother Cancer* 2024;12. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38862251>.

93. Seifert KE, Dinkel H, Huberth L, et al. Response assessment of patients with locally advanced renal cell carcinoma receiving prior systemic therapy. *BMC Cancer* 2026;26:129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41545991>.

94. Haas NB, Manola J, Uzzo RG, et al. Adjuvant sunitinib or sorafenib for high-risk, non-metastatic renal-cell carcinoma (ECOG-ACRIN E2805): a double-blind, placebo-controlled, randomised, phase 3 trial. *Lancet*



NCCN Guidelines Version 2.2026 Kidney Cancer

2016;387:2008–2016. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26969090>.

95. Motzer RJ, Haas NB, Donskov F, et al. Randomized phase III trial of adjuvant pazopanib versus placebo after nephrectomy in patients with localized or locally advanced renal cell carcinoma. *J Clin Oncol* 2017;35:3916–3923. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28902533>.

96. Gross-Goupil M, Kwon TG, Eto M, et al. Axitinib versus placebo as an adjuvant treatment of renal cell carcinoma: results from the phase III, randomized ATLAS trial. *Ann Oncol* 2018;29:2371–2378. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30346481>.

97. Ravaud A, Motzer RJ, Pandha HS, et al. Adjuvant sunitinib in high-risk renal-cell carcinoma after nephrectomy. *N Engl J Med* 2016;375:2246–2254. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27718781>.

98. Motzer RJ, Ravaud A, Patard JJ, et al. Adjuvant sunitinib for high-risk renal cell carcinoma after nephrectomy: Subgroup analyses and updated overall survival results. *Eur Urol* 2018;73:62–68. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28967554>.

99. Riaz IB, Siddiqi R, Islam M, et al. Adjuvant tyrosine kinase inhibitors in renal cell carcinoma: A concluded living systematic review and meta-analysis. *JCO Clin Cancer Inform* 2021;5:588–599. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/34043431>.

100. Laukhtina E, Quhal F, Mori K, et al. Adjuvant therapy with tyrosine kinase inhibitors for localized and locally advanced renal cell carcinoma: an updated systematic review and meta-analysis. *Urol Oncol* 2021;39:764–773. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/34400065>.

101. Choueiri TK, Tomczak P, Park SH, et al. Adjuvant pembrolizumab after nephrectomy in renal-cell carcinoma. *N Engl J Med* 2021;385:683–694. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34407342>.

102. Powles T, Tomczak P, Park SH, et al. Pembrolizumab versus placebo as post-nephrectomy adjuvant therapy for clear cell renal cell carcinoma (KEYNOTE-564): 30-month follow-up analysis of a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2022;23:1133–1144. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/36055304>.

103. Choueiri TK, Tomczak P, Park SH, et al. Overall survival with adjuvant pembrolizumab in renal-cell carcinoma. *N Engl J Med* 2024;390:1359–1371. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/38631003>.

104. Wei H, Miao J, Cui J, et al. The prognosis and clinicopathological features of different distant metastases patterns in renal cell carcinoma: analysis based on the SEER database. *Sci Rep* 2021;11:17822. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34497343>.

105. Eggener SE, Yossepowitch O, Pettus JA, et al. Renal cell carcinoma recurrence after nephrectomy for localized disease: predicting survival from time of recurrence. *J Clin Oncol* 2006;24:3101–3106. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16809736>.

106. Stewart SB, Thompson RH, Psutka SP, et al. Evaluation of the National Comprehensive Cancer Network and American Urological Association renal cell carcinoma surveillance guidelines. *J Clin Oncol* 2014;32:4059–4065. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25403213>.

107. Dabestani S, Beisland C, Stewart GD, et al. Long-term outcomes of follow-up for initially localised clear cell renal cell carcinoma: RECUR database analysis. *Eur Urol Focus* 2019;5:857–866. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29525381>.

108. Mucksavage P, Kutikov A, Magerfleisch L, et al. Comparison of radiographical imaging modalities for measuring the diameter of renal masses: is there a sizeable difference? *BJU Int* 2011;108:E232–236. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21348913>.



NCCN Guidelines Version 2.2026 Kidney Cancer

109. Herr HW. Partial nephrectomy for incidental renal cell carcinoma. *Br J Urol* 1994;74:431–433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7820418>.
110. Morgan WR, Zincke H. Progression and survival after renal-conserving surgery for renal cell carcinoma: experience in 104 patients and extended followup. *J Urol* 1990;144:852–857; discussion 857–858. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2398558>.
111. Lam JS, Shvarts O, Leppert JT, et al. Postoperative surveillance protocol for patients with localized and locally advanced renal cell carcinoma based on a validated prognostic nomogram and risk group stratification system. *J Urol* 2005;174:466–472; discussion 472; quiz 801. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16006866>.
112. Correa AF, Jegede OA, Haas NB, et al. Predicting Disease Recurrence, Early Progression, and Overall Survival Following Surgical Resection for High-risk Localized and Locally Advanced Renal Cell Carcinoma. *Eur Urol* 2021;80:20–31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33707112>.
113. Motzer RJ, Bacik J, Murphy BA, et al. Interferon-alfa as a comparative treatment for clinical trials of new therapies against advanced renal cell carcinoma. *J Clin Oncol* 2002;20:289–296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11773181>.
114. Heng DY, Xie W, Regan MM, et al. Prognostic factors for overall survival in patients with metastatic renal cell carcinoma treated with vascular endothelial growth factor-targeted agents: results from a large, multicenter study. *J Clin Oncol* 2009;27:5794–5799. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19826129>.
115. Mekhail TM, Abou-Jawde RM, Boucher G, et al. Validation and extension of the Memorial Sloan-Kettering prognostic factors model for survival in patients with previously untreated metastatic renal cell carcinoma. *J Clin Oncol* 2005;23:832–841. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15681528>.
116. Heng DY, Xie W, Regan MM, et al. External validation and comparison with other models of the International Metastatic Renal-Cell Carcinoma Database Consortium prognostic model: a population-based study. *Lancet Oncol* 2013;14:141–148. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23312463>.
117. Culp SH, Tannir NM, Abel EJ, et al. Can we better select patients with metastatic renal cell carcinoma for cytoreductive nephrectomy? *Cancer* 2010;116:3378–3388. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20564061>.
118. Choueiri TK, Xie W, Kollmannsberger C, et al. The impact of cytoreductive nephrectomy on survival of patients with metastatic renal cell carcinoma receiving vascular endothelial growth factor targeted therapy. *J Urol* 2011;185:60–66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21074201>.
119. Mejean A, Ravaud A, Thezenas S, et al. Sunitinib alone or after nephrectomy in metastatic renal-cell carcinoma. *N Engl J Med* 2018;379:417–427. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29860937>.
120. Motzer RJ, Russo P. Cytoreductive Nephrectomy - Patient Selection Is Key. *N Engl J Med* 2018;379:481–482. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29860908>.
121. Mejean A, Ravaud A, Thezenas S, et al. Sunitinib alone or after nephrectomy for patients with metastatic renal cell carcinoma: Is there still a role for cytoreductive nephrectomy? *Eur Urol* 2021;80:417–424. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34187771>.
122. Hannan R, Christensen M, Hammers H, et al. Phase II Trial of Stereotactic Ablative Radiation for Oligoprogressive Metastatic Kidney Cancer. *Eur Urol Oncol* 2022;5:216–224. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34986993>.
123. Tang C, Msaouel P, Hara K, et al. Definitive radiotherapy in lieu of systemic therapy for oligometastatic renal cell carcinoma: a single-arm,



NCCN Guidelines Version 2.2026 Kidney Cancer

single-centre, feasibility, phase 2 trial. *Lancet Oncol* 2021;22:1732–1739. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34717797>.

124. Palma DA, Olson R, Harrow S, et al. Stereotactic ablative radiotherapy versus standard of care palliative treatment in patients with oligometastatic cancers (SABR-COMET): a randomised, phase 2, open-label trial. *Lancet* 2019;393:2051–2058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30982687>.

125. Stinauer MA, Kavanagh BD, Schefter TE, et al. Stereotactic body radiation therapy for melanoma and renal cell carcinoma: impact of single fraction equivalent dose on local control. *Radiat Oncol* 2011;6:34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21477295>.

126. Zaorsky NG, Lehrer EJ, Kothari G, et al. Stereotactic ablative radiation therapy for oligometastatic renal cell carcinoma (SABR ORCA): a meta-analysis of 28 studies. *Eur Urol Oncol* 2019;2:515–523. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31302061>.

127. Ali M, Mooi J, Lawrentschuk N, et al. The role of stereotactic ablative body radiotherapy in renal cell carcinoma. *Eur Urol* 2022;82:613–622. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35843777>.

128. Siva S, Ellis RJ, Ponsky L, et al. Consensus statement from the International Radiosurgery Oncology Consortium for Kidney for primary renal cell carcinoma. *Future Oncol* 2016;12:637–645. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26837701>.

129. Siva S, Louie AV, Warner A, et al. Pooled analysis of stereotactic ablative radiotherapy for primary renal cell carcinoma: A report from the International Radiosurgery Oncology Consortium for Kidney (IROCK). *Cancer* 2018;124:934–942. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29266183>.

130. Meyer E, Pasquier D, Bernadou G, et al. Stereotactic radiation therapy in the strategy of treatment of metastatic renal cell carcinoma: A study of the Getug group. *Eur J Cancer* 2018;98:38–47. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29864737>.

131. Rini BI, Plimack ER, Stus V, et al. Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med* 2019;380:1116–1127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30779529>.

132. Powles T, Plimack ER, Soulieres D, et al. Pembrolizumab plus axitinib versus sunitinib monotherapy as first-line treatment of advanced renal cell carcinoma (KEYNOTE-426): extended follow-up from a randomised, open-label, phase 3 trial. *Lancet Oncol* 2020;21:1563–1573. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33284113>.

133. Plimack ER, Powles T, Stus V, et al. Pembrolizumab plus axitinib versus sunitinib as first-line treatment of advanced renal cell carcinoma: 43-month follow-up of the phase 3 KEYNOTE-426 study. *Eur Urol* 2023;84:449–454. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37500340>.

134. Rini BI, Plimack ER, Stus V, et al. Pembrolizumab plus axitinib versus sunitinib for advanced clear cell renal cell carcinoma: 5-year survival and biomarker analyses of the phase 3 KEYNOTE-426 trial. *Nat Med* 2025;31:3475–3484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40750932>.

135. Choueiri TK, Powles T, Burotto M, et al. Nivolumab plus cabozantinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med* 2021;384:829–841. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33657295>.

136. Motzer RJ, Powles T, Burotto M, et al. Nivolumab plus cabozantinib versus sunitinib in first-line treatment for advanced renal cell carcinoma (CheckMate 9ER): long-term follow-up results from an open-label, randomised, phase 3 trial. *Lancet Oncol* 2022;23:888–898. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35688173>.

137. Cella D, Motzer RJ, Suarez C, et al. Patient-reported outcomes with first-line nivolumab plus cabozantinib versus sunitinib in patients with advanced renal cell carcinoma treated in CheckMate 9ER: an open-label, randomised, phase 3 trial. *Lancet Oncol* 2022;23:292–303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35032437>.



NCCN Guidelines Version 2.2026

Kidney Cancer

138. Motzer RJ, Escudier B, Burotto M, et al. Final analysis of nivolumab plus cabozantinib for advanced renal cell carcinoma from the randomized phase III CheckMate 9ER trial. *Ann Oncol* 2026;37:33–43. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40998092>.
139. Motzer RJ, Tannir NM, McDermott DF, et al. Nivolumab plus ipilimumab versus sunitinib in advanced renal-cell carcinoma. *N Engl J Med* 2018;378:1277–1290. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29562145>.
140. Motzer RJ, McDermott DF, Escudier B, et al. Conditional survival and long-term efficacy with nivolumab plus ipilimumab versus sunitinib in patients with advanced renal cell carcinoma. *Cancer* 2022;128:2085–2097. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35383908>.
141. Tannir NM, Albiges L, McDermott DF, et al. Nivolumab plus ipilimumab versus sunitinib for first-line treatment of advanced renal cell carcinoma: extended 8-year follow-up results of efficacy and safety from the phase III CheckMate 214 trial. *Ann Oncol* 2024;35:1026–1038. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39098455>.
142. Motzer R, Alekseev B, Rha SY, et al. Lenvatinib plus pembrolizumab or everolimus for advanced renal cell carcinoma. *N Engl J Med* 2021;384:1289–1300. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33616314>.
143. Choueiri TK, Eto M, Motzer R, et al. Lenvatinib plus pembrolizumab versus sunitinib as first-line treatment of patients with advanced renal cell carcinoma (CLEAR): extended follow-up from the phase 3, randomised, open-label study. *Lancet Oncol* 2023;24:228–238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36858721>.
144. Motzer RJ, Porta C, Eto M, et al. Lenvatinib plus pembrolizumab versus sunitinib in first-line treatment of advanced renal cell carcinoma: Final prespecified overall survival analysis of CLEAR, a phase III study. *J Clin Oncol* 2024;42:1222–1228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38227898>.

145. Choueiri TK, Halabi S, Sanford BL, et al. Cabozantinib Versus Sunitinib As Initial Targeted Therapy for Patients With Metastatic Renal Cell Carcinoma of Poor or Intermediate Risk: The Alliance A031203 CABOSUN Trial. *J Clin Oncol* 2017;35:591–597. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28199818>.
146. Chen RC, Choueiri TK, Feuille M, et al. Quality-adjusted survival with first-line cabozantinib or sunitinib for advanced renal cell carcinoma in the CABOSUN randomized clinical trial (Alliance). *Cancer* 2020;126:5311–5318. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33022096>.
147. Motzer RJ, Penkov K, Haanen J, et al. Avelumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med* 2019;380:1103–1115. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30779531>.
148. Choueiri TK, Motzer RJ, Rini BI, et al. Updated efficacy results from the JAVELIN Renal 101 trial: first-line avelumab plus axitinib versus sunitinib in patients with advanced renal cell carcinoma. *Ann Oncol* 2020;31:1030–1039. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32339648>.
149. Choueiri TK, Penkov K, Uemura H, et al. Avelumab + axitinib versus sunitinib as first-line treatment for patients with advanced renal cell carcinoma: final analysis of the phase III JAVELIN Renal 101 trial. *Ann Oncol* 2025;36:387–392. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39706335>.
150. Haanen J, Larkin J, Choueiri TK, et al. Extended follow-up from JAVELIN Renal 101: subgroup analysis of avelumab plus axitinib versus sunitinib by the International Metastatic Renal Cell Carcinoma Database Consortium risk group in patients with advanced renal cell carcinoma. *ESMO Open* 2023;8:101210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37104931>.
151. Choueiri TK, Larkin J, Pal S, et al. Efficacy and correlative analyses of avelumab plus axitinib versus sunitinib in sarcomatoid renal cell carcinoma: post hoc analysis of a randomized clinical trial. *ESMO Open* 2021;6:100101. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33901870>.



NCCN Guidelines Version 2.2026 Kidney Cancer

152. Sternberg CN, Davis ID, Mardiak J, et al. Pazopanib in locally advanced or metastatic renal cell carcinoma: results of a randomized phase III trial. *J Clin Oncol* 2010;28:1061–1068. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20100962>.

153. Sternberg CN, Hawkins RE, Wagstaff J, et al. A randomised, double-blind phase III study of pazopanib in patients with advanced and/or metastatic renal cell carcinoma: final overall survival results and safety update. *Eur J Cancer* 2013;49:1287–1296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23321547>.

154. Motzer RJ, Hutson TE, Cella D, et al. Pazopanib versus sunitinib in metastatic renal-cell carcinoma. *N Engl J Med* 2013;369:722–731. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23964934>.

155. Motzer RJ, Hutson TE, McCann L, et al. Overall survival in renal-cell carcinoma with pazopanib versus sunitinib. *N Engl J Med* 2014;370:1769–1770. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24785224>.

156. Motzer RJ, Hutson TE, Tomczak P, et al. Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med* 2007;356:115–124. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17215529>.

157. Motzer RJ, Michaelson MD, Redman BG, et al. Activity of SU11248, a multitargeted inhibitor of vascular endothelial growth factor receptor and platelet-derived growth factor receptor, in patients with metastatic renal cell carcinoma. *J Clin Oncol* 2006;24:16–24. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16330672>.

158. Chow LQ, Eckhardt SG. Sunitinib: from rational design to clinical efficacy. *J Clin Oncol* 2007;25:884–896. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17327610>.

159. Faivre S, Delbaldo C, Vera K, et al. Safety, pharmacokinetic, and antitumor activity of SU11248, a novel oral multitarget tyrosine kinase inhibitor, in patients with cancer. *J Clin Oncol* 2006;24:25–35. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16314617>.

160. Motzer RJ, Hutson TE, Tomczak P, et al. Overall survival and updated results for sunitinib compared with interferon alfa in patients with metastatic renal cell carcinoma. *J Clin Oncol* 2009;27:3584–3590. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19487381>.

161. Rini BI, Dorff TB, Elson P, et al. Active surveillance in metastatic renal-cell carcinoma: a prospective, phase 2 trial. *Lancet Oncol* 2016;17:1317–1324. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27498080>.

162. Harrison MR, Costello BA, Bhavsar NA, et al. Active surveillance of metastatic renal cell carcinoma: Results from a prospective observational study (MaRCC). *Cancer* 2021;127:2204–2212. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33765337>.

163. Bex A. Increasing the evidence for surveillance of metastatic renal cancer. *Cancer* 2021;127:2184–2186. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33765328>.

164. Rini BI, Escudier B, Tomczak P, et al. Comparative effectiveness of axitinib versus sorafenib in advanced renal cell carcinoma (AXIS): a randomised phase 3 trial. *Lancet* 2011;378:1931–1939. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22056247>.

165. Hutson TE, Lesovoy V, Al-Shukri S, et al. Axitinib versus sorafenib as first-line therapy in patients with metastatic renal-cell carcinoma: a randomised open-label phase 3 trial. *Lancet Oncol* 2013;14:1287–1294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24206640>.

166. Pal SK, Albiges L, Tomczak P, et al. Atezolizumab plus cabozantinib versus cabozantinib monotherapy for patients with renal cell carcinoma after progression with previous immune checkpoint inhibitor treatment (CONTACT-03): a multicentre, randomised, open-label, phase 3 trial. *Lancet* 2023;402:185–195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37290461>.

167. Choueiri TK, Albiges L, Barthelemy P, et al. Tivozanib plus nivolumab versus tivozanib monotherapy in patients with renal cell carcinoma following an immune checkpoint inhibitor: results of the phase 3 TiNivo-2



NCCN Guidelines Version 2.2026

Kidney Cancer

Study. Lancet 2024;404:1309–1320. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/39284329>.

168. Choueiri TK, Escudier B, Powles T, et al. Cabozantinib versus everolimus in advanced renal-cell carcinoma. N Engl J Med 2015;373:1814–1823. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26406150>.

169. Choueiri TK, Escudier B, Powles T, et al. Cabozantinib versus everolimus in advanced renal cell carcinoma (METEOR): final results from a randomised, open-label, phase 3 trial. Lancet Oncol 2016;17:917–927. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27279544>.

170. Motzer RJ, Escudier B, Powles T, et al. Long-term follow-up of overall survival for cabozantinib versus everolimus in advanced renal cell carcinoma. Br J Cancer 2018;118:1176–1178. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29576624>.

171. Obeng-Kusi M, Kreutzfeldt JJ, Estrada-Mendizabal RJ, et al. Network meta-analysis of second line and beyond treatment options in metastatic clear cell renal cell carcinoma. Urol Oncol 2024;42:32 e31–32 e38. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38216444>.

172. Tannir NM, Agarwal N, Porta C, et al. Efficacy and safety of telaglenastat plus cabozantinib vs placebo plus cabozantinib in patients with advanced renal cell carcinoma: The CANTATA randomized clinical trial. JAMA Oncol 2022;8:1411–1418. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/36048457>.

173. Motzer RJ, Hutson TE, Glen H, et al. Lenvatinib, everolimus, and the combination in patients with metastatic renal cell carcinoma: a randomised, phase 2, open-label, multicentre trial. Lancet Oncol 2015;16:1473–1482. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26482279>.

174. Motzer RJ, Hutson TE, Ren M, et al. Independent assessment of lenvatinib plus everolimus in patients with metastatic renal cell carcinoma. Lancet Oncol 2016;17:e4–5. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26758760>.

175. Wiele AJ, Bathala TK, Hahn AW, et al. Lenvatinib with or Without Everolimus in Patients with Metastatic Renal Cell Carcinoma After Immune Checkpoint Inhibitors and Vascular Endothelial Growth Factor Receptor-Tyrosine Kinase Inhibitor Therapies. Oncologist 2021;26:476–482. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33792094>.

176. Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus everolimus in advanced renal-cell carcinoma. N Engl J Med 2015;373:1803–1813. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26406148>.

177. Motzer RJ, Escudier B, George S, et al. Nivolumab versus everolimus in patients with advanced renal cell carcinoma: Updated results with long-term follow-up of the randomized, open-label, phase 3 CheckMate 025 trial. Cancer 2020;126:4156–4167. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/32673417>.

178. Motzer RJ, Escudier B, Tomczak P, et al. Axitinib versus sorafenib as second-line treatment for advanced renal cell carcinoma: overall survival analysis and updated results from a randomised phase 3 trial. Lancet Oncol 2013;14:552–562. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23598172>.

179. Dizman N, Austin M, Considine B, et al. Outcomes with combination pembrolizumab and axitinib in second and further line treatment of metastatic renal cell carcinoma. Clin Genitourin Cancer 2023;21:221–229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36681606>.

180. Apolo AB, Nadal R, Girardi DM, et al. Phase I study of cabozantinib and nivolumab alone or with ipilimumab for advanced or metastatic urothelial carcinoma and other genitourinary tumors. J Clin Oncol 2020;38:3672–3684. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/32915679>.

181. Apolo AB, Girardi DM, Niglio SA, et al. Final results from a phase I trial and expansion cohorts of cabozantinib and nivolumab alone or with ipilimumab for advanced/metastatic genitourinary tumors. J Clin Oncol 2024;42:3033–3046. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/38954785>.



NCCN Guidelines Version 2.2026 Kidney Cancer

182. Hammers HJ, Plimack ER, Infante JR, et al. Safety and efficacy of nivolumab in combination with ipilimumab in metastatic renal cell carcinoma: The CheckMate 016 study. *J Clin Oncol* 2017;35:3851–3858. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28678668>.

183. Grimm MO, Esteban E, Barthelemy P, et al. Tailored immunotherapy approach with nivolumab with or without nivolumab plus ipilimumab as immunotherapeutic boost in patients with metastatic renal cell carcinoma (TITAN-RCC): a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2023;24:1252–1265. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37844597>.

184. Choueiri TK, Kluger H, George S, et al. FRACTION-RCC: nivolumab plus ipilimumab for advanced renal cell carcinoma after progression on immuno-oncology therapy. *J Immunother Cancer* 2022;10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36328377>.

185. Jonasch E, Bauer TM, Papadopoulos KP, et al. Phase I LITESPARK-001 study of belzutifan for advanced solid tumors: Extended 41-month follow-up in the clear cell renal cell carcinoma cohort. *Eur J Cancer* 2023;196:113434. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38008031>.

186. Choueiri TK, Powles T, Peltola K, et al. Belzutifan versus everolimus for advanced renal-cell carcinoma. *N Engl J Med* 2024;391:710–721. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39167807>.

187. Lee CH, Shah AY, Rasco D, et al. Lenvatinib plus pembrolizumab in patients with either treatment-naïve or previously treated metastatic renal cell carcinoma (Study 111/KEYNOTE-146): a phase 1b/2 study. *Lancet Oncol* 2021;22:946–958. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34143969>.

188. Lee C-H, Yogesh Shah A, Rao A, et al. Final database lock results of the phase 2 cohort of lenvatinib + pembrolizumab for progressive disease after a PD-1/PD-L1-containing therapy in metastatic clear cell renal cell carcinoma. *The Oncologist* 2023;28:S3–S4. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10445556/>.

189. Rini BI, Pal SK, Escudier BJ, et al. Tivozanib versus sorafenib in patients with advanced renal cell carcinoma (TIVO-3): a phase 3, multicentre, randomised, controlled, open-label study. *Lancet Oncol* 2020;21:95–104. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31810797>.

190. Beckermann KE, Asnis-Alibozek AG, Atkins MB, et al. Long-term survival in patients with relapsed/refractory advanced renal cell carcinoma treated with tivozanib: Analysis of the Phase III TIVO-3 trial. *Oncologist* 2024;29:254–262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38262444>.

191. Zugman M, McDermott DF, Escudier BJ, et al. Updated overall survival in patients with prior checkpoint inhibitor therapy in the phase III TIVO-3 study. *Oncologist* 2025;30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39912344>.

192. Szarek M, Needle MN, Rini BI, et al. Q-TWiST analysis of tivozanib versus sorafenib in patients with advanced renal cell carcinoma in the TIVO-3 study. *Clin Genitourin Cancer* 2021;19:468 e461–468 e465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33980467>.

193. Hainsworth JD, Rubin MS, Arrowsmith ER, et al. Pazopanib as second-line treatment after sunitinib or bevacizumab in patients with advanced renal cell carcinoma: a Sarah Cannon Oncology Research Consortium Phase II Trial. *Clin Genitourin Cancer* 2013;11:270–275. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23665131>.

194. Motzer RJ, Rini BI, Bukowski RM, et al. Sunitinib in patients with metastatic renal cell carcinoma. *JAMA* 2006;295:2516–2524. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16757724>.

195. Motzer RJ, Escudier B, Oudard S, et al. Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial. *Lancet* 2008;372:449–456. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18653228>.

196. Motzer RJ, Escudier B, Oudard S, et al. Phase 3 trial of everolimus for metastatic renal cell carcinoma : final results and analysis of prognostic



NCCN Guidelines Version 2.2026

Kidney Cancer

factors. Cancer 2010;116:4256–4265. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20549832>.

197. Yang JC, Haworth L, Sherry RM, et al. A randomized trial of bevacizumab, an anti-vascular endothelial growth factor antibody, for metastatic renal cancer. N Engl J Med 2003;349:427–434. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/12890841>.

198. de Velasco G, McKay RR, Lin X, et al. Comprehensive analysis of survival outcomes in non-clear cell renal cell carcinoma patients treated in clinical trials. Clin Genitourin Cancer 2017;15:652–660 e651. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28410911>.

199. Pal SK, Tangen C, Thompson IM, Jr., et al. A comparison of sunitinib with cabozantinib, crizotinib, and savolitinib for treatment of advanced papillary renal cell carcinoma: a randomised, open-label, phase 2 trial. Lancet 2021;397:695–703. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/33592176>.

200. Barata P, Tangen C, Plets M, et al. Final Overall Survival Analysis of S1500: A Randomized, Phase II Study Comparing Sunitinib With Cabozantinib, Crizotinib, and Savolitinib in Advanced Papillary Renal Cell Carcinoma. J Clin Oncol 2024;42:3911–3916. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/39255440>.

201. Lee CH, Voss MH, Carlo MI, et al. Phase II trial of cabozantinib plus nivolumab in patients with non-clear-cell renal cell carcinoma and genomic correlates. J Clin Oncol 2022;40:2333–2341. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/35298296>.

202. Fitzgerald KN, Lee CH, Voss MH, et al. Cabozantinib plus nivolumab in patients with non-clear cell renal cell carcinoma: Updated results from a phase 2 trial. Eur Urol 2024;86:90–94. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/38782695>.

203. Albiges L, Gurney H, Atduiev V, et al. Pembrolizumab plus lenvatinib as first-line therapy for advanced non-clear-cell renal cell carcinoma (KEYNOTE-B61): a single-arm, multicentre, phase 2 trial. Lancet Oncol

2023;24:881–891. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/37451291>.

204. Voss MH, Gurney H, Atduiev V, et al. First-line Pembrolizumab Plus Lenvatinib for Advanced Non-clear-cell Renal Cell Carcinoma: Updated Results from the Phase 2 KEYNOTE-B61 Trial. Eur Urol 2025;88:614–624. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40707309>.

205. Menko FH, Maher ER, Schmidt LS, et al. Hereditary leiomyomatosis and renal cell cancer (HLRCC): renal cancer risk, surveillance and treatment. Fam Cancer 2014;13:637–644. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25012257>.

206. Srinivasan R, Gurram S, Singer EA, et al. Bevacizumab and erlotinib in hereditary and sporadic papillary kidney cancer. N Engl J Med 2025;392:2346–2356. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/40532152>.

207. Hutson TE, Michaelson MD, Kuzel TM, et al. A single-arm, multicenter, phase 2 study of lenvatinib plus everolimus in patients with advanced non-clear cell renal cell carcinoma. Eur Urol 2021;80:162–170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33867192>.

208. Koshkin VS, Barata PC, Zhang T, et al. Clinical activity of nivolumab in patients with non-clear cell renal cell carcinoma. J Immunother Cancer 2018;6:9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29378660>.

209. McKay RR, Bosse D, Xie W, et al. The clinical activity of PD-1/PD-L1 inhibitors in metastatic non-clear cell renal cell carcinoma. Cancer Immunol Res 2018;6:758–765. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29748390>.

210. McDermott DF, Lee JL, Ziobro M, et al. Open-label, single-arm, phase II study of pembrolizumab monotherapy as first-line therapy in patients with advanced non-clear cell renal cell carcinoma. J Clin Oncol 2021;39:1029–1039. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/33529058>.



NCCN Guidelines Version 2.2026 Kidney Cancer

211. Armstrong AJ, Halabi S, Eisen T, et al. Everolimus versus sunitinib for patients with metastatic non-clear cell renal cell carcinoma (ASPEN): a multicentre, open-label, randomised phase 2 trial. *Lancet Oncol* 2016;17:378–388. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26794930>.

212. Tannir NM, Jonasch E, Albiges L, et al. Everolimus versus sunitinib prospective evaluation in metastatic non-clear cell renal cell carcinoma (ESPN): A randomized multicenter phase 2 trial. *Eur Urol* 2016;69:866–874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26626617>.

213. Ciccarese C, Iacovelli R, Brunelli M, et al. Addressing the best treatment for non-clear cell renal cell carcinoma: A meta-analysis of randomised clinical trials comparing VEGFR-TKIs versus mTORi-targeted therapies. *Eur J Cancer* 2017;83:237–246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28756136>.

214. Park I, Lee SH, Lee JL. A multicenter phase II trial of axitinib in patients with recurrent or metastatic non-clear-cell renal cell carcinoma who had failed prior treatment with temsirolimus. *Clin Genitourin Cancer* 2018;16:e997–e1002. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29903415>.

215. Blank CU, Bono P, Larkin JMG, et al. Safety and efficacy of everolimus in patients with non-clear cell renal cell carcinoma refractory to VEGF-targeted therapy: Subgroup analysis of REACT [abstract]. *J Clin Oncol* 2012;30 (5_suppl):Abstract 402. Available at: http://ascopubs.org/doi/abs/10.1200/jco.2012.30.5_suppl.402.

216. Koh Y, Lim HY, Ahn JH, et al. Phase II trial of everolimus for the treatment of nonclear-cell renal cell carcinoma. *Ann Oncol* 2013;24:1026–1031. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23180114>.

217. Escudier B, Molinie V, Bracarda S, et al. Open-label phase 2 trial of first-line everolimus monotherapy in patients with papillary metastatic renal cell carcinoma: RAPTOR final analysis. *Eur J Cancer* 2016;69:226–235. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27680407>.

218. Voss MH, Molina AM, Chen YB, et al. Phase II trial and correlative genomic analysis of everolimus plus bevacizumab in advanced non-clear cell renal cell carcinoma. *J Clin Oncol* 2016;34:3846–3853. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27601542>.

219. Tykodi SS, Gordan LN, Alter RS, et al. Safety and efficacy of nivolumab plus ipilimumab in patients with advanced non-clear cell renal cell carcinoma: results from the phase 3b/4 CheckMate 920 trial. *J Immunother Cancer* 2022;10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35210307>.

220. Bergmann L, Albiges L, Ahrens M, et al. Prospective randomized phase-II trial of ipilimumab/nivolumab versus standard of care in non-clear cell renal cell cancer - results of the SUNNIFORECAST trial. *Ann Oncol* 2025;36:796–806. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40180121>.

221. Hakimi AA, Koi PT, Milhoua PM, et al. Renal medullary carcinoma: the Bronx experience. *Urology* 2007;70:878–882. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18068443>.

222. Watanabe IC, Billis A, Guimaraes MS, et al. Renal medullary carcinoma: report of seven cases from Brazil. *Mod Pathol* 2007;20:914–920. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17643096>.

223. Shah AY, Karam JA, Malouf GG, et al. Management and outcomes of patients with renal medullary carcinoma: a multicentre collaborative study. *BJU Int* 2017;120:782–792. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27860149>.

224. Srigley JR, Delahunt B. Uncommon and recently described renal carcinomas. *Mod Pathol* 2009;22 Suppl 2:S2–S23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19494850>.

225. Tokuda N, Naito S, Matsuzaki O, et al. Collecting duct (Bellini duct) renal cell carcinoma: a nationwide survey in Japan. *J Urol* 2006;176:40–43; discussion 43. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16753362>.



NCCN Guidelines Version 2.2026 Kidney Cancer

226. Karakiewicz PI, Trinh QD, Rioux-Leclercq N, et al. Collecting duct renal cell carcinoma: a matched analysis of 41 cases. *Eur Urol* 2007;52:1140–1145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17336449>.

227. Gupta R, Billis A, Shah RB, et al. Carcinoma of the collecting ducts of Bellini and renal medullary carcinoma: clinicopathologic analysis of 52 cases of rare aggressive subtypes of renal cell carcinoma with a focus on their interrelationship. *Am J Surg Pathol* 2012;36:1265–1278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22895263>.

228. Oudard S, Banu E, Vieillefond A, et al. Prospective multicenter phase II study of gemcitabine plus platinum salt for metastatic collecting duct carcinoma: results of a GETUG (Groupe d'Etudes des Tumeurs Uro-Genitales) study. *J Urol* 2007;177:1698–1702. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17437788>.

229. Wilson NR, Wiele AJ, Surasi DS, et al. Efficacy and safety of gemcitabine plus doxorubicin in patients with renal medullary carcinoma. *Clin Genitourin Cancer* 2021;19:e401–e408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34625389>.

230. Roubaud G, Gross-Goupil M, Wallerand H, et al. Combination of gemcitabine and doxorubicin in rapidly progressive metastatic renal cell carcinoma and/or sarcomatoid renal cell carcinoma. *Oncology* 2011;80:214–218. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21720184>.

231. Sahgal A, Myrehaug SD, Siva S, et al. Stereotactic body radiotherapy versus conventional external beam radiotherapy in patients with painful spinal metastases: an open-label, multicentre, randomised, controlled, phase 2/3 trial. *Lancet Oncol* 2021;22:1023–1033. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34126044>.

232. Taunk NK, Spratt DE, Bilsky M, Yamada Y. Spine radiosurgery in the management of renal cell carcinoma metastases. *J Natl Compr Canc Netw* 2015;13:801–809; quiz 809. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26085394>.

233. Kwon YS, Hsu E, Stein M, et al. Extracranial palliative radiation therapy for renal cell carcinoma. *Adv Radiat Oncol* 2025;10:101870. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40979684>.

234. Dudani S, de Velasco G, Wells JC, et al. Evaluation of clear cell, papillary, and chromophobe renal cell carcinoma metastasis sites and association with survival. *JAMA Netw Open* 2021;4:e2021869. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33475752>.

235. Mulvenna P, Nankivell M, Barton R, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): results from a phase 3, non-inferiority, randomised trial. *Lancet* 2016;388:2004–2014. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27604504>.

236. Huang JF, Shen J, Li X, et al. Incidence of patients with bone metastases at diagnosis of solid tumors in adults: a large population-based study. *Ann Transl Med* 2020;8:482. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32395526>.

237. Hunter GK, Balagamwala EH, Koyfman SA, et al. The efficacy of external beam radiotherapy and stereotactic body radiotherapy for painful spinal metastases from renal cell carcinoma. *Pract Radiat Oncol* 2012;2:e95–e100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24674192>.

238. Zelefsky MJ, Greco C, Motzer R, et al. Tumor control outcomes after hypofractionated and single-dose stereotactic image-guided intensity-modulated radiotherapy for extracranial metastases from renal cell carcinoma. *Int J Radiat Oncol Biol Phys* 2012;82:1744–1748. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21596489>.

239. Lipton A, Zheng M, Seaman J. Zoledronic acid delays the onset of skeletal-related events and progression of skeletal disease in patients with advanced renal cell carcinoma. *Cancer* 2003;98:962–969. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12942563>.



NCCN Guidelines Version 2.2026 Kidney Cancer

240. Rosen LS, Gordon D, Tchekmedyian NS, et al. Long-term efficacy and safety of zoledronic acid in the treatment of skeletal metastases in patients with nonsmall cell lung carcinoma and other solid tumors: a randomized, Phase III, double-blind, placebo-controlled trial. *Cancer* 2004;100:2613–2621. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15197804>.

241. Henry DH, Costa L, Goldwasser F, et al. Randomized, double-blind study of denosumab versus zoledronic acid in the treatment of bone metastases in patients with advanced cancer (excluding breast and prostate cancer) or multiple myeloma. *J Clin Oncol* 2011;29:1125–1132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21343556>.

242. Henry D, Vadhan-Raj S, Hirsh V, et al. Delaying skeletal-related events in a randomized phase 3 study of denosumab versus zoledronic acid in patients with advanced cancer: an analysis of data from patients with solid tumors. *Support Care Cancer* 2014;22:679–687. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24162260>.

243. Maher ER. Hereditary renal cell carcinoma syndromes: diagnosis, surveillance and management. *World J Urol* 2018;36:1891–1898. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29680948>.

244. Rednam SP, Erez A, Druker H, et al. Von hippel-lindau and hereditary pheochromocytoma/paraganglioma syndromes: Clinical features, genetics, and surveillance recommendations in childhood. *Clin Cancer Res* 2017;23:e68–e75. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28620007>.

245. Tufton N, Sahdev A, Akker SA. Radiological surveillance screening in asymptomatic succinate dehydrogenase mutation carriers. *J Endocr Soc* 2017;1:897–907. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29264540>.

246. Eijkelenkamp K, Osinga TE, de Jong MM, et al. Calculating the optimal surveillance for head and neck paraganglioma in SDHB-mutation carriers. *Fam Cancer* 2017;16:123–130. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27573198>.

247. Louise MBM, Smerdel M, Borgwadt L, et al. von Hippel-Lindau disease: Updated guideline for diagnosis and surveillance. *Eur J Med Genet* 2022;65:104538. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35709961>.

248. Northrup H, Aronow ME, Bebin EM, et al. Updated international tuberous sclerosis complex diagnostic criteria and surveillance and management recommendations. *Pediatr Neurol* 2021;123:50–66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34399110>.

249. Menko FH, van Steensel MA, Giraud S, et al. Birt-Hogg-Dube syndrome: diagnosis and management. *Lancet Oncol* 2009;10:1199–1206. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19959076>.

250. Star P, Goodwin A, Kapoor R, et al. Germline BAP1-positive patients: the dilemmas of cancer surveillance and a proposed interdisciplinary consensus monitoring strategy. *Eur J Cancer* 2018;92:48–53. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29413689>.

251. Ornstein DK, Lubensky IA, Venzon D, et al. Prevalence of microscopic tumors in normal appearing renal parenchyma of patients with hereditary papillary renal cancer. *J Urol* 2000;163:431–433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10647647>.

252. Pavlovich CP, Grubb RL, 3rd, Hurley K, et al. Evaluation and management of renal tumors in the Birt-Hogg-Dube syndrome. *J Urol* 2005;173:1482–1486. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15821464>.

253. Shuch B, Singer EA, Bratslavsky G. The surgical approach to multifocal renal cancers: hereditary syndromes, ipsilateral multifocality, and bilateral tumors. *Urol Clin North Am* 2012;39:133–148, v. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22487757>.

254. Gill AJ, Hes O, Papathomas T, et al. Succinate dehydrogenase (SDH)-deficient renal carcinoma: a morphologically distinct entity: a clinicopathologic series of 36 tumors from 27 patients. *Am J Surg Pathol* 2014;38:1588–1602. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25025441>.



NCCN Guidelines Version 2.2026 Kidney Cancer

255. Singer EA, Vourganti S, Lin KY, et al. Outcomes of patients with surgically treated bilateral renal masses and a minimum of 10 years of followup. J Urol 2012;188:2084–2088. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23083858>.

256. Bissler JJ, Kingswood JC, Radzikowska E, et al. Everolimus for angiomyolipoma associated with tuberous sclerosis complex or sporadic lymphangioleiomyomatosis (EXIST-2): a multicentre, randomised, double-blind, placebo-controlled trial. Lancet 2013;381:817–824. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23312829>.

257. Bissler JJ, Kingswood JC, Radzikowska E, et al. Everolimus long-term use in patients with tuberous sclerosis complex: Four-year update of the EXIST-2 study. PLoS One 2017;12:e0180939. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28792952>.

258. Bissler JJ, McCormack FX, Young LR, et al. Sirolimus for angiomyolipoma in tuberous sclerosis complex or lymphangioleiomyomatosis. N Engl J Med 2008;358:140–151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18184959>.

259. Jonasch E, Donskov F, Iliopoulos O, et al. Belzutifan for renal cell carcinoma in von Hippel-Lindau disease. N Engl J Med 2021;385:2036–2046. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34818478>.

260. Srinivasan R, Iliopoulos O, Beckermann KE, et al. Belzutifan for von Hippel-Lindau disease-associated renal cell carcinoma and other neoplasms (LITESPARK-004): 50 months follow-up from a single-arm, phase 2 study. Lancet Oncol 2025;26:571–582. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40228516>.

261. Jonasch E, McCutcheon IE, Gombos DS, et al. Pazopanib in patients with von Hippel-Lindau disease: a single-arm, single-centre, phase 2 trial. Lancet Oncol 2018;19:1351–1359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30236511>.