



National Comprehensive
Cancer Network®


NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

Version 2.2026 — June 16, 2026

NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.





NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

***Aaron T. Gerds, MD, MS/Chair ‡ † P**
Case Comprehensive Cancer Center/University
Hospitals Seidman Cancer Center and
Cleveland Clinic Taussig Cancer Institute

***Jason Gotlib, MD, MS/Vice-Chair ‡**
Stanford Cancer Institute

Peter Abdelmessieh, DO, MSc ξ
Fox Chase Cancer Center

Haris Ali, MD ‡ ξ
City of Hope National Medical Center

Mariana Castells, MD, PhD P σ
Dana-Farber/Brigham and
Women's Cancer Center

Brian J. Chernak, MD ‡ †
Memorial Sloan Kettering Cancer Center

Manuel R. Espinoza-Gutarra, MD ‡ ξ
O'Neal Comprehensive Cancer Center at UAB

Ruth Fein Revell, BS ¥
Patient advocate

Steven Green, MD †
Roswell Park Comprehensive Cancer Center

Krishna Gundabolu, MBBS, MS ‡
Fred & Pamela Buffett Cancer Center

Elizabeth Hexner, MD, MS ‡ ξ
Abramson Cancer Center
at the University of Pennsylvania

NCCN

Mary Anne Bergman
Cindy Hochstetler, PhD

[NCCN Guidelines Panel Disclosures](#)

Tania Jain, MBBS †
Johns Hopkins Kimmel Cancer Center

Catriona H. Jamieson, MD, PhD ‡
UC San Diego Moores Cancer Center

Irum Khan, MD ‡ †
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Andrew T. Kuykendall, MD ‡ P
Moffitt Cancer Center

Yazan F. Madanat, MD ‡ P
UT Southwestern Simmons
Comprehensive Cancer Center

Lucia Masarova, MD ‡
The University of Texas
MD Anderson Cancer Center

Brandon McMahon, MD ‡
University of Colorado Cancer Center

Sanjay R. Mohan, MD, MSCI ‡
Vanderbilt-Ingram Cancer Center

Kalyan V. Nadiminti, MD ‡
University of Wisconsin Carbone Cancer Center

Stephen Oh, MD, PhD ‡
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

Jeanne Palmer ‡
Mayo Clinic Comprehensive Cancer Center

Ami B. Patel, MD ‡
Huntsman Cancer Institute
at the University of Utah

Anand A. Patel, MD ‡
The UChicago Medicine
Comprehensive Cancer Center

Nikolai Podoltsev, MD, PhD ‡
Yale Cancer Center/Smilow Cancer Hospital

Lindsay Rein, MD ‡
Duke Cancer Institute

Rachel Salit, MD ‡
Fred Hutchinson Cancer Center

Kelly Schoenbeck, MD ‡
UCSF Helen Diller
Family Comprehensive Cancer Center

Mary Sehl, MD †
UCLA Jonsson Comprehensive Cancer Center

Moshe Talpaz, MD †
University of Michigan Rogel Cancer Center

Martha Wadleigh, MD ‡ †
Dana-Farber/Brigham and
Women's Cancer Center

Sarah A. Wall, MD, MPH ‡
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

σ Allergy/Immunology
‡ Hematology/Hematologic oncology
P Internal medicine
† Medical oncology
≠ Pathology
¥ Patient advocacy
ξ Transplantation
* Discussion Writing Committee Member



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

[NCCN Myeloproliferative Neoplasms Panel Members](#)

[NCCN Evidence Blocks Definitions \(EB-DEF\)](#)

Myeloproliferative Neoplasms:

- [Workup \(MPN-1\)](#)
- [Diagnosis and Risk Stratification \(MPN-2\)](#)

Myelofibrosis:

- [Treatment for Lower-Risk Myelofibrosis \(MF-1\)](#)
- [Treatment for Higher-Risk Myelofibrosis \(MF-2\)](#)
- [Management of MF-Associated Anemia \(MF-3\)](#)
- [Risk Stratification for Patients with Myelofibrosis \(MF-A\)](#)
- [2013 IWG-MRT AND ELN Response Criteria for MF \(MF-B\)](#)
- [Special Considerations for Allogeneic Hematopoietic Cell Transplant \(MF-C\)](#)

Polycythemia Vera:

- [Treatment for Low-Risk Polycythemia Vera \(PV-1\)](#)
- [Treatment for High-Risk Polycythemia Vera \(PV-2\)](#)
- [2013 IWG-MRT and ELN Response Criteria for PV \(PV-A\)](#)
- [Overall Survival Risk Stratification for Patients with Polycythemia Vera \(PV-B\)](#)

Essential Thrombocythemia:

- [Treatment for Very-Low-Risk or Low-Risk or Intermediate-Risk Essential Thrombocythemia \(ET-1\)](#)
- [Treatment for High-Risk Essential Thrombocythemia \(ET-2\)](#)
- [2013 IWG-MRT AND ELN Response Criteria for Essential Thrombocythemia \(ET-A\)](#)
- [Overall Survival Risk Stratification for Patients with Essential Thrombocythemia \(ET-B\)](#)

Myeloproliferative Neoplasms:

- [Accelerated/Blast Phase MPN \(MPN-AP/BP-1\)](#)
- [WHO and International Consensus Classification \(ICC\) Diagnostic Criteria for Primary Myelofibrosis \(MPN-A 1 of 2\)](#)
- [Grading of Myelofibrosis \(MPN-A 2 of 2\)](#)
- [WHO and ICC Diagnostic Criteria for Polycythemia Vera and Post-PV Myelofibrosis \(MPN-B\)](#)
- [WHO and ICC Diagnostic Criteria for Essential Thrombocythemia and Post-ET Myelofibrosis \(MPN-C\)](#)

[NCCN Guidelines for Patients®](#) [NCCN Guidelines Navigator™](#)

- [Prognostic Significance of Mutations in Myelofibrosis \(MPN-D\)](#)
- [Assessment of Symptom Burden \(MPN-E 1 of 2\)](#)
- [Myeloproliferative Neoplasm Symptom Assessment Form: Total Symptom Score \(MPN-SAF TSS\) \(MPN-E 2 of 2\)](#)
- [Supportive Care for Patients with MPN \(MPN-F\)](#)
- [Special Considerations for the Use of JAK Inhibitors \(MPN-G 1 of 2\)](#)
- [Special Considerations for the Use of Interferons and Anagrelide \(MPN-G 2 of 2\)](#)
- [Special Considerations in the Treatment of Myeloproliferative Neoplasms \(MPN-H\)](#)
- [Definition of Resistance/Intolerance to Hydroxyurea \(MPN-I\)](#)
- [Evidence Blocks](#)
- [Abbreviations \(ABBR-1\)](#)

Myelodysplastic/Myeloproliferative Neoplasms (MDS/MPN): [NCCN Guidelines for Myelodysplastic Syndromes](#)

Find an NCCN Member Institution:

<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus:

All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference:

All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS

5					
4					
3					
2					
1					

E S Q C A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

Example Evidence Block

5					
4					
3					
2					
1					

E S Q C A

E = 4
S = 4
Q = 3
C = 4
A = 3

Efficacy of Regimen/Agent

5	Highly effective: Cure likely and often provides long-term survival advantage
4	Very effective: Cure unlikely but sometimes provides long-term survival advantage
3	Moderately effective: Modest impact on survival, but often provides control of disease
2	Minimally effective: No, or unknown impact on survival, but sometimes provides control of disease
1	Palliative: Provides symptomatic benefit only

Safety of Regimen/Agent

5	Usually no meaningful toxicity: Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
4	Occasionally toxic: Rare significant toxicities or low-grade toxicities only; little interference with ADLs
3	Mildly toxic: Mild toxicity that interferes with ADLs
2	Moderately toxic: Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
1	Highly toxic: Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

Note: For significant chronic or long-term toxicities, score decreased by 1

Quality of Evidence

5	High quality: Multiple well-designed randomized trials and/or meta-analyses
4	Good quality: One or more well-designed randomized trials
3	Average quality: Low quality randomized trial(s) or well-designed non-randomized trial(s)
2	Low quality: Case reports or extensive clinical experience
1	Poor quality: Little or no evidence

Consistency of Evidence

5	Highly consistent: Multiple trials with similar outcomes
4	Mainly consistent: Multiple trials with some variability in outcome
3	May be consistent: Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
2	Inconsistent: Meaningful differences in direction of outcome between quality trials
1	Anecdotal evidence only: Evidence in humans based upon anecdotal experience

Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)

5	Very inexpensive
4	Inexpensive
3	Moderately expensive
2	Expensive
1	Very expensive



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

WORKUP

Suspicion of
myeloproliferative
neoplasms (MPN)^a

- History and physical (H&P), including spleen size by palpation, evaluation of thrombotic/hemorrhagic events and cardiovascular risk factors
- Complete blood count (CBC) with differential and examination of blood smear
- Comprehensive metabolic panel with uric acid, lactate dehydrogenase (LDH), and liver function tests (LFTs)
- Fluorescence in situ hybridization (FISH) or multiplex reverse transcriptase polymerase chain reaction (RT-PCR) (if available) for *BCR::ABL1* to exclude the diagnosis of chronic myeloid leukemia (CML); if *BCR::ABL1*-positive, see [NCCN Guidelines for Chronic Myeloid Leukemia](#)
- Bone marrow aspirate with iron stain; bone marrow biopsy with trichrome and reticulin stain^{b,c,d}
- Bone marrow cytogenetics (if inaspirable, metaphase karyotype on peripheral blood or second core or use FISH for MDS- or MPN-associated abnormalities on peripheral blood)^{b,c,d}
- Molecular testing (blood or bone marrow) with multigene panel testing to detect driver mutations (*JAK2*, *CALR*, *MPL*)^e and other pathogenic variants for diagnosis and prognostic risk stratification
- If there is evidence of mast cell aggregates in the bone marrow, see [NCCN Guidelines for Systemic Mastocytosis](#) for diagnostic workup
- Assessment of symptom burden using MPN Symptom Assessment Form Total Symptom Score (MPN-SAF TSS; [MPN-E 1 of 2](#))
- Documentation of transfusion/medication history
- Human leukocyte antigen (HLA) testing, if considering allogeneic hematopoietic cell transplant (HCT)^f
- Serum erythropoietin (EPO) level
- Serum iron studies
- In patients with elevated platelet count undergoing surgical procedures, or those with unexplained bleeding, consider testing for acquired von Willebrand syndrome (VWS):^g
 - ▶ Plasma von Willebrand factor antigen (VWFA) measurement
 - ▶ Von Willebrand activity

Diagnosis
and
Risk
stratification

→ [MPN-2](#)

^a See Workup in the [NCCN Guidelines for Systemic Mastocytosis](#) and [Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions](#).

^b See WHO and International Consensus Classification (ICC) diagnostic criteria for primary myelofibrosis (PMF) ([MPN-A](#)), polycythemia vera (PV) ([MPN-B](#)), and essential thrombocythemia (ET) ([MPN-C](#)).

^c WHO and ICC Diagnostic Criteria for PV and Post-PV MF. See [MPN-B](#).

^d WHO and ICC Diagnostic Criteria for ET and Post-ET MF. See [MPN-C](#).

^e Prognostic models incorporating other mutations have been proposed to identify patients with myelofibrosis (MF) as well as PV and ET to better estimate overall survival (OS), MF-free survival (PV and ET), and rates of leukemic transformation. Next-generation sequencing (NGS) is useful to establish clonality.

See [MPN-D](#) for a list of somatic mutations with prognostic significance in patients with MF.

^f See [MF-1](#) and [MF-2](#).

^g An expanded panel including VWFA, factor VIII activity, and VWS multimers may be useful under certain circumstances.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

DIAGNOSIS^{h,i}

PROGNOSTIC RISK MODEL

RISK STRATIFICATION

Myelofibrosis
(MF)

Primary myelofibrosis (PMF)^j
• Preferred: [MIPSS-70](#) or [MIPSS-70+ Version 2.0](#)
• [DIPSS](#) (if recent karyotyping is not available)
• [DIPSS-Plus](#) (if molecular testing is not available)
Post-PV or Post-ET MF^{c,d}
• Preferred: [MIPSS-70+ Version 2.0](#)
• [DIPSS-Plus](#)
• [MYSEC-PM](#)

[Lower-risk \(MF-1\)](#)

- [MIPSS-70](#): ≤3
- [MIPSS-70+ Version 2.0](#): ≤3
- [DIPSS-Plus](#): ≤1
- [DIPSS](#): ≤2
- [MYSEC-PM](#): <14

[Higher-risk \(MF-2\)](#)

- [MIPSS-70](#): ≥4
- [MIPSS-70+ Version 2.0](#): ≥4
- [DIPSS-Plus](#): >1
- [DIPSS](#): >2
- [MYSEC-PM](#): ≥14

Polycythemia vera
(PV)^c

Conventional risk model^k

[Low-risk \(PV-1\)](#)

- Age <60 years and no prior history of thrombosis

[High-risk \(PV-2\)](#)

- Age ≥60 years and/or prior history of thrombosis

[Very-low-risk \(ET-1\)](#)

- Age ≤60 years, no *JAK2* mutation, no prior history of thrombosis

[Low-risk \(ET-1\)](#)

- Age ≤60 years, with *JAK2* mutation, no prior history of thrombosis

[Intermediate-risk \(ET-1\)](#)

- Age >60 years, no *JAK2* mutation, no prior history of thrombosis

[High-risk \(ET-2\)](#)

- History of thrombosis at any age or age >60 years with *JAK2* mutation

Essential
thrombocythemia
(ET)^d

IPSET-thrombosis (revised)^l

[MPN-AP/BP-1](#)

Accelerated/blast phase MPN

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Footnotes on
MPN-2A](#)



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

FOOTNOTES

^c WHO and ICC Diagnostic Criteria for PV and Post-PV MF. See [MPN-B](#).

^d WHO and ICC Diagnostic Criteria for ET and Post-ET MF. See [MPN-C](#).

^h The diagnosis of MPN is based on the 2022 WHO criteria and ICC criteria.

ⁱ Referral to specialized centers with expertise in the management of MPN is strongly recommended for all patients diagnosed with MF, PV, or ET.

^j See WHO and ICC Diagnostic Criteria for PMF ([MPN-A](#)).

^k Marchioli R, et al. J Clin Oncol 2005;23:2224-2232.

^l The revised International Prognostic Score of Thrombosis for Essential Thrombocythemia (IPSET-thrombosis) is preferred for the risk stratification of ET (Haider M, et al. Am J Hematol 2016;91:390-394; Barbui T, et al. Blood Cancer J 2015;5:e369).


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™



[See Evidence Blocks on EB-1](#)

- ^a Evaluation for allogeneic HCT is recommended for patients with low platelet counts or complex cytogenetics. Identification of higher-risk mutations may be helpful in the decision-making regarding allogeneic HCT for patients with MF.
- ^b [Prognostic Significance of Mutations in Myelofibrosis \(MPN-D\)](#).
- ^c When counseling patients about transplant, the Myelofibrosis Transplant Scoring System (MTSS) can be helpful in predicting post-transplant survival.
- ^d [Supportive Care for Patients with MPN \(MPN-F\)](#).
- ^e In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate (see [MPN-G 2 of 2](#)).

^f Bone marrow aspirate and biopsy with NGS and karyotyping should be performed at diagnosis and as clinically indicated (if supported by increased symptoms and signs of progression). Additional molecular testing using multigene NGS panel should be considered to evaluate for higher-risk mutations associated with disease progression in patients with MF.

^g Response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the [2013 IWG-MRT and ELN Response Criteria for MF \(MF-B\)](#). Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician. RR6 may also be used to gauge response. Continuation of JAK inhibitors is recommended based on the discretion of the clinician.

^h For ruxolitinib, use RR6 model to assess. Maffioli M, et al. Blood Adv 2022;6:1855-1864.

ⁱ [Special Considerations for the Use of JAK Inhibitors \(MPN-G\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

TREATMENT FOR HIGHER-RISK MYELOFIBROSIS



[See Evidence Blocks on EB-1](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

[Footnotes on MF-2A](#)



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

TREATMENT FOR HIGHER-RISK MYELOFIBROSIS

FOOTNOTES

^b [Prognostic Significance of Mutations in Myelofibrosis \(MPN-D\)](#).

^c When counseling patients about transplant, the MTSS can be helpful in predicting post-transplant survival.

^d [Supportive Care for Patients with MPN \(MPN-F\)](#).

^f Bone marrow aspirate and biopsy with NGS and karyotyping should be performed at diagnosis and as clinically indicated (if supported by increased symptoms and signs of progression). Additional molecular testing using multigene NGS panel should be considered to evaluate for higher-risk mutations associated with disease progression in patients with MF.

^g Response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the [2013 IWG-MRT and ELN Response Criteria for MF \(MF-B\)](#). Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician. RR6 may also be used to gauge response. Continuation of JAK inhibitors is recommended based on the discretion of the clinician.

^h For ruxolitinib, use RR6 model to assess. Maffioli M, et al. Blood Adv 2022;6:1855-1864.

ⁱ [Special Considerations for the Use of JAK Inhibitors \(MPN-G\)](#).

^j [Special Considerations for Allogeneic Hematopoietic Cell Transplant \(MF-C\)](#).

^k Donor selection and conditioning should be evaluated on a case-by-case basis. See [NCCN Guidelines for Hematopoietic Cell Transplantation](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

MANAGEMENT OF MF-ASSOCIATED ANEMIA^d



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

RISK STRATIFICATION FOR PATIENTS WITH MYELOFIBROSIS

PMF

DIPSS	(MF-A, 2 of 5)
DIPSS-PLUS	(MF-A, 2 of 5)
MIPSS-70	(MF-A, 3 of 5)
MIPSS-70+ Version 2.0	(MF-A, 4 of 5)

Post-PV AND Post-ET MF	
MYSEC-PM	(MF-A, 5 of 5)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

RISK STRATIFICATION FOR PATIENTS WITH PMF

DYNAMIC INTERNATIONAL PROGNOSTIC SCORING SYSTEM (DIPSS)¹

Prognostic Variable	Points		
	0	1	2
Age, y	≤65	>65	
White blood cell count, x10 ⁹ /L	≤25	>25	
Hemoglobin, g/dL	≥10		<10
Peripheral blood blast, %	<1	≥1	
Constitutional symptoms, Y/N	N	Y	

Risk Group	Points
Low	0
Intermediate-1 (INT-1)	1 or 2
Intermediate-2 (INT-2)	3 or 4
High	5 or 6

Online calculator for DIPSS score can be found at
https://qxmd.com/calculate/calculator_187/dipss-prognosis-in-myelofibrosis

DIPSS-PLUS²

Prognostic Variable	Points
DIPSS low-risk	0
DIPSS intermediate-risk 1 (INT-1)	1
DIPSS intermediate-risk 2 (INT-2)	2
DIPSS high-risk	3
Platelets <100 x 10 ⁹ /L	1
Transfusion need	1
Unfavorable karyotype*	1

*Unfavorable karyotype: complex karyotype or sole or two abnormalities that include trisomy 8, -7/7q-, i(17q), -5/5q-, 12p-, inv(3), or 11q23 rearrangement.

Risk Group	Points
Low	0
Intermediate-1 (INT-1)	1
Intermediate-2 (INT-2)	2 or 3
High	4 to 6

Online calculator for DIPSS-PLUS score can be found at
https://qxmd.com/calculate/calculator_315/dipss-plus-score-for-prognosis-in-myelofibrosis

¹ Passamonti F, Cervantes F, Vannucchi AM, et al. A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). Blood 2010;115:1703-1708.

² Gangat N, Caramazza D, Vaidya R, et al. DIPSS plus: a refined Dynamic International Prognostic Scoring System for primary myelofibrosis that incorporates prognostic information from karyotype, platelet count, and transfusion status. J Clin Oncol 2011;29:392-397.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

RISK STRATIFICATION FOR PATIENTS WITH PMF

MUTATION-ENHANCED IPSS (MIPSS-70) FOR PATIENTS WITH PMF AGE ≤70 YEARS³

Prognostic Variable	Points
Hemoglobin <10 g/dL	1
Leukocytes >25 x 10 ⁹ /L	2
Platelets <100 x 10 ⁹ /L	2
Circulating blasts ≥2%	1
Bone marrow fibrosis grade ≥2	1
Constitutional symptoms	1
CALR type-1 unmutated genotype	1
High-molecular-risk (HMR) mutations ^a	1
≥2 HMR mutations	2

Risk Group	Points
Low	0–1
Intermediate	2–4
High	≥5

Online calculator for MIPSS-70 can be found at <http://www.mipss70score.it>.

Footnote

^a Presence of a mutation in any of the following genes: *ASXL1*, *EZH2*, *SRSF2*, or *IDH1/2*.

Reference

³ Guglielmelli P, Lasho TL, Rotunno G, et al. MIPSS-70: Mutation-Enhanced International Prognostic Score System for Transplantation-Age Patients with Primary Myelofibrosis. *J Clin Oncol* 2018;36:310-318.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

RISK STRATIFICATION FOR PATIENTS WITH PMF MUTATION AND KARYOTYPE-ENHANCED IPSS (MIPSS-70+ VERSION 2.0) FOR PATIENTS WITH PMF^{4,5}

Prognostic Variable	Points
Severe anemia (Hemoglobin <8 g/dL in women and <9 g/dL in men)	2
Moderate anemia (Hemoglobin 8–9.9 g/dL in women and 9–10.9 g/dL in men)	1
Circulating blasts ≥2%	1
Constitutional symptoms	2
Absence of CALR type 1 mutation	2
HMR mutations ^b	2
≥2 HMR mutations	3
Unfavorable karyotype ^c	3
Very-high-risk (VHR) karyotype ^d	4

Online calculator for MIPSS-70+ Version 2.0 can be found at <http://www.mipss70score.it>.

Risk Group	Points
Very low	0
Low	1–2
Intermediate	3–4
High	5–8
Very high	≥9

Footnotes

^b Presence of a mutation in any of the following genes: *ASXL1*, *EZH2*, *SRSF2*, *U2AF1* Q157, or *IDH1/2*.

^c Unfavorable karyotype: any abnormal karyotype other than normal karyotype or sole abnormalities of 20q–, 13q–, +9, chromosome 1 translocation/duplication, or –Y or sex chromosome abnormality other than –Y.

^d VHR karyotype: single/multiple abnormalities of –7, i(17q), inv(3)/3q21, 12p–/12p11.2, 11q–/11q23, or other autosomal trisomies not including + 8/+9 (eg, +21, +19).

References

⁴ Tefferi A, Guglielmelli P, Lasho TL, et al. MIPSS70 + Version 2.0: Mutation and Karyotype-Enhanced International Prognostic Scoring System for Primary Myelofibrosis. *J Clin Oncol* 2018;36:1769-1770.

⁵ Tefferi A, Nicolosi M, Mudireddy M, et al. Revised cytogenetic risk stratification in primary myelofibrosis: analysis based on 1002 informative patents. *Leukemia* 2018;32:1189-1199.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

RISK STRATIFICATION FOR PATIENTS WITH POST-PV AND POST-ET MF

MYELOFIBROSIS SECONDARY TO PV AND ET-PROGNOSTIC MODEL (MYSEC-PM)⁶

Prognostic Variable	Points
Age at diagnosis	0.15 per patient's year of age
Hemoglobin <11 g/dL	2
Circulating blasts ≥3%	2
Absence of CALR type 1 mutation	2
Platelets <150 x 10 ⁹ /L	1
Constitutional symptoms	1

Risk Group	Points
Low	<11
Intermediate-1 (INT-1)	≥11 and <14
Intermediate-2 (INT-2)	≥14 and <16
High	≥16

Online calculator for MYSEC-PM can be found at <http://mysec-pm.eu>.

⁶ Passamonti F, Giorgino T, Mora B, et al. A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis. Leukemia 2017;31:2726-2731.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

IWG-MRT AND ELN RESPONSE CRITERIA FOR MYELOFIBROSIS

- Tefferi A, Barosi G, Passamonti F, et al. Proposals for revised International Working Group-European LeukemiaNet criteria for anemia response in myelofibrosis. Blood 2024;144:1813-1820.
- Tefferi A, Cervantes F, Mesa R, et al. Revised response criteria for myelofibrosis: International Working Group-Myeloproliferative Neoplasms Research and Treatment (IWG-MRT) and European LeukemiaNet (ELN) consensus report. Blood 2013;122:1395-1398.
- These response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the IWG-MRT Response Criteria. Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician.
- Patients undergoing HCT may be evaluated for response assessment using other response criteria.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



SPECIAL CONSIDERATIONS FOR ALLOGENEIC HEMATOPOIETIC CELL TRANSPLANT^{1,2}

- Referral to an HCT expert for allogeneic HCT evaluation is recommended for all patients stage DIPSS-Plus-intermediate-1 or MIPSS-intermediate or higher.
- The selection of patients for allogeneic HCT is based on age, performance status, major comorbid conditions, psychosocial status, patient preference, and availability of caregiver.
- Transplant is recommended for patients with DIPSS-Plus or MYSEC intermediate-2 or high-risk disease or MIPSS70 or MIPSS70+ high risk.
- Transplant can also be considered in selective cases – red blood cell (RBC) transfusion dependence, high-risk mutations (ie, ASXL1, RAS, TP53), or loss of response to JAK inhibitor therapy.
- The Myelofibrosis Transplant Scoring System (MTSS) can be used to optimize patient selection, low and intermediate risk being optimal.³
- JAK inhibitors should be considered for use in patients for at least 2 months prior to transplant in patients with splenomegaly and/or constitutional symptoms even if the patient would otherwise be already transplant eligible.
- JAK inhibitors can be tapered prior to or during conditioning to be completed before cell infusion.
- In patients with massive splenomegaly not responding to JAK inhibition, alternative measures to reduce spleen size may be considered prior to transplant (eg, splenic radiation, splenic artery embolization, splenectomy).
- For MF with >10% blasts in the peripheral blood, consider azacitidine with or without a JAK inhibitor or prior to allogeneic HCT to reduce the blast percentage.
- An HLA-matched sibling is the preferred donor; however, an HLA-matched unrelated donor, 7/8-mismatched unrelated donor, or haploidentical donor are also appropriate.
- When counseling patients about transplant, the MTSS can be helpful in predicting post-transplant survival.
- Post-transplant monitoring for measurable residual disease by driver mutations and chimerism is recommended.⁴

¹ Kröger N, Bacigalupo A, Barbui T, et al. Indication and management of allogeneic haematopoietic stem-cell transplantation in myelofibrosis: updated recommendations by the EBMT/ELN International Working Group. *Lancet Haematol* 2024;11:e62-e74.

² Polverelli N, Hernández-Boluda JC, Czerw T, et al. Splenomegaly in patients with primary or secondary myelofibrosis who are candidates for allogeneic hematopoietic cell transplantation: a Position Paper on behalf of the Chronic Malignancies Working Party of the EBMT. *Lancet Haematol* 2023;10:e59-e70.

³ Gagelmann N, Ditschkowski M, Bogdanov R, et al. Comprehensive clinical-molecular transplant scoring system for myelofibrosis undergoing stem cell transplantation. *Blood* 2019;133:2233-2242.

⁴ Gagelmann N, Quarder M, Badbaran A, et al. Clearance of driver mutations after transplantation for myelofibrosis. *N Engl J Med* 2025;392:150-160.


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

NCCN Guidelines Version 2.2026

Polycythemia Vera

NCCN Evidence Blocks™

TREATMENT FOR LOW-RISK POLYCYTHEMIA VERA^a



[See Evidence Blocks on EB-3](#)

^a [Special Considerations in the Treatment of MPN \(MPN-H\)](#).

^b Cytoreductive therapy is not recommended as initial treatment.

^c Landolfi R, et al. N Engl J Med 2004;350:114-124.

^d Aspirin twice daily may be considered for patients with refractory symptoms (Dillinger JG, et al. Thromb Res 2012;129:91-94; Pascale S, et al. Blood 2012;119:3595-3603).

^e Hematocrit <45% is based on the data from the CYTO-PV study (Marchioli R, et al. N Engl J Med 2013;368:22-33). There may be situations in which a lower hematocrit cutoff may be appropriate and it should be individualized (eg, progressive symptoms).

^f [Supportive Care for Patients with MPN \(MPN-F\)](#).

^g While normalization of blood counts after initiation of treatment is usually a goal in clinical practice, it is not associated with long-term clinical benefit and there are no evidence-based data to recommend a target white blood cell (WBC) or platelet count for patients receiving cytoreductive therapy. In selected patients with a severe thrombotic event or other disease-related symptoms, normalization of blood counts might be a goal of treatment.

^h Barbui T, et al. Leukemia 2018;32:1057-1069.

ⁱ [WHO and ICC Diagnostic Criteria for Post-PV MF \(MPN-B\)](#).

^j Peginterferon alfa-2a is an option for younger patients or in pregnant patients in need of cytoreductive therapy (MPN-G 2 of 2).

^k In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropoginterferon alfa-2b-njft) is appropriate (MPN-G 2 of 2).


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF
All recommendations are category 2A unless otherwise indicated.

NCCN Guidelines Version 2.2026

Polycythemia Vera

NCCN Evidence Blocks™

TREATMENT FOR HIGH-RISK POLYCYTHEMIA VERA^a



^a [Special Considerations in the Treatment of MPN \(MPN-H\)](#).

^c Landolfi R, et al. N Engl J Med 2004;350:114-124.

^d Aspirin twice daily may be considered for patients with refractory symptoms (Dillinger JG, et al. Thromb Res 2012;129:91-94; Pascale S, et al. Blood 2012;119:3595-3603).

^e Hematocrit <45% is based on the data from the CYTO-PV study (Marchioli R, et al. N Engl J Med 2013;368:22-33). There may be situations in which a lower hematocrit cutoff may be appropriate and it should be individualized (eg, progressive symptoms).

^f [Supportive Care for Patients with MPN \(MPN-F\)](#).

^g While normalization of blood counts after initiation of treatment is usually a goal in clinical practice, it is not associated with long-term clinical benefit and there are no evidence-based data to recommend a target WBC or platelet count for patients receiving cytoreductive therapy. In selected patients with a severe thrombotic event or other disease-related symptoms, normalization of blood counts might be a goal of treatment.

^h Barbui T, et al. Leukemia 2018;32:1057-1069.

ⁱ [WHO and ICC Diagnostic Criteria for Post-PV MF \(MPN-B\)](#).

^j Peginterferon alfa-2a is an option for younger patients or in pregnant patients in need of cytoreductive therapy ([MPN-G 2 of 2](#)).

^k In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate ([MPN-G 2 of 2](#)).

^l Response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the [2013 IWG-MRT and ELN Response Criteria for PV \(PV-A\)](#). Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician.

^m Definition of intolerance/resistance to hydroxyurea ([MPN-I](#)).

ⁿ Ruxolitinib is FDA approved for the treatment of patients with PV who have had an inadequate response to or are intolerant of hydroxyurea. Ruxolitinib may have activity after inadequate response or loss of response to other agents besides hydroxyurea. See [Discussion](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myelofibrosis

NCCN Evidence Blocks™

2013 IWG-MRT AND ELN RESPONSE CRITERIA FOR POLYCYTHEMIA VERA

- Barosi G, Mesa R, Finazzi G, et al. Revised response criteria for polycythemia vera and essential thrombocythemia: an ELN and IWG-MRT consensus project. Blood 2013;121:4778-4781.
- These response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the IWG-MRT Response Criteria. Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Polycythemia Vera

NCCN Evidence Blocks™

OVERALL SURVIVAL RISK STRATIFICATION FOR PATIENTS WITH POLYCYTHEMIA VERA^a

MIPSS-PV

Prognostic Variable	Points
Thrombosis history	1
Leukocyte count $\geq 15 \times 10^9/L$	1
Age > 67	2
Adverse mutations (SRSF2)	3

Risk Group	Points
Low	0–1
Intermediate	2–3
High	≥ 4

^a Tefferi A, Guglielmelli P, Lasho TL, et al. Mutation-enhanced international prognostic systems for essential thrombocythaemia and polycythaemia vera. Br J Haematol 2020;189:291-302.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Essential Thrombocythemia

NCCN Evidence Blocks™

TREATMENT FOR VERY-LOW-RISK OR LOW-RISK OR INTERMEDIATE-RISK ESSENTIAL THROMBOCYTHEMIA^a



^a [Special Considerations in the Treatment of MPN \(MPN-H\)](#).

^b Harrison CN, et al. N Engl J Med 2005;353:33-45.

^c Aspirin should be used with caution in patients with acquired VWS. Higher-dose aspirin may be appropriate in selected patients as clinically indicated. The risks and benefits of higher-dose aspirin (>100 mg) must be weighed based on the presence of vasomotor symptoms versus the risk of bleeding.

^d Aspirin twice daily may be considered for patients with refractory symptoms or cardiovascular risk factors (Dillinger JG, et al. Thromb Res 2012;129:91-94; Pascale S, et al. Blood 2012;119:3595-3603; Rocca B, et al. Am J Hematol 2024;99:1462-1474).

^e [Supportive Care for Patients with MPN \(MPN-F\)](#).

^f Bone marrow aspirate and biopsy should be performed to rule out disease progression to MF prior to the initiation of cytoreductive therapy.

^g While normalization of blood counts after initiation of treatment is usually a goal in clinical practice, it is not associated with long-term clinical benefit and there are no evidence-based data to recommend a target WBC or platelet count for patients receiving cytoreductive therapy. In selected patients with a severe thrombotic event or other disease-related symptoms, normalization of blood counts might be a goal of treatment.

^h Barbui T, et al. Leukemia 2018;32:1057-1069.

ⁱ [WHO and ICC Diagnostic Criteria for Post-ET MF \(MPN-C\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Essential Thrombocythemia

NCCN Evidence Blocks™

TREATMENT FOR HIGH-RISK ESSENTIAL THROMBOCYTHEMIA^a



[See Evidence Blocks on EB-4](#)

^a [Special Considerations in the Treatment of MPN \(MPN-H\)](#).

^c Aspirin should be used with caution in patients with acquired VWS. Higher-dose aspirin may be appropriate in selected patients as clinically indicated. The risks and benefits of higher-dose aspirin (>100 mg) must be weighed based on the presence of vasomotor symptoms versus the risk of bleeding.

^d Aspirin twice daily may be considered for patients with refractory symptoms or cardiovascular risk factors (Dillinger JG, et al. Thromb Res 2012;129:91-94; Pascale S, et al. Blood 2012;119:3595-3603; Rocca B, et al. Am J Hematol 2024;99:1462-1474).

^e [Supportive Care for Patients with MPN \(MPN-F\)](#).

^f Bone marrow aspirate and biopsy should be performed to rule out disease progression to MF if clinical/laboratory suspicion of MF.

^g While normalization of blood counts after initiation of treatment is usually a goal in clinical practice, it is not associated with long-term clinical benefit and there are no evidence-based data to recommend a target WBC or platelet count for patients receiving cytoreductive therapy. In selected patients with a severe thrombotic event or other disease-related symptoms, normalization of blood counts might be a goal of treatment.

ⁱ [WHO and ICC Diagnostic Criteria for Post-ET MF \(MPN-C\)](#).

^j Response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the [2013 IWG-MRT and ELN Response Criteria for ET \(ET-A\)](#). Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician.

^k Peginterferon alfa-2a can be considered for patients in need of cytoreductive therapy who are younger or pregnant or who defer hydroxyurea.

^l In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Essential Thrombocythemia

NCCN Evidence Blocks™

TREATMENT FOR HIGH-RISK ESSENTIAL THROMBOCYTHEMIA^a

Inadequate response
or
Loss of response

Potential indications for change of cytoreductive therapy^h:

- Intolerance or resistance to hydroxyurea^m or peginterferon alfa-2a^{k,l} or anagrelide
- New thrombosis, acquired VWS, and/or disease-related major bleeding
- Splenomegaly
- Thrombocytosis
- Leukocytosis
- Disease-related symptoms (eg, pruritus, night sweats, fatigue)
- Vasomotor/microvascular disturbances not responsive to aspirin (eg, headaches/chest pain, erythromelalgia)

Preferred:

- Clinical trial
or
- [Ropeginterferon alfa-2b-njft](#) (category 1)
or
- Hydroxyurea if not previously used

Other Recommended:

- [Peginterferon alfa-2a](#) if not previously used^{k,l}

Useful in Certain Circumstances:

- [Anagrelide](#) if not previously used
or
- Plateletpheresis (for emergent situations, eg, severe thrombocytosis-related neurologic complications)ⁿ
or
- Ruxolitinib

[See Evidence Blocks on EB-2](#)

^a [Special Considerations in the Treatment of MPN \(MPN-H\)](#).

^h Barbui T, et al. Leukemia 2018;32:1057-1069.

^k Peginterferon alfa-2a can be considered for patients in need of cytoreductive therapy who are younger or pregnant or who defer hydroxyurea.

^l In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate.

^m [Definition of Resistance/Intolerance to Hydroxyurea \(MPN-I\)](#).

ⁿ Padmanabhan A, et al. J Clin Apher 2019;34:171-354.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Essential Thrombocythemia

NCCN Evidence Blocks™

2013 IWG-MRT AND ELN RESPONSE CRITERIA FOR ESSENTIAL THROMBOCYTHEMIA

- Barosi G, Mesa R, Finazzi G, et al. Revised response criteria for polycythemia vera and essential thrombocythemia: an ELN and IWG-MRT consensus project. Blood 2013;121:4778-4781.
- These response criteria were developed mainly for use in clinical trials. Clinical benefit may not reach the threshold of the IWG-MRT Response Criteria. Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Essential Thrombocythemia

NCCN Evidence Blocks™

OVERALL SURVIVAL RISK STRATIFICATION FOR PATIENTS WITH ESSENTIAL THROMBOCYTHEMIA^a

MIPSS-ET

Prognostic Variable	Points
Male sex	1
Leukocyte count $\geq 11 \times 10^9/L$	1
Adverse mutations (SRSF2, SF3B1, U2AF1, TP53)	2
Age >60	4

Risk Group	Points
Low	0–1
Intermediate	2–5
High	≥ 6

^a Tefferi A, Guglielmelli P, Lasho TL, et al. Mutation-enhanced international prognostic systems for essential thrombocythaemia and polycythaemia vera. Br J Haematol 2020;189:291-302.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

WORKUP



MPN-accelerated phase (blasts 10%–19% in peripheral blood or bone marrow)

MPN blast phase (blasts ≥20% in peripheral blood or bone marrow, or myeloid sarcoma)

Transplant candidate^{a,b}

Not a candidate for transplant^a

TREATMENT

- Clinical trial or
- Bridging therapy^c followed by allogeneic HCT. Bridging therapy options include:
 - ▶ Hypomethylating agent (HMA)
 - ▶ HMA + JAK inhibitors^d
 - ▶ HMA + venetoclax^e
 - ▶ Intensive induction chemotherapy^{e,f} (see [NCCN Guidelines for AML](#))

- Clinical trial or
- HMA or
- HMA + JAK inhibitors^d or
- HMA + venetoclax^e or
- Lower-intensity chemotherapy or targeted therapy^{e,f} (see [NCCN Guidelines for AML](#))

^a The selection of patients for allogeneic HCT should be based on age, performance status, major comorbid conditions, psychosocial status, patient preference, and availability of caregiver. Early referral to transplant team is recommended for planning purposes and to discuss the role of bridging therapy.

^b JAK inhibitors may be continued near to the start of conditioning therapy for the improvement of splenomegaly and other disease-related symptoms ([MPN-G](#)). See [Discussion](#) for the use of JAK inhibitors prior to transplant.

^c Some patients in accelerated phase may proceed to transplant directly without bridging therapy (Gagelmann N, et al. Blood Adv 2022;6:1222-1231).

^d JAK inhibitors can be used in combination with HMA (azacitidine or decitabine) for the palliation of splenomegaly or other disease-related symptoms. There are very limited data regarding the use of fedratinib, momelotinib, or pacritinib with HMAs.

^e AML-type induction chemotherapy regimens and HMA + venetoclax may be used for the management of disease progression of MPN. However, these regimens typically result in low response rates and are associated with significant toxicities. Based on NGS panel results, targeted therapy (eg, IDH1/2 and FLT3 inhibitors) alone or in combination with HMAs can be considered.

^f Consider prophylaxis for tumor lysis syndrome (TLS). See [Supportive Care for Patients with MPN \(MPN-F\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

WHO¹ AND INTERNATIONAL CONSENSUS CLASSIFICATION (ICC)² DIAGNOSTIC CRITERIA FOR PRIMARY MYELOFIBROSIS

PMF, prefibrotic stage (pre-PMF)	PMF, fibrotic stage
Major criteria 1. Megakaryocytic proliferation and atypia, without reticulin fibrosis grade >1, ^a accompanied by increased age-adjusted bone marrow cellularity, granulocytic proliferation, and (often) decreased erythropoiesis 2. Not meeting diagnostic criteria for chronic myeloid leukemia, polycythemia vera, essential thrombocythemia, myelodysplastic neoplasms, or other defined myeloid neoplasms 3. JAK2, CALR, or MPL mutation or presence of another clonal marker ^b or absence of reactive bone marrow fibrosis ^c	Major criteria 1. Megakaryocytic proliferation and atypia, accompanied by reticulin and/or collagen fibrosis grade 2 or 3 ^a 2. Not meeting diagnostic criteria for chronic myeloid leukemia, polycythemia vera, essential thrombocythemia, myelodysplastic neoplasms, or other defined myeloid neoplasms ^d 3. JAK2, CALR, or MPL mutation or presence of another clonal marker ^e or absence of reactive bone marrow myelofibrosis ^f
Minor criteria • Anemia not attributed to a comorbid condition • Leukocytosis $\geq 11 \times 10^9/L$ • Splenomegaly detected clinically and/or by imaging • LDH level above the upper limit of the institutional reference range	Minor criteria • Anemia not attributed to a comorbid condition • Leukocytosis $\geq 11 \times 10^9/L$ • Splenomegaly detected clinically and/or by imaging • LDH level above the upper limit of the institutional reference range • Leukoerythroblastosis
The diagnosis of prefibrotic-PMF or overt PMF requires all 3 major criteria and at least 1 minor criterion confirmed in 2 consecutive determinations	The diagnosis of overt primary myelofibrosis requires all 3 major criteria and at least 1 minor criterion to be met in 2 consecutive determinations.

^a See [MPN-A 2 of 2](#) for the WHO grading system for myelofibrosis.

^b In the absence of any of the three major clonal mutations, a search for other mutations associated with myeloid neoplasms (eg, *ASXL1*, *EZH2*, *TET2*, *IDH1*, *IDH2*, *SRSF2*, and *SF3B1* mutations) may be of help in determining the clonal nature of the disease.

^c Minor (grade 1) reticulin fibrosis secondary to infection, autoimmune disorder or other chronic inflammatory conditions, hairy cell leukemia or another lymphoid neoplasm, metastatic malignancy, or toxic (chronic) myelopathies.

^d Myeloproliferative neoplasms (MPNs) can be associated with monocytosis, or patients can develop it during the course of the disease. These cases may mimic chronic myelomonocytic leukemia; in these rare instances, a history of MPN excludes chronic myelomonocytic leukemia, whereas the presence of MPN features in the bone marrow and/or MPN-associated mutations (in *JAK2*, *CALR*, or *MPL*) tends to support the diagnosis of MPN with monocytosis rather than chronic myelomonocytic leukemia.

^e In the absence of any of the three major clonal mutations, a search for other mutations associated with myeloid neoplasms (eg, *ASXL1*, *EZH2*, *TET2*, *IDH1*, *IDH2*, *SRSF2*, and *SF3B1* mutations) may be of help in determining the clonal nature of the disease.

^f Bone marrow fibrosis secondary to infection, autoimmune disorder or another chronic inflammatory condition, hairy cell leukemia or another lymphoid neoplasm, metastatic malignancy, or toxic (chronic) myelopathy.

¹ Adapted with permission from Kanagal-Shamanna R, Naresh KN, Dave SS, et al. Primary myelofibrosis. In: WHO Classification of Tumours Editorial Board. Haematolymphoid tumours [Internet]. Lyon (France): International Agency for Research on Cancer; 2024 [cited 2024 November 15]. (WHO classification of tumours series, 5th ed; vol. 11). Available from: <https://tumourclassification.iarc.who.int/chapters/63>.

² Arber DA, Orazi A, Hasserjian RP, et al. International Consensus Classification of myeloid neoplasms and acute leukemias: Integrating morphologic, clinical and genomic data. Blood 2022;140:1200-1228. from Arber DA, et al. Blood 2022;140:1200-1228.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

GRADING OF MYELOFIBROSIS^{3,4}

Grade	Grade definition ^a	Collagen pattern ^b	Osteosclerosis ^c
MF-0	Scattered linear reticulin with no intersections (crossovers) corresponding to normal bone marrow	Perivascular collagen only (normal)	Regular bone trabecula (distinct paratrabecular borders)
MF-1	Loose network of reticulin with many intersections, especially in perivascular areas	Focal paratrabecular or central collagen deposition with no connecting meshwork	Focal budding, hooks, spikes, or paratrabecular apposition of new bone
MF-2	Diffuse and dense increase in reticulin with extensive intersections, occasionally with focal bundles of thick fibers mostly consistent with collagen, and/or focal osteosclerosis	Paratrabecular or central deposition of collagen with focally connecting meshwork or generalized paratrabecular apposition of collagen	Diffuse paratrabecular formation of new bone with thickening of trabecula, occasionally with focal interconnections
MF-3	Diffuse and dense increase in reticulin with extensive intersections and coarse bundles of thick fibers consistent with collagen, usually associated with osteosclerosis	Diffuse (complete) connecting meshwork of collagen in >30% of marrow spaces	Extensive interconnecting meshwork of new bone with overall effacement of marrow spaces

^a Reticulin and collagen fiber density should be assessed only in hematopoietic areas; if the pattern of reticulin fibrosis, collagen deposition, and/or osteosclerosis is heterogeneous, the final grade is determined by the highest grade present in ≥30% of the marrow area.

^b Using trichrome staining (Masson trichrome or Martius Scarlet Blue); the stain is recommended for grades MF-2 and MF-3.

^c Best performed on a bone marrow core biopsy of sufficient length, taken at a right angle from the cortical bone without significant fragmentation.

³ Adapted with permission from Kanagal-Shamanna R, Naresh KN, Dave SS, et al. Primary myelofibrosis. In: WHO Classification of Tumours Editorial Board. Haematolymphoid tumours [Internet]. Lyon (France): International Agency for Research on Cancer; 2024 [cited 2024 November 15]. (WHO classification of tumours series, 5th ed; vol 11). Available from: <https://tumourclassification.iarc.who.int/chapters/63>.

⁴ Kvasnicka HM, Beham-Schmid C, Bob R, et al. Problems and pitfalls in grading of bone marrow fibrosis, collagen deposition and osteosclerosis - a consensus-based study. Histopathology 2016;68:905-915.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

WHO¹ AND ICC² DIAGNOSTIC CRITERIA FOR POLYCYTHEMIA VERA AND POST-PV MYELOFIBROSIS

PV	Post-PV MF
Major criteria 1. Elevated hemoglobin concentration (>16.5 g/dL in men, >16.0 g/dL in women) or elevated hematocrit (>49%^a in men, >48% in women) 2. Bone marrow biopsy showing age-adjusted hypercellularity with trilineage proliferation (panmyelosis), including prominent erythroid, granulocytic, and megakaryocytic proliferation with pleomorphic, mature megakaryocytes (differences in size)^b 3. Presence of JAK2 p.V617F or JAK2 exon 12 mutation Minor criterion • Subnormal serum erythropoietin level	Required criteria 1. Documentation of a previous diagnosis of WHO-defined polycythemia vera 2. Bone marrow fibrosis of grade 2–3 on a scale of 0–3 Additional criteria (two are required): 1. Anemia (below the reference range given age, sex, and altitude considerations) or sustained loss of requirement of either phlebotomy (in the absence of cytoreductive therapy) or cytoreductive treatment for erythrocytosis 2. Leukoerythroblastosis 3. Increasing splenomegaly, defined as either an increase in palpable splenomegaly of >50 mm from baseline (distance from the left costal margin) or the development of a newly palpable splenomegaly 4. Development of any 2 (or all 3) of the following constitutional symptoms: >10% weight loss in 6 months, night sweats, unexplained fever (>37.5°C)
The diagnosis of PV requires either all 3 major criteria or the first 2 major criteria plus the minor criterion.	The diagnosis of post-PV MF is established by all required criteria and at least 2 additional criteria

^a Hematocrit for diagnosis in the presence of a *JAK2* mutation. In the absence of a *JAK2* mutation, a higher hematocrit target (eg, 52%) could be considered in men before further investigation is required.

^b Major criterion 2 (bone marrow biopsy) may not be required in patients with sustained absolute erythrocytosis (hemoglobin concentrations of >18.5 g/dL in men or >16.5 g/dL in women, or hematocrit values of >55.5% in men or >49.5% in women) if major criterion 3 and the minor criterion are present.

¹ Adapted with permission from Harrison CN, Godley LA, Naresh KN, et al. Polycythaemia vera. In: WHO Classification of Tumours Editorial Board. Haematolymphoid tumours [Internet]. Lyon (France): International Agency for Research on Cancer; 2024 [cited 2024 November 15]. (WHO classification of tumours series, 5th ed; vol 11). Available from: <https://tumourclassification.iarc.who.int/chapters/63>.

² Arber DA, Orazi A, Hasserjian RP, et al. International Consensus Classification of Myeloid Neoplasms and Acute Leukemias: Integrating morphologic, clinical, and genomic data. Blood 2022;140:1200-1228.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

WHO¹ AND ICC² DIAGNOSTIC CRITERIA FOR ESSENTIAL THROMBOCYTHEMIA AND POST-ET MYELOFIBROSIS

ET	Post-ET MF
Major criteria 1. Platelet count $\geq 450 \times 10^9/L$ 2. Bone marrow biopsy showing proliferation mainly of the megakaryocytic lineage, with increased numbers of enlarged, mature megakaryocytes with hyperlobulated nuclei, no significant increase or left shift in neutrophil granulopoiesis or erythropoiesis; very rarely a minor (grade 1) increase in reticulin fibers 3. WHO criteria for BCR::ABL1-positive CML, PV, PMF, and other myeloid neoplasms are not met 4. JAK2, CALR, or MPL mutation Minor criteria • Presence of a clonal marker OR • Exclusion of reactive thrombocytosis	Required criteria 1. Documentation of a previous diagnosis of WHO-defined essential thrombocythemia 2. Bone marrow fibrosis of grade 2–3 on a scale of 0–3 Additional criteria 1. Anemia (below the reference range given age, sex, and altitude considerations) and a >2 g/dL decrease from baseline hemoglobin concentration 2. Leukoerythroblastosis 3. Increasing splenomegaly, defined as either an increase in palpable splenomegaly of >50 mm from baseline (distance from the left costal margin, or on imaging) or the development of a newly palpable splenomegaly 4. Elevated LDH level above the reference range 5. Development of any 2 (or all 3) of the following constitutional symptoms: $>10\%$ weight loss in 6 months, night sweats, unexplained fever ($>37.5^\circ C$)
The diagnosis of ET requires either all the major criteria or the first 3 major criteria plus a minor criterion to be met	The diagnosis of post-ET MF requires both of the required criteria and at least 2 additional criteria

¹ Adapted with permission from Harrison CN, Godley LA, Naresh KN, et al. Essential thrombocythemia. In: WHO Classification of Tumours Editorial Board. Haematolymphoid tumours [Internet]. Lyon (France): International Agency for Research on Cancer; 2024 [cited 2024 November 15]. (WHO classification of tumours series, 5th ed; vol 11). Available from: <https://tumourclassification.iarc.who.int/chapters/63>.

² Arber DA, Orazi A, Hasserjian RP, et al. International Consensus Classification of myeloid neoplasms and acute leukemias: Integrating morphologic, clinical and genomic data. Blood 2022;140:1200-1228.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

PROGNOSTIC SIGNIFICANCE OF MUTATIONS IN MYELOFIBROSIS

To be used in conjunction with the prognostic scoring systems^a

Mutated Gene	Driver Mutations ^b
JAK2 V617F	Intermediate prognosis and higher risk of thrombosis compared to patients with CALR type 1 mutation ¹
MPL W515L/K	Intermediate prognosis and higher risk of thrombosis compared to patients with CALR type 1 mutation ¹
CALR type 1	Improved overall survival (OS) and lower risk of thrombosis compared to JAK2 mutation and "triple-negative" PMF. ¹⁻⁴ Improved OS compared to CALR type 2 mutation ⁵⁻⁸
CALR type 2	Decreased OS compared to CALR type 1 mutation ⁵⁻⁸
Mutated Gene	Other Somatic Mutations
ASXL1	Associated with inferior OS ^c and leukemia-free survival (LFS) and LFS following HCT ^{9,10}
EZH2	Associated with inferior OS ⁹
RAS	Associated with inferior OS ¹¹
IDH1/2	Associated with inferior LFS and lower progression-free survival (PFS) following HCT ^{9,10}
SRSF2	Associated with inferior OS and LFS ⁹
TP53	Associated with increased risk of leukemic transformation ¹²
U2AF1	Associated with inferior OS following HCT ¹³⁻¹⁵ U2AF1 Q157 is associated with inferior OS compared to patients with U2AF1 S34 mutated or U2AF1 unmutated MF
DNMT3A	Associated with inferior OS following HCT ^{14,15}
CBL	Associated with inferior OS following HCT ^{14,15}

[References on MPN-D \(2 of 2\)](#)

^a Various clinical outcomes are worse with multiple versus one somatic mutation identified.¹⁶

^b Driver mutation negative "triple negative" MF is associated with inferior OS and LFS.¹⁻³

^c ASXL1 mutation retains prognostic significance for inferior OS independent of IPSS or DIPSS-Plus risk score.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

PROGNOSTIC SIGNIFICANCE OF MUTATIONS IN MYELOFIBROSIS

REFERENCES

- ¹ Rumi E, Pietra D, Pascutto C, et al. Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood* 2014;124:1062-1069.
- ² Tefferi A, Lasho TL, Finke CM, et al. CALR vs JAK2 vs MPL-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia* 2014;28:1472-1477.
- ³ Tefferi A, Guglielmelli P, Larson DR, et al. Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. *Blood* 2014;124:2507-2513.
- ⁴ Klampfl T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med* 2013;369:2379-2390.
- ⁵ Guglielmelli P, Rotunno G, Fanelli T, et al. Validation of the differential prognostic impact of type 1/type-1 like versus type 2/type 2-like CALR mutations in myelofibrosis. *Blood Cancer J* 2015;5:e360.
- ⁶ Tefferi A, Lasho TL, Tischer A, et al. The prognostic advantage of calreticulin mutations in myelofibrosis might be confined to type 1 or type 1-like CALR variants. *Blood* 2014;124:2465-2466.
- ⁷ Tefferi A, Lasho TL, Finke CM, et al. Type 1 vs type 2 calreticulin mutations in primary myelofibrosis: differences in phenotype and prognostic impact. *Leukemia* 2014;28:1568-1570.
- ⁸ Li B, Xu J, Wang J, et al. Calreticulin mutations in Chinese with primary myelofibrosis. *Haematologica* 2014;99:1697-1700.
- ⁹ Vannucchi AM, Lasho TL, Guglielmelli P, et al. Mutations and prognosis in primary myelofibrosis. *Leukemia* 2013;27:1861-1869.
- ¹⁰ Kröger N, Panagiotou V, Badbaran A, et al. Impact of molecular genetics on outcome in myelofibrosis patients after allogeneic stem cell transplantation. *Biol Blood Marrow Transplant* 2017;23:1095-1101.
- ¹¹ Santos FPS, Getta B, Masarova, et al. Prognostic impact of RAS-pathway mutations in patients with myelofibrosis. *Leukemia* 2020;34:799-810.
- ¹² Rampal R, Ahn J, Abdel-Wahab O, et al. Genomic and functional analysis of leukemic transformation of myeloproliferative neoplasms. *Proc Natl Acad Sci USA* 2014;111:E5401-E5410.
- ¹³ Tefferi A, Finke CM, Lasho TL, et al. U2AF1 mutation types in primary myelofibrosis: phenotypic and prognostic distinctions. *Leukemia* 2018;32:2274-2278.
- ¹⁴ Tamari R, Rapaport F, Zhang N, et al. Impact of high-molecular-risk mutation in transplantation outcomes in patients with myelofibrosis. *Biol Blood Marrow Transplant* 2019;25:1142-1151.
- ¹⁵ Ali H, Aldoss I, Yang D, et al. MIPSS70+ v2.0 predicts long-term survival in myelofibrosis after allogeneic HCT with the Flu/Mel conditioning regimen. *Blood Adv* 2019;3:83-95.
- ¹⁶ Guglielmelli P, Lasho TL, Rotunno G, et al. The number of prognostically detrimental mutations and prognosis in primary myelofibrosis: An international study of 797 patients. *Leukemia* 2014;28:1804-1810.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

ASSESSMENT OF SYMPTOM BURDEN

- Assessment of symptoms (in provider's office) at baseline and monitoring symptom status (stable, improved, or worsening) during the course of treatment is recommended for all patients.
- Myeloproliferative Neoplasm Symptom Assessment Form Total Symptom Score (MPN-SAF TSS) is recommended for the assessment of symptom burden at baseline and monitoring symptom status during the course of treatment ([MPN-E 2 of 2](#)).
- MPN-SAF TSS is assessed by the patients themselves. Scoring is from 0 (absent/as good as it can be) to 10 (worst imaginable/as bad as it can be) for each item. The MPN-SAF TSS is the summation of all the individual scores (0–100 scale).
- Symptom response requires ≥50% reduction in the MPN-SAF TSS. A symptom response <50% may be clinically meaningful and justify continued use of JAK inhibitors.
- Changes in symptom status could be a sign of disease progression. Therefore, change in symptom status should prompt evaluation of treatment efficacy and/or disease status.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

MYELOPROLIFERATIVE NEOPLASM SYMPTOM ASSESSMENT FORM: TOTAL SYMPTOM SCORE (MPN-SAF TSS)¹

(Recommended for monitoring symptoms during the course of treatment)

Symptom	1 to 10 (0 if absent) ranking 1 is most favorable and 10 least favorable
Please rate your fatigue (weariness, tiredness) by circling the one number that best describes your WORST level of fatigue during past 24 hours	(No Fatigue) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)

Circle the one number that describes, during the past week, how much difficulty you have had with each of the following symptoms

Filling up quickly when you eat (early satiety)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Abdominal discomfort	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Inactivity	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Problems with concentration-compared to prior to my MPD	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Night sweats	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Itching (pruritus)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Bone pain (diffuse not joint pain or arthritis)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)
Fever (>100 F)	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Daily)
Unintentional weight loss last 6 months	(Absent) 0 1 2 3 4 5 6 7 8 9 10 (Worst Imaginable)

¹ Reproduced with permission from Emanuel RM, Dueck AC, Geyer HL, et al. Myeloproliferative neoplasm (MPN) symptom assessment form total symptom score: prospective international assessment of an abbreviated symptom burden scoring system among patients with MPNs. J Clin Oncol 2012;30:4098-4103.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SUPPORTIVE CARE FOR PATIENTS WITH MPN

MYELOFIBROSIS

- Transfusion support
 - ▶ RBC transfusions for symptomatic anemia
 - ▶ Platelet transfusions for thrombocytopenic bleeding or a platelet count $<10,000/\text{mm}^3$
 - ▶ In transplant candidates, use leukocyte-reduced blood products to prevent HLA alloimmunization and reduce the risk of cytomegalovirus (CMV) transmission.
- Consider antifibrinolytic agents for bleeding that is refractory to transfusions.
- Iron chelation could be considered for patients who have received >20 transfusions and/or ferritin >2500 ng/mL in patients with lower-risk MF. However, the role of iron chelation remains unclear.
- In patients with massive splenomegaly not responding to JAK inhibition, alternative measures to reduce spleen size may be considered (eg, splenic radiation, splenic artery embolization, splenectomy).
- In patients who have had a splenectomy, vaccinations and antibiotic prophylaxis should be given per [IDSA Guidelines](#).
- Vaccinations: See [NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).
 - ▶ Consider recombinant (killed) zoster vaccine for patients on, or prior to, treatment with a JAK inhibitor.
- Hematopoietic growth factor support
 - ▶ ESAs: See [\(MF-3\)](#) for the management of MF-associated anemia. ESAs are generally less effective for patients with transfusion-dependent anemia.
 - ▶ Consider granulocyte colony-stimulating factor (G-CSF) or granulocyte-macrophage colony-stimulating factor (GM-CSF) for recurrent infections in patients with neutropenia. However, these should be used with caution in patients with an enlarged spleen since the use of G-CSF or GM-CSF has been associated with splenic rupture. See [NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).
- Consider cytoreductive therapy (eg, hydroxyurea) for hyperproliferative manifestations of PMF (thrombocytosis or leukocytosis).
- Consider prophylaxis for tumor lysis syndrome (TLS) for patients undergoing induction therapy for advanced-stage MF or disease progression to AML. See [NCCN Guidelines for Acute Myeloid Leukemia](#).
- Counseling at baseline and throughout disease course for assessment for, identification of, and decreasing cardiovascular risk factors (eg, smoking, diet, exercise, hypertension, diabetes mellitus, lipid management), and thrombotic and hemorrhagic risk factors.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



SUPPORTIVE CARE FOR PATIENTS WITH MPN

SYMPTOM MANAGEMENT IN PATIENTS WITH MPN

Disease-related symptoms commonly contribute to decreased quality of life in patients with MPN.¹ While JAK inhibitors have been shown to broadly improve disease-related symptoms, their use is not indicated in all patients with symptomatic MPN, and the presence of specific symptoms often requires a targeted approach. Fatigue, pruritus, bone pain, and headaches/tinnitus occur across all MPN (albeit with some disease preference) and greatly impact quality of life. The optimal management of these symptoms in the setting of MPN has not been established and is based on subset analysis of large trials, small pilot studies, anecdotal evidence, extrapolation from other disease states, and expert opinion.

MPNs are a group of chronic blood cancers that typically progress over time. The stress and uncertainty of long-term care should be acknowledged. Referral to a mental health professional and other supportive services can be invaluable, including integrative health specialists.

• Pruritus²⁻¹¹

- ▶ Initial efforts to improve pruritus should include sensitive skin care practices (ie, short showers, mild soap, moisturizing), optimized antihistamine therapy (ie, cetirizine, diphenhydramine), and topical steroids.
- ▶ Ruxolitinib was shown to improve pruritus in patients with ET, PV, and MF in the MAJIC-ET, RESPONSE, and COMFORT-I trials, respectively.
- ▶ Small pilot studies have shown selective serotonin reuptake inhibitors (SSRIs) and narrow-band ultraviolet B to be effective in treating pruritic symptoms.
- ▶ Additional options include peginterferon alfa-2a, gabapentin, aprepitant, and immunosuppressant agents such as cyclosporine, methotrexate, azathioprine, mycophenolate mofetil, or dupilumab. In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate.
- ▶ A risk-stratified, step-wise approach should be utilized with the specific therapeutic option chosen based on strength of evidence, side effect profile, cost/benefit analysis, and concomitant disease-related symptoms.

• Bone Pain¹²⁻¹⁴

- ▶ Close evaluation to distinguish disease-related bone pain from arthralgias should be undertaken in order to identify symptoms that may be amenable to local therapies.
- ▶ Ruxolitinib was shown to stabilize bone/muscle pain in patients with MF in the COMFORT-1 study.
- ▶ Loratadine and nonsteroidal anti-inflammatory drugs (NSAIDs) (naproxen) have been used in MPN-associated bone pain due to their efficacy in the treatment of growth-factor-related bone pain.
- ▶ Single-fraction radiation has been effective in temporarily relieving MPN-associated bone pain.

• Headache/Tinnitus¹⁵⁻²²

- ▶ Given the increased risk of vascular complications in patients with MPN (ie, stroke, retinal artery or vein thrombosis, cerebral venous thrombosis), all patients with new onset of neurologic symptoms including headache and tinnitus or with progressive refractory symptoms should undergo appropriate and indicated workup to assess for thrombosis.
- ▶ Low-dose aspirin (80–100 mg/day) has been shown to improve vasomotor symptoms including headache in patients with MPN. In patients with aspirin-resistant symptoms, consider a twice-daily rather than once-daily regimen of low-dose aspirin or alternative anti-platelet agents (clopidogrel 75 mg/day) as monotherapy or in combination with aspirin.
- ▶ Cyto reduction or phlebotomy (if PV with elevated hematocrit) is recommended when aspirin is ineffective at relieving symptoms.
- ▶ The use of ruxolitinib improves headache in patients with PV and associated iron deficiency, more so in patients with baseline iron deficiency.
- ▶ NSAIDs should be used with caution (given concurrent aspirin use).
- ▶ Consider treatment/prophylaxis with triptans or topiramate for migraine headaches.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References on MPN-F \(3 of 3\)](#)

MPN-F
2 OF 3



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SUPPORTIVE CARE FOR PATIENTS WITH MPN

REFERENCES

- ¹ Mesa RA, Niblack J, Wadleigh M, et al. The burden of fatigue and quality of life in myeloproliferative disorders (MPDs): an international internet-based survey of 1179 MPD patients. *Cancer* 2007;109:68-76.
- ² Saini KS, Patnaik MM, Tefferi A. Polycythemia vera-associated pruritus and its management. *Eur J Clin Invest* 2010;40:828-834.
- ³ Vaa BE, Wolanskyj AP, Roeker L, et al. Pruritus in primary myelofibrosis: Clinical and laboratory correlates. *Am J Hematol* 2012;87:136-138.
- ⁴ Tefferi A, Fonseca R. Selective serotonin reuptake inhibitors are effective in the treatment of polycythemia vera-associated pruritus. *Blood* 2002;99:267.
- ⁵ Baldo A, Sammarco E, Plaitano R, et al. Narrowband (TL-01) ultraviolet B phototherapy for pruritus in polycythaemia vera. *Br J Dermatol* 2002;147:979-981.
- ⁶ Vaa BE, Tefferi A. Pruritus in primary myelofibrosis: Treatment attempts, response and outcomes. *Blood* 2013;122:4081.
- ⁷ Vaa BE, Tefferi A, Gangat N, et al. Pruritus in primary myelofibrosis: management options in the era of JAK inhibitors. *Ann Hematol* 2016;97:1185-1189.
- ⁸ Diehn F, Tefferi A. Pruritus in polycythaemia vera: prevalence, laboratory correlates and management. *Br J Haematol* 2001;115:619-621.
- ⁹ Vannucchi AM, Kiladjan JJ, Griesshammer M, et al. Ruxolitinib versus standard therapy for the treatment of polycythemia vera. *N Engl J Med* 2015;372:426-435.
- ¹⁰ Ishii T, Wang J, Zhang W, et al. Pivotal role of mast cells in pruritogenesis in patients with myeloproliferative disorders. *Blood* 2009;113:5942-5950.
- ¹¹ Harrison CN, Mead AJ, Panchal A, et al. Ruxolitinib vs best available therapy for ET intolerant or resistant to hydroxycarbamide. *Blood* 2017;130:1889-1897.
- ¹² Kirshner JJ, McDonald MC, Kruter F, et al. NOLAN: a randomized, phase 2 study to estimate the effect of prophylactic naproxen or loratadine vs no prophylactic treatment on bone pain in patients with early-stage breast cancer receiving chemotherapy and pegfilgrastim. *Support Care Cancer* 2018;26:1323-1334.
- ¹³ Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med* 2012;366:799-807.
- ¹⁴ Neben-Wittich MA, Brown PD, Tefferi A. Successful treatment of severe extremity pain in myelofibrosis with low-dose single-fraction radiation therapy. *Am J Hematol* 2010;85:808-810.
- ¹⁵ Frewin R, Dowson A. Headache in essential thrombocythaemia. *Int J Clin Pract* 2012;66:976-983.
- ¹⁶ Michiels JJ, Berneman Z, Schroyens, et al. Platelet-mediated erythromelalgic, cerebral, ocular and coronary microvascular ischemic and thrombotic manifestations in patients with essential thrombocythemia and polycythemia vera: a distinct aspirin-responsive and coumadin-resistant arterial thrombophilia. *Platelets* 2006;17:528-544.
- ¹⁷ Pascale S, Petrucci G, Dragani A, et al. Aspirin-insensitive thromboxane biosynthesis in essential thrombocythemia is explained by accelerated renewal of the drug target. *Blood* 2012;119:3595-3603.
- ¹⁸ Tefferi A, Barbui T. Essential thrombocythemia and polycythemia vera: Focus on clinical practice. *Mayo Clin Proc* 2015;90:1283-1293.
- ¹⁹ Michiels JJ. Erythromelalgia and vascular complications in polycythemia vera. *Semin Thromb Hemost* 1997;23:441-454.
- ²⁰ Verstovsek S, Harrison CN, Kiladjan JJ, et al. Markers of iron deficiency in patients with polycythemia vera receiving ruxolitinib or best available therapy. *Leuk Res* 2017;56:52-59.
- ²¹ Mesa R, Verstovsek S, Kiladjan JJ, et al. Changes in quality of life and disease-related symptoms in patients with polycythemia vera receiving ruxolitinib or standard therapy. *Eur J Haematol* 2016;97:192-200.
- ²² Billot S, Kouroupi EG, Le Guilloux J, et al. Neurological disorders in essential thrombocythemia. *Haematologica* 2011;96:1866-1869.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SPECIAL CONSIDERATIONS FOR THE USE OF JAK INHIBITORS¹

- JAK inhibitors are fedratinib, momelotinib, pacritinib, and ruxolitinib.
 - ▶ A withdrawal syndrome including fever, respiratory distress, hypotension, and multi-organ failure may occur following interruption or discontinuation of treatment. Consider tapering the dose gradually rather than discontinuing abruptly for reasons other than thrombocytopenia or neutropenia. Corticosteroids may also be considered for withdrawal syndrome.
 - ▶ Patients should have a dermatology exam at least yearly due to increased risk of non-melanoma skin cancers.
 - ▶ Check lipid panel every 6 months.
 - ▶ JAK inhibitors are associated with an increased risk of infections, including viral reactivations, and atypical organisms.

Lymphoma Risk with JAK Inhibitors in Patients with MPN:

Both low- and high-grade lymphoid neoplasms may be diagnosed concurrently with MPNs or may develop during the natural history of PV, ET, or MF. Although one report indicated an increased risk of lymphomas with JAK inhibitor therapy,² other studies found no evidence of increased lymphoma risk in patients treated with a JAK inhibitor.³⁻⁶

¹ Please refer to package insert for full prescribing information available at <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

² Porpaczy E, Tripolt S, Hoelbl-Kovacic A, et al. Aggressive B-cell lymphomas in patients with myelofibrosis receiving JAK1/2 inhibitor therapy. *Blood* 2018;132:694-706.

³ Pemmaraju N, Kantarjian H, Nastoupil L, et al. Characteristics of patients with myeloproliferative neoplasms with lymphoma, with or without JAK inhibitor therapy. *Blood* 2019;133:2348-2351.

⁴ Rumi E, Zibellini S, Boveri E, et al. Ruxolitinib treatment and risk of B-cell lymphomas in myeloproliferative neoplasms. *Am J Hematol* 2019;94:E185-E188.

⁵ Barbui T, Ghirardi A, Masciulli A, et al. Second cancer in Philadelphia negative myeloproliferative neoplasms (MPN-K). A nested case-control study. *Leukemia* 2019;33:1996-2005.

⁶ Polverelli N, Elli EM, Abruzzese E, et al. Second primary malignancy in myelofibrosis in patients treated with ruxolitinib. *Br J Haematol* 2021;193:356-368.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SPECIAL CONSIDERATIONS FOR THE USE OF INTERFERONS AND ANAGRELIDE

Special Considerations for the Use of Interferons in MPNs

- Pegylated interferon alfa-2a: Starting dose 45 mcg/week. Escalate by 45 mcg every 4 weeks (see package insert) if needed to a maximum of 180 mcg.⁷
- Ropeginterferon alfa-2b-njft: Starting dose 100 mcg every 2 weeks (50 mcg/week in patients transitioning from hydroxyurea). Escalate by 50 mcg every 2 weeks (see package insert).
 - ▶ Screen patients for a history of psychiatric disorder (eg, depression) and monitor for mood symptoms including depression and anxiety during treatment.
 - ▶ Monitor for development or exacerbation of autoimmune disorders.
 - ▶ Monitor thyroid function at baseline and annually.
 - ▶ Monitor liver enzymes at baseline and during treatment.
 - ▶ Eye examination at baseline. Patients with preexisting ophthalmologic disorders (eg, diabetic or hypertensive retinopathy) should receive periodic eye exams during treatment.

Special Considerations for the Use of Anagrelide in MPNs

- Starting dose: 0.5 mg BID, increase by 0.5 mg dose every week.⁸
- Baseline electrocardiogram (see package insert).
- Common side effects: tachycardia, headache, and dizziness are probably due to phosphodiesterase III-inhibiting properties of anagrelide. Patients with tachycardia or palpitations should be advised to reduce caffeine intake and, if possible, to increase physical exercise.
- Worsening of cardiac insufficiency has been reported, and caution should be used in patients with a previous history of cardiac failure.
- Monitor for development of post-ET myelofibrosis.

⁷ Mascarenhas J, Kosiorek HE, Prchal JT, et al. A randomized phase 3 trial of interferon- α vs hydroxyurea in polycythemia vera and essential thrombocythemia. *Blood* 2022;139:2931-2941.

⁸ Gisslinger H, Gotic M, Holowiecki J, et al. Anagrelide compared with hydroxyurea in WHO-classified essential thrombocythemia: The ANAHYDRET study, a randomized controlled trial. *Blood* 2013;121:1720-1728.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SPECIAL CONSIDERATIONS IN THE TREATMENT OF MYELOPROLIFERATIVE NEOPLASMS

Vascular Events

• Thrombosis

- ▶ The use of clinically appropriate anticoagulant therapy (eg, low-molecular-weight heparin [LMWH], direct oral anticoagulant, warfarin) is recommended for patients with active thrombosis. The initial use of anticoagulant therapy for the prevention and treatment of thrombosis should be based on the current American College of Chest Physicians (ACCP) Guidelines.^{1,2}
- ▶ Consider aspirin for patients with cardiovascular risk factors. The risks and benefits of aspirin plus anticoagulation need to be individualized on a case-by-case basis.
- ▶ There are no data to guide the selection or appropriate duration of anticoagulation with or without antiplatelet therapy in patients with PV or ET. The use of anticoagulant therapy in combination with aspirin is associated with an increased risk of bleeding compared with aspirin alone. Caution is required when using antiplatelet agents with anticoagulants for the treatment of thrombosis in patients with PV.³ The duration of anticoagulant therapy is dependent on the severity of the thrombotic event (eg, abdominal vein thrombosis vs. deep vein thrombosis), degree of disease control, and assessment of likelihood of recurrence after cessation of anticoagulant therapy.
- ▶ Low-dose aspirin can be considered for patients at high risk of thrombosis based on the arterial thrombosis score.⁴
- ▶ Assess the need for cytoreductive therapy (if not done before) and initiate cytoreductive therapy (to maintain hematocrit <45% in patients with PV) if necessary. In the presence of inadequate response, consider intensification of therapy or switch to an alternate agent. The value of cytoreduction in reducing future vascular events has not been studied in prospective, randomized controlled trials in PV.
- ▶ Plateletpheresis may be indicated in patients with ET presenting with acute life-threatening thrombosis or severe bleeding.

• Bleeding

- ▶ Rule out other potential causes and treat coexisting causes as necessary.
- ▶ Aspirin should be withheld until bleeding is under control. Consider the use of appropriate cytoreductive therapy to optimize platelet counts while minimizing hematologic and non-hematologic toxicities; see [Discussion](#).
- ▶ Coagulation tests to evaluate for acquired VWS and/or other coagulopathies are recommended for patients undergoing high-risk surgical procedures and those with elevated platelet count and/or splenomegaly or unexplained bleeding ([MPN-1](#)).
- ▶ In unanticipated gastrointestinal (GI) bleeding, particularly in the setting of splenomegaly, portal hypertension, and gastric varices, special consultation (for endoscopic evaluation) with a hepatologist or a GI specialist is recommended.

Surgery

- Multidisciplinary management with surgical and perioperative medical teams (eg, review of bleeding and thrombosis history; medication list) is recommended.
- Emergency surgery should be performed as necessary with close postoperative surveillance for the symptoms of arterial or venous thrombosis and bleeding.
- Patients with MPN are at higher risk for bleeding despite optimal management. The thrombotic and bleeding risk of the surgical procedure (eg, orthopedic and cardiovascular surgery) should be strongly considered prior to elective surgery.

[References on MPN-H \(4 of 4\)](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SPECIAL CONSIDERATIONS IN THE TREATMENT OF MYELOPROLIFERATIVE NEOPLASMS^a

Surgery (continued)

- Thrombosis and bleeding risk should be well controlled (normalization or near-normalization CBC without causing prohibitive cytopenias) prior to performing elective surgery (particularly for orthopedic surgeries or any surgical procedures associated with prolonged immobilization) with the use of appropriate anticoagulant prophylaxis and cytoreductive therapy. If surgery is associated with a high risk for venous thromboembolism (VTE) (eg, cancer surgery, splenectomy, orthopedic and cardiovascular surgery), extended prophylaxis with anticoagulation should be considered. Prophylaxis with aspirin may be considered following vascular surgery.
- In patients with PV, hematocrit should be controlled for 3 months before elective surgery (normalization or near-normalization of CBC). Additional phlebotomy may also be necessary to maintain hematocrit <45% prior to performing elective surgery.
- Aspirin should be discontinued 1 week prior to surgical procedure and restarted 24 hours after surgery or when considered acceptable depending on the bleeding risk.
- Anticoagulant therapy should be withheld (based on the half-life/type of agent) prior to surgery and restarted after surgery when considered acceptable depending on the bleeding risk.
- Cytoreductive therapy could be continued throughout the perioperative period, unless there are unique contraindications expressed by the surgical team.

Pregnancy

General Pregnancy-Related Recommendations

- Pregnancy pre-conception meeting and evaluation by high-risk obstetrician is recommended.
- All females with PV should maintain hematocrit, ideally, below the gestational range (<41% trimester 1, <38% trimester 2, <39% trimester 3).
- Hydroxyurea should be discontinued in anticipation or in the event of pregnancy. If cytoreductive therapy is needed during pregnancy, peginterferon alfa-2a^b could be considered.⁵ Potential indications include those with prior pregnancy loss or complications (pre-eclampsia), or uncontrolled leukocytosis/thrombocytosis. There are no sufficient data to establish the use of peginterferon alfa-2a in pregnancy. It should be used only if benefits outweigh potential risk to the fetus.⁵
- Direct oral anticoagulants should be avoided in breastfeeding females. Unfractionated heparin, LMWH, warfarin, and fondaparinux are all safe options in females who require anticoagulation and are breastfeeding.⁶
- Hydroxyurea is excreted in breastmilk and should be avoided in females who are breastfeeding.

^a NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. On this page, the terms males and females refer to sex assigned at birth.

^b In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) can be considered. The data for ropeginterferon alfa-2b-njft use in pregnancy are very limited. Two case reports have described the use of ropeginterferon alfa-2b-njft in pregnant women with polycythemia vera, with no adverse events reported (Okikiolu J, et al. Leuk Res Rep 2022;19:100360; Bang SY, et al. Hematol Rep 2023;15:172-179). However, safety data remain insufficient to recommend its routine use during pregnancy. According to the FDA label for ropeginterferon alfa-2b-njft, available human data do not establish the risk of birth defects, miscarriage, or adverse maternal or fetal outcomes, and reproductive toxicity studies in animals have not been conducted (refer to package insert for full prescribing information, available at: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF
All recommendations are category 2A unless otherwise indicated.

References on MPN-H (4 of 4)

MPN-H
2 OF 4



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SPECIAL CONSIDERATIONS IN THE TREATMENT OF MYELOPROLIFERATIVE NEOPLASMS

Recommendations for the Management of Standard-Risk Pregnancy

- Patients with PV or ET who become pregnant and do not have the high-risk features (standard-risk pregnancy, listed below) should take low-dose aspirin daily throughout the pregnancy and should receive prophylactic LMWH alone during the first 6 weeks of the postpartum period (unless there are patient-specific contraindications). If they were taking low-dose aspirin prior to pregnancy, it can be resumed once the postpartum course of LMWH is complete.

Definition of High-Risk Pregnancy in PV or ET⁷

Pregnancy in the setting of PV or ET is considered high risk if any of the following is present:

- Previous venous and/or arterial thrombosis with/without pregnancy
- Previous hemorrhage due to PV or ET (see management section)
- Previous pregnancy complications:
 - ▶ Unexplained death of a morphologically normal fetus ≥ 10 weeks of gestation. Premature delivery of a morphologically normal fetus at < 34 weeks of gestation due to:
 - ◊ Severe preeclampsia or eclampsia defined according to standard criteria
 - ◊ Recognized features of placental insufficiency
 - ▶ ≥ 3 unexplained consecutive miscarriages at < 10 weeks of gestation, without anatomic, hormonal, or chromosomal abnormalities
 - ▶ Unexplained intrauterine growth restriction
 - ▶ Significant antepartum or postpartum hemorrhage

Recommendations for the Management of High-Risk Pregnancy⁸⁻¹⁵

Treatment to start (continue) when pregnancy test is positive, and pregnancy is considered high risk:

- Low-dose aspirin daily
- Prophylactic LMWH throughout pregnancy and for 6 weeks postpartum^c
- Cytoreductive therapy with interferon or peginterferon alfa-2a.^b See Table 1. Mastocytosis Treatments and Pregnancy/Lactation Risk in the [NCCN Guidelines for Systemic Mastocytosis](#).

^b In the event that peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) can be considered. The data for ropeginterferon alfa-2b-njft use in pregnancy are very limited. Two case reports have described the use of ropeginterferon alfa-2b-njft in pregnant women with polycythemia vera, with no adverse events reported (Okikiolu J, et al. Leuk Res Rep 2022;19:100360; Bang SY, et al. Hematol Rep 2023;15:172-179). However, safety data remain insufficient to recommend its routine use during pregnancy. According to the FDA label for ropeginterferon alfa-2b-njft, available human data do not establish the risk of birth defects, miscarriage, or adverse maternal or fetal outcomes, and reproductive toxicity studies in animals have not been conducted (refer to package insert for full prescribing information available at: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>).

^c Administration of LMWH should be modified based on renal function, body weight, and medical history. Prophylactic LMWH should be avoided among patients with a history of MPN-related bleeding. Therapeutic anticoagulation with LMWH should be continued during pregnancy by patients receiving anticoagulation therapy after venous and/or arterial thrombotic events prior to pregnancy. Timing of LMWH and aspirin discontinuation prior to epidural and delivery and re-initiation of medications after delivery should be discussed with a high-risk obstetrician and obstetric anesthesiologist.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References on MPN-H \(4 of 4\)](#)

MPN-H
3 OF 4



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

SPECIAL CONSIDERATIONS IN THE TREATMENT OF MYELOPROLIFERATIVE NEOPLASMS

REFERENCES

- ¹ Guyatt GH, Akl EA, Crowther M, et al. Executive summary: Antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2012;141(2 Suppl):7S-47S.
- ² Kearon C, Akl EA, Ornelas J, et al. Antithrombotic therapy for VTE disease: CHEST Guideline and Expert Panel Report. Chest 2016;149:315-352.
- ³ Zwicker JJ, Paranagama D, Lessen DS, et al. Hemorrhage in patients with polycythemia vera receiving aspirin with an anticoagulant: A prospective, observational study. Haematologica 2022;107:1106-1110.
- ⁴ Pasquer H, Daltro de Oliveira R, Vasseur L, et al. Distinct clinico-molecular arterial and venous thrombosis scores for myeloproliferative neoplasms risk stratification. Leukemia 2024;38:326-339.
- ⁵ Beauverd Y, Radia D, Cargo C, et al. Pegylated interferon alpha-2a for essential thrombocythemia during pregnancy: outcome and safety. A case series. Haematologica 2016;101:e182-e184.
- ⁶ Bates SM, Rajasekhar A, Middeldorp S, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: venous thromboembolism in the context of pregnancy. Blood Adv 2018;2:3317-3359.
- ⁷ Lishner M, Avivi I, Apperley JF, et al. Hematologic malignancies in pregnancy: Management guidelines from an international consensus meeting. J Clin Oncol 2016;34:501-508.
- ⁸ Griesshammer M, Struve S, Barbui T. Management of Philadelphia negative chronic myeloproliferative disorders in pregnancy. Blood Rev 2008;22:235-245.
- ⁹ Alimam S, Bewley S, Chappell LC, et al. Pregnancy outcomes in myeloproliferative neoplasms: UK prospective cohort study. Br J Haematol 2016;175:31-36.
- ¹⁰ Barbui T, Tefferi A, Vannucchi AM, et al. Philadelphia chromosome-negative classical myeloproliferative neoplasms: revised management recommendations from European LeukemiaNet. Leukemia 2018;32:1057-1069.
- ¹¹ Maze D, Kazi S, Gupta V, et al. Association of treatments for myeloproliferative neoplasms during pregnancy with birth rates and maternal outcomes. JAMA Netw Open 2019;2:e1912666.
- ¹² Lapoirie J, Contis A, Guy A, et al. Management and outcomes of 27 pregnancies in women with myeloproliferative neoplasms. J Matern Fetal Neonatal Med 2018;1-8.
- ¹³ Bertozzi I, Rumi E, Cavalloni C, et al. Pregnancy outcome and management of 25 pregnancies in women with polycythemia vera. Am J Hematol 2018;93:E234-E235.
- ¹⁴ McMullin MFF, Mead AJ, Ali S, et al. A guideline for the management of specific situations in polycythaemia vera and secondary erythrocytosis: A British Society for Haematology Guideline. Br J Haematol 2019;184:161-175.
- ¹⁵ Skeith L, Carrier M, Robinson SE, et al. Risk of venous thromboembolism in pregnant women with essential thrombocythemia: a systematic review and meta-analysis. Blood 2017;129:934-939.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

DEFINITION OF RESISTANCE/INTOLERANCE TO HYDROXYUREA¹

Myeloproliferative Neoplasm	Definition of Resistance/Intolerance to Hydroxyurea
Polycythemia vera	1. Need for phlebotomy to keep hematocrit <45% after 3 months of at least 2 g/d of hydroxyurea, OR 2. Uncontrolled myeloproliferation (ie, platelet count >400 x 10⁹/L AND WBC count >10 x 10⁹/L) after 3 months of at least 2 g/d of hydroxyurea, OR 3. Failure to reduce massive* splenomegaly by >50% as measured by palpation OR failure to completely relieve symptoms related to splenomegaly after 3 months of at least 2 g/d of hydroxyurea, OR 4. Absolute neutrophil count <1.0 x 10⁹/L OR platelet count <100 x 10⁹/L OR hemoglobin <10 g/dL at the lowest dose of hydroxyurea required to achieve a complete or partial clinicohematologic response,[†] OR 5. Presence of leg ulcers or other unacceptable hydroxyurea-related nonhematologic toxicities, such as mucocutaneous manifestations, GI symptoms, pneumonitis, or fever at any dose of hydroxyurea
Essential thrombocythemia	1. Platelet count >600 x 10⁹/L after 3 months of at least 2 g/d of hydroxyurea (2.5 g/d in patients with a body weight >80 kg), OR 2. Platelet count >400 x 10⁹/L and WBC count <2.5 x 10⁹/L at any dose of hydroxyurea, OR 3. Platelet count >400 x 10⁹/L and hemoglobin <10 g/dL at any dose of hydroxyurea, OR 4. Presence of leg ulcers or other unacceptable mucocutaneous manifestations at any dose of hydroxyurea, OR 5. Hydroxyurea-related fever

*Organ extending by >10 cm from the costal margin.

[†]Complete response is defined as hematocrit less than 45% without phlebotomy, platelet count ≤400 x 10⁹/L, WBC count ≤10 x 10⁹/L, and no disease-related symptoms. Partial response is defined as hematocrit less than 45% without phlebotomy or response in three or more of other criteria.

¹ Reproduced with permission from Barbui T, Barosi G, Birgegard G, et al. Philadelphia-negative classical myeloproliferative neoplasms: Critical concepts and management recommendations from European LeukemiaNet. J Clin Oncol 2011;29:761-770.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

EVIDENCE BLOCKS FOR THE TREATMENT OF PRIMARY MF OR POST-PV OR POST-ET MF

Symptomatic Lower-Risk		
	First-Line Therapy	Subsequent-Line Therapy
	Useful in Certain Circumstances	
Hydroxyurea		
Pacritinib		
Peginterferon alfa-2a		
Ruxolitinib		
Momelotinib		

Higher-Risk (Platelets <50 x 10⁹/L; and not a transplant candidate or transplant not currently feasible)	
Preferred Regimen	
Pacritinib	
Other Recommended Regimen	
Momelotinib	

Higher-Risk (Platelets $\geq 50 \times 10^9/L$)		
Regimens	Not a transplant candidate or transplant not currently feasible	
	First-Line Therapy	Subsequent-Line Therapy
Fedratinib		
Ruxolitinib		
Momelotinib		
Pacritinib		

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

EVIDENCE BLOCKS FOR THE MANAGEMENT OF MF-ASSOCIATED ANEMIA

Regimens	Splenomegaly and constitutional symptoms well controlled on current JAK inhibitor	Ongoing symptomatic splenomegaly and/or constitutional symptoms
Preferred Regimen		
Momelotinib	—	
Other Recommended Regimens		
Pacritinib	—	
Ruxolitinib/luspatercept-aamt		
Ruxolitinib/epoetin alfa		
Ruxolitinib/darbepoetin alfa		
Ruxolitinib/danazol		
Fedratinib/luspatercept-aamt		—
Fedratinib/epoetin alfa		—
Fedratinib/darbepoetin alfa		—
Fedratinib/danazol		—
Momelotinib/luspatercept-aamt		—
Momelotinib/epoetin alfa		—
Momelotinib/darbepoetin alfa		—
Momelotinib/danazol		—

Regimens	Splenomegaly and constitutional symptoms well controlled on current JAK inhibitor	Ongoing symptomatic splenomegaly and/or constitutional symptoms
Other Recommended Regimens (continued)		
Pacritinib/luspatercept-aamt		—
Pacritinib/epoetin alfa		—
Pacritinib/darbepoetin alfa		—
Pacritinib/danazol		—
Useful in Certain Circumstances		
Momelotinib		—
Pacritinib		—

Regimens	No splenomegaly or constitutional symptoms
Other Recommended Regimens	
Danazol	
Darbepoetin alfa	
Epoetin alfa	
Luspatercept-aamt	
Momelotinib	
Pacritinib	
Useful in Certain Circumstances	
Lenalidomide/prednisone	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

EVIDENCE BLOCKS FOR THE TREATMENT OF POLYCYTHEMIA VERA

Management of Polycythemia Vera

Regimen	Low-Risk	High-Risk
Aspirin		

Cytoreductive Treatment

	Symptomatic Low-Risk		High-Risk	
	First-Line Therapy	Subsequent-Line Therapy	First-Line Therapy	Subsequent-Line Therapy
Preferred Regimens				
Hydroxyurea	—	—		—
Ropeginterferon alfa-2b-njft		—		—
Ruxolitinib	—		—	
Other Recommended Regimens				
Hydroxyurea			—	
Peginterferon alfa-2a				
Ropeginterferon alfa-2b-njft	—		—	
Useful in Certain Circumstances				
Ruxolitinib	—	—		—

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)

EVIDENCE BLOCKS FOR THE TREATMENT OF ESSENTIAL THROMBOCYTHEMIA

Management of Essential Thrombocythemia

Regimen	Very Low-Risk	Low-Risk	Intermediate-Risk	High-Risk
Aspirin				

Cytoreductive Treatment

	Symptomatic Very Low-Risk		Symptomatic Low-Risk		Symptomatic Intermediate-Risk		High-Risk	
	First-Line Therapy	Subsequent-Line Therapy	First-Line Therapy	Subsequent-Line Therapy	First-Line Therapy	Subsequent-Line Therapy	First-Line Therapy	Subsequent-Line Therapy
Preferred Regimen								
Ropeginterferon alfa-2b-njft	—		—		—		—	
Hydroxyurea								
Other Recommended Regimens								
Peginterferon alfa-2a								
Anagrelide		—		—		—		—
Useful in Certain Circumstances								
Anagrelide	—		—		—		—	
Ruxolitinib	—		—		—		—	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

ABBREVIATIONS

ACCP	American College of Chest Physicians	ICC	International Consensus Classification	NGS	next-generation sequencing
AML	acute myeloid leukemia	IPSET	International Prognostic Score of Thrombosis for Essential Thrombocythemia	NSAID	nonsteroidal anti-inflammatory drug
CBC	complete blood count	IPSS	International Prognostic Scoring System	OS	overall survival
CML	chronic myeloid leukemia	IWG-MRT	International Working Group-Myeloproliferative Neoplasms Research and Treatment	PFS	progression-free survival
CMV	cytomegalovirus			PMF	primary myelofibrosis
DIPSS	Dynamic International Prognostic Scoring System			PV	polycythemia vera
ELN	European LeukemiaNet	LDH	lactate dehydrogenase	RBC	red blood cell
ESA	erythropoiesis-stimulating agent	LFT	liver function test	RT-PCR	reverse transcriptase polymerase chain reaction
ET	essential thrombocythemia	LFS	leukemia-free survival		
		LMWH	low-molecular-weight heparin		
FISH	fluorescence in situ hybridization	MDS	myelodysplastic syndrome	SSRI	selective serotonin reuptake inhibitor
		MF	myelofibrosis		
		MIPSS	Mutation-Enhanced International Prognostic Scoring System	TLS	tumor lysis syndrome
G-CSF	granulocyte colony-stimulating factor	MPD	myeloproliferative disorder		
GI	gastrointestinal	MPN	myeloproliferative neoplasm(s)	VHR	very high risk
GM-CSF	granulocyte-macrophage colony-stimulating factor	MPN-SAF TSS	Myeloproliferative Neoplasm Symptom Assessment Form Total Symptom Score	VTE	venous thromboembolism
				VWFA	von Willebrand factor antigen
				VWS	von Willebrand syndrome
H&P	history and physical	MTSS	Myelofibrosis Transplant Scoring System		
HCT	hematopoietic cell transplant	MYSEC-PM	Myelofibrosis Secondary to Polycythemia Vera and Essential Thrombocythemia-Prognostic Model	WBC	white blood cell
HLA	human leukocyte antigen				
HMA	hypomethylating agent				
HMR	high molecular risk				



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

NCCN Evidence Blocks™

NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Discussion

This discussion corresponds to the NCCN Guidelines for Myeloproliferative Neoplasms. Last updated: June 16, 2026.

Table of Contents

Overview	MS-2
Guidelines Update Methodology	MS-2
Literature Search Criteria.....	MS-2
Sensitive/Inclusive Language Usage.....	MS-3
Molecular Abnormalities in MPN	MS-3
Myelofibrosis.....	MS-3
Polycythemia Vera and Essential Thrombocythemia.....	MS-4
Diagnostic Classification	MS-5
Workup of Suspected MPN	MS-7
Assessment of Symptom Burden	MS-8
Symptom Management in Patients with MPN.....	MS-8
Management of Myelofibrosis.....	MS-9
Risk Stratification	MS-9
Treatment Options	MS-11
Treatment Recommendations Based on Symptom Assessment and Risk Stratification	MS-24
Treatment Response Criteria	MS-29
Monitoring Response and Follow-up Therapy for Lower- and Higher-Risk MF	MS-30

Supportive Care	MS-32
Management of Polycythemia Vera and Essential Thrombocythemia	MS-32
Risk Stratification	MS-32
Treatment Options.....	MS-34
Treatment Recommendations Based on Risk Stratification	MS-39
Treatment Response Criteria.....	MS-41
Monitoring Response and Follow-up Therapy	MS-41
Special Considerations in the Management of MPN	MS-43
Management of Thrombosis	MS-43
Management of Bleeding.....	MS-43
Surgery	MS-43
Pregnancy.....	MS-44
Accelerated/Blast Phase MPN.....	MS-45
Treatment Options.....	MS-46
Treatment Recommendations Based on Eligibility for Transplant	MS-47
Summary.....	MS-48
References	MS-49



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Overview

Myelofibrosis (MF), polycythemia vera (PV), and essential thrombocythemia (ET) are a group of heterogeneous disorders of the hematopoietic system collectively known as Philadelphia chromosome-negative myeloproliferative neoplasms (MPN). In 2023, the estimated incidence of MF in the United States was 1.56 (95% CI, 1.38–1.77) per 100,000 person-years.¹ In a survey that assessed the incidence rates (IRs) of different subtypes of MPN in the United States (2002–2016), the age-adjusted IRs per 100,000 person-years were highest for PV (IR = 1.57) and ET (IR = 1.55) and was 0.44 for primary MF (PMF).²

MPN are characterized by a complicated symptom profile; the symptom profile varies within and between each MPN subtype, but often includes fatigue, pruritus, weight loss, and symptoms from splenomegaly.^{3–6} Variable laboratory abnormalities are observed depending on the type of MPN, including erythrocytosis, thrombocytosis, and leukocytosis, and sometimes myeloid immaturity, especially in progressive MF.⁶ A SEER-Medicare database analysis showed that patients with MPN have substantially inferior survival compared to matched controls, and the survival for patients with MF is worse than that of patients with ET or PV and significantly worse than matched controls.⁷ In addition, MPN also have the propensity for disease transformation into blast phase (MPN-BP), which is akin to acute myeloid leukemia (AML), and is associated with a poor prognosis.^{8,9}

The diagnosis and comprehensive care of patients with MPN has increased since the identification of JAK-STAT “driver” mutations (*JAK2*, *CALR*, and *MPL* mutations), and the development of therapies targeting the JAK-STAT pathway has resulted in significant improvements in disease-related symptoms and quality of life.¹⁰ However, certain aspects of clinical management regarding the diagnosis, assessment of symptom

burden, and selection of appropriate symptom-directed therapies continue to present challenges for hematologists and oncologists.¹¹

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Myeloproliferative Neoplasms were developed as a result of meetings convened by a multidisciplinary Panel with expertise in MPN, with the aim of providing recommendations for the comprehensive care of adults with these diseases. The NCCN Guidelines® for Myeloproliferative Neoplasms include recommendations for the diagnostic workup, risk stratification, treatment, and supportive care strategies for the management of MF, PV, and ET.

Guidelines Update Methodology

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Literature Search Criteria

Prior to the update of this version of the NCCN Guidelines for Myeloproliferative Neoplasms, an electronic search of the PubMed database was performed to obtain key literature in Myeloproliferative Neoplasms published since the previous Guidelines update using the following search terms: myeloproliferative neoplasms, myelofibrosis, polycythemia vera, and essential thrombocythemia. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.¹²

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Practice Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies. The data from key PubMed articles as well as articles from additional sources



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

deemed as relevant to these Guidelines as discussed by the Panel during the Guidelines update have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel's review of lower-level evidence and expert opinion.

Sensitive/Inclusive Language Usage

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation.¹³ NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-weight biased; and inclusive of individuals of all sexual orientations and gender identities. NCCN Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will continue to use the terms men, women, female, and male when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Molecular Abnormalities in MPN

JAK2 V617F mutations account for most patients with PV (>95%) and 60% of patients with ET or MF.¹⁴⁻¹⁶ *JAK2* V617F can be identified in hematopoietic stem and progenitor cells.¹⁷ The V617F mutation occurs in exon 14; however, rare insertions and deletions have been found in exon

12. *JAK2* exon 12 mutations have been described in 2% to 3% of patients with PV.^{18,19}

Activating mutations in the thrombopoietin receptor gene (*MPL* W515L/K) are reported in approximately 5% to 8% of all patients with MF and 1% to 4% of all patients with ET.²⁰⁻²²

Frameshift mutations in exon 9 of *CALR* are reported in approximately 20% to 35% of all patients with ET and MF (accounting for approximately 60%–80% of patients with *JAK2/MPL*-negative ET and MF).^{23,24} Type 1 (52 base pair deletion) and type 2 (5 base pair insertion) mutations are the most frequent *CALR* variants. *CALR* type 1 mutations are more frequent in patients with MF and *CALR* type 2 mutations are preferentially associated with ET.²⁵⁻²⁷ Type-1 like and type-2 like *CALR* mutations, which are respectively similar to the canonical 52 base pair deletion and 5 base pair deletion mutations, have been identified in patients with ET and MF.²⁸

Mutations in several other genes that are involved in signal transduction (*CBL* and *LNK/SH2B3*), chromatin modification (*TET2*, *EZH2*, *IDH1/2*, *ASXL1*, and *DNM3TA*), RNA splicing (*SF3B1*, *SRSF2*, *U2AF1*, and *ZRSR2*), and tumor suppressor function (*TP53*) have also been reported in patients with MPN.^{29,30}

Myelofibrosis

Mutations in *CALR* are associated with better overall survival (OS) than the *JAK2* V617F or *MPL* W515 mutations and the survival advantage is more pronounced in patients with a type 1/type 1-like mutation.^{9,26,31,32} In a study of 617 patients with PMF, the median OS was 18 years for those with *CALR* mutations versus 9 years for those with *JAK2* V617F mutation or *MPL* mutation and 3 years for patients with triple-negative MF.³¹ *CALR* mutations retained their prognostic significance for better OS compared to *JAK2* V617F mutation ($P = .019$) or triple-negative status ($P < .001$) in a multivariate analysis corrected for age. The 10-year cumulative incidence



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

of leukemic transformation was also lower (9%) for patients with *CALR* mutation compared to 19% for those with *JAK2* V617F mutation, 17% for those with *MPL* mutation, and 34% for those with triple-negative disease. In a study that evaluated the prognostic impact of the two different types of *CALR* mutations in 396 patients with PMF, the median survival was significantly higher for patients with type 1/type 1-like mutation than for those with type 2/type 2-like mutation or *JAK2* V617F mutation (26 vs 7 years; $P < .0001$).³² The rate of leukemic transformation was also higher among patients with type 2/type 2-like mutation than for those with type 1/type 1-like and *JAK2* V617F mutation. The presence of *JAK2* V617F mutations in patients with PMF increases the risk of thrombosis.³³ See discussion of the arterial/venous thrombosis score in the discussion section.

MPL mutations are associated with lower hemoglobin levels at diagnosis and increased risk of transfusion dependence in patients with MF.³⁴ The triple-negative mutation status (lack of all three “driver” mutations—*JAK2*, *CALR*, or *MPL*), which occurs in approximately 10% of patients, is associated with a worse prognosis in patients with MF.^{35,36}

ASXL1, *EZH2*, *SRSF2*, *TP53*, *IDH1*, *IDH2*, or *U2AF1* mutations are considered as high molecular risk (HMR) mutations and are associated with significantly shorter OS and leukemia-free survival (LFS) in patients with PMF.³⁷⁻⁴² *ASXL1*, *EZH2*, *SRSF2*, and *RAS* mutations are predictive of OS, while *ASXL1*, *SRSF2*, and *IDH1* or *IDH2* are predictive of leukemic transformation in patients with PMF.^{37-40,43,44} *TET2* or *TP53* mutations have also been associated with a worsened overall prognosis and an increased rate of leukemic transformation.^{30,41,45,46} *U2AF1* mutations have also been associated with inferior survival in patients with PMF.⁴²

In a study that evaluated the prognostic significance of somatic mutations in 879 patients with PMF, the median survival was significantly shorter (81 vs 148 months; $P < .0001$) in patients with at least one mutation in the

prognostically significant genes (*ASXL1*, *EZH2*, *SRSF2*, *IDH1*, or *IDH2*) compared with those with no mutation in any of these genes.³⁹ However, only *ASXL1* mutations retained prognostic significance after accounting for known prognostic factors. The results of a subsequent analysis that evaluated the additional prognostic value of the “number” of mutated genes in 797 patients with PMF confirmed that patients harboring ≥ 2 HMR mutations had significantly reduced OS and LFS compared not only to patients with no mutations but also to those presenting with only one HMR mutation.⁴⁰ The median OS was 3 years for patients with ≥ 2 HMR mutations compared to 7 years and 12 years, respectively, for those with one HMR mutation and no mutations. The corresponding LFS was 7 years, 11 years, and 27 years, respectively.

An analysis that assessed the impact of both *CALR* and *ASXL1* mutations on OS in 570 patients with PMF identified *CALR*(-)/*ASXL1*(+) mutational status as the most significant adverse risk factor for survival.⁴⁷ *CALR*(+)/*ASXL1*(-) was associated with the longest median OS (10.4 years) and *CALR*(-)/*ASXL1*(+) was associated with shortest median OS (2.3 years); this prognostic significance was independent of the Dynamic International Prognostic Scoring System (DIPSS)-plus risk score.

The prognostic significance of these HMR mutations, perhaps with the exception of *SRSF2* mutations, has not yet been established in patients with post-PV or post-ET MF.⁴⁸

Polycythemia Vera and Essential Thrombocythemia

JAK2 exon 12-mutated PV is characterized by significantly higher hemoglobin level and lower platelet and leukocyte counts at diagnosis compared to *JAK2* V617F-mutated PV.^{49,50} However, both *JAK2* V617F and *JAK2* exon 12 mutations are associated with similar rates of thrombosis, transformation to MF or leukemia, and death.⁴⁹ Another study reported similar findings but found no difference in hemoglobin level.⁵¹ The



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

number of deaths was also significantly higher in patients with *JAK2* V617F mutation.

CALR-mutated ET is characterized by younger age, male sex, higher platelet count, lower hemoglobin, lower leukocyte count, and lower risk of thrombosis than *JAK2*-mutated ET, whereas the presence of *MPL* mutations might be associated with a higher risk of fibrotic transformation.⁵²⁻⁵⁶ Compared to patients with *MPL*-mutated ET, patients with *CALR*-mutated ET had a lower risk of thrombosis but similar hemoglobin levels and leukocyte and platelet counts.⁵²

However, *CALR* mutations have no impact on OS or myelofibrotic or leukemic transformation.^{54,57} *CALR* mutation status also did not have a significant impact on the International Prognostic Score of Thrombosis for Essential Thrombocythemia (IPSET-thrombosis) prognostic score for predicting the risk of thrombosis.⁵⁸

Next-generation sequencing (NGS) has identified adverse variants/mutations in several other genes and may be useful to identify a minority of patients with PV and ET with increased risk of leukemic transformation.^{36,59-61} In one report, the presence of at least one of the three non-driver mutations (*ASXL1*, *SRSF2*, and *IDH2*) was associated with inferior OS and MF-free survival but it did not significantly affect the LFS in patients with PV.⁵⁹ In the multivariable analysis, *ASXL1* and *SRSF2* retained the prognostic significance for OS, and *ASXL1* was prognostic of MF-free survival. *SH2B3*, *IDH2*, *U2AF1*, *SF3B1*, *EZH2*, and *TP53* mutations were identified as significant risk factors for inferior OS, MF-free survival, and LFS in patients with ET. Multivariable analysis confirmed the individual prognostic significance of *U2AF1* mutation for OS and MF-free survival and *TP53* mutation for LFS. In one report, myelofibrotic transformation was more frequent in patients with *SF3B1* and *IDH1/2* mutations, although a persistently high or a progressive increase of the

JAK2 V617F allele burden while receiving cytoreductive therapy was the strongest predictor of myelofibrotic transformation.⁶⁰

Diagnostic Classification

The World Health Organization (WHO) classification of myeloid neoplasms was first published in 2001 and was updated in 2008 to refine the diagnostic criteria for previously described neoplasms based on new scientific and clinical information and to introduce newly recognized disease entities.^{62,63} It was revised in 2017 and once again in 2022 to incorporate new clinical, prognostic, morphologic, immunophenotypic, and genetic data that have emerged since the publication of the 2008 WHO classification.⁶⁴⁻⁶⁸

The 2022 WHO diagnostic criteria include molecular testing for *JAK2*, *CALR*, and *MPL* mutations for PMF and ET and molecular testing for *JAK2* V617F or *JAK2* exon 12 mutations for PV.⁶⁸ In the absence of *JAK2*, *CALR*, and *MPL* mutations, the presence of another clonal marker is included as one of the major diagnostic criteria for PMF. Additional mutations in *ASXL1*, *EZH2*, *TET2*, *IDH1*, *IDH2*, *SRSF2*, and *SF3B1* genes are noted to be of use in determining the clonal nature of the disease.^{39,40}

MF can present as a de novo disorder (PMF) or it can develop from the progression of PV and ET (post-PV MF or post-ET MF).⁶⁹ Prefibrotic/early-stage PMF is characterized by an increase in atypical megakaryocytes, reduced erythropoiesis, and increased age-matched bone marrow cellularity. However, overt bone marrow fibrosis might be absent in early-stage/prefibrotic PMF, leading to a diagnosis of ET.⁷⁰ The revised 2022 WHO and the 2022 International Consensus Criteria (ICC) diagnostic criteria include separate criteria for prefibrotic/early-stage PMF and overt fibrotic-stage PMF in order to differentiate true ET from prefibrotic/early PMF by the morphologic findings of the bone marrow biopsy, including the lack of reticulin fibrosis at onset.^{66,68} The 2017 WHO



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

diagnostic criteria for prefibrotic/early-stage PMF and overt fibrotic-stage PMF have also been validated in a large series of patients with pre-PMF and overt PMF.⁷¹⁻⁷³

In the International Working Group for Myeloproliferative Neoplasms Research and Treatment (IWG-MRT) study that reevaluated 1104 patients with a diagnosis of ET, central pathology review revealed a diagnosis (as defined by the WHO criteria) of ET in 891 patients (81%) and early/prefibrotic PMF in 180 patients (16%). The remaining 33 patients (3%) were unevaluable.⁷⁰ The frequency of grade 1 bone marrow fibrosis was greater in patients with early/prefibrotic PMF. In addition, leukocyte count, platelet count, serum lactate dehydrogenase (LDH) level, and the incidence of palpable splenomegaly were greater in patients with early/prefibrotic PMF, whereas hemoglobin level was greater in patients with ET. The long-term clinical outcomes were significantly worse for patients with early-stage/prefibrotic PMF. The 15-year rates of OS, leukemic transformation, and fibrotic progression were 59%, 12%, and 17%, respectively, for patients with early-stage/prefibrotic PMF. The corresponding rates were 80%, 2%, and 9%, respectively, for patients with ET. In a multivariate analysis, bone marrow histopathology remained prognostically significant for survival ($P = .03$), leukemic transformation ($P = .007$), and overt fibrotic progression ($P = .019$). Therefore, accurate evaluation of bone marrow morphology is essential to distinguish early-stage/prefibrotic PMF from ET, especially since the long-term clinical outcomes are significantly better for patients with ET than for those with prefibrotic MF.

In the 2017 WHO criteria, the diagnostic criteria for PV have been refined to differentiate masked PV from ET (recognizing the utility of bone marrow biopsy in patients with hemoglobin levels <18.5 g/dL in men and <16.5 g/dL in women).⁶⁵ In an international study of 397 patients with *JAK2* V617F or a *JAK2* exon 12 mutation and WHO-defined PV morphology,

257 patients were diagnosed with overt PV that met the full 2008 WHO diagnostic criteria for PV. The remaining 140 patients were classified as having masked PV with hemoglobin levels at diagnosis of <18.5 g/dL in males (range, 16.0–18.4 g/dL) and <16.5 g/dL in females (range, 15.0–16.4 g/dL) and frequent presence of subnormal erythropoietin (EPO) levels.⁷⁴ In a multivariate analysis, the diagnosis of masked PV was an independent predictor of poor survival in patients aged >65 years with a leukocyte count $>10 \times 10^9/L$. In the absence of these risk factors, the outcome of patients with masked PV was similar to that of patients with overt PV, suggesting that a fraction of patients with lower hemoglobin levels should still be considered as overt PV. The results of a study also showed that the OS, rates of thrombosis and major bleeding, and probability of transformation were similar among patients with masked and overt PV.⁷⁵ Thus, the major diagnostic criteria for PV have been refined to include hemoglobin levels (>16.5 g/dL in men and >16.0 g/dL in women) or hematocrit $>49\%$ in men and $>48\%$ in women and a bone marrow biopsy to confirm the age-adjusted hypercellularity.⁶⁸ Bone marrow biopsy may not be required to make the diagnosis in patients with sustained absolute erythrocytosis (hemoglobin levels >18.5 g/dL in men [hematocrit $>55.5\%$] or >16.5 g/dL in women [hematocrit $>49.5\%$]) and *JAK2* V617F or *JAK2* exon 12 mutations and subnormal serum EPO levels. However, a bone marrow biopsy in these patients can still provide helpful prognostic information.

The diagnosis of MPN should be based on the 2022 WHO and ICC diagnostic criteria.⁶⁶⁻⁶⁸ The diagnosis of PMF requires meeting all three major criteria and at least one minor criterion confirmed in two consecutive determinations as outlined in the 2022 WHO or ICC criteria.⁶⁶⁻⁶⁸ The diagnosis of PV requires meeting either all three major criteria or the first two major criteria and the minor criterion, whereas the diagnosis of ET requires meeting all four major criteria or the first three major criteria and



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

the minor criterion as outlined in the 2022 WHO or ICC criteria.^{66-68,76} See *WHO and International Consensus Classification Diagnostic Criteria for Primary Myelofibrosis*, *WHO and ICC Diagnostic Criteria for Polycythemia Vera and Post-PV Myelofibrosis*, and *WHO and ICC Diagnostic Criteria for Essential Thrombocythemia and Post-ET Myelofibrosis* in the algorithm for the lists of major and minor criteria. The diagnosis of post-PV MF or post-ET MF is based on the 2022 WHO and 2022 ICC diagnostic criteria. The 2022 ICC criteria require the documentation of a previous diagnosis of PV or ET as defined by the WHO criteria and the development of European bone marrow fibrosis grade MF-2 to MF-3 (or 3–4+, depending on the scale) and at least two additional criteria.⁶⁶

Workup of Suspected MPN

Initial evaluation of patients with suspected MPN should include a history and physical examination, palpation of spleen, evaluation of thrombotic and hemorrhagic events, cardiovascular risk factors, as well as transfusion and medication history. Laboratory evaluations should include complete blood count (CBC) with differential, microscopic examination of the peripheral blood smear, comprehensive metabolic panel with serum uric acid, serum LDH, liver function tests, serum EPO level, and serum iron studies. Human leukocyte antigen (HLA) typing should be performed for patients with MF for whom allogeneic hematopoietic cell transplant (HCT) would be considered.

Fluorescence in situ hybridization (FISH) or a multiplex reverse transcriptase polymerase chain reaction (RT-PCR), if available, on peripheral blood to detect *BCR::ABL1* transcripts and exclude the diagnosis of CML is especially recommended for patients with left-shifted leukocytosis and/or thrombocytosis with basophilia.⁶⁸ FISH may have better specificity than RT-PCR in the detection of unusual breakpoints. Some institutions prefer FISH to detect cryptic translocations while others prefer RT-PCR. The preferred method may vary based on institutional

expertise. Molecular testing on blood or bone marrow with multigene panel testing to detect driver mutations is recommended as part of initial workup for all patients.⁶⁸ The application of an NGS-based 28-gene panel in patients with MPN identified significantly more mutated splicing genes (*SF3B1*, *SRSF2*, and *U2AF1*) in patients with PMF compared to those with ET, and no mutations in splicing genes were found in patients with PV.⁷⁷ NGS is useful to establish clonality and can also identify additional mutations that hold prognostic significance.

Bone marrow aspirate with iron stain and biopsy with trichrome and reticulin stains and bone marrow cytogenetics (if inaspirable, metaphase karyotype on peripheral blood or second core or FISH for MDS- or MPN-associated abnormalities should be used on peripheral blood) are necessary to accurately distinguish the bone marrow morphologic features between the disease subtypes (early or prefibrotic PMF, ET, and masked PV).^{68,70,74} Bone marrow histology shows hypercellularity and megakaryocytic proliferation. In the case of MF, bone marrow fibrosis is demonstrated on the reticulin stain and an additional trichrome stain is recommended to distinguish grade MF-1 from MF-2 or MF-3, as outlined in the 2022 WHO diagnostic criteria.^{68,78} Progression of PV or ET to MF can only be detected by performing a bone marrow biopsy. If there is evidence of mast cell aggregates, the diagnostic workup should be performed according to the NCCN Guidelines for Systemic Mastocytosis (available at www.NCCN.org).

MPN are associated with an increased risk of major bleeding and thrombosis compared to the general population, and these events contribute considerably to morbidity and mortality in patients with MPN.^{79,80} Acquired von Willebrand syndrome (VWS) is associated with a variety of hematologic disorders, being particularly frequent in lymphoproliferative (48%) and myeloproliferative disorders (15%). Among MPN, the frequency of acquired VWS is more common among patients with ET (11%–17%)



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

but can also be seen in patients with PV.⁸¹ Testing for acquired VWS (plasma von Willebrand factor antigen [VWFA] measurement, von Willebrand activity) should be considered for patients with an elevated platelet count undergoing surgical procedures or for those with unexplained bleeding. An expanded panel including VWFA, factor VIII activity, and VWF multimers may be useful under certain circumstances.⁸²

Assessment of Symptom Burden

MPN are characterized by a complicated symptom profile resulting in reductions in quality of life, functional status, and activities of daily living.³⁻⁵ Constitutional symptoms (fever, night sweats, and weight loss) are more frequently reported in patients with MF compared to those with PV or ET.^{3,5} In a landmark survey that evaluated the symptom burden experienced by patients with MPN, disease-related symptoms were reported ≥ 1 years before diagnosis in 49% of patients with MF, 61% of patients with PV, and 58% of patients with ET.⁴ In an online survey of 669 patients with MPN, fatigue was the most frequent symptom observed in 54% of patients with MF, 45% of patients with PV, and 64% of patients with ET.⁵ Abdominal discomfort, night sweats, difficulty sleeping, shortness of breath, pruritus, bruising, loss of concentration, and dizziness were the other common symptoms and the incidences varied by disease type.

Various tools have been developed and validated in a large cohort of patients with MPN for the assessment of symptom burden.⁸³⁻⁸⁷

The Myelofibrosis Symptom Assessment Form (MF-SAF) is a 20-item tool used for the assessment of MF-associated symptoms, including fatigue, symptoms associated with splenomegaly (early satiety, abdominal pain or discomfort, inactivity, and cough), constitutional symptoms (night sweats, itching, bone pain, fever, and weight loss), and quality of life.⁸³ MF-SAF was subsequently expanded to a 27-item tool, Myeloproliferative

Neoplasm Symptom Assessment Form (MPN-SAF), to include the assessment of additional symptoms that are relevant to ET and PV (insomnia, headaches, concentration, dizziness, vertigo, lightheadedness, numbness or tingling, depression, and sexual desire dysfunction).⁸⁵

MPN-SAF was further simplified to a concise and abbreviated tool, MPN-SAF Total Symptom Score (MPN-SAF TSS; MPN-10), which is used for the assessment of the 10 most relevant symptoms in patients with MPN (fatigue, concentration, early satiety, inactivity, night sweats, itching, bone pain, abdominal discomfort, weight loss, and fever) in both clinical practice and clinical trial settings.⁸⁶ The MPN-10 score is influenced by *JAK2*, *CALR*, and triple-negative mutation status, which can be helpful for predicting survival and disease progression in patients with MPN.⁸⁸

All three symptom assessment tools are coadministered with the Brief Fatigue Inventory and symptom severity is rated by patients on a scale of 1 to 10. Assessment of symptom burden at baseline and during the course of treatment with MPN-SAF TSS (MPN-10) is recommended for all patients.^{85,86} Blood counts should also be assessed at baseline for lower- and higher-risk MF, if not previously done.

Symptom Management in Patients with MPN

Disease-related symptoms commonly contribute to decreased quality of life in patients with MPN.⁸⁹ While JAK inhibitors have been shown to broadly improve disease-related symptoms,⁹⁰⁻⁹⁸ their use is not indicated in all patients with symptomatic MPN, and the presence of specific symptoms often requires a targeted approach. Fatigue, pruritus, bone pain, headaches, and tinnitus occur across all MPN, albeit with some disease preference, and greatly impact quality of life. The optimal management of these symptoms in the setting of MPN has not been established and recommendations for symptom management as outlined in the guidelines (see *Supportive Care for Patients with MPN: Symptom*



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Management in Patients with MPN in the algorithm) are based on the subset analysis of large trials, small pilot studies, anecdotal evidence, extrapolation from other disease states, and expert opinion.

Management of Myelofibrosis

The treatment approach is currently identical for PMF and post-PV or post-ET MF. Referral to specialized centers with expertise in the management of MPN is strongly recommended for all patients diagnosed with MF.

Risk Stratification

Primary Myelofibrosis

DIPSS, DIPSS-Plus, Mutation-Enhanced International Prognostic Scoring System 70 (MIPSS-70), and MIPSS-70-Plus are prognostic scoring systems used for the risk stratification of patients with MF.⁹⁹⁻¹⁰² MIPSS-70 and MIPSS-70-Plus version 2.0 incorporate cytogenetic information and mutational status and have been developed to refine the risk stratification.¹⁰²

DIPSS is a dynamic model and has been validated for use at any point over the course of disease.¹⁰⁰ MIPSS-70 or MIPSS-70-Plus version 2.0 is preferred for the prognostic risk stratification of patients with PMF.^{102,103} Additionally, DIPSS-Plus is recommended for risk stratification at the time of treatment if molecular testing is not available¹⁰¹ and DIPSS can be used if recent karyotyping is not available.¹⁰⁰

DIPSS

In a subsequent analysis that evaluated the impact of each adverse factor on survival during follow-up after treatment, all variables retained statistical significance. However, the development of anemia over time significantly affected survival (hazard ratio [HR] was approximately double that of other adverse factors).¹⁰⁰ Thus, a modified risk stratification system (DIPSS)

was developed using the same prognostic variables as in IPSS (age >65 years, presence of constitutional symptoms, hemoglobin level <10 g/dL, leukocyte count >25 x 10⁹/L, and circulating blast cells ≥1%), but two points were assigned for hemoglobin <10 g/dL. The DIPSS can be applied at any point during the disease course to stratify patients into four different risk groups: low risk (0 adverse points), intermediate-1 risk (1 or 2 points), intermediate-2 risk (3 or 4 points), and high risk (5 or 6 points) with the median survival rates of not reached, 14 years, 4 years, and 1.5 years, respectively.¹⁰⁰

DIPSS-Plus

In subsequent reports, the need for red blood cell (RBC) transfusion, platelet count, and unfavorable karyotype have been identified as additional IPSS- and DIPSS-independent prognostic factors for inferior OS and LFS in patients with PMF.¹⁰⁴⁻¹⁰⁷ The median survival of patients with DIPSS low-risk disease with thrombocytopenia or unfavorable karyotype was 6.5 years compared to >15 years in the absence of these two additional risk factors.¹⁰¹ Similarly, the median survival was <1.5 years for patients with DIPSS high-risk disease with ≥1 of these additional prognostic factors compared to approximately 3 years for those patients without these prognostic factors.¹⁰¹

DIPSS was modified into DIPSS-Plus by the incorporation of platelet count <100 x 10⁹/L, RBC transfusion need, and unfavorable karyotype [complex karyotype or one or two abnormalities that include trisomy 8, del(7/7q), i(17q), del(5/5q), del(12p), inv(3), or 11q23 rearrangement].¹⁰¹ DIPSS-Plus also stratifies patients into four risk groups based on the aforementioned eight risk factors: low risk (no risk factors), intermediate-1 risk (one risk factor), intermediate-2 risk (two or three risk factors), and high risk (≥4 risk factors) with respective median survival rates of 15.4, 6.5, 2.9, and 1.3 years, respectively. To calculate the DIPSS-Plus score, clinicians must first calculate the DIPSS score. Points are assigned as follows: 0 for



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

DIPSS low risk, 1 for DIPSS intermediate-1 risk, 2 for DIPSS intermediate-2 risk, and 3 for DIPSS high risk. One point each is then added for platelets $<100 \times 10^9/L$, RBC transfusion need, and unfavorable karyotype.

MIPSS-70 and MIPSS-70-Plus

In a study of 805 patients with PMF (aged ≤ 70 years), in a multivariate analysis, hemoglobin level <10 g/dL, leukocyte count $>25 \times 10^9/L$, platelet count $<100 \times 10^9/L$, circulating blast cells $\geq 2\%$, bone marrow fibrosis grade $\geq$ MF-2, constitutional symptoms, absence of *CALR* type-1 mutation, and presence of ≥ 2 HMR mutations (*ASXL1*, *EZH2*, *SRSF2*, and *IDH1/2*) were identified as independent predictors of inferior OS.¹⁰² This mutation-informed (MIPSS-70) prognostic model (without the cytogenetic information) stratified patients into three risk categories (low risk, intermediate risk, and high risk) with a median OS of 28 years, 7 years, and 2 years, respectively. The 5-year OS rates were 95%, 70%, and 29%, respectively. The MIPSS-70-Plus prognostic model, which included cytogenetic information but omitted bone marrow fibrosis grade and leukocyte and platelet counts, stratified patients into four risk categories (low risk, intermediate risk, high risk, and very high risk) with 5-year OS rates of 91%, 66%, 42%, and 7%, respectively. The MIPSS-70-Plus version 2.0 prognostic model accounted for very-high-risk (VHR) karyotype, included *U2AF1* Q157 as an HMR mutation, and specified new hemoglobin thresholds with adjustments for sex and severity.¹⁰³ It stratified patients into five risk categories (very low risk, low risk, intermediate risk, high risk, and very high risk) with a median OS of not reached, 10.3 years, 7.0 years, 3.5 years, and 1.8 years, respectively, for patients of all ages. The 10-year survival rates were 86%, 50%, 30%, 10%, and $<3\%$, respectively.

Post-PV MF and Post-ET MF

MIPSS-70-Plus version 2.0 (preferred), DIPSS-Plus, and Myelofibrosis Secondary to PV and ET-Prognostic Model (MYSEC-PM) are recommended for the risk stratification of post-PV or post-ET MF.¹⁰⁸⁻¹¹¹

The MYSEC-PM is a prognostic model that stratifies patients with post-PV or post-ET MF into four risk groups, with distinct survival outcomes (low risk, intermediate-1, intermediate-2, and high risk) based on age, hemoglobin level (<11 g/dL), circulating blasts ($\geq 3\%$), *CALR* mutation status, platelet count ($<150 \times 10^9/L$), and constitutional symptoms.¹¹⁰ The median survival was not reached, 9 years, 4 years, and 2 years, respectively. Palandri et al¹¹¹ validated the MYSEC-PM model in post-PV and post-ET MF. The model was successfully used to stratify patients into different risk categories, while the IPSS could not. Spleen responses and hematologic toxicities also differed based on the predicted risk. In a retrospective analysis of cytogenetic data from 376 patients with post-PV and post-ET MF, a significant association was uncovered between abnormal karyotypes and higher MYSEC-PM risk categories ($P = .006$).¹¹² However, patients with a monosomal karyotype had a lower chance of survival that was independent of the MYSEC-PM stratification.

The inclusion of HMR mutations and unfavorable karyotype as prognostic variables can improve the risk prognostication.^{108,109} Using Harrell's C-index, Guerra et al compared PMF prognostic scoring systems in patients with post-PV or post-ET MF.¹⁰⁸ Both MIPSS-70 and MIPSS-70-Plus version 2.0 had a high performance score (C-index of 0.731 and 0.794, respectively), with MIPSS-70-Plus version 2.0 outperforming MYSEC-PM (C-index, 0.731). IPSS (C-index, 0.725) and DIPSS (C-index, 0.694) had lower performance scores compared to MYSEC-PM. Another study showed that DIPSS-Plus also outperformed MYSEC-PM in patients with post-PV or post-ET MF (C-index of 0.887 vs 0.806, respectively).¹⁰⁹ Unlike the previous study, which reported a lower performance of IPSS and



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

DIPSS compared to MYSEC-PM, this study reported a higher performance of IPSS and DIPSS over MYSEC-PM (C-index of 0.858 and 0.849 vs 0.806, respectively).

Treatment Options

Interferons

Interferons may demonstrate activity in low-risk MF¹¹³⁻¹¹⁵ but they are generally not recommended for higher-risk disease.

In a retrospective study of 62 patients with early MF treated with peginterferon alfa-2a, improvement in constitutional symptoms and complete resolution of thrombocytosis and leukocytosis were observed in 82%, 83%, and 69% of patients, respectively, and a reduction of splenomegaly was seen in 47% of patients.¹¹³ Ianotto et al¹¹⁴ reported an improved OS compared to the reference cohorts used to determine DIPSS scores (intermediate-2: 6.9 vs 4 years and high risk: 4.58 vs 1.5 years). A reduction of >50% in the *JAK2* V617F allele burden was observed in 58.8% of patients; the presence of ≥1 additional mutation(s) was associated with worse OS and LFS.

In a prospective trial of 30 patients (21 patients with PMF, 7 patients with post-PV MF, and 2 patients with post-ET MF), treatment with interferon alfa-2b or peginterferon alfa-2a resulted in an overall response rate (ORR) of 73% (7% complete response [CR], 30% partial response [PR], 13% clinical improvement, and 23% of patients had stable disease [SD]).¹¹⁵ The corresponding response rates were 3%, 27%, 6%, and 13%, respectively, for patients with low-risk disease. Among patients with marked splenomegaly, spleen response (≥50% reduction in spleen size) was observed in 40% of patients (4 out of 10) and 60% of patients (6 out of 10) had either a slight decrease in spleen size or stable spleen size. Among the 25 patients with evaluable bone marrow biopsies, reduction in bone marrow cellularity and reductions of reticulin fibrosis were

observed in 12 patients and 5 patients, respectively, after a median treatment duration of 6 years. The presence of HMR mutations or ≥3 mutations was associated with inferior response rates and the survival rates were better for patients without *ASXL1* mutation; the 5-year progression-free survival (PFS) and OS rates were 88% and 92%, respectively.

The combination of interferons with JAK inhibitors is under investigation in clinical trials. The phase II COMBI study, which evaluated the efficacy of combined ruxolitinib and low-dose pegylated interferon alfa-2 in 32 patients with PV and 18 patients with primary or secondary MF, reported a remission rate of 31% in patients with PV and 44% in patients with primary or secondary MF at 2 years, as determined by the 2013 European LeukemiaNet (ELN) and IWG-MRT response criteria.¹¹⁶ Forty-six patients previously had disease that was intolerant of, or refractory to pegylated interferon alfa-2. Reductions in symptom burden (22 to 15) as assessed by the MPN-SAF TSS and in the median *JAK2* V617F allele burden (47% to 12%) were also obtained. The main grade 3–4 hematologic adverse events reported were anemia (14.0%), thrombocytopenia (4.0%), and leukopenia (2.0%) and the main grade 3–4 nonhematologic adverse events were pneumonia (12.0%), hypertension (6%), and gastrointestinal bleeding (6%). Data from the phase I/II RUXOPEG trial demonstrated reduction of ≥50% in spleen length in 70% of patients within 24 weeks in the intention-to-treat population in patients with MF treated with ruxolitinib and pegylated interferon alfa-2a.¹¹⁷ A reduction in the *JAK2* V617F allele burden was also reported (mean of 84% at baseline to 65% and 53% after 6 and 12 months, respectively).

Fedratinib

Fedratinib is a potent and selective JAK2 and FLT3 inhibitor approved by the U.S. Food and Drug Administration (FDA) for the treatment of



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

intermediate-2 or high-risk MF as determined by IPSS, based on the results of the randomized phase III JAKARTA trial, as well as the non-randomized phase II JAKARTA-2 trial, which evaluated efficacy in patients with ruxolitinib-resistant or ruxolitinib-intolerant intermediate-1, intermediate-2, or high-risk MF.^{93,98}

The phase III JAKARTA trial randomized patients with intermediate-2-risk or high-risk MF (PMF, post-PV MF, or post-ET MF) with platelet counts $\geq 50 \times 10^9/L$ to once-daily fedratinib 400 mg (n = 96) or placebo (n = 96).⁹⁸ Patients with progressive disease (PD) were permitted to cross over from placebo to fedratinib. The proportion of patients achieving the primary endpoint (spleen response; $\geq 35\%$ reduction in spleen volume as assessed by MRI or CT scan at 24 weeks and confirmed 4 weeks later) was significantly higher ($P < .0001$) in the 400 mg fedratinib group (37% [95% CI, 27%–46%]) than in the placebo group (1% [95% CI, 0%–3%]). The symptom response rates at 24 weeks ($\geq 50\%$ reduction in the MF-SAF-TSS from baseline) in evaluable patients were 40% (95% CI, 30%–51%) and 9% (95% CI, 3%–15%), respectively, for the 400 mg and placebo groups.

Seventy-four percent of patients initially in the placebo group crossed over to fedratinib during the study.¹¹⁸ The median OS was not reached in either group (HR, 0.57; 95% CI, 0.30–1.10; $P = .094$). The survival rates at 1 year and 18 months were 92% and 87%, respectively, for the fedratinib group, and 86% and 80%, respectively, for the placebo group. Patients treated with fedratinib had significantly longer median PFS (23.2 vs 17.5 months) (HR, 0.42; 95% CI, 0.23–0.76; $P = .004$). The PFS rates were 83% for the fedratinib group and 67% for the placebo group at 1 year. A subsequent analysis of the JAKARTA study showed that the baseline platelet count did not significantly impact the rate of spleen response ($P = .37$), which was 36% in patients with platelet count 50 to $<100 \times 10^9/L$ (N = 14) and 49% in patients with platelet count $\geq 100 \times 10^9/L$ (N = 82) who

were treated with 400 mg daily fedratinib at 24 weeks.¹¹⁹ Similar results were obtained for the rates of symptom response (33% in the first group and 42% in the second group; $P = .57$).

The phase II non-randomized JAKARTA-2 trial (n = 97) showed that fedratinib 400 mg was also effective in reducing splenomegaly and symptom burden in patients with ruxolitinib-resistant or ruxolitinib-intolerant intermediate-1-risk or intermediate-2-risk/high-risk MF (PMF, post-PV MF, or post-ET MF, palpable splenomegaly [≥ 5 cm below the left costal margin], and platelet count $\geq 50 \times 10^9/L$).⁹³ Patients were assigned by treating investigators as resistant or intolerant to ruxolitinib. Spleen response ($\geq 35\%$ reduction in spleen volume as assessed by MRI or CT scan at 24 weeks; 83 evaluable patients) and symptom response ($\geq 50\%$ reduction in the MF-SAF-TSS at 24 weeks; 90 evaluable patients) were achieved in 55% (53% in the ruxolitinib-resistant group and 63% in the ruxolitinib-intolerant group) and 26% (21% in the ruxolitinib-resistant group and 32% in the ruxolitinib-intolerant group) of patients, respectively. Another analysis of the JAKARTA-2 study reported the efficacy data in three different cohorts of patients (intent-to-treat population, n = 97; stringent criteria cohort, n = 79; and sensitivity analysis cohort, 66 patients treated with 6 cycles of fedratinib or discontinued before cycle 6 for reasons other than study closure) by using updated criteria for ruxolitinib failure and intolerance.¹²⁰ The spleen response rates were 31%, 30%, and 36%, respectively, for these three cohorts. The corresponding symptom response rates were 27%, 27%, and 32%, respectively. At the end of the study, 81% of patients were censored for survival.¹¹⁸ The median OS was not reached (95% CI, 17.1 months–not reached) and the survival rates at 1 year and 18 months were 84% and 67%, respectively. The median PFS was 13.3 months (95% CI, 8.4–17.1 months) and the PFS rate at 1 year was 59%. A subgroup analysis of the JAKARTA2 study showed the baseline platelet count did not significantly impact the rate of spleen response ($P = .41$), which was 36% in patients with platelet count 50 to



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

<100 x 10⁹/L (N = 33) and 28% in patients with platelet count ≥100 x 10⁹/L (N = 64) at 24 weeks.¹¹⁹ The rate of symptom response was 39% in the former group and 20% in the latter group (*P* = .06). Post hoc analyses from the JAKARTA and JAKARTA2 trials determined that treatment with fedratinib (400 mg daily) was not associated with clinically significant weight gain or an increase in the body mass index.¹²¹

The phase IIIb FREEDOM trial evaluated the efficacy and safety of fedratinib at a dose of 400 mg daily in patients with DIPSS intermediate- or high-risk PMF or post-PV/ET MF who were previously treated with ruxolitinib.¹²² At the end of cycle 6, 25.7% of evaluable patients achieved the primary endpoint of ≥35% reduction in spleen volume and 44.4% achieved the secondary endpoint of ≥50% reduction in total symptom score.

In the phase III FREEDOM2 trial, patients with intermediate-2 or high-risk MF that relapsed or that was refractory or intolerant to ruxolitinib were randomized to receive fedratinib (n = 134) (or best available therapy [BAT]; n = 67).¹²³ Patients were allowed to cross over from the BAT group to the fedratinib group after response assessment following cycle 6 or if there was confirmed progressive disease. 36% of patients in the fedratinib arm achieved the primary endpoint of ≥35% spleen volume reduction compared to 6% of patients in the control arm (*P* < .0001). At the completion of cycle 6, 34% of patients treated with fedratinib had a ≥50% decrease in total symptom score vs. 17% of those treated with BAT (*P* = .0033).

Toxicity

Anemia and thrombocytopenia were the most common hematologic toxicities associated with fedratinib.^{93,98,122,123} In the JAKARTA trial, ≥grade 3 anemia was reported in 30% of patients.⁹⁸ In an analysis of the JAKARTA-2 trial, grade 3 or 4 anemia was reported in 46% of patients and thrombocytopenia in 24% of patients.⁹³ A pooled analysis of the

JAKARTA/JAKARTA2/ARD11936 cohorts revealed a higher percentage of grade 3–4 treatment-emergent thrombocytopenia (40% for platelet count 50 to <100 x 10⁹/L [N = 48] and 5% for platelet count ≥100 x 10⁹/L [N = 155]) in patients treated with 400 mg daily fedratinib.¹¹⁹

Diarrhea, vomiting, and nausea were the most common nonhematologic toxicities and usually abated after the first 28-day cycle.^{93,98} Fedratinib has demonstrated inhibition of FLT3, which has been implicated in the occurrence of these gastrointestinal toxicities.^{124,125} Elevation of liver enzymes or creatinine levels were more frequent with fedratinib than with placebo.⁹⁸ Fedratinib was also associated with a higher rate of infections (42% for fedratinib 400 mg compared to 27% in the placebo group).¹²⁶

In the phase IIIb FREEDOM trial, grade 3/4 anemia and thrombocytopenia occurred in 39.5% and 23.7% of patients, respectively.¹²² Grade 3 gastrointestinal adverse events were also reported in 15.8% of patients. Data also suggest that early treatment with gastrointestinal prophylactic agents may help to mitigate the rates of gastrointestinal adverse events. No cases of Wernicke encephalopathy (WE) were observed.

In the phase III FREEDOM2 trial, 40% of patients in the fedratinib arm experienced grade ≥3 treatment-related adverse events compared to 12% in the control group, with anemia (9% in both arms) and thrombocytopenia (12% in the fedratinib arm vs 3% in the control arm) being the most common adverse events.¹²³ Mitigation strategies with prophylactic agents helped to manage gastrointestinal adverse events and thiamine deficiency. One case of suspected grade 1 WE occurred in the third cycle in the fedratinib arm; however, this was resolved following thiamine supplementation.

In August 2017, the FDA removed the clinical hold on the fedratinib development program, which was initially placed in 2013 because 8 out of 670 patients in fedratinib clinical trials experienced symptoms suggestive



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

of WE, which is a neurological disorder that develops in the setting of thiamine deficiency.¹²⁷ A subsequent report showed that fedratinib does not increase the risk of thiamine deficiency beyond its potential to worsen malnutrition, which could be due to poor management of preventable gastrointestinal adverse events.¹²⁷ In the JAKARTA2 study, only one case of encephalopathy was reported, which was subsequently determined to be related to hepatic encephalopathy and inconsistent with WE.¹²⁰ In 670 patients enrolled in clinical trials evaluating fedratinib in patients with MPN or solid tumors, the overall prevalence of WE was observed in <1% of treated patients,¹²⁷ and thus was not found to be clearly different than the 1% to 2% prevalence of WE in the general U.S. population.¹²⁸

As a result of these updated analyses, the FDA approved fedratinib in 2019 for the treatment of patients with intermediate-2-risk or high-risk MF (PMF, post-PV MF, or post-ET MF). The prescribing information for fedratinib includes a boxed warning regarding the potential risk of encephalopathy, including WE. See the prescribing information for monitoring of thiamine levels.

Ruxolitinib

Ruxolitinib is a potent and selective JAK1 and JAK2 inhibitor that is FDA-approved for the treatment of intermediate-risk or high-risk MF as determined by IPSS, based on the results of phase III studies (COMFORT-I and COMFORT-II).^{90,129} The COMFORT studies did not include patients with low-risk or intermediate-1-risk MF, and the use of ruxolitinib in this patient population is based on the evidence from retrospective analysis and non-randomized clinical studies as discussed below.¹³⁰⁻¹³³

Lower-Risk MF

The efficacy of ruxolitinib in low-risk MF has not been evaluated in prospective clinical trials. The results from a retrospective analysis suggest that ruxolitinib may be an appropriate treatment option for

symptomatic patients with low-risk MF.¹³⁰ In this retrospective analysis of 108 patients (25 patients with low-risk MF and 83 patients with intermediate-1-risk MF) treated with ruxolitinib, patients with low-risk MF experienced a substantial improvement in splenomegaly and constitutional symptoms. The proportion of patients with moderate to severe splenomegaly reduced from 64% at the time of diagnosis to 16% at the time of best response to ruxolitinib. The proportion of patients with moderate or severe fatigue decreased from 90% at the time of diagnosis to 37% at the time of best response to ruxolitinib.

The safety and efficacy of ruxolitinib in patients with intermediate-1-risk MF have been demonstrated in a retrospective analysis¹³⁰ and nonrandomized studies.¹³¹⁻¹³³ In the retrospective analysis (discussed above), among the 83 patients with intermediate-1-risk MF, the proportion of patients with moderate or severe splenomegaly decreased from 53% at the time of diagnosis to 10% at the time of best response to ruxolitinib, and the proportion of patients with moderate or severe fatigue decreased from 76% at the time of diagnosis to 42% at the time of best response to ruxolitinib.¹³⁰

The ROBUST trial is an open-label phase II trial that evaluated the efficacy of ruxolitinib in patients with intermediate-1-risk MF (48 patients; 14 patients with intermediate-1-risk MF along with 13 patients with intermediate-2-risk MF and 21 patients with high-risk MF).¹³¹ The primary composite endpoint was the achievement of treatment success at 48 weeks after ruxolitinib therapy (≥50% reduction in palpable spleen length and/or a ≥50% decrease in MF-SAF). At 48 weeks, 47% of the overall population achieved a reduction in mean palpable spleen length and the effect was seen across all risk groups (52% of patients with intermediate-1-risk, 37% of patients with intermediate-2-risk, and 49% of patients with high-risk disease). A ≥50% reduction in MF-SAF at 48 weeks was achieved in 20.8% of patients in the overall population and



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

across all risk groups (intermediate-1 risk, 21%; intermediate-2 risk, 23%; high risk, 19%). Improvements in MF-SAF were seen in 80%, 73%, and 72% of patients with intermediate-1-risk, intermediate-2-risk, and high-risk disease, respectively.

JUMP is an expanded-access phase III study designed to assess the safety and efficacy of ruxolitinib in patients with intermediate-2-risk or high-risk MF with or without splenomegaly or intermediate-1-risk MF with a palpable spleen (≥ 5 cm from the costal margin).¹³⁴ The JUMP study comprised 2087 patients with platelet count $\geq 100 \times 10^9/L$ and 138 patients with platelet count $< 100 \times 10^9/L$. A primary analysis revealed that at 24, 48, and 96 weeks, 56.5%, 61.4%, and 66.5% of evaluable patients achieved a $\geq 50\%$ reduction from baseline in palpable spleen length, respectively. At the same time points, 23.3%, 18.9%, and 14.3% of patients had a 25% to $< 50\%$ reduction from baseline in palpable spleen length, respectively. Of evaluable patients with platelet count $< 100 \times 10^9/L$, 38.4% and 31.9% achieved a $\geq 50\%$ reduction from baseline in palpable spleen length at 24 and 48 weeks, respectively. The most common grade 3 or 4 hematologic adverse events were anemia and thrombocytopenia in patients with platelet count $\geq 100 \times 10^9/L$ (34.7% and 17.1%, respectively) and in patients with platelet count $< 100 \times 10^9/L$ (35.5% and 54.3%, respectively). The most common grade 3 or 4 nonhematologic adverse events were pneumonia (4.6%), pyrexia (2.3%), and asthenia (2.2%) in patients with platelet count $\geq 100 \times 10^9/L$ and pneumonia (5.8%), pyrexia (3.6%), and dyspnea (3.6%) in patients with platelet count $< 100 \times 10^9/L$. At 96 weeks, the estimated OS and PFS (per IWG-MRT criteria) probabilities were 87% (95% CI, 85%–89%) and 81% (95% CI, 78%–83%), respectively. Treatment with ruxolitinib also led to the amelioration of symptoms. A multivariate analysis determined that IPSS low/intermediate-1 risk category (43.1% vs 30.6% for IPSS intermediate-2/high-risk category; adjusted odds ratio [AOR], 0.65; 95% CI, 0.44–0.95), use of ruxolitinib in the first-line setting (40.2% vs 31.5%

for use in subsequent-line setting; AOR, 0.53; 95% CI, 0.38–0.75), and a total daily dose of > 20 mg/day at 12 weeks (41.3% vs 30.4% for < 20 mg/day; AOR, 0.47; 95% CI, 0.33–0.68) were associated with a higher spleen response rate.¹³⁵ However, no association was found with symptom response rate.

In another study that evaluated efficacy and safety of ruxolitinib in 70 patients with intermediate-1-risk MF, the rates of spleen and symptom response at 6 months were 55% and 80%, respectively. The majority of patients (83%) were still on therapy after a median follow-up of 27 months.¹³³

Higher-Risk MF

The results of COMFORT-I^{90,136,137} and COMFORT-II^{129,138,139} studies demonstrated that continuous ruxolitinib therapy was associated with significant clinical benefits in patients with MF in terms of reduction in spleen size, amelioration of disease-related symptoms, and improvement in quality of life and OS compared to either placebo or BAT for patients with intermediate-2-risk or high-risk MF (PMF, post-PV MF, or post-ET MF).

The COMFORT-I trial randomized 259 patients with intermediate-2-risk or high-risk MF to twice-daily ruxolitinib ($n = 155$) or placebo ($n = 154$).⁹⁰ The starting dose of ruxolitinib was based on the baseline platelet count (15 mg twice daily for a platelet count $100 \times 10^9/L$ to $200 \times 10^9/L$ and 20 mg twice daily for $> 200 \times 10^9/L$) and patients with protocol-defined worsening splenomegaly were permitted to cross over from placebo to ruxolitinib. The primary endpoint ($\geq 35\%$ reduction in spleen volume as assessed by MRI at 24 weeks) was reached in 42% of patients in the ruxolitinib group as compared with 0.7% in the placebo group ($P < .001$). An improvement of $\geq 50\%$ in the MF-SAF at 24 weeks was seen in 46% of patients treated with ruxolitinib as compared with 5% of patients who received placebo ($P < .001$). Long-term follow-up results confirmed the safety and durable



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

efficacy of ruxolitinib for the treatment of patients with intermediate-2-risk or high-risk MF.^{136,137} The 5-year follow-up data showed that patients treated with ruxolitinib had prolonged median OS compared to placebo (not reached compared to 200 weeks for patients randomized to placebo; HR, 0.69; 95% CI, 0.50–0.96; $P = .025$).¹³⁷ Spleen response ($\geq 35\%$ reduction from baseline in spleen volume) was achieved in 59% of patients randomized to ruxolitinib and the median duration of spleen response was 168 weeks. At the time of this analysis, 111 patients from the placebo group had crossed over to ruxolitinib (median time to crossover was 40 weeks). The subgroup analyses showed that clinical benefit of ruxolitinib was seen across all patient subgroups including PMF, post-ET MF or post-PV MF, IPSS risk groups, and *JAK* mutation status (positive or negative), and there was also a non-significant trend toward longer OS for patients with IPSS intermediate-2-risk and high-risk MF treated with ruxolitinib. However, this study was not designed or powered to detect treatment efficacies between treatment arms within each subgroup.^{137,140}

In the COMFORT-II study, 219 patients with intermediate-2-risk or high-risk MF were randomized to ruxolitinib ($n = 146$) or BAT ($n = 73$).¹²⁹ The primary endpoint was at least a 35% reduction in spleen volume as assessed with MRI or CT scan at 48 weeks. The starting dose of ruxolitinib was based on the baseline platelet count (15 mg twice daily if the platelet count was $\leq 200 \times 10^9/L$ and 20 mg twice daily if the platelet count was $> 200 \times 10^9/L$). A total of 28% of the patients in the ruxolitinib arm had a $\geq 35\%$ reduction in spleen volume at 48 weeks compared with 0% in the group receiving BAT ($P < .001$). The median duration of response among patients treated with ruxolitinib was not reached, with 80% of patients still having a response at a median follow-up of 12 months.¹²⁹ Patients receiving ruxolitinib had improved quality of life and role functioning as well as significant reductions in disease-related symptoms compared to those receiving BAT. Long-term follow-up results

confirmed that ruxolitinib is associated with durable efficacy and survival benefit compared to BAT for patients with intermediate-2-risk or high-risk MF.^{138,139} At the time of the 5-year final analysis, 53% of patients in the ruxolitinib arm achieved a $\geq 35\%$ reduction in spleen volume at any time on treatment, and spleen volume reductions of $\geq 35\%$ were sustained with long-term therapy (median duration, 3 years).¹³⁹ The median OS was not reached for patients in the ruxolitinib arm, and it was 4 years for those in the BAT arm.

The pooled analysis of COMFORT-I and COMFORT-II studies showed that patients with intermediate-2-risk or high-risk MF treated with ruxolitinib had prolonged OS, and the OS of patients with high-risk disease in the ruxolitinib group was similar to that of patients with intermediate-2-risk MF in the control group.¹⁴¹ Larger spleen size at baseline was associated with shortened survival, whereas any spleen volume reductions ($> 10\%$ reduction in spleen size) and a palpable spleen length reduction of $\geq 25\%$ correlated with longer survival. Verstovsek et al¹⁴² also determined that compared to patients who had a decrease of $< 25\%$ in spleen length, those with a $\geq 50\%$ decrease had significantly improved survival (HR, 0.223; 95% CI, 0.097–0.512; $P = .0001$).

The European Registry for Myeloproliferative Neoplasms: Toward a Better Understanding of Epidemiology, Survival, and Treatment (ERNEST) study enrolled patients with PMF or post-PV/ET MF.¹⁴³ At enrollment, 10.7% of patients had received treatment with ruxolitinib and 48.2% of patients had received treatment with hydroxyurea only. Sixty-four percent of patients treated with ruxolitinib had received treatment with hydroxyurea. Analysis of the real-world data revealed an improved median OS with ruxolitinib compared to those treated with hydroxyurea (6.7 vs 5.1 years; $P = .001$). A propensity score matching analysis also demonstrated an improved median OS in patients treated with ruxolitinib (7.7 years) as first-line



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

therapy or second-line therapy after hydroxyurea compared to those treated with hydroxyurea only (3.4 years; $P = .002$).

Toxicity

Anemia and thrombocytopenia were the most common hematologic toxicities associated with ruxolitinib, consistent with its mechanism of action, and the incidences of grade 3/4 anemia or thrombocytopenia were higher during the first 8 to 12 weeks of treatment.^{90,129,132} In the COMFORT-I study, ecchymosis, dizziness, and headache were the most frequent nonhematologic toxicities associated with ruxolitinib, and diarrhea was the most frequent nonhematologic adverse event associated with ruxolitinib in the COMFORT-II study.^{90,129} In general, the incidences of nonhematologic toxicities decreased with long-term therapy.^{136,139} Anemia associated with ruxolitinib treatment may not share the inferior prognosis of disease-related anemia as ruxolitinib can overcome the inferior prognosis of disease-induced anemia.¹⁴⁴ A study by Cervantes et al¹⁴⁵ suggests that an alternative dosing strategy for ruxolitinib consisting of a dose of 10 mg twice daily for 12 weeks and titrating up to a dose of 25 mg twice daily was well-tolerated and effective in patients with PMF or post-PV/ET MF and anemia.

Management of Treatment-Related Anemia and Thrombocytopenia

In the COMFORT-I and COMFORT-II studies, anemia and thrombocytopenia were managed with dose modifications and RBC transfusions.^{90,129} Patients enrolled in the COMFORT trials were required to have a baseline platelet count $\geq 100 \times 10^9/L$, and the initial starting dose of ruxolitinib was dependent on the patient's baseline platelet counts.^{90,129} The results of a phase II study suggest that a lower initial dose of ruxolitinib (5 mg twice daily with optional escalation up to 15 mg twice a day) may be appropriate in patients with baseline platelet count 50 to $100 \times 10^9/L$.¹⁴⁶ In the dose-finding phase Ib EXPAND study, ruxolitinib was tolerated at a maximum safe starting dose of 10 mg twice daily in patients

with MF with platelet count 50 to $74 \times 10^9/L$ or 75 to $99 \times 10^9/L$.¹⁴⁷ Patients with platelet count of 75 to $99 \times 10^9/L$ displayed higher tolerability. At 48 weeks, 33.3% of patients with platelet count 75 to $99 \times 10^9/L$ demonstrated a spleen response compared to 30% of patients with platelet count 50 to $74 \times 10^9/L$. See the prescribing information for dose modifications for the management of hematologic toxicities.

Other Toxicities

Ruxolitinib is associated with a potentially increased risk of opportunistic infections and viral reactivations.^{148,149} Non-melanoma skin cancers and pre-cancerous lesions have been reported in patients treated with ruxolitinib.¹⁵⁰ Lymphoid neoplasms may be diagnosed concurrently with MPN or may develop during the natural history of MF, PV, or ET.¹⁵¹⁻¹⁵⁴ Although one report indicated that *JAK* inhibitor therapy may be associated with an increased risk of aggressive B-cell lymphomas in patients with MF,¹⁵⁵ other studies found no evidence of increased lymphoma risk in patients treated with a *JAK* inhibitor.¹⁵⁶⁻¹⁵⁹

Impact of Mutational Status and Response to Ruxolitinib

In the COMFORT-II study, ruxolitinib was associated with clinical efficacy and survival improvement across different molecular subsets of patients with MF.¹⁶⁰ HMR mutations (*ASXL1*, *EZH2*, *SRSF2*, *IDH1*, or *IDH2*) were identified in 33%, 7%, 3%, <1%, and 0% of patients, respectively, and these frequencies were comparable in ruxolitinib and BAT arms. Responses in splenomegaly (>35% spleen volume reduction), symptomatic improvement, and the risk of ruxolitinib-associated anemia and thrombocytopenia were observed at similar frequencies across different mutation profiles. Ruxolitinib improved survival and reduced the risk of death in patients harboring HMR mutations (*ASXL1*, *EZH2*, *SRSF2*, *IDH1*, or *IDH2*) with an HR of 0.57.¹⁶⁰ The use of ruxolitinib did not appreciably influence the acquisition of additional mutations during treatment compared to the use of hydroxyurea.¹⁶¹ A decrease in the *JAK2*



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

V617F variant allele frequency was associated with the duration of the spleen volume response. An increase in the variant allele frequency of any initial mutation or the acquisition of ≥ 1 non-driver mutations during treatment was associated with increased rates of treatment discontinuation.

The results of another analysis of 95 patients with MF treated with ruxolitinib in a single institution also showed that *ASXL1*, *EZH2*, and *IDH1/2* mutations are associated with poor outcomes and patients with ≥ 3 mutations in *ASXL1*, *EZH2*, or *IDH1/2* had shorter time to treatment discontinuation and OS.¹⁶² However, in contrast to the findings of the COMFORT-II study, patients with ≥ 1 mutations in *ASXL1*, *EZH2*, or *IDH1/2* were significantly less likely to have a spleen response. Patients with ≥ 3 mutations had the worst outcomes, suggesting that multigene profiling may be useful for treatment planning in patients with MF.

Response Prediction to Ruxolitinib

The response to ruxolitinib after 6 months (RR6) model can be used to gauge response to ruxolitinib. This prognostic model takes into account the ruxolitinib dose, spleen response, and RBC transfusion needs for 6 months following the initiation of ruxolitinib therapy in order to predict OS.¹⁶³ The results of a multivariable analysis revealed several risk factors: a ruxolitinib dose of < 20 mg twice a day at baseline and at 3 and 6 months, a $\leq 30\%$ reduction in spleen length from baseline at 3 and 6 months, an RBC transfusion requirement at 3 and/or 6 months, and an RBC transfusion requirement at baseline and at 3 and 6 months. The model stratified patients into 3 risk categories: low-risk (median OS, not reached), intermediate-risk (median OS, 61 months; 95% CI, 43–80 months), and high-risk (median OS, 33 months; 95% CI, 21–50 months). The model's predictive potential was validated and confirmed in an external cohort of 40 patients with MF.

Momelotinib

Momelotinib, a potent and selective JAK1/2 inhibitor and ACVR1/ALK2 inhibitor, is FDA-approved for the treatment of intermediate- or high-risk MF in patients with anemia based on the results of the randomized phase III MOMENTUM trial, as well as subgroup data from the randomized phase III SIMPLIFY-1 trial.^{94,95,164} Momelotinib was also evaluated in randomized phase III studies in patients with intermediate-1 (symptomatic), intermediate-2, or high-risk MF who were not previously treated with a JAK inhibitor, as well as in those who were previously treated with ruxolitinib.^{164,165}

In the phase III MOMENTUM trial, patients with PMF or post-PV/ET MF with DIPSS intermediate-1, intermediate-2, or high-risk disease were randomized 2:1 to receive treatment with momelotinib or danazol.⁹⁴ The patients had symptomatic disease, anemia, and had previously received treatment with a JAK inhibitor. At 24 weeks, a significantly higher percentage of patients in the momelotinib arm had a total symptom score response rate of $\geq 50\%$ (25% vs 9%; $P = .0095$). Following week 24, all patients who remained in the study were treated with momelotinib.⁹⁵ At 48 weeks, among those who were evaluable for total symptom score, 45% of patients treated with momelotinib from the start of the study had a response, compared to 50% of patients treated with danazol who crossed over.

The phase III SIMPLIFY-1 study randomized 432 patients with intermediate-1 (symptomatic), intermediate-2, or high-risk MF with no prior treatment with a JAK inhibitor to receive momelotinib 200 mg once daily or ruxolitinib 20 mg twice daily (or according to the label) for 24 weeks.¹⁶⁴ Following this time period, all patients could cross over to the momelotinib arm. At 24 weeks, the data showed that momelotinib was noninferior to ruxolitinib. In the momelotinib arm, 26.5% of patients achieved the primary endpoint of a spleen response, defined as a $\geq 35\%$ decrease in the spleen



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

volume, compared to 29% of patients in the ruxolitinib arm ($P = .011$). However, it did not improve the total symptom score response rate ($P = .98$). As the symptoms benefit did not achieve statistical significance, all subsequent results, including transfusion results, were considered nominal. Mometotinib treatment led to an improvement in transfusion burden (transfusion rate, median of 0 units/month vs 0.4 units/month; $P < .001$; transfusion independence, 66.5% vs 49.3%; $P < .001$; transfusion dependence, 30.2% vs 40.1%; $P = .019$). At 2 years, the OS and LFS were 81.6% (HR, 1.02; 95% CI, 0.73–1.43) and 80.7% (HR, 1.08; 95% CI, 0.78–1.50), respectively, in patients treated with momelotinib compared to 80.6% and 79.3%, respectively, in patients initially treated with ruxolitinib who crossed over to the momelotinib group.¹⁶⁶

The phase III SIMPLIFY 2 trial randomized patients with intermediate-1 (symptomatic), intermediate-2, or high-risk MF who received prior ruxolitinib treatment 2:1 to receive momelotinib or BAT for 24 weeks.¹⁶⁵ In an intention-to-treat analysis, 7% of patients in the momelotinib group met the primary endpoint of a $\geq 35\%$ reduction in spleen volume, compared to 6% of patients in the BAT group ($P = .90$). At 24 weeks, 26% of those treated with momelotinib achieved a $\geq 50\%$ decrease in MPN Symptom Assessment Form TSS (MPN-SAF TSS) from baseline compared to 6% of those treated with BAT ($P = .0006$). At 2 years, the OS and LFS were 65.8% (HR, 0.98; 95% CI, 0.59–1.62) and 64.2% (HR, 0.97; 95% CI, 0.59–1.60), respectively, in patients treated with momelotinib compared to 61.2% and 59.7%, respectively, in patients initially treated with BAT who crossed over to the momelotinib group.¹⁶⁶ Data from both SIMPLIFY trials showed that treatment with momelotinib led to a clinically meaningful symptomatic benefit in patients with MF.¹⁶⁷

Toxicity

In the MOMENTUM study, at 24 weeks, anemia and thrombocytopenia were the most common grade ≥ 3 treatment-emergent hematologic

adverse events and were observed in 61% and 28%, respectively, of patients receiving momelotinib, and in 75% and 26%, respectively, of patients receiving danazol.⁹⁴ At 48 weeks, anemia and thrombocytopenia were reported in 11% and 19% of patients treated with momelotinib, including those who crossed over.⁹⁵ Acute kidney injury (mometotinib, 3%; danazol, 9%) and pneumonia (mometotinib, 2%; danazol, 9%) were the most common grade ≥ 3 nonhematologic treatment-emergent adverse events at 24 weeks.⁹⁴ Asthenia (mometotinib, 3%; danazol, 2%), dyspnea (mometotinib, 3%; danazol, 0%), and fatigue (mometotinib, 3%; danazol, 7%) were the most common grade ≥ 3 nonhematologic adverse events at 48 weeks.⁹⁵ In the SIMPLIFY-1 trial, anemia and thrombocytopenia were the most frequent hematologic abnormalities in both groups.¹⁶⁴ Seven percent of patients in the momelotinib group and 3% of patients in the ruxolitinib group had grade ≥ 3 infections. Similarly, anemia (mometotinib group, 14%; BAT group, 14%) and thrombocytopenia (mometotinib group, 7%; BAT group, 6%) were the most frequent grade ≥ 3 treatment-emergent adverse events in the SIMPLIFY-2 trial, with the most common nonhematologic treatment-emergent adverse events being asthenia (5%) in the momelotinib group and abdominal pain (6%) in the BAT group.¹⁶⁵

Pacritinib

Pacritinib, a JAK2, FLT3, and IRAK1 inhibitor, was evaluated in patients with intermediate-1, intermediate-2, and high-risk MF.^{96,97,168} Pacritinib is FDA-approved for the treatment of intermediate or high-risk MF with platelet count $< 50 \times 10^9/L$.^{97,168}

The phase II PAC203 trial reported that 200 mg pacritinib twice daily showed clinical activity and had a manageable safety profile in patients with ruxolitinib-resistant or ruxolitinib-intolerant intermediate-1, intermediate-2, or high-risk MF with platelet count $< 50 \times 10^9/L$.¹⁶⁸ At 24 weeks, the spleen response rate ($\geq 35\%$ reduction in spleen volume) was 9.3% in the overall cohort versus 16.7% in those with platelet count $< 50 \times$



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

$10^9/L$ and the total symptom score response rate ($\geq 50\%$ reduction in total symptom score based on the MPN-SAF TSS 2.0) was 7.4% in the overall cohort versus 8.3% in those with platelet count $< 50 \times 10^9/L$.

In the phase III PERSIST-1 trial, patients with intermediate-1, intermediate-2, or high-risk MF with palpable splenomegaly (≥ 5 cm below the left costal margin) were randomized 2:1 to receive pacritinib ($n = 220$), 400 mg once daily, or BAT ($n = 107$) (excluding JAK2 inhibitors).⁹⁶

Patients were allowed to cross over to pacritinib at 24 weeks or if their disease progressed. In the BAT group, 84% of the patients crossed over to the pacritinib group at a median time point of 6.3 months. Nineteen percent of patients receiving pacritinib met the primary endpoint ($\geq 35\%$ spleen volume reduction, as determined by MRI or CT, in the intention-to-treat population) compared to 5% of patients receiving BAT ($P = .0003$) at 24 weeks. At the same time point, the percentage of patients with a total symptom score reduction of $\geq 50\%$, as determined using the MPN-SAF TSS 2.0, was similar in the pacritinib and BAT study arms (19% vs 10%; $P = .24$). At 48 weeks, a significantly higher percentage of patients in the pacritinib study arm achieved this reduction (15% vs 0%; $P = .0027$). OS did not differ between the two groups (HR, 1.36; 95% CI, 0.89–2.09; $P = .16$) prior to week 24.

The phase III PERSIST-2 trial randomized patients with intermediate-1, intermediate-2, or high-risk MF with platelet count $\leq 100 \times 10^9/L$ 1:1:1 to receive once-daily pacritinib 400 mg, twice-daily pacritinib 200 mg, or BAT.⁹⁷ Patients had palpable splenomegaly (≥ 5 cm below the left costal margin) and platelet count $\leq 100 \times 10^9/L$. Forty-eight percent of patients were previously treated with ruxolitinib. Among the BAT group, 45% of patients received ruxolitinib. Patients were allowed to cross over to pacritinib at 24 weeks or if splenomegaly progressed. At 24 weeks, in the intention-to-treat population, the proportion of patients achieving the co-primary endpoint of $\geq 35\%$ reduction in spleen volume, as assessed by

MRI/CT, was significantly higher in the pacritinib groups (15% [95% CI, 7.6%–24.7%; $P = .02$] for 400 mg once daily and 22% [95% CI, 12.9%–32.7%; $P = .001$] for 200 mg twice daily) than in the BAT group (3% [95% CI, 0.3%–9.7%]). Seventeen percent (95% CI, 9.6%–27.8%; $P = .65$) of patients receiving once-daily 400 mg pacritinib and 32% (95% CI, 22.0%–44.3%; $P = .01$) of patients receiving twice-daily 200 mg pacritinib met the co-primary endpoint of $\geq 50\%$ reduction in total symptom score (MPN-SAF TSS 2.0), as opposed to 14% (95% CI, 6.9%–24.1%) of patients receiving BAT. OS was similar across all three groups (HR, 1.18; 95% CI, 0.57–2.44; and HR, 0.68; 95% CI, 0.30–1.53 for pacritinib 400 mg once daily and 200 mg twice daily, respectively, when compared to BAT). A post hoc analysis determined that the achievement of spleen volume reduction with pacritinib treatment (200 mg twice daily) was associated with a significant improvement in OS compared to treatment with BAT in patients with platelet counts $\leq 100 \times 10^9/L$.¹⁶⁹

An analysis of pooled data from the PERSIST-1 and PERSIST-2 studies demonstrated the efficacy of pacritinib irrespective of the platelet and hemoglobin counts at baseline.¹⁷⁰ Pacritinib elicited a spleen response across all subgroups. Symptoms response, as assessed by TSS, was comparable across the platelet subgroups. However, among the subgroups of patients with anemia, those with baseline hemoglobin < 8 g/dL had the most instances of $\geq 50\%$ reduction in TSS compared to those with baseline hemoglobin ≥ 8 to < 10 g/dL and > 10 g/dL.

Toxicity

The phase II PAC203 trial reported thrombocytopenia (33.3%), anemia (20.4%), and neutropenia (5.6%) as the most common grade 3 or 4 treatment-emergent hematologic events in patients with MF resistant to or intolerant of ruxolitinib who received twice-daily pacritinib 200 mg.¹⁶⁸ Pneumonia (9.3%) as well as diarrhea, abdominal pain, and hyperuricemia (5.6% each) were the most common nonhematologic grade 3 or 4



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

treatment-emergent adverse events. Like fedratinib, pacritinib also exhibits FLT3 inhibition, which has been implicated in gastrointestinal toxicity.^{124,125}

In the PERSIST-1 trial, the most frequent grade 3 or 4 adverse events in the pacritinib study arm were anemia (17%), thrombocytopenia (12%), and diarrhea (5%) and in the BAT arm, they were anemia (15%), thrombocytopenia (11%), dyspnea (3%), and hypotension (3%).⁹⁶ One percent of patients in the pacritinib group had an infection compared to none in patients receiving BAT. In the PERSIST-2 trial, the most frequent grade 3 or 4 treatment-emergent adverse events in patients receiving once-daily pacritinib 400 mg, twice-daily pacritinib 200 mg, or BAT were thrombocytopenia (31%, 32%, and 18%, respectively) and anemia (27%, 22%, and 14%, respectively).⁹⁷

In 2016, the FDA placed a clinical hold on the development of pacritinib while evaluating deaths related to intracerebral hemorrhage and cardiovascular events. In 2017, the FDA lifted the clinical hold and in 2022, the drug was approved for the treatment of intermediate- or high-risk MF (PMF, post-PV MF, or post-ET MF) for patients with platelet count $<50 \times 10^9/L$.

Allogeneic Hematopoietic Cell Transplant

Allogeneic HCT is the only potentially curative treatment option resulting in long-term remissions for patients with MF. Donor selection and conditioning should be evaluated on a case-by-case basis (See NCCN Guidelines for Hematopoietic Cell Transplantation, available at www.NCCN.org). One study showed that the cumulative relapse incidence at 1 year was 6% in those with mutation clearance at 30 days post-transplant versus 21% in those without mutation clearance at the same time point.¹⁷¹ At 6 years, patients with mutation clearance at 30 days post-transplant had a higher disease-free survival (61% vs 41%) and overall survival (74% vs 60%) rates compared to those without mutation clearance. As the clearance of driver mutations at 30 days post-

transplant appears to influence relapse and survival, the Panel recommends post-transplant monitoring for measurable residual disease by driver mutations and chimerism.^{171,172}

Myeloablative conditioning and reduced-intensity conditioning (RIC) are relatively similar in terms of OS.¹⁷³ The use of RIC is associated with a lower rate of non-relapse mortality (NRM), but it is also associated with a higher risk of relapse compared to myeloablative conditioning.¹⁷⁴⁻¹⁸¹ Comparison studies of RIC also do not show a difference in OS,^{181,182} although one study reported a trend towards lower NRM (HR, 0.52; 95% CI, 0.26–1.05; $P = .068$) and a higher relapse rate (HR, 9.21; 95% CI, 1.81–46.9; $P = .008$) with regimens that use the combination of busulfan and fludarabine.¹⁸¹ Another study also determined a higher relapse rate but the difference was not statistically significant ($P = .21$).¹⁸² No statistically significant difference was obtained for NRM ($P = .32$).

Patients with MPN are at particularly high risk for hepatobiliary toxicities related to transplant, including sinusoidal obstructive syndrome (SOS). Approaches to reduce SOS and NRM using specialized myeloablative conditioning have been used and may be helpful.^{183,184} The estimated OS and NRM rates for myeloablative conditioning at 3 to 5 years range from 30% to 61% and 35% to 50%, respectively.¹⁸⁵ In a retrospective registry analysis of 289 patients with MF, allogeneic HCT resulted in long-term OS in approximately one third of patients, but the probability of long-term survival and NRM was dependent on the source of stem cells.¹⁸⁶ The 5-year post-transplant OS rates were 37%, 40%, and 30%, respectively, for HLA-matched sibling donor transplant, other related donor transplant, and unrelated donor (URD) transplant, respectively. The corresponding 5-year disease-free survival rates were 33%, 22%, and 27%, respectively. The NRM rate at 5 years was higher for URD transplant (50% compared to 35% and 38% for HLA-matched sibling donor transplant and other related donor transplant, respectively).



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

In a prospective, multicenter study that evaluated allogeneic HCT with RIC in 103 patients with MF, the cumulative incidence of NRM at 1 year was 16% and the cumulative incidence of relapse at 3 years was 22%.¹⁷⁵ The estimated 5-year event-free survival (EFS) and OS rates were 51% and 67%, respectively. The NRM was significantly lower for patients with a completely matched donor (12% vs 38%; $P = .003$). Other large retrospective registry analyses have also reported similar outcomes.^{178,179} In the Center for International Blood and Marrow Transplant Research (CIBMTR) analysis that included 233 patients who underwent allogeneic HCT using RIC for PMF, the probabilities of OS and PFS at 5 years were 47% and 27%, respectively.¹⁷⁸ The cumulative incidence of NRM and relapse/progression at 5 years were 24% and 48%, respectively. In the European Bone Marrow Transplantation Registry (EBMTR) analysis that included 193 patients who underwent transplantation for post-PV or post-ET MF, the 3-year OS rate, incidence of relapse, and NRM were 55%, 32%, and 28%, respectively.¹⁷⁹ Another study that included 2459 patients with MF who underwent allogeneic HCT reported an OS rate of 41% (95% CI, 39%–44%) and a disease-free survival rate of 32% (95% CI, 30%–35%) at 10 years.¹⁸⁷ In 1055 patients who were disease-free at 2 years, the 10-year OS and disease-free survival rates were 74% (71%–78%) and 64% (60%–68%), respectively.

Age (>55 years) and donor type (HLA-identical sibling donor transplant vs HLA-well-matched URD transplant or partially/mismatched URD transplant) have been the most important prognostic factors of OS and NRM. Among patients who underwent allogeneic HCT with RIC for PMF, the 5-year survival rates following HLA-identical sibling donor transplant, HLA-well-matched URD transplant, and partially/mismatched URD transplant were 56%, 48%, and 34%, respectively ($P = .002$) and the relative risk of NRM was also the lowest for HLA-identical sibling donor transplant (1%) compared to 3% and 9% for HLA-well-matched URD

transplant and partial/mismatched URD transplant, respectively.¹⁷⁸ In patients who underwent allogeneic HCT with RIC for post-PV MF or post-ET MF, the overall 3-year cumulative incidence of NRM was significantly higher in patients >55 years (35% vs 20% for younger patients; $P = .032$) and in those who underwent URD transplant (34% vs 18% for those who had a related donor transplant; $P = .034$).¹⁷⁹

The results of a retrospective study by the European Society for Blood and Marrow Transplantation with patients with MF who underwent allogeneic HCT from an HLA-identical sibling or an URD identified age ≥ 60 years, Karnofsky performance status of <90% at the time of transplant, graft failure, acute graft-versus-host disease (GVHD) (grades III–IV), and disease progression or relapse as factors that were independently associated with a higher mortality rate.¹⁸⁸ These factors, along with HCT-specific Comorbidity Index ≥ 3 and extensive chronic GVHD, were associated with higher NRM. The DIPSS risk score was not a prognostic factor.

Another retrospective multicenter study of 69 patients with chronic phase MF who were treated with allogeneic blood or marrow transplantation from a haploidentical donor and received cyclophosphamide post-transplantation reported an OS of 72% (95% CI, 59%–81%), a relapse-free survival (RFS) of 44% (95% CI, 29%–59%), and a GVHD-free RFS of 30% (95% CI, 17%–43%) at 3 years.¹⁸⁹ A cumulative incidence of 10% was obtained for grade 3–4 acute GVHD and 8% for extensive chronic GVHD.

A few studies have shown that larger spleen size may be associated with inferior outcomes after transplant, possibly reflecting an aggressive disease biology.^{189–191} A spleen size ≥ 22 cm or a prior splenectomy (HR, 6.37; 95% CI, 2.02–20.1; $P = .002$) and bone marrow grafts (HR, 4.92; 95% CI, 1.68–14.4; $P = .004$) were associated with a higher incidence of relapse.¹⁸⁹ A univariate analysis determined that a spleen size ≥ 17 cm or



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

a prior splenectomy was associated with worse RFS (HR, 3.50; 95% CI, 1.18–10.37; $P = .02$) and a higher relapse rate (subdistribution HR not calculable; $P = .01$).¹⁹⁰ The results of a multivariate analysis by Polverelli et al¹⁹¹ demonstrated that splenectomy was associated with reduced NRM (HR, 0.64; 95% CI, 0.44–0.93; $P = .018$) and a higher risk of relapse (HR, 1.43; 95% CI, 1.01–2.02; $P = .042$), but no effect on OS (HR, 0.86; 95% CI, 0.67–1.12; $P = .274$).

In another study, DIPSS risk score has been shown to predict outcome after transplant.^{178,192} In the aforementioned CIBMTR analysis, there was a trend towards lower mortality rates in patients with low- or intermediate-1-risk disease, and higher NRM in patients with intermediate-2 or high-risk disease.¹⁷⁸ In another retrospective analysis of 170 patients with MF who received HCT, DIPSS risk score significantly correlated with mortality risk and NRM (HR for post-transplant mortality was 4.11 for high-risk disease compared to 3.15, 1.97, and 1, respectively, for intermediate-2, intermediate-1, and low-risk disease; the corresponding HRs for NRM were 3.41, 3.19, 1.41, and 1, respectively).¹⁹² The association of DIPSS risk score with relapse was not significant, although patients with higher-risk disease experienced more relapses than those with lower-risk disease.

DIPSS risk scores prior to HCT have also been shown to correlate with OS following allogeneic HCT.^{178,193,194} However, in one retrospective analysis, the differences in OS between patients with intermediate-1 and intermediate-2-risk disease were not significantly different. In a multivariate analysis, only *JAK2* wild-type, age ≥ 57 years, and the presence of constitutional symptoms were independent predictors of OS. The 5-year OS rates were 90%, 74%, and 50% for the presence of 0, 1, and 2 risk factors.¹⁹³ In another retrospective analysis that evaluated the impact of allogeneic HCT on survival in patients < 65 years of age at the time of diagnosis of PMF ($n = 438$; 190 patients received allogeneic HCT

and 248 patients received conventional therapy), the relative risk of death after allogeneic HCT was 5.6 for patients with DIPSS low-risk disease, 1.6 for patients with intermediate-1-risk disease, 0.55 for patients with intermediate-2-risk disease, and 0.37 for patients with high-risk disease.¹⁹⁴

These findings suggest that outcomes following allogeneic HCT are better for patients with low- or intermediate-1-risk MF.^{178,192} However, since HCT is associated with a significant rate of transplant-related complications and morbidity that may not otherwise occur with non-transplant therapies in this group of patients, the overall benefit may be with non-transplant therapies.¹⁹⁵ Allogeneic HCT is associated with a clear benefit in patients with intermediate-2 or high-risk MF. A retrospective study of 544 patients with MF investigated the different prognostic models (IPSS, DIPSS, and DIPSS-Plus) and determined that the IPSS and DIPSS-plus models were most able to differentiate between the intermediate-1 and intermediate-2-risk categories.¹⁹⁶

The Myelofibrosis Transplant Scoring System (MTSS) is a model that takes into account clinical (age ≥ 57 years, Karnofsky performance status $< 90\%$, platelet count $< 150 \times 10^9/L$, and leukocyte count $> 25 \times 10^9/L$), molecular (presence of *ASXL1* mutation and absence of *CALR* and *MPL* mutations), and transplant-specific factors (HLA-mismatched URD), and is designed to assess prognosis after allogeneic transplant in patients with primary and post-ET/PV MF.¹⁹⁷ It stratifies patients into four risk categories: low, intermediate, high, and very high. Validated in a cohort of 156 patients, the survival rates for these categories were 83% (95% CI, 71%–95%), 64% (95% CI, 53%–75%), 37% (95% CI, 17%–57%), and 22% (95% CI, 4%–39%), respectively ($P < .001$). Another study evaluating the performance of the MTSS model concluded that it may need to be refined as it did not distinctly stratify patients into four risk categories.¹⁹⁸ However, the authors note that it still has clinical value. When the risk



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

levels were combined to give two new categories, standard (low and intermediate) and high (high and very high), the MTSS was better able to distinguish risk ($P < .001$). The OS at 3 years for the standard- and high-risk levels were 62% (95% CI, 49%–72%) and 25% (95% CI, 9%–45%), respectively. Further validation studies are needed to confirm these findings.

Impact of Mutational Status

CALR mutation is associated with higher OS rates and lower rate of NRM following allogeneic HCT in patients with PMF as well as post-PV or post-ET MF.^{199,200} Identification of HMR mutations (*ASXL1*, *EZH2*, *SRSF2*, *TP53*, *IDH1*, or *IDH2* mutations) may be helpful in decision-making regarding allogeneic HCT in patients with MF.^{30,39-41,200} *CBL*, *DNMT3A*, and *U2AF1* were associated with worse OS in patients with MF undergoing allogeneic HCT.^{201,202} The results from another study also suggest inferior OS with *ASXL1* mutations (subdistribution HR, 2.36; 95% CI, 0.85–6.6; $P = .09$).¹⁹⁰

In a study of 133 patients who underwent allogeneic HCT for PMF ($n = 97$) or post-ET/post-PV MF ($n = 36$), the 4-year OS rate was 82% for patients with *CALR* mutations compared to 56% for patients without *CALR* mutations (*CALR* wild-type). The NRM rate was also significantly lower in patients with *CALR* mutations compared with those who were *CALR* wild-type (4-year NRM rates were 7% and 31%, respectively; $P = .024$).¹⁹⁹ In another study that evaluated the impact of molecular genetics on the outcome after allogeneic HCT in patients with MF (PMF, $n = 110$; post-PV or ET MF, $n = 46$; and MF in transformation, $n = 13$), the results of a multivariate analysis showed that *CALR* mutation was an independent factor for lower NRM and improved PFS and OS.²⁰⁰ *ASXL1* and *IDH2* mutations were independent risk factors for lower PFS, whereas no impact was observed for patients with triple-negative disease. As discussed earlier, *CALR*(-)/*ASXL1*(+) is associated with a poor prognosis

(independent of the DIPSS-Plus risk score) in patients with PMF and this subset of patients should be considered for allogeneic HCT earlier in the disease course.⁴⁷

A small study with 18 patients with primary or post-ET MF found that *MPL* mutations were associated with a favorable outcome following allogeneic HCT with an OS rate and an RFS rate of 83.5% (95% CI, 65.9%–100%) at 5 years and a relapse rate of 5.5%.²⁰³

The addition of mutational status to DIPSS-Plus can help improve the prediction of transplantation outcome.²⁰⁴ Patients with ≥ 3 mutations along with *CALR* or *JAK2* mutations had higher NRM and risk of relapse following transplant compared to those with fewer mutations.

Treatment Recommendations Based on Symptom Assessment and Risk Stratification

The selection of appropriate treatment should be based on the risk score, the presence of symptoms, and the disease stage. A clinical trial or consideration of a clinical trial is recommended for all patients with MF who require treatment with the aim of reducing bone marrow fibrosis, improving cytopenias and symptom burden, restoring transfusion independence, and/or preventing/delaying progression to AML.

Lower-Risk MF

Patients with asymptomatic lower-risk MF should be observed and monitored for signs and symptoms of disease progression with MPN-SAF TSS (MPN-10). Enrollment in a clinical trial is also an option. Hydroxyurea has been shown to be an effective treatment option for the hyperproliferative manifestations of lower-risk MF (thrombocytosis or leukocytosis). In a small study of 40 patients with symptomatic MF (constitutional symptoms, splenomegaly, thrombocytosis, leukocytosis, pruritus, and bone pain), treatment with hydroxyurea (500 mg/day, subsequently adjusted to the individual efficacy and tolerability) resulted in



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

clinical improvement in 40% of patients.²⁰⁵ Anemia induced by hydroxyurea was manageable with concomitant treatment. The Panel has included hydroxyurea as an option for symptomatic lower-risk MF, if the use of cytoreductive therapy would be symptomatically beneficial in selected patients with high platelet counts. Hydroxyurea, pacritinib (if platelets $<50 \times 10^9/L$), peginterferon alfa-2a,¹¹⁵ ruxolitinib,¹³⁰⁻¹³² and momelotinib (category 2B) are listed as useful in certain circumstances options for the first-line treatment of patients with symptomatic lower-risk MF. Clinical trials are always strongly encouraged. For MF, if peginterferon alfa-2a is unavailable, the use of other available pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate. Data from the PROUD-PV and CONTINUATION-PV studies support the use of ropeginterferon alfa-2b-njft in PV while data from the SURPASS-ET study supports its use in hydroxyurea-resistant or -intolerant ET.^{206,207} For patients with anemia, see *Management of MF-Associated Anemia* in the algorithm.

Although the outcomes following allogeneic HCT are better for patients with lower-risk MF, due to the high transplantation-related morbidity and mortality, treatment decisions regarding allogeneic HCT should be individualized.^{178,192,194} Allogeneic HCT should be considered for lower-risk MF in patients with refractory, transfusion-dependent anemia, circulating blast cells $>2\%$ in peripheral blood, adverse cytogenetics, or molecular abnormalities.²⁰⁸ Evaluation for allogeneic HCT is recommended for patients with low platelet counts or complex cytogenetics. The MTSS can be helpful in predicting post-transplant survival when counseling patients about transplant.¹⁹⁷

Higher-Risk MF

Referral to an HCT expert for allogeneic HCT evaluation is recommended for all patients with higher-risk MF that is DIPSS-Plus Intermediate-1 or MIPSS-intermediate or higher. Transplant is recommended for patients with higher-risk MF that is DIPSS-Plus or MYSEC Intermediate-2 or high-

risk disease or MIPSS70 or MIPSS 70+ high-risk.¹⁹² Transplant can also be considered in selective cases: RBC transfusion dependence, high-risk mutations (ie, *ASXL1*, *RAS*, *TP53*), or loss of response to JAK inhibitor therapy. The selection of patients for allogeneic HCT should be based on age, performance status, major comorbid conditions, psychosocial status, patient preference, and availability of caregiver. The MTSS can be helpful in predicting post-transplant survival when counseling patients about transplant and can be used to optimize patient selection, with low- and intermediate-risk being optimal.¹⁹⁷

For patients with $>10\%$ blasts in the peripheral blood, azacitidine with or without a JAK inhibitor may be considered prior to allogeneic HCT to reduce the blast percentage.²⁰⁹ The results of several studies suggest that prior exposure to ruxolitinib may improve outcomes after allogeneic HCT.^{184,210-212} The guidelines recommend that JAK inhibitors should be considered for use at least 2 months prior to transplant in patients with splenomegaly and/or constitutional symptoms even if the patient would otherwise be transplant eligible.^{184,210,211} JAK inhibitors can be tapered prior to or during conditioning to be completed before cell infusion. In patients with massive splenomegaly that does not respond to JAK inhibitors, alternative measures to reduce spleen size may be considered prior to transplant (eg, splenic radiation, splenic artery embolization, splenectomy).^{213,214}

In a prospective phase II trial, 28 patients with MF were treated with ruxolitinib for at least 8 weeks prior to HCT and followed a taper schedule that ended 4 days before donor cell infusion.¹⁸⁴ Twenty-three patients underwent myeloablative conditioning while the remaining five underwent RIC. After termination of treatment with ruxolitinib, cytokine release syndrome was not observed, and engraftment was successful in all patients. Following transplant, the 2-year OS was 86% (95% CI, 61%–96%). Shanavas and colleagues²¹¹ examined data from 100 patients with



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

MF who were treated with JAK inhibitors prior to HCT. Sixty-six patients continued ruxolitinib therapy until transplant. Most of the observed symptoms were consistent with symptoms associated with MF and were mild or moderate. Two patients had a severe adverse occurrence and, as a result, HCT was delayed. Patients who displayed clinical improvement with the use of a JAK inhibitor also had more favorable outcomes post-transplant. At 2 years, the OS was 61% (95% CI, 49%–71%).

Similarly, a study by Chhabra et al²¹⁰ reported that treatment with ruxolitinib and management of splenomegaly with splenic irradiation prior to transplant, along with fludarabine/busulfan-based conditioning, led to more favorable outcomes. At 3 years, the OS was 81.1% (95% CI, 64.4%–90.5%) and the RFS was 78.4% (95% CI, 61.4%–88.5%). Another study assessing the use of ruxolitinib prior to RIC and transplant in patients with MF found that treatment with ruxolitinib significantly reduced symptom burden.²¹² Patients did not experience significant side effects while tapering off ruxolitinib and HCT was not delayed. A retrospective study with 551 patients with MF who underwent HCT determined that the NRM at 1 year (HR, 0.80; $P = .32$), and the EFS (HR, 0.81; $P = .19$) and OS (HR, 0.81; $P = .21$) rates at 2 years did not differ between patients who received ruxolitinib prior to transplant versus those who did not.²¹⁵ However, patients with ruxolitinib pretreatment who had an ongoing spleen response at the time of transplant had a decreased risk of relapse (HR, 0.34; $P = .04$) and an improved 2-year EFS (HR, 0.61; $P = .02$).

Pacritinib has demonstrated significant activity resulting in $\geq 35\%$ spleen volume reductions and symptom improvement, even in patients with severe baseline cytopenias,^{96,97} and is a category 1, preferred option for patients with higher-risk MF with platelet count $< 50 \times 10^9/L$ who are not transplant candidates or for whom transplant is not currently feasible. Momelotinib is a category 2B, other recommended regimen for these patients.¹⁶⁴ Enrollment in an appropriate clinical trial is always

encouraged. The use of ruxolitinib at a lower dose (5 mg twice daily) has shown some efficacy, resulting in some reductions in spleen volume and improvement in total symptom score even in patients with low platelet counts at baseline ($50\text{--}100 \times 10^9/L$).¹⁴⁶

Enrollment in a clinical trial, fedratinib⁹⁸ (category 1 for initial treatment), ruxolitinib^{90,129,136–138} (category 1 for initial treatment), momelotinib,^{95,164,165} or pacritinib^{96,97} (category 2B) are options for patients with higher-risk MF with symptomatic splenomegaly and/or constitutional symptoms and with platelet count $\geq 50 \times 10^9/L$ who are not candidates for transplant or for whom transplant is not currently feasible. A study by Hernandez-Boluda²¹⁶ reported that patients with severe thrombocytopenia (platelet count $< 50 \times 10^9/L$) were in a higher risk category and had more instances of anemia and leukopenia. Patients with platelet count $< 50 \times 10^9/L$ experience a greater symptom burden and might benefit from symptomatically guided treatment options.²¹⁷ For patients with anemia, see *Management of MF-Associated Anemia* in the algorithm.

Management of MF-Associated Anemia

Anemia is considered a negative prognostic risk factor for survival in patients with MF.⁹⁹ Symptomatic anemia is observed in 35% to 54% of patients at the time of diagnosis.^{218,219} It is essential to assess for, and treat the most common alternative causes of anemia (ie, bleeding, nutritional deficiencies, hemolysis) before considering other treatment options.

Momelotinib, pacritinib, danazol, EPO-stimulating agents (ESAs), luspatercept-aamt, and immunomodulatory agents (lenalidomide, thalidomide, and pomalidomide) have also been evaluated for the management of MF-associated anemia.

In a post hoc analysis of the phase III SIMPLIFY-1 trial, treatment with momelotinib led to a greater transfusion independence rate at 24 weeks in



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

both the subgroup with moderate/severe anemia (<10 g/dL) (mometotinib, 46.5%; ruxolitinib, 26.6%) and the subgroup with mild anemia (≥ 10 to <12 g/dL) (mometotinib, 80.8%; ruxolitinib, 50.7%).²²⁰ Among patients with moderate/severe anemia who did not require transfusions at baseline, 72.0% of those treated with mometotinib maintained transfusion independence at 24 weeks, compared to 34.1% of those treated with ruxolitinib. In the subgroup with mild anemia, 86.4% of those treated with mometotinib maintained transfusion independence, compared to 57.9% of those treated with ruxolitinib.

In the phase III SIMPLIFY 2 trial, a higher percentage of patients achieved transfusion independence at 24 weeks when treated with mometotinib compared to BAT (43% vs 21%, respectively; $P = .0012$).¹⁶⁵ Fifty percent of those treated with mometotinib required transfusions at 24 weeks compared to 64% of those treated with BAT ($P = .10$). Additionally, 40% of those in the mometotinib arm did not require transfusions during the treatment period versus 27% in the BAT arm. Subgroup analyses demonstrated a higher transfusion independence rate at 24 weeks with mometotinib as opposed to BAT in patients with a baseline hemoglobin level <10 g/dL (33.3% vs 12.8%) and those with non-transfusion independent MF at baseline (34.7% vs 3.0%).²²¹ A longitudinal analysis also showed that treatment with mometotinib led to a reduction in transfusion burden.²²²

Data from the phase III MOMENTUM trial showed that at 24 weeks, treatment with mometotinib resulted in a significantly higher transfusion independence rate (31% vs 20%; $P = .0064$), and a spleen volume reduction of $\geq 35\%$ (23% vs 3%; $P = .0006$) compared to treatment with danazol.⁹⁴ At 48 weeks, the transfusion independence rate and the proportion of patients achieving a $\geq 35\%$ spleen volume reduction were 57% and 43%, respectively, in the mometotinib group and 60% and 13%, respectively, in the danazol group who crossed over.⁹⁵

Data from the PERSIST-2 study demonstrated the anemia benefit of pacritinib.²²³ A greater number of patients treated with pacritinib compared to those treated with BAT had transfusion-independent disease over any 12-week period in the first 24 weeks of treatment (37% vs 7%; $P = .001$ based on Gale criteria). Similarly, a greater proportion of patients treated with pacritinib had a $\geq 50\%$ decrease in transfusion requirement over any 12-week period in the first 24 weeks of treatment (49% vs 9%; $P < .0001$).

In the COMFORT-II study, anemia was managed with packed RBC transfusions.²²⁴ In a small number of patients (13 out of 146 patients) who received both ruxolitinib and an ESA, the use of an ESA with ruxolitinib was well tolerated and did not impact the effectiveness of ruxolitinib. Another study that assessed the use of ESAs along with ruxolitinib ($n = 9$) or the addition of ESAs after treatment with ruxolitinib for a median of 4 months ($n = 50$) in patients with MF also showed that the concomitant use of an ESA with ruxolitinib was effective for the management of anemia in patients with MF.²²⁵ Fifty-four percent of patients achieved an anemia response (per IWG-MRT criteria) and, at 5 years, a response was observed in 76% of patients. Spleen reduction was reported in 78% of patients. These findings support the feasibility of administering ESAs for the management of anemia in patients receiving ruxolitinib. However, ESAs are less effective for the management of transfusion-dependent anemia.²²⁶

In a study of 50 patients with MF and anemia, danazol therapy resulted in an anemia response in 30% of patients, and responses were less frequent in patients with transfusion dependency (19% compared to 44% in patients without transfusion requirements).²²⁷ As danazol is an androgen, it is contraindicated in patients with androgen-dependent tumors.²²⁸ It can also cause liver toxicity. As such, prostate cancer screening and monitoring of liver function tests are recommended.²²⁷ Caution is advised



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

with concomitant use of statins over concerns for increased risk of rhabdomyolysis in patients receiving danazol.²²⁸

The use of recombinant human EPO or darbepoetin alfa has resulted in anemia responses (transfusion independence with normal hemoglobin levels, sustained increase in hemoglobin levels [>2 g/dL] within 12 weeks, or $>50\%$ reduction in transfusion requirements within 12 weeks) in 45% to 60% of patients with MF.²²⁹⁻²³¹ Lower serum EPO levels (<125 mU/mL), smaller spleen size, and low RBC transfusion requirements have been associated with favorable responses.

Because MF can be characterized by increased transforming growth factor beta (TGF- β) signaling and anemia related to increased TGF- β can be alleviated by inhibition of TGF- β signaling,²³² luspatercept has garnered significant attention in the MF field and is the subject of a randomized phase III clinical trial for patients with RBC transfusion-dependent MF on JAK2 inhibitor therapy (NCT04717414). The phase II open-label ACE-536-MF-001 clinical trial assessed the safety and efficacy of luspatercept for MF-related anemia.²³³ Patients were divided into four groups: no transfusion dependence and no ruxolitinib treatment; transfusion dependence and no ruxolitinib treatment; no transfusion dependence and ruxolitinib treatment; and transfusion dependence and ruxolitinib treatment. Anemia response rate, defined as a ≥ 1.5 g/dL rise in hemoglobin from baseline in the non-transfusion dependent group and transfusion independence in the transfusion-dependent group, over 12 consecutive weeks in the primary treatment period was the primary endpoint. In the group with no transfusion dependence, 13.6% of patients who did not receive ruxolitinib achieved an anemia response, while 14.3% of patients who received ruxolitinib achieved an anemia response. In the group with transfusion dependence, 9.5% of patients with no ruxolitinib treatment achieved an anemia response, while 26.3% of patients who received ruxolitinib achieved an anemia response. All groups had a

decrease in the total symptom score; patients with no transfusion dependence who received ruxolitinib had the highest decrease. Overall, hypertension was the most common treatment-related adverse event. Complete results from the phase III INDEPENDENCE study are forthcoming.²³⁴ Luspatercept-aamt is FDA-approved for the treatment of anemia without previous ESA use in adults with very low- to intermediate-risk MDS who may require regular RBC transfusions; and for the treatment of anemia refractory or intolerant to prior ESA treatment that requires ≥ 2 RBC transfusions over 8 weeks in adults with very-low- to intermediate-risk myelodysplastic syndromes (MDS) with ring sideroblasts or with myelodysplastic/MPN with ring sideroblasts and thrombocytosis.

Thalidomide (in escalating daily doses of 100–800 mg) has demonstrated very minimal efficacy, resulting in anemia response rates of 0% to 29%, and is also poorly tolerated.²³⁵⁻²⁴¹ A lower dose of thalidomide (50 mg/day), when used in combination with prednisone, is better tolerated, leading to improved anemia response rates (62%) compared to high-dose thalidomide monotherapy in the management of MF-associated symptomatic anemia (hemoglobin level <10 g/dL or symptomatic splenomegaly).²⁴² Lenalidomide, alone or in combination with prednisone, has also demonstrated modest efficacy in the management of MF-associated anemia, resulting in response rates of 19% to 32%, with myelosuppression being the most common grade ≥ 3 hematologic toxicity.²⁴³⁻²⁴⁶ Lenalidomide is more likely to induce better response rates in patients with isolated chromosome 5q deletion.²⁴⁷

In an analysis that reassessed the efficacy of thalidomide and lenalidomide in 125 patients with MF treated in three consecutive phase II trials, the combination of lenalidomide and prednisone was more effective and safer than single-agent thalidomide or lenalidomide.²⁴⁸ After a median follow-up of 42 months, the ORR was 38% for the combination of lenalidomide and prednisone compared to 34% and 16%, respectively, for



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

lenalidomide and thalidomide. There was also a trend for a higher efficacy in patients receiving lenalidomide-based therapy ($P = .06$), and in a multivariate analysis the lenalidomide-based regimen was the only factor independently associated with a higher response rate.

Pomalidomide has also been evaluated as a treatment option for MF-associated anemia.^{249,250} In one phase II study, pomalidomide (with or without prednisone) resulted in similar response rates (39%) in patients with MF and anemia and/or thrombocytopenia and/or neutropenia, with a median response duration of 13 months.²⁴⁹ However, in another randomized study that evaluated pomalidomide in patients with MF and RBC transfusion dependence, the RBC transfusion independence response rates were similar for patients treated with pomalidomide and placebo.²⁵⁰

Studies are ongoing to evaluate the combination treatment of ruxolitinib with thalidomide or pomalidomide in patients with MF (NCT03069326 and NCT01644110).^{251,252} A response rate of 55% was obtained in a phase II study investigating the combination of ruxolitinib and lenalidomide in patients with PMF or post-PV/ET MF with anemia.²⁵³ However, a dose interruption was needed in 75% of patients due to toxicity and the study was terminated early due to lack of efficacy.

For patients with MF-associated anemia and ongoing symptomatic splenomegaly and/or constitutional symptoms, enrollment in a clinical trial and momelotinib (category 1) are preferred options. Pacritinib, as well as ruxolitinib combination, are listed as other recommended regimens. Luspatercept-aamt, danazol (category 2B), or ESAs (epoetin alfa or darbepoetin alfa) (if serum EPO <500 mU/mL) (category 2B) can be added to ruxolitinib. An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.

For patients with MF-associated anemia and no symptomatic splenomegaly or constitutional symptoms, a clinical trial is preferred. Danazol, ESAs (if serum EPO <500 mU/mL), luspatercept-aamt, momelotinib (category 2B), and pacritinib (category 2B) are other recommended regimens. Lenalidomide with prednisone for del(5q) is a category 2B, useful in certain circumstances option. This regimen should start as a combination followed by tapering of prednisone over 3 months.

For patients with MF-associated anemia with splenomegaly and constitutional symptoms that are well controlled on a current JAK inhibitor, enrollment in a clinical trial is preferred. JAK inhibitor combinations are other recommended regimens. ESAs (if serum EPO <500 mU/mL), luspatercept-aamt, or danazol (category 2B) can be added to JAK inhibitors. Changing to momelotinib or pacritinib may be useful in certain circumstances. JAK inhibitors (fedratinib, momelotinib, pacritinib, or ruxolitinib) may be continued for the improvement of splenomegaly and other disease-related symptoms.

Treatment Response Criteria

In 2006, the IWG-MRT first published the response criteria for MF, and the responses were categorized as CR, PR, clinical improvement, PD, SD, and relapse.²⁵⁴ In 2013, these response criteria were revised by IWG-MRT and ELN to include MPN-SAF TSS as a quantifiable tool to assess changes in disease-related symptoms and stricter definitions of RBC transfusion dependency and independency.²⁵⁵ In 2024, the response criteria for anemia were updated.²⁵⁶ These response criteria were developed mainly for use in clinical trials.

In addition to CR, PR, and clinical improvement, three other response categories (anemia response, spleen response, and symptoms response) were included in the revised 2013 IWG-MRT and ELN response criteria to quantify treatment-induced improvements in



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

symptom burden, particularly anemia, splenomegaly, and constitutional symptoms.²⁵⁵ The revised response criteria recommend that symptoms should be evaluated by the MPN-SAF TSS and that symptom response requires $\geq 50\%$ reduction in the TSS.⁸⁶ The revised 2013 IWG-MRT and ELN response criteria also require that a $\geq 35\%$ reduction in spleen volume should be confirmed by MRI or CT scan; volumetric imaging of the spleen is typically included in clinical trials to adjudicate this endpoint.²⁵⁵ In addition, a $\geq 35\%$ reduction in spleen volume by MRI or CT scan constitutes a spleen response regardless of that reported by physical examination. Additional criteria are also included for PD, SD, and relapse.

Morphologic response in bone marrow is required for CR. The criteria for PR require morphologic response in the peripheral blood (but not necessarily in the bone marrow). Patients meeting criteria for CR with inadequate blood count recovery are also included in the PR category to capture those patients who have achieved CR with persistent drug-induced cytopenia despite a morphologically normal bone marrow. The revised response criteria also include response categories for cytogenetic and molecular response. However, these are not required for CR assignment.

In the 2024 update, the hemoglobin threshold was updated to take gender into consideration.²⁵⁶ The definition for transfusion-dependent anemia and the response criteria for anemia were updated. Criteria for minor response, duration of anemia response, progressive anemia, and stable anemia were introduced. The definitions for major versus minor responses are based on whether the patient is transfusion-dependent or transfusion-independent at baseline.

Monitoring Response and Follow-up Therapy for Lower- and Higher-Risk MF

The goal of treatment is to reduce symptom burden and minimize the risk of leukemic transformation. Changes in symptom status could be a sign of disease progression. Therefore, change in symptom status should prompt evaluation of treatment efficacy and/or disease status. Evaluation of treatment efficacy should include CBC to assess normalization of blood counts, monitoring symptom status using MPN-SAF TSS, and monitoring spleen size either by palpation or imaging.²⁵⁵

The guidelines recommend monitoring for intolerance, response (anemia response, spleen response, and symptom response), signs, and symptoms of disease progression as clinically indicated during the course of treatment. Bone marrow aspirate and biopsy with NGS and karyotyping should be performed as clinically indicated (if supported by increased symptoms and signs of progression). Additional molecular testing using a multi-gene NGS panel to evaluate for HMR mutations associated with disease progression should be considered for patients with MF.^{39,40}

Continuation of JAK inhibitors is recommended for patients achieving response to initial treatment. In the COMFORT-I study, the majority of patients (91%) treated with ruxolitinib experienced significant improvements in individual MF-related symptoms ($\geq 50\%$ improvement in total symptom score as assessed by MF-SAF) and quality of life; most importantly, patients with a lesser degree of symptom improvement ($< 50\%$ improvement in total symptom score) also achieved improvements over placebo on these measures and other patient-reported outcomes.⁸⁷ The Panel acknowledges that clinical benefit may not reach the threshold of the 2013 IWG-MRT and ELN Response Criteria (ie, symptom response requires $\geq 50\%$ reduction in the MPN-SAF TSS) in patients receiving treatment with JAK inhibitors. Response assessment should be done based on the improvement of disease-related symptoms at the discretion



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

of the clinician. The RR6 model may also be used to gauge response to ruxolitinib.¹⁶³ Continuation of JAK inhibitors is also recommended based on the discretion of the clinician, since a symptom response of <50%, as well as spleen volume reduction that does not meet the threshold of >35% (reduction in palpable splenomegaly of <50%), may be clinically meaningful.

Disease-related symptoms may return to pretreatment levels over a period of approximately 1 week following discontinuation or interruption of ruxolitinib.²⁵⁷ Low platelet counts (at initiation or completion of therapy) and clonal evolution (acquisition of new mutations while on treatment with ruxolitinib) were associated with a significantly shorter survival after discontinuation of ruxolitinib.²⁵⁸ In a study that evaluated the outcomes of ruxolitinib discontinuation in patients with MF, after a median follow-up of 32 months, the median survival was 14 months among 42 patients who had molecular data at baseline; during follow-up, clonal evolution was seen in 14 patients (33%; *ASXL1* mutation in 60% of patients).²⁵⁸ RBC transfusion dependence at baseline was the only clinical variable associated with clonal evolution; survival after discontinuation of ruxolitinib was 6 months for patients with clonal evolution compared to 16 months for those without clonal evolution. A population-based analysis of 290 patients with MF found that 50% of patients developed cytopenias after terminating treatment with ruxolitinib.²⁵⁹ The median OS after discontinuation was 11.1 months (95% CI, 8.4–14.5 months) and the median PFS was 6.0 months (95% CI, 4.4–8.3 months).

For patients with symptomatic lower-risk MF with intolerance, no response, or loss of response following initial treatment, an alternate option not used for initial treatment is recommended (clinical trial, ruxolitinib, peginterferon alfa-2a, hydroxyurea [if cytoreduction would be symptomatically beneficial], pacritinib [if platelets <50 x 10⁹/L], or

momelotinib [category 2B]). If anemia is present, see *Management of MF-Associated Anemia* in the algorithm.

For patients with higher-risk MF with platelet count ≥50 x 10⁹/L who are not candidates for transplant and who have intolerance, no response or loss of response following initial treatment, enrollment in a clinical trial or an alternate JAK inhibitor (ruxolitinib, fedratinib, momelotinib, or pacritinib [category 2B]) not used before is recommended.^{93,97,120,165} If anemia is present, see *Management of MF-Associated Anemia* in the algorithm.

JAK2 V617F Allele Burden

Reductions in *JAK2* V617F allele burden have been observed in patients with MF with long-term fedratinib²⁶⁰ or ruxolitinib therapy.^{139,261} In the COMFORT-I study, a >50% reduction in *JAK2* V617F allele burden were observed in 12% of patients (28 patients); 20 of these patients met the criteria for partial molecular response (PMR) and six patients had *JAK2* V617F allele burden values below the quantifiable limit, meeting the criteria for complete molecular response (CMR).²⁶¹ The median times to PMR and CMR were 22 months and 28 months, respectively. *JAK2* V617F allele burden reductions also correlated with spleen volume reductions. Achievement of *JAK2* V617F negativity or *JAK2* V617F allele burden reduction after allogeneic HCT has also been associated with a decreased incidence of relapse.^{262,263}

However, at the present time, the utility of *JAK2* V617F allele burden reduction as a predictor of treatment efficacy remains unclear. In the 2013 IWG-MRT and ELN response criteria, cytogenetic and molecular responses are not required for CR assignment.²⁵⁵ Therefore, measurement of the *JAK2* V617F allele burden is not currently recommended for use in routine clinical practice to guide treatment decisions.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Supportive Care

Supportive care for disease-related symptoms should be an integral part of clinical management during the course of treatment. This should include assessment and monitoring of symptom status and counseling for identification, assessment, and management of cardiovascular risk factors (eg, smoking, diet, exercise, hypertension, diabetes mellitus, lipid management) and thrombotic and hemorrhagic risk factors.

Transfusion support should include platelet transfusions for thrombocytopenic bleeding or platelet count $<10 \times 10^9/L$ and RBC transfusions for symptomatic anemia.²⁶⁴ The use of leukocyte-reduced blood products is recommended in transplant candidates to prevent HLA alloimmunization and reduce the risk of cytomegalovirus transmission. Antifibrinolytic agents should be considered for bleeding that is refractory to transfusions. Iron chelation could be considered for patients who have received >20 transfusions and/or ferritin >2500 ng/mL in patients with lower-risk disease. However, the role of iron chelation remains unclear.

In patients with massive splenomegaly who do not respond to JAK inhibitors, alternative measures to reduce spleen size may be considered (eg, splenic radiation, splenic artery embolization, splenectomy).^{213,214} Vaccinations are recommended as outlined in the NCCN Guidelines for the Prevention and Treatment of Cancer-Related Infections (available at www.NCCN.org). A recombinant (killed) zoster vaccine may be considered for patients on, or prior to, treatment with a JAK inhibitor. In patients who have had a splenectomy, vaccinations and antibiotic prophylaxis should be given per the Infectious Diseases Society of America (IDSA) Guidelines. Growth factor support (granulocyte colony-stimulating factor [G-CSF] or granulocyte-macrophage colony-stimulating factor [GM-CSF]) should be considered for recurrent infections with neutropenia. However, these should be used with caution in patients with an enlarged spleen since the use of G-CSF or GM-CSF has been associated with splenic rupture.²⁶⁵

Cytoreductive therapy (eg, hydroxyurea) could be considered for the management of hyperproliferative manifestations of PMF (thrombocytosis or leukocytosis).²⁰⁵ Prophylaxis for tumor lysis syndrome should be considered for patients undergoing induction chemotherapy for advanced-stage MF or disease progression to AML.

Management of Polycythemia Vera and Essential Thrombocythemia

Referral to specialized centers with expertise in the management of MPN is strongly recommended for all patients diagnosed with PV or ET.

Risk Stratification

Studies have shown that leukocytosis at diagnosis is associated with higher risk of thrombosis and major hemorrhage in patients with PV and ET.²⁶⁶⁻²⁷¹ Data from some studies suggest that the prognostic significance of leukocytosis for the risk of recurrent thrombosis may be significant only in patients <60 years of age,^{272,273} and other studies have reported that leukocytosis at diagnosis is not associated with the risk of subsequent thrombosis.²⁶⁷ Thrombocytosis (platelet count $>1000 \times 10^9/L$) has been associated with an immediate risk of major hemorrhage but not with the risk of thrombosis in patients with ET.²⁷⁰ In fact, some studies have reported that elevated platelet counts at diagnosis ($>1000 \times 10^9/L$) are associated with significantly lower rate of thrombosis; this association was significant even in patients with JAK2-mutated ET.^{268,269} The potential benefit of initiation of cytoreductive therapy based on elevated blood counts (leukocytosis or thrombocytosis) at the time of diagnosis has not been evaluated in prospective studies.

Arterial/Venous Thrombosis Score

In a multivariable analysis of 1057 patients with MPN (training cohort), prior arterial thrombosis, age >60 years, presence of cardiovascular risk factors, and presence of *TET2* or *DNMT3A* mutations were identified as



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

independent predictors of arterial thrombosis and were used to calculate an arterial thrombosis score.²⁷⁴ This model stratified patients into two risk categories: low risk (0.37% patients-year) and high risk (1.19% patients-year), and was deemed to be superior when compared to the two-tiered conventional risk model in the training cohort, regardless of the MPN subtype. The arterial thrombosis score model was also validated in two external cohorts. A prior history of venous thrombosis and the presence of *JAK2* V617F mutation with VAF $\geq 50\%$ were identified as independent predictors of venous thrombosis and were used to calculate a venous thrombosis score. This model stratified patients into three risk categories: low risk, intermediate risk, and high risk. However, this model had a low prediction potential and was found to be similar to the two-tiered conventional risk model.

Polycythemia Vera

Conventional Risk Model

Advanced age (ie, >60 years) and history of thrombosis are the most consistent risk factors associated with the risk of thrombosis.²⁷⁵ In a cohort of 1638 patients with PV who were screened for inclusion in the ECLAP trial, age >65 years and a previous history of thrombosis were the two most important prognostic factors associated with an increasing risk of cardiovascular events resulting in the identification of three different risk groups: low risk (age <65 years and no prior history of thrombosis), intermediate risk (age <65 years with prior thrombosis or age ≥ 65 years without prior thrombosis), and high risk (age ≥ 65 years with prior thrombosis). There is a consensus to use age ≥ 60 years or history of thrombosis as prognostic factors for the risk of thrombosis.^{276,277}

MIPSS-PV

In a study of 336 patients with PV, the presence of *SRSF2* mutation, age >67 years, leukocyte count $\geq 15 \times 10^9/L$, and history of thrombosis were identified as independent risk factors for survival.²⁷⁸ Based on these

findings, MIPSS-PV was developed. Patients were stratified into three risk categories: low risk, intermediate risk, and high risk, with a median OS of 24 years, 13.1 years, and 3.2 years, respectively. Further studies are needed to validate these findings.

Essential Thrombocythemia

IPSET-Thrombosis

In an analysis of 867 patients with ET, age ≥ 60 years, leukocyte count $\geq 11 \times 10^9/L$, and prior thrombosis were significantly associated with inferior survival.²⁷⁹ Based on these findings, IPSET was developed to stratify patients at the time of diagnosis into three risk categories: low risk, intermediate risk, and high risk. The median survival was not reached for the low-risk group and the median survival was 24 years and 14 years, respectively, for the intermediate-risk and high-risk groups. In a subsequent analysis of 891 patients with ET, age >60 years, history of thrombosis, cardiovascular risk factors, and presence of *JAK2* V617F mutation retained their prognostic significance regarding thrombosis risk in multivariable analysis.²⁸⁰ Thus, a modified prognostic model (IPSET-thrombosis) including cardiovascular risk factors and presence of *JAK2* V617F mutation status as additional risk factors was developed to stratify patients into the same three groups with significantly different thrombosis-free survival: 87% after 15-year follow-up for patients with low-risk disease and 50% after 7-year follow-up for patients with high-risk disease.²⁸⁰ In the intermediate-risk group, the thrombosis-free survival rate for the first 10 years was closer to that of the low-risk group and then progressively reached the high-risk survival rate in the subsequent 5 years.

Further analysis of the IPSET-thrombosis showed that among the patients with low-risk disease, the risk of thrombosis was significantly lower in patients with *JAK2*-negative/unmutated ET in the absence of cardiovascular risk factors (0.44%) compared to the risk of thrombosis in



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

patients with *JAK2* unmutated ET in the presence of cardiovascular risk factors (1%).²⁸¹ The risk of thrombosis in the presence of *JAK2* mutation without cardiovascular risk factors and in the presence of both *JAK2* mutation and cardiovascular risk factors was 2% and 3%, respectively. These findings led to the development of revised IPSET-thrombosis that stratifies patients into four different risk groups: very low risk (age ≤60 years, no *JAK2* mutation, and no prior history of thrombosis); low risk (age ≤60 years, *JAK2* mutation, and no prior history of thrombosis); intermediate risk (age >60 years, no *JAK2* mutation, and no prior history of thrombosis); and high risk (history of thrombosis at any age; or age >60 years with *JAK2* mutation). The revised IPSET-thrombosis has also been validated in an independent cohort of 585 patients.^{281,282}

CALR mutation status, however, did not have a significant impact on the IPSET-thrombosis prognostic score for predicting the risk of thrombosis.⁵⁸ While the incidences of thrombosis were slightly lower in patients with *CALR*-mutated ET than in those with *JAK2*-mutated ET, in multivariable analysis, *CALR* mutation status did not retain the association with the risk of thrombosis in low-risk and intermediate-risk groups. In part, this may be explained by the fact that *CALR* mutation status tended to cluster with other lower-risk features. The significance of *CALR* mutations and the risk of thrombosis could not be evaluated in the high-risk group since there was a lower proportion of patients with the *CALR* mutation in this group.

MIPSS-ET

In a study of 451 patients with ET, the presence of adverse mutations (ie, *SF3B1*, *SRSF2*, *TP53*, *U2AF1*), age >60 years, male sex, and leukocyte count ≥11 x 10⁹/L were identified as independent risk factors for survival.²⁷⁸ Based on these findings, MIPSS-ET was developed. Patients were stratified into three risk categories: low risk, intermediate risk, and high risk, with a median OS of 34.4 years, 14.1 years, and 7.9 years, respectively. Further studies are needed to validate these findings.

“Triple A” Risk Model

The prognostic AAA risk model for ET is based on age, absolute neutrophil count, and absolute lymphocyte count. Data from 598 patients were used and this model was validated in an external cohort of 485 patients. Four risk categories were identified: low risk (47-year median survival), intermediate-1 risk (20.7-year median survival; HR, 3.8), intermediate-2 risk (13.5-year median survival; HR, 12.7), and high risk (8-year median survival; HR, 30.1). Age ≥50 years, ANC ≥8 x 10⁹/L, absolute lymphocyte count <1.7 x 10⁹/L were identified as independent predictors of survival.

Treatment Options

Antiplatelet Therapy

The safety and efficacy of low-dose aspirin for the prevention of thrombotic complications in PV were established in a multicenter trial in patients with no contraindication to aspirin therapy and no history of a thrombotic event (ECLAP study; 518 patients).²⁸³ The use of aspirin resulted in a significant reduction (60%) of combined risk of nonfatal myocardial infarction, nonfatal stroke, pulmonary embolism, major venous thrombosis, or death from cardiovascular causes (*P* = .03) and the incidence of major bleeding was not significantly increased in the aspirin group. The role of maintaining the hematocrit level below 45% in patients receiving treatment was established in the CYTO-PV study.²⁸⁴ In this randomized study of 365 patients with PV treated with phlebotomy and/or hydroxyurea, the hematocrit target of <45% resulted in a significantly lower rate of cardiovascular death and major thrombotic events (primary endpoint) than a hematocrit target of 45% to 50%.²⁸⁴ After a median follow-up of 31 months, death from cardiovascular causes or major thrombotic events was recorded in 3% (5 of 182 patients) of patients with a hematocrit level of <45% compared to 10% (18 of 183 patients) of patients with a hematocrit level of 45% to 50% (*P* = .007).



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

The efficacy of low-dose aspirin for the prevention of thrombosis in patients with ET has not been evaluated in randomized clinical trials. The data supporting the use of aspirin in patients with ET is based on the extrapolation of results from the ECLAP study that evaluated the efficacy of aspirin in patients with PV and the results of retrospective analyses.^{285,286} Results from one retrospective analysis suggest that aspirin may be effective for the prevention of thrombosis in patients with low-risk *JAK2*-mutated ET and in those with cardiovascular risk factors.²⁸⁵ Observation may be appropriate for all other patients with low-risk ET. In this retrospective analysis of 300 patients with low-risk ET managed with aspirin (n = 198) or observation (n = 102), the incidences of venous thrombosis were higher for those with *JAK2* V617F-positive ET not receiving any antiplatelet therapy; patients with cardiovascular risk factors had increased rates of arterial thrombosis while on observation.²⁸⁵

Cytoreductive Therapy

Hydroxyurea,^{284,287,288} peginterferon alfa,²⁸⁹⁻²⁹² and ropeginterferon alfa-2b-njft^{206,293} have been shown to be effective for the prevention of thrombotic complications in patients with PV.

Hydroxyurea for the Treatment of PV

In a nonrandomized study of 51 patients with PV, the use of hydroxyurea along with phlebotomy as needed significantly reduced the risk of thrombosis compared to a historical control of patients treated with phlebotomy alone.²⁸⁷ Long-term follow-up of this study (after a median follow-up of 9 years) showed that prolonged use of hydroxyurea was associated with leukemic transformation (6% compared to 2% for phlebotomy).²⁹⁴ However, an analysis from the ECLAP study identified older age and the use of other alkylating agents (eg, P32, busulfan, pipobroman) but not hydroxyurea alone as an independent risk factor for leukemic transformation.²⁹⁵ In the randomized trial that compared hydroxyurea and pipobroman as first-line therapy in 285 patients with PV

<65 years of age, the cumulative incidence of leukemic transformation was significantly higher with pipobroman than with hydroxyurea.²⁸⁸ At a median follow-up of 15 years the incidences of leukemic transformation were 17% and 34%, respectively, for hydroxyurea and pipobroman.

Peginterferon alfa-2a for the Treatment of PV

In a phase II multicenter study of 40 patients with PV, peginterferon alfa-2a resulted in high rates of complete hematologic response (CHR; 95%) and CMR (24%) with limited toxicity.²⁹⁰ At a median follow-up of 31 months, 36 patients with a response remained phlebotomy free. A phase II trial that included 43 patients with PV reported a CHR rate of 77% and a CMR rate of 20% after a median follow-up of 83 months.²⁹² The duration of response was longer among patients with CMR (70 months) than for those with CHR (65 months). The presence of *TET2*, *ASXL1*, *EZH2*, *DNMT3A*, and *IDH1/2* mutations was associated with non-achievement of CMR.²⁹¹ Patients with both *JAK2* V617F and *TET2* mutations at initiation of treatment had a less significant reduction in *JAK2* V617F allele burden compared to those with *JAK2*-mutated/*TET2* wild-type disease.

A phase III study comparing hydroxyurea to peginterferon alfa-2a in patients with high-risk PV or ET reported no significant difference in CR rates at 12 months (37% vs 35%; *P* = .80).²⁹⁶ However, the authors note that with prolonged treatment, hydroxyurea elicited a greater number of histopathologic responses, while peginterferon-alfa-2a resulted in a greater reduction in the *JAK2* V617F mutation burden. Grade ≥3 adverse events, irrespective of cause, also occurred more frequently with peginterferon alfa-2a treatment (46% vs 28% for hydroxyurea).

Rpeginterferon alfa-2b-njft for the Treatment of PV

Rpeginterferon alfa-2b-njft was FDA-approved in 2021 for the treatment of adult patients with PV. In the phase II Low-PV trial comprising 127 patients, a higher proportion of patients treated with ropeginterferon-



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

alfa2b-njft in addition to phlebotomy achieved the primary endpoint, defined as the maintenance of a median hematocrit level of $\leq 45\%$ over 12 months in the absence of disease progression, when compared to those treated with phlebotomy alone (81% vs 51%; $P < .001$).²⁹³ At 24 months, the response rates were maintained (ropeginterferon-alfa-2b-njft with phlebotomy, 83%; phlebotomy alone, 59%; $P = .02$).

In the phase III PROUD-PV trial, patients received either ropeginterferon alfa-2b-njft ($n = 127$) or hydroxyurea ($n = 127$).²⁰⁶ The composite primary endpoint was the achievement of CHR and normal spleen size by imaging. At 12 months, noninferiority was not demonstrated ($P = .23$) with 21% of patients in the ropeginterferon alfa-2b-njft group and 28% in the hydroxyurea group achieving the composite primary endpoint. Not accounting for the spleen, 43% of patients achieved CHR in the ropeginterferon alfa-2b-njft group compared to 46% in the hydroxyurea group ($P = .63$). At the end of the 12-month PROUD-PV trial, patients were eligible to enter the CONTINUATION-PV extension study. Patients taking ropeginterferon alfa-2b-njft ($n = 95$) remained on the drug and those taking hydroxyurea received BAT ($n = 76$), chosen by the investigator. The co-primary endpoints were the achievement of CHR and normal spleen size as well as CHR accompanied by improved disease burden. Patient response to ropeginterferon alfa-2b-njft improved over time and, at 36 months, was significantly higher as CHR with improved disease burden was reported in 53% of patients, compared to 38% in the hydroxyurea group ($P = .044$). However, there was no significant difference at 36 months in terms of CHR with normal spleen size, with a response rate of 42% in the ropeginterferon alfa-2b-njft group and 30% in the hydroxyurea group ($P = .16$). Not accounting for the spleen, CHR was reported in 71% of patients in the ropeginterferon alfa-2b-njft group and in 51% of patients in the hydroxyurea group ($P = .012$). Across both studies, the most common grade 3 and 4 adverse occurrences for patients on ropeginterferon alfa-2b-njft were increased γ -glutamyltransferase and

alanine aminotransferase and for those on hydroxyurea were leucopenia and thrombocytopenia.

Data from the PROUD-PV and CONTINUATION-PV trials at 5 years revealed a CHR rate and molecular response rate of 55.8% and 69.1%, respectively, in patients treated with ropeginterferon alfa-2b-njft compared to 44.0% (rate ratio, 1.30; $P = .0974$) and 21.6% (rate ratio, 3.04; $P < .0001$), respectively, in patients treated with BAT, which was mostly hydroxyurea.²⁹⁷ In the ropeginterferon alfa-2b-njft group, the median JAK2 V617F allele burden decreased from 37.3% at baseline to 8.5% at 60 months whereas in the BAT group, a decrease was observed at 12 months (38.1% at baseline to 18.2%) but at 60 months, the percentage was at 44.4% ($P < .0001$). The rates of treatment-related adverse events were similar in both groups, irrespective of prior treatment with hydroxyurea.

At 72 months, patients treated with ropeginterferon alfa-2b-njft maintained a higher CHR rate compared to those treated with BAT (54.5% vs 34.9%; $P = .02$).²⁹⁸ After 6 years, 66.0% of patients in the ropeginterferon alfa-2b-njft arm had a molecular response compared to 19.4% in the control arm ($P < .0001$), with a median JAK2 V617F allele burden of 8.5% and 50.4%, respectively ($P < .0001$), at 72 months. Patients in the former group also had a significantly higher probability of EFS (0.94 vs 0.82 in the control group; $P = .04$), with 5.3% of patients in the former group having a risk event (thromboembolic event, 2; MF, 1; death, 2) versus 16.2% in the control group (thrombotic event, 5; MF, 2; acute leukemia, 2; death, 2).

Ruxolitinib for the Treatment of PV

A futility analysis of the phase IIb RuxoBEAT study showed that in patients with PV with no prior treatment, ruxolitinib resulted in a decrease in the median hematocrit, the median number of phlebotomies received per year, and the median pruritus scores at 6 months.²⁹⁹ Adverse events were reported in 24 out of 28 patients.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

The results of the phase III randomized trial (RESPONSE) confirmed that ruxolitinib is superior to BAT (hydroxyurea, interferon or pegylated interferon, pipobroman, anagrelide, lenalidomide, thalidomide, or observation with the use of aspirin) at controlling hematocrit and improving splenomegaly and symptoms in patients with PV.^{91,300,301}

In this study, 222 patients with PV who are phlebotomy-dependent with splenomegaly and whose disease had an inadequate response to or was intolerant of hydroxyurea were randomized to receive ruxolitinib (110 patients) or BAT (112 patients).⁹¹ The primary endpoint was hematocrit control without phlebotomy and at least a 35% reduction in spleen volume (as assessed by imaging) by 32 weeks. Patients randomized to BAT were eligible to cross over to ruxolitinib after 32 weeks if the primary endpoint was not met or if there were signs of disease progression. After 32 weeks, hematocrit control was achieved in 60% of patients treated with ruxolitinib compared to 20% of patients treated with BAT. A reduction in spleen volume ($\geq 35\%$), CHR, and at least a 50% reduction in symptom burden were achieved in 38%, 24%, and 49% of patients, respectively, in the ruxolitinib group and in 1%, 9%, and 5% of patients, respectively, in the BAT group. The incidences of grade 3/4 anemia and herpes zoster infection were higher among patients treated with ruxolitinib (occurring in 2% and 6% of patients, respectively, compared to 0% of patients treated with BAT). The 80-week follow-up data confirmed the long-term efficacy of ruxolitinib, and the probability of maintaining CHR for ≥ 80 weeks was 69%.³⁰⁰ Ruxolitinib was also associated with a lower rate of thromboembolic events (1.8% and 4.1%, respectively, for patients originally randomized to ruxolitinib and for those receiving ruxolitinib after crossover compared to 8.2% for those receiving BAT). The 5-year follow-up of the RESPONSE study further confirmed the safety and efficacy of ruxolitinib as a long-term option for patients with PV that is resistant to or intolerant of hydroxyurea.³⁰² By week 80, patients who did not cross over to the ruxolitinib arm discontinued the study. The probability of maintaining

the primary endpoint response, complete hematologic remission, and overall clinicohematologic response at 5 years was 74% (95% CI, 51%–88%), 55% (95% CI, 32%–73%), and 67% (95% CI, 54%–77%), respectively. Compared to the BAT study arm, the patients in the ruxolitinib study arm experienced fewer thromboembolic and nonhematologic adverse events.

In a subsequent phase IIIb study (RESPONSE-2), ruxolitinib was shown to be effective for the treatment of PV with an inadequate response to hydroxyurea in patients without splenomegaly.³⁰³ A follow-up study performed 80 weeks later revealed sustained CHR in 24% of patients receiving ruxolitinib compared to 3% of patients receiving BAT.³⁰⁴ Of those receiving BAT, 77% crossed over to the ruxolitinib arm after week 28. At 80 weeks, patients discontinued BAT.³⁰⁵ At 5 years, durable hematocrit control was reported in 22% of patients in the ruxolitinib group.³⁰⁵ The results of another phase III study showed that ruxolitinib was also effective and resulted in improvements in symptoms (although non-significant) compared to hydroxyurea in patients with well-controlled PV; however, other disease-associated symptoms were reported.³⁰⁶

Results from the phase II MAJIC-PV study demonstrated the benefit of ruxolitinib over BAT in patients with PV that is resistant or intolerant to hydroxyurea.³⁰⁷ Forty-three percent of patients treated with ruxolitinib achieved a CR within 1 year compared to 26% of patients treated with BAT (OR, 2.12; 90% CI, 1.25–3.60; $P = .02$). Ruxolitinib treatment also led to more frequent molecular responses, which were associated with improved PFS, EFS, and OS. The presence of additional driver mutations negatively impacted EFS.

Hydroxyurea,³⁰⁸⁻³¹⁰ peginterferon alfa-2a,^{291,292} and possibly anagrelide³⁰⁹⁻³¹² have been shown to be effective for the prevention of venous thrombotic complications in patients with high-risk ET.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Hydroxyurea for the Treatment of ET

In a study of 114 patients with high-risk ET (>60 years and high risk of thrombosis) randomized to receive hydroxyurea ($n = 56$), which was administered to maintain the platelet count $<600 \times 10^9/L$ or no myelosuppressive therapy ($n = 58$), the incidences of thrombotic episodes were significantly lower in patients treated with hydroxyurea (3.6% compared to 24% in patients with no myelosuppressive therapy; $P = .003$).³⁰⁸ In another randomized study of 809 patients with high-risk ET, hydroxyurea plus low-dose aspirin was superior to anagrelide plus low-dose aspirin.³⁰⁹ Patients in the hydroxyurea arm initially received the drug at a dose of 0.5 to 1 g daily, while those in the anagrelide arm received the drug at a dose of 0.5 mg twice daily. The dose of the drugs was adjusted subsequently to keep the platelet count at $<400 \times 10^9/L$. After a median follow-up of 39 months, the long-term control of platelet counts was equivalent in both groups and anagrelide plus aspirin was better in the prevention of venous thrombosis ($P = .006$). However, the incidences of arterial thrombosis ($P = .004$), serious hemorrhage ($P = .008$), and transformation to MF ($P = .01$) were higher with anagrelide plus aspirin. In addition, treatment discontinuation rate was also significantly higher with anagrelide plus aspirin. The diagnosis of ET in this trial was based on the Polycythemia Vera Study Group criteria.

Anagrelide for the Treatment of ET

A phase III randomized study showed that anagrelide was not inferior to hydroxyurea as first-line therapy for the prevention of thrombotic complications in patients with high-risk ET diagnosed according to the WHO criteria.³¹⁰ In this study, 259 patients were randomized to either hydroxyurea ($n = 122$) or anagrelide ($n = 137$). The dose of the drugs was increased until platelet counts were maintained at a normal level ($\leq 450 \times 10^9/L$) or close to it ($>450 \times 10^9/L$ to $600 \times 10^9/L$). After a total observation time of 730 patient-years, there was no significant difference between anagrelide and hydroxyurea in the incidences of arterial or venous

thrombotic events, severe bleeding, or rates of discontinuation. Another study showed that over a median period of 10 years, patients taking anagrelide experienced fewer minor arterial events ($P < .001$), had more major arterial events ($P = .049$), and had improved OS ($P = .001$) and PFS ($P = .004$) compared to patients taking hydroxyurea and aspirin.³¹²

Peginterferon alfa-2a for the Treatment of ET

In a phase II trial that included 40 patients with ET, peginterferon alfa-2a induced a CHR rate of 73% and a CMR rate of 9% after a median follow-up of 83 months.²⁹² The presence of *TET2*, *ASXL1*, *EZH2*, *DNMT3A*, and *IDH1/2* mutations was associated with non-achievement of CMR.²⁹¹ Patients with both *JAK2* V617F and *TET2* mutations at initiation of treatment had a less significant reduction in *JAK2* V617F allele burden compared to those with *JAK2*-mutated or *TET2* wild-type disease. The phase II Myeloproliferative Disorders Research Consortium-111 study consisted of patients with high-risk ET ($n = 65$) or PV ($n = 50$) that is resistant or intolerant to hydroxyurea. Treatment with peginterferon alfa-2a resulted in a 12-month ORR of 69% and 60%, respectively.³¹³ Patients with ET who have a *CALR* mutation had increased CHR rates compared to those without a *CALR* mutation. Fourteen percent of patients discontinued treatment due to adverse events.

Ropeginterferon alfa-2b-njft for the Treatment of ET

In the phase III SURPASS ET study, patients with high-risk ET that is intolerant or resistant to hydroxyurea and with leukocytosis were randomly assigned to be treated with ropeginterferon alfa-2b-njft ($n = 91$) or anagrelide ($n = 83$).²⁰⁷ The results showed the superiority of ropeginterferon alfa-2b-njft over anagrelide, with 43% of those in the ropeginterferon alfa-2b-njft arm achieving the primary endpoint of durable modified ELN response criteria at months 9 and 12, compared to 6% of those in the anagrelide arm ($P = .0001$). Treatment with ropeginterferon alfa-2b-njft also resulted in significantly higher response rates in the other



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

parameters of the primary endpoint ($P < .0001$): peripheral blood count response (56% vs 6%), WBC response (74% vs 13%), platelet response (56% vs 22%), improvement or non-progression of splenomegaly (88% vs 54%), symptom response based on the MPN-SAF-TSS (71% vs 34%), and a lack of hemorrhagic or thrombotic events (85% vs 52%).

Additionally, of the evaluable patients treated with ropeginterferon alfa-2b-njft, there was a 21.3% decrease in the *JAK2* V617F VAF compared to a 12.3% increase in patients treated with anagrelide ($P < .0001$). Twenty-three percent of those in the former group had grade ≥ 3 treatment-emergent adverse events compared to 34% in the latter group, with infections being the most frequent adverse event (9% in the ropeginterferon alfa-2b-njft arm vs 6% in the anagrelide arm). Nervous system disorders were the most frequent grade ≥ 3 treatment-emergent adverse events (8% vs 1% in the ropeginterferon alfa-2b-njft arm).

Ruxolitinib for the Treatment of ET

The phase II MAJIC-ET trial investigated the efficacy of ruxolitinib versus BAT in patients with ET that is resistant or intolerant to hydroxyurea.⁹² The CR rates at 1 year, as well as occurrence of thrombosis, hemorrhage, and disease transformation at 2 years were similar in both groups. Ruxolitinib use was associated with a decrease in some disease-related symptoms, with a median total symptom score reduction of 32%, compared to 0% for patients receiving BAT ($P = .03$). An expanded analysis of the trial revealed that the presence of *TP53* and splicing factor mutations led to poorer transformation-free survival.³¹⁴ Treatment with ruxolitinib did not modify disease transformation. Another phase II study found that long-term treatment with ruxolitinib in patients with ET that is refractory to or intolerant of hydroxyurea led to lasting reductions in platelet counts and amelioration of ET-related symptoms.³¹⁵

Treatment Recommendations Based on Risk Stratification

Treatment options should be individualized based on age and history of thrombosis for patients with PV,²⁷⁵ and the revised IPSET-thrombosis is recommended for the risk stratification of patients with ET.^{281,282}

Polycythemia Vera

Low Risk (Age <60 years and no prior history of thrombosis)

Aspirin (81–100 mg/day), phlebotomy (to maintain hematocrit <45%), and the management of cardiovascular risk factors are recommended for all patients with low-risk PV.^{283,284} In the CYTO-PV study, the hematocrit target was the same for both males and females. No thrombotic event was observed in the 66 females with hematocrit of <45% compared to nine events reported in the 72 females with a hematocrit target of 45% to 50%.²⁸⁴ However, normal hematocrit levels vary in males (42%–54%) and females (38%–46%). While the target hematocrit level of <45% may be adequate for the majority of patients, there may be situations in which a lower hematocrit cutoff may be appropriate and it should be individualized (eg, for patients with progressive symptoms).

High Risk (Age ≥ 60 years and/or prior history of thrombosis)

In addition to aspirin and phlebotomy, cytoreductive therapy is also used to reduce the risk of thrombotic complications in patients with high-risk PV. Management of cardiovascular risk factors is recommended. Cytoreductive therapy with aspirin (81–100 mg/day) for vascular symptoms and phlebotomy (to maintain hematocrit <45%) is recommended. Cytoreductive therapy options comprise hydroxyurea (preferred regimen), ropeginterferon alfa-2b-njft (preferred regimen), peginterferon alfa-2a (other recommended regimen), and ruxolitinib (useful in certain circumstances). Peginterferon alfa-2a is an option for younger patients or in pregnant patients in need of cytoreductive therapy. If



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

peginterferon alfa-2a is unavailable, the use of other pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate.

Essential Thrombocytopenia

Very-Low Risk (Age ≤60 years without JAK2 mutation and no prior history of thrombosis), Low Risk (age ≤60 years with JAK2 mutation and no prior history of thrombosis), or Intermediate Risk (age >60 years, no JAK2 mutation, and no prior history of thrombosis)

As discussed above, the efficacy and safety of low-dose aspirin in patients with ET has not been evaluated in randomized clinical trials. The results of a systematic review also suggest that the risks and benefits of antiplatelet therapy in patients with ET remain highly uncertain.³¹⁶ Observation is appropriate for patients with very-low-risk, low-risk, and intermediate-risk ET. Aspirin (81–100 mg/day) is an option for patients with very-low-risk (with vasomotor/microvascular disturbances), low-risk, or intermediate-risk ET. Aspirin should be used with caution in patients with acquired VWS who have an increased risk of bleeding.

A report from a retrospective analysis suggests that the use of low-dose aspirin may not be beneficial in patients with low-risk *CALR*-mutated ET.²⁸⁶ In an analysis that evaluated the benefit-to-risk ratio of low-dose aspirin in 433 patients with low-risk ET (271 patients with a *CALR* mutation and 162 patients with a *JAK2* V617F mutation) who were on antiplatelet therapy or observation, low-dose aspirin did not affect the risk of thrombosis but was associated with a higher incidence of bleeding in patients with *CALR*-mutated ET.²⁸⁶ These findings have to be confirmed in prospective clinical trials.

In carefully selected patients, twice-daily aspirin at a 100-mg dose may be more effective than once-daily aspirin (100 mg).³¹⁷⁻³¹⁹ The results of a randomized phase II trial also showed that 100 mg of aspirin twice daily was more effective than once daily aspirin in reducing thromboxane levels.^{319,320} Another study that compared once-daily aspirin (75 mg) to

twice-daily aspirin (37.5 mg per dose) found that the twice-daily schedule led to improved platelet inhibition as measured by platelet aggregation studies as well.³²¹ These studies did not, and were not powered to, show a difference in thrombosis rates between the two dosing strategies.

Aspirin twice daily may be considered for patients with refractory symptoms or cardiovascular risk factors.³¹⁷⁻³¹⁹ At the present time, the risks and benefits of higher dose aspirin (>100 mg) must be weighed based on the presence of vasomotor symptoms versus the risk of bleeding. It may be appropriate in carefully selected patients as clinically indicated.

In one study, patients with ET and no high-risk factors for thrombosis or extreme thrombocytosis were given either aspirin alone (n = 176) or aspirin with hydroxyurea (n = 182).³²² The dose of hydroxyurea was adjusted in order to maintain platelet count between 200 x 10⁹/L and 400 x 10⁹/L. The results showed that the addition of hydroxyurea to aspirin did not decrease the incidence of vascular events and myelofibrotic or leukemic transformation.

High Risk (History of thrombosis at any age; or age >60 years with JAK2 mutation)

Cytoreductive therapy (clinical trial [preferred regimen], hydroxyurea [preferred regimen], peginterferon alfa-2a [based on other patient-specific variables] [other recommended regimen], or anagrelide [other recommended regimen]) with aspirin (81–100 mg/day) is recommended as initial treatment. Peginterferon alfa-2a can be considered for patients in need of cytoreductive therapy who are younger or pregnant or who defer hydroxyurea. If peginterferon alfa-2a is unavailable, the use of other pegylated interferons (eg, ropeginterferon alfa-2b-njft) is appropriate. Aspirin twice daily may be considered for patients with refractory symptoms or cardiovascular risk factors.³¹⁷⁻³¹⁹



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Treatment Response Criteria

The IWG-MRT and ELN treatment response criteria for PV and ET were first published in 2009 and were revised in 2013.³²³ Responses are categorized as CR, PR, no response, and PD. The revised response criteria recommend that symptoms should be evaluated by the MPN-SAF TSS. The evaluation of CR or PR includes four categories: 1) resolution of disease-related signs and symptoms including palpable splenomegaly and large symptom improvement (≥ 10 point decrease in MPN-SAF TSS); 2) peripheral blood count response (platelet count $\leq 400 \times 10^9/L$, white blood cell [WBC] count $< 10 \times 10^9/L$, absence of leukoerythroblastosis, and hematocrit $< 45\%$ without phlebotomies); 3) absence of signs of PD and absence of any hemorrhagic or thrombotic events; and 4) histologic response in bone marrow. Molecular response is not required for the assignment of CR or PR and the revised IWG-MRT and ELN treatment response criteria do not provide a definition of molecular response.

JAK2 V617F Allele Burden

Long-term ruxolitinib therapy has been shown to reduce *JAK2 V617F* allele burden in patients with PV that is resistant to hydroxyurea.³²⁴ High *JAK2 V617F* allele burden has also been reported as a risk factor for myelofibrotic transformation and higher incidences of thrombotic events in patients with PV and ET.³²⁵⁻³²⁷ These findings suggest that monitoring *JAK2 V617F* allele burden could be useful to identify patients at higher risk of myelofibrotic transformation. It could also be a useful adjunctive evaluation to assess the impact of cytoreduction on molecular response. However, the utility of *JAK2 V617F* allele burden reduction as a predictor of clinical outcome is not well-established. In addition, in patients with other mutations in addition to a *JAK2* mutation, a remission of one mutated clone is not always accompanied by remission of other mutated clones.³²³

Monitoring Response and Follow-up Therapy

The goal of therapy is to prevent thrombotic and hemorrhagic complications without increasing the risk of bleeding. Monitoring for new thrombosis or bleeding is recommended for patients with PV. Monitoring for new thrombosis, acquired VWS, and/or disease-related major bleeding is recommended for patients with ET. After initiation of low-dose aspirin (and phlebotomy for patients with PV), the guidelines recommend monitoring for new thrombosis or bleeding, symptom status using MPN-SAF TSS, signs/symptoms of disease progression, and evaluation for potential indications for cytoreductive therapy as clinically indicated. For ET, bone marrow aspirate and biopsy should be performed to rule out disease progression to MF if there is clinical and/or laboratory suspicion of MF. The development of new thrombosis or disease-related major bleeding, frequent phlebotomy or intolerance to phlebotomy, splenomegaly, progressive thrombocytosis and/or leukocytosis, or disease-related symptoms are considered as potential indications for cytoreductive therapy. Acquired VWS and vasomotor/microvascular disturbances not responsive to aspirin (eg, headaches/chest pain, erythromelalgia) are additional potential indications for cytoreductive therapy for patients with ET.

In one retrospective study, the need for ≥ 3 phlebotomies per year was associated with a significantly higher rate of thrombosis in patients with PV treated with hydroxyurea (21% at 3 years compared to 5% at 3 years for those receiving ≤ 2 phlebotomies per year; $P < .0001$).³²⁸ However, these findings could not be confirmed by other investigators.^{329,330} The development of cytopenia (one of the ELN-defined criteria for resistance or intolerance to hydroxyurea) at the lowest dose of hydroxyurea is an adverse prognostic factor associated with higher risk of death and transformation to AML.^{331,332} Patients with high-risk PV or ET treated with cytoreductive therapy as initial treatment should also be monitored for intolerance or resistance to hydroxyurea.³³³



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

The Panel acknowledges that the IWG-MRT and ELN treatment response criteria were developed mainly for use in clinical trials and that clinical benefit may not reach the threshold of the IWG-MRT and ELN response criteria. Response criteria are not defined for patients treated with low-dose aspirin. Available evidence from retrospective studies that have evaluated these response criteria in patients with PV and ET treated with cytoreductive therapy suggests that achievement of CR as outlined in the response criteria did not correlate with a lower incidence of thrombosis or improvement in thrombosis-free survival.^{331,334-336} While normalization of blood counts after initiation of treatment is usually a goal in clinical practice, it is not associated with long-term clinical benefit and there are no evidence-based data to recommend a target WBC or platelet count for patients receiving cytoreductive therapy. In selected patients with a severe thrombotic event, normalization of blood counts might be a goal of treatment. Response assessment should be done based on the improvement of disease-related symptoms at the discretion of the clinician, and target WBC or platelet counts should be individualized to prevent new thrombosis or bleeding in each patient depending on the presence of risk factors.

Continuation of prior treatment is recommended for patients with asymptomatic disease (low-risk PV and very-low-risk, low-risk, or intermediate-risk ET) with no potential indications for cytoreductive therapy and for patients with high-risk PV or ET with adequate response to initial cytoreductive therapy. Initiation of cytoreductive therapy is recommended for patients with symptomatic disease with potential indications for cytoreductive therapy. For symptomatic low-risk PV, enrollment in a clinical trial and ropeginterferon alfa-2b-njft are preferred regimens, while hydroxyurea and peginterferon alfa-2a are other recommended regimens. Peginterferon alfa-2a is an option for younger patients or in pregnant patients in need of cytoreductive therapy. Subsequent recommendations are the same as those for high-risk PV. The cytoreductive therapy

regimens for symptomatic very-low-risk, low-risk, and intermediate-risk ET are the same as for those with high-risk ET.

Potential indications for change of cytoreductive therapy for patients with PV include intolerance or resistance to prior cytoreductive treatment, new thrombosis or disease-related major bleeding, frequent phlebotomy or intolerance to phlebotomy, splenomegaly, progressive thrombocytosis and/or leukocytosis, and disease-related symptoms (eg, pruritus, night sweats, fatigue).³³⁷ For patients with an inadequate response or a loss of response, a clinical trial and ruxolitinib^{91,300,301,303,306} (category 1 for high-risk PV for hydroxyurea resistance or intolerance) are preferred regimens. Ruxolitinib is FDA approved for the treatment of patients with PV who have had an inadequate response to or are intolerant of hydroxyurea.^{91,300,303,306} and may have activity after inadequate response or loss of response to other agents besides hydroxyurea.³⁰¹ If not previously used, hydroxyurea,^{284,287,288} peginterferon alfa-2,²⁸⁹⁻²⁹² and ropeginterferon alfa-2b-njft,^{206,293} are other recommended regimens. Ropiginterferon alfa-2b-njft is FDA-approved for the treatment of patients with PV.^{206,293}

Potential indications for change of cytoreductive therapy for patients with ET include intolerance or resistance to hydroxyurea, peginterferon alfa-2a, or anagrelide; new thrombosis; acquired VWS and/or disease-related major bleeding; splenomegaly; thrombocytosis; leukocytosis; disease-related symptoms; and vasomotor/microvascular disturbances not responsive to aspirin (eg, headaches, chest pain, erythromelalgia).³³⁷ For patients with an inadequate response or a loss of response, a clinical trial, ropeginterferon alfa-2b-njft (category 1),²⁰⁷ and hydroxyurea (if not previously used),³⁰⁸⁻³¹⁰ are preferred regimens. If not previously used, peginterferon alfa-2a^{289,291,292} is an other recommended regimen. Peginterferon alfa-2a can be considered for patients in need of cytoreductive therapy who are younger or pregnant or who defer



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

hydroxyurea. Anagrelide (if not previously used),³⁰⁹⁻³¹¹ plateletpheresis³³⁸ (for emergent situations such as severe thrombocytosis-related neurologic complications), and ruxolitinib^{92,315} are options that are useful in certain circumstances.

Special Considerations in the Management of MPN

Management of Thrombosis

The use of clinically appropriate anticoagulant therapy (eg, low-molecular-weight heparin [LMWH], direct oral anticoagulants [DOACs], warfarin) is recommended for patients with active thrombosis.³³⁹⁻³⁴¹ The initial use of anticoagulant therapy for the prevention and treatment of thrombosis should be based on the current American College of Chest Physicians Guidelines.³³⁹ Aspirin may be considered for patients with cardiovascular risk factors. There are no evidence-based data to guide the selection or appropriate duration of anticoagulation with or without antiplatelet therapy in patients with PV or ET. The duration of anticoagulant therapy is dependent on the severity of the thrombotic event, degree of disease control, and assessment of likelihood of recurrence after cessation of anticoagulant therapy.³⁴⁰ Aspirin may be considered for patients with cardiovascular risk factors. Low-dose aspirin may also be considered for patients at high risk of thrombosis based on the arterial thrombosis score.²⁷⁴ The risks and benefits of aspirin plus anticoagulation need to be individualized on a case-by-case basis. Plateletpheresis may be indicated in patients with ET presenting with acute life-threatening thrombosis or severe bleeding.

Management of Bleeding

It is essential to rule out other potential causes and treat any coexisting causes, as necessary. Aspirin should be withheld until bleeding is under control and the use of appropriate cytoreductive therapy should be considered to optimize platelet counts while minimizing hematologic and nonhematologic toxicities. Testing for acquired VWS (plasma VWFA

measurement, von Willebrand activity) should be considered for patients with elevated platelet count undergoing surgical procedures or for those with unexplained bleeding. In unanticipated gastrointestinal bleeding, particularly in the setting of splenomegaly, portal hypertension, and gastric varices, special consultation (for endoscopic evaluation) with a hepatologist or a gastrointestinal specialist is recommended.

Surgery

The thrombotic and bleeding risk of the surgical procedure should be strongly considered prior to elective surgery since patients with MPN are at higher risk for bleeding despite optimal management. A matched population-based study demonstrated a higher rate of major bleeding, arterial events, and venous events in patients with MF compared to matched controls (HR, 3.78, 1.73, and 2.75, respectively).³⁴² In a retrospective analysis that evaluated the postoperative outcomes in patients with PV (n = 105) and ET (n = 150), although the majority of patients (74%) were treated with cytoreductive therapy and phlebotomy prior to surgery and antithrombotic prophylaxis, a significant proportion of surgeries was complicated by vascular occlusion (8%) or major hemorrhage (7%). Arterial thrombotic events were more frequent in patients with ET (5% vs 2%; $P = .08$) and venous thrombotic events were more frequent in PV (8% vs 1%; $P = .002$).³⁴³

Multidisciplinary management with careful review of bleeding and thrombosis history is recommended prior to surgery for all patients. Emergency surgery should be performed as necessary with close postoperative surveillance for the symptoms of arterial or venous thrombosis and bleeding. Thrombosis and bleeding should be well-controlled without causing prohibitive cytopenias prior to performing elective surgery (particularly for orthopedic surgeries or any surgical procedures associated with prolonged immobilization) with the use of appropriate antiplatelet therapy, anticoagulant prophylaxis, and



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

cytoreductive therapy. In patients with PV, hematocrit should be controlled for 3 months before elective surgery with the use of additional phlebotomy if necessary to maintain hematocrit <45% prior to performing elective surgery. Prophylaxis with aspirin may be considered following vascular surgery. Extended prophylaxis with anticoagulation should be considered if surgery is associated with a high risk for venous thromboembolism.

Pregnancy

Pregnancy is considered a high-risk clinical situation in patients with PV and ET.³⁴⁴ The presence of a *JAK2* V617F mutation is an adverse prognostic factor for pregnancy outcome, and pregnancy complications are associated with a higher risk of subsequent thrombotic events in patients with ET.³⁴⁵⁻³⁴⁸ The use of aspirin has been reported to be effective in reducing pregnancy complications, especially in patients with *JAK2*-mutated ET.³⁴⁹⁻³⁵¹ In a study that investigated 129 pregnancies in 78 patients with ET, among patients with *JAK2*-mutated ET, complications occurred in 36% of patients receiving aspirin compared to 68% of patients not receiving aspirin.³⁵⁰ In another study of 63 pregnancies among 36 females with ET, the rate of pregnancy loss was 21% among patients receiving aspirin during the first trimester compared to 75% among those not receiving aspirin ($P = .002$).³⁴⁹ The results of a UK prospective cohort study (58 women with a diagnosis of MPN; 47 had a diagnosis of ET) suggest that maternal MPN is associated with higher incidences of maternal complications, preterm delivery, and small-for-gestational-age infants compared to the general population.³⁵² The majority of women (88%) received aspirin and 38% of females additionally received a prophylactic dose of LMWH. Preeclampsia was the most common antenatal complication reported in 9% of women, and 22% of neonates were below the 10th percentile for growth. A systematic review and meta-analysis of pregnant patients with MPN also reported preeclampsia as the most common adverse occurrence (3.1% incidence). Aspirin and interferon use were linked to increased odds of a successful pregnancy.³⁵³

Aggressive intervention for the control of hematocrit, the use of aspirin, and LMWH were associated with significantly better pregnancy outcome in patients with PV.³⁵⁴

Evaluation by a high-risk obstetrician is recommended prior to conception. Hydroxyurea should be discontinued in anticipation or in the event of pregnancy. Peginterferon alfa-2a should be considered if cytoreductive therapy is needed during pregnancy.³⁵⁵ Potential indications include those with prior pregnancy loss or complications (pre-eclampsia) or uncontrolled leukocytosis/thrombocytosis. There are no sufficient data to establish the use of peginterferon alfa-2a in pregnancy and it should be used only if benefits outweigh potential risk to the fetus.³⁵⁵ In the setting of pregnancy and if peginterferon alfa-2a is unavailable, the use of other pegylated interferons (eg, ropeginterferon alfa-2b-njft) can be considered. The data for the use of ropeginterferon alfa-2b-njft in pregnancy are very limited. Two case reports have described the use of ropeginterferon alfa-2b-njft in pregnant women with PV, with no adverse events reported.^{356,357} However, safety data remain insufficient to recommend its routine use during pregnancy. DOACs should be avoided in breastfeeding females. Although large studies do not exist, some reports have shown that DOACs are in the breastmilk of patients who are taking them.³⁵⁸ Unfractionated heparin, LMWH, warfarin, and fondaparinux are all safe options in patients who require anticoagulation and are breastfeeding as they are unlikely to transfer into milk in clinically significant amounts, and are not absorbed from the infant's gastrointestinal tract due to large molecular weight.³⁵⁹ Hydroxyurea is excreted in breastmilk and should be avoided in females who are breastfeeding. Patients with PV or ET who become pregnant and do not have high-risk features (standard-risk pregnancy) should take low-dose aspirin daily throughout the pregnancy and should receive prophylactic LMWH alone during the first 6 weeks of the postpartum period unless there are patient-specific contraindications. If they were



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

taking low-dose aspirin prior to pregnancy, it can be resumed once the postpartum course of LMWH is complete.

For patients with PV or ET whose pregnancies are considered high risk, the recommended treatments to start or continue once a positive pregnancy test is obtained are: 1) low-dose aspirin daily; 2) prophylactic LMWH throughout pregnancy and for 6 weeks postpartum; and 3) cytoreductive therapy with interferon or peginterferon alfa-2a.^{337,344,352,353,360-363} The administration of LMWH should be modified based on renal function, body weight, and medical history. Prophylactic LMWH should be avoided among patients with a history of MPN-related bleeding. Therapeutic anticoagulation with LMWH should be continued during pregnancy by patients receiving anticoagulation therapy after venous and/or arterial thrombotic events prior to pregnancy. The timing of LMWH and aspirin discontinuation prior to the epidural and delivery and the re-initiation of medications after delivery should be discussed with a high-risk obstetrician and obstetric anesthesiologist.

Accelerated/Blast Phase MPN

MPN in accelerated phase is characterized by the presence of 10% to 19% blasts in the peripheral blood or bone marrow, while blast phase is defined as the presence of ≥20% blasts or the presence of myeloid sarcoma.⁶⁶

Patients with a blast percentage of 5% to 9% in the peripheral blood or bone marrow and those with 10% to 19% share similar clinical characteristics.³⁶⁴ In patients with <10% blasts in the peripheral blood or bone marrow, treatment with ruxolitinib improved survival.^{364,365} Patients with a blast percentage of ≥4% in the peripheral blood or ≥5% in the bone marrow have unfavorable outcomes. OS was not significantly different in patients with 5% to 9% blasts in the peripheral blood or bone marrow and in patients with 10% to 19% blasts (24 vs 13 months; $P = .19$).³⁶⁴ Similarly,

OS was comparable in patients with 5% to 9% blasts in the bone marrow and in patients with ≥10% blasts (22 vs 14 months; $P = .73$).³⁶⁵

The incidence of transformation to AML is significantly higher for patients with MF than for those with PV and ET, although the risk is very low in patients who remain in chronic phase MF.^{366,367} Among patients who present with chronic phase MF, development of accelerated phase features during follow-up was associated with short median survival times.³⁶⁶

Treatment with hydroxyurea has been associated with increased risk of transformation to AML in some studies.^{288,368} These findings, however, were not confirmed in subsequent reports.³⁶⁹⁻³⁷¹ In a large cohort analysis (n = 11,039; 162 patients with transformation to AML/MDS) that evaluated treatment-related risk factors for transformation to AML/MDS in patients with MPN, the use of alkylating agents or a combination of ≥2 cytoreductive agents—but not treatment with hydroxyurea alone—was significantly associated with an increased risk of transformation to AML.³⁶⁹ The results of another large analysis (649 patients with PMF, post-PV MF, or post-ET MF) identified bone marrow blasts ≥10% and high-risk karyotypes as independent prognostic factors for the transformation to AML.³⁷¹ Hydroxyurea, however, was not an independent risk factor for transformation to AML.

Mutations in several genes (*ASXL1*, *EZH2*, *TP53*, *SRSF2*, and *IDH1* or *IDH2*) and other chromosomal abnormalities (eg, aberrations in chromosomes 1q and 9p) have been associated with transformation to AML.^{30,39,41,372} Broad-based NGS testing that includes AML-associated mutations is recommended as part of initial workup for patients with accelerated/blast phase MPN (MPN-AP/BP).



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Treatment Options

In a retrospective analysis of 91 patients with MF that had transformed to AML, the median OS after transformation to AML was 3 months. Among patients who were treated with AML-type induction chemotherapy, reversal to chronic phase without an increase in the blast percentage occurred in 41% of patients.³⁷³ However, it was also associated with a treatment-related mortality (TRM) rate of 33%. The median OS was 4 months, which was comparable to that observed in patients treated with supportive care or low-intensity chemotherapy (2 months and 3 months, respectively).

Hypomethylating agents (HMAs) (azacitidine or decitabine) have been evaluated in a few small studies as a treatment option for MPN that has transformed to AML.³⁷⁴⁻³⁷⁶ In a small series of 11 patients with MF-BP/AML, decitabine was associated with improved survival in patients who were not eligible for allogeneic HCT.³⁷⁴ At a median follow-up of 9 months, 67% of the patients treated with decitabine were alive. In another series of 54 patients with MPN-BP/AML (21 patients with ET, 21 patients with PV, 7 patients with PMF, and 5 patients with unclassified MPN), first-line therapy with azacitidine resulted in an ORR of 52% (24% CR, 11% PR, 8% bone marrow CR or CR with incomplete recovery of cytopenias, and 9% hematologic improvement).³⁷⁵ The median duration of response and the median OS were 9 months and 11 months, respectively. In a retrospective analysis of 21 patients with MPN-BP/AML and 13 patients with MPN-AP treated with decitabine, the ORRs were 62% (8 out of 13 patients) and 29% (6 out of 21 patients), respectively, for patients with MPN-AP and MPN-BP/AML.³⁷⁶ The median OS was significantly higher in patients with disease that responded to decitabine (12 vs 5 months, respectively, for patients with MPN-AP; 11 vs 4 months, respectively, for patients with MPN-BP/AML).

In a small study of 21 patients with MPN-AP/BP, decitabine in combination with dose-escalated ruxolitinib resulted in an ORR (by protocol-defined criteria) of 53% (95% CI, 27.8%–77.0%) and a median OS of 7.9 months (95% CI, 4.1 months–not reached).³⁷⁷ The results of a phase II study with 25 patients with MPN-AP/BP demonstrated an ORR of 44% (95% CI, 24.4%–65.1%) and a median OS of 9.5 months (95% CI, 4.3–12.0 months) in patients treated with the combination of ruxolitinib and decitabine.³⁷⁸ A phase 1/2 trial investigating the combination of ruxolitinib and decitabine in patients with post-MPN AML reported an ORR of 45% in the intention-to-treat population and 61% in patients who received the recommended phase II dose.³⁷⁹ A phase II study (NCT04282187) is examining the combination of decitabine with ruxolitinib or fedratinib in patients with MPN-AP/BP prior to HCT.

The role of venetoclax combinations has not yet been established in MPN-BP/AML. In one study, the complete remission rate (with or without count recovery) was 44% in patients with MPN-BP treated with venetoclax in combination with azacitidine or decitabine.³⁸⁰ The authors note that 6 patients with complete remission (with or without count recovery) proceeded to transplant. In another study, venetoclax-based combination therapy resulted in an ORR of 50% and 53%, respectively, in MPN-AP/BP.³⁸¹ However, significant toxicities were also reported. In a systematic review and meta-analysis, treatment with HMAs in combination with venetoclax resulted in a higher CR/CR with incomplete count recovery (CRi) rate compared to treatment with HMA monotherapy (36% vs 19%; $P = .0204$).³⁸² In addition, this combination also yielded an improved CR rate compared to HMAs in combination with ruxolitinib (22% vs 8%; $P = .0313$). Treatment with HMAs in combination with ruxolitinib elicited an improved ORR compared to HMA monotherapy (45% vs 30%; $P = .0395$). However, there was no difference in CR/CRi.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

There is a paucity of data regarding the use of targeted therapies for MPN-AP/BP. In a retrospective analysis of 8 patients with MPN-AP/BP with *IDH2* mutations treated with enasidenib, an ORR of 37.5% was obtained based on the 2017 ELN criteria.³⁸³ Another study also demonstrated the clinical benefit of *IDH1/2* inhibitor-based combinations in patients with post-MPN AML with *IDH1/2* mutations.³⁸⁴ Of the 12 patients that were evaluated, 3 achieved CR, with a median OS of 10 months. The addition of midostaurin, a *FLT3* inhibitor, to chemotherapy resulted in improved median OS (74.7 vs 25.6 months; HR, 0.78; $P = .009$) in patients with AML with a *FLT3* mutation, when compared to placebo.³⁸⁵ Similarly, the use of quizartinib, another *FLT3* inhibitor, in combination with chemotherapy led to higher median OS (31.9 vs 15.1 months for placebo; HR, 0.78; $P = .032$) in patients with *FLT3*-ITD AML.³⁸⁶

Allogeneic HCT remains the only curative option resulting in long-term disease control in selected transplant-eligible patients who achieve a CR to induction chemotherapy.³⁸⁷⁻³⁹⁰ Early referral to a transplant team is recommended for planning purposes and to discuss the role of bridging therapy. Some patients in accelerated phase may proceed to transplant directly without bridging therapy.³⁹¹ In one retrospective analysis of 75 patients with MPN-BP, patients who were treated with curative intent (induction chemotherapy with or without allogeneic HCT) had significantly improved survival compared with those treated with non-curative intent (non-intensive chemotherapy or supportive care).³⁸⁹ The 2-year OS rates were 26% and 3%, respectively, and the median survival was 9 months and 2 months, respectively ($P < .0001$). Among patients treated with curative intent, the ORR to induction chemotherapy was 46% and reversal to chronic phase was observed in 31% of patients, with 17 patients undergoing allogeneic HCT. The OS rate was significantly higher for patients who underwent allogeneic HCT following induction chemotherapy (2-year OS rate was 47% compared with 15% for those who did not undergo allogeneic HCT; $P = .03$). In another retrospective analysis of 46

patients who received allogeneic HCT for MF-BP, the 3-year PFS and OS rates following transplant were 26% and 33%, respectively. The response status prior to transplant (CR vs no CR) was a significant predictor of OS (69% for CR vs 22% for no CR; $P = .008$) and PFS (55% and 19%, respectively; $P = .02$).³⁹⁰ The cumulative incidence of TRM was 28% at 1 year and the absence of CR before allogeneic HCT was also associated with significantly increased TRM (35% vs 0%; $P = .053$). Another study reported a median OS of 2.30 years in patients with MPN-AP/BP who underwent allogeneic HCT.³⁹²

Treatment Recommendations Based on Eligibility for Transplant

The selection of patients for allogeneic HCT should be based on age, performance status, major comorbid conditions, psychosocial status, patient preference, and availability of caregiver(s). Patients may be taken immediately to transplant or bridging therapy can be used to decrease marrow blasts to an acceptable level prior to transplant.

Enrollment in a clinical trial or bridging therapy followed by allogeneic HCT are treatment options for patients who are candidates for transplant. Bridging therapy options include HMAs (azacitidine or decitabine), HMAs with JAK inhibitors (fedratinib, momelotinib, pacritinib, or ruxolitinib), HMAs with venetoclax, or intensive AML-type induction chemotherapy.^{374,389,390} Enrollment in a clinical trial, HMAs (azacitidine or decitabine), HMAs with JAK inhibitors (fedratinib, momelotinib, pacritinib, or ruxolitinib), HMAs with venetoclax, lower-intensity AML-type chemotherapy, or targeted therapy is recommended for those who are not candidates for transplant. AML-type induction chemotherapy regimens and HMAs in combination with venetoclax may be used for the management of disease progression of MPN. However, these regimens typically result in low response rates and are associated with significant toxicities. Based on NGS panel results (eg, if NGS shows particular mutations such as *IDH1*, *IDH2*, or *FLT3*), targeted therapy (eg, *IDH1/2* and *FLT3* inhibitors) alone or in combination with



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

HMA can be considered.³⁸³⁻³⁸⁶ HMAs (azacitidine or decitabine) can be used in combination with a JAK inhibitor (fedratinib, momelotinib, pacritinib) for the palliation of splenomegaly or other disease-related symptoms.^{377,379} However, the Panel notes that there are very limited data regarding the use of fedratinib, momelotinib, or pacritinib with HMAs. While the combination of an HMA and pacritinib has been evaluated in AML and chronic myelomonocytic leukemia (CMML),^{393,394} the combination has not been evaluated in MPN-AP/BP. Fedratinib, momelotinib, pacritinib, or ruxolitinib may be continued near to the start of conditioning therapy for the improvement of splenomegaly and other disease-related symptoms in patients who are transplant candidates.^{184,210-212}

Summary

MPN are characterized by a significant symptom burden and a propensity for disease transformation to accelerated and blast phases. The goal of treatment is to reduce symptom burden and the risk of developing vascular complications. Regular monitoring of disease-related symptoms, assessment of need for cytoreductive therapy, and appropriate evaluation for disease progression should be an integral part of the comprehensive care of patients with MPN. Referral to a mental health professional and other supportive services can be invaluable, including integrative health specialists.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

References

1. Gerds A, Rein L, Engel-Nitz N, et al. Incidence, prevalence, and clinical outcomes of myelobrosis with and without cytopenia in the United States [abstract]. Blood 2025;146:5576. Available at: <https://ashpublications.org/blood/article/146/Supplement%201/5576/556730/Incidence-prevalence-and-clinical-outcomes-of>.
2. Verstovsek S, Yu J, Scherber RM, et al. Changes in the incidence and overall survival of patients with myeloproliferative neoplasms between 2002 and 2016 in the United States. Leuk Lymphoma 2022;63:694–702. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34689695>.
3. Anderson LA, James G, Duncombe AS, et al. Myeloproliferative neoplasm patient symptom burden and quality of life: Evidence of significant impairment compared to controls. Am J Hematol 2015;90:864–870. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26113113>.
4. Mesa R, Miller CB, Thyne M, et al. Myeloproliferative neoplasms (MPNs) have a significant impact on patients' overall health and productivity: the MPN Landmark survey. BMC Cancer 2016;16:167. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26922064>.
5. Harrison CN, Koschmieder S, Foltz L, et al. The impact of myeloproliferative neoplasms (MPNs) on patient quality of life and productivity: Results from the international MPN Landmark survey. Ann Hematol 2017;96:1653–1665. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28780729>.
6. Geyer HL, Scherber RM, Dueck AC, et al. Distinct clustering of symptomatic burden among myeloproliferative neoplasm patients: Retrospective assessment in 1470 patients. Blood 2014;123:3803–3810. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24553173>.
7. Price GL, Davis KL, Karve S, et al. Survival patterns in United States (US) medicare enrollees with non-CML myeloproliferative neoplasms (MPN). PLoS One 2014;9:e90299. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24618579>.
8. Tam CS, Nussenzveig RM, Popat U, et al. The natural history and treatment outcome of blast phase BCR-ABL- myeloproliferative neoplasms. Blood 2008;112:1628–1637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18566326>.
9. Tefferi A, Guglielmelli P, Larson DR, et al. Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. Blood 2014;124:2507–2513. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25037629>.
10. Tefferi A. Myeloproliferative neoplasms: A decade of discoveries and treatment advances. Am J Hematol 2016;91:50–58. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26492355>.
11. Stein BL, Gotlib J, Arcasoy M, et al. Historical views, conventional approaches, and evolving management strategies for myeloproliferative neoplasms. J Natl Compr Canc Netw 2015;13:424–434. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25870379>.
12. PubMed Overview. Available at: <https://pubmed.ncbi.nlm.nih.gov/about/>. Accessed July 26, 2022.
13. Freedman-Cass DA, Fischer T, Alpert AB, et al. The value and process of inclusion: Using sensitive, respectful, and inclusive language and images in NCCN content. J Natl Compr Canc Netw 2023;21:434–441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37156485>.
14. Baxter EJ, Scott LM, Campbell PJ, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. Lancet 2005;365:1054–1061. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15781101>.
15. Kralovics R, Passamonti F, Buser AS, et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. N Engl J Med 2005;352:1779–1790. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15858187>.
16. Levine RL, Wadleigh M, Cools J, et al. Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia,



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

and myeloid metaplasia with myelofibrosis. *Cancer Cell* 2005;7:387–397. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15837627>.

17. Jamieson CH, Gotlib J, Durocher JA, et al. The JAK2 V617F mutation occurs in hematopoietic stem cells in polycythemia vera and predisposes toward erythroid differentiation. *Proc Natl Acad Sci U S A* 2006;103:6224–6229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16603627>.

18. Scott LM, Tong W, Levine RL, et al. JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med* 2007;356:459–468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17267906>.

19. Pietra D, Li S, Brisci A, et al. Somatic mutations of JAK2 exon 12 in patients with JAK2 (V617F)-negative myeloproliferative disorders. *Blood* 2008;111:1686–1689. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17984312>.

20. Pardanani AD, Levine RL, Lasho T, et al. MPL515 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood* 2006;108:3472–3476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16868251>.

21. Pikman Y, Lee BH, Mercher T, et al. MPLW515L is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia. *PLoS Med* 2006;3:e270. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16834459>.

22. Beer PA, Campbell PJ, Scott LM, et al. MPL mutations in myeloproliferative disorders: analysis of the PT-1 cohort. *Blood* 2008;112:141–149. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18451306>.

23. Klampfl T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med* 2013;369:2379–2390. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24325356>.

24. Nangalia J, Massie CE, Baxter EJ, et al. Somatic CALR mutations in myeloproliferative neoplasms with nonmutated JAK2. *N Engl J Med*

2013;369:2391–2405. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24325359>.

25. Tefferi A, Lasho TL, Finke C, et al. Type 1 vs type 2 calreticulin mutations in primary myelofibrosis: differences in phenotype and prognostic impact. *Leukemia* 2014;28:1568–1570. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24569778>.

26. Tefferi A, Lasho TL, Tischer A, et al. The prognostic advantage of calreticulin mutations in myelofibrosis might be confined to type 1 or type 1-like CALR variants. *Blood* 2014;124:2465–2466. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25301336>.

27. Pietra D, Rumi E, Ferretti VV, et al. Differential clinical effects of different mutation subtypes in CALR-mutant myeloproliferative neoplasms. *Leukemia* 2016;30:431–438. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26449662>.

28. How J, Hobbs GS, Mullally A. Mutant calreticulin in myeloproliferative neoplasms. *Blood* 2019;134:2242–2248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31562135>.

29. Nangalia J, Green TR. The evolving genomic landscape of myeloproliferative neoplasms. *Hematology Am Soc Hematol Educ Program* 2014;2014:287–296. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25696868>.

30. Rampal R, Ahn J, Abdel-Wahab O, et al. Genomic and functional analysis of leukemic transformation of myeloproliferative neoplasms. *Proc Natl Acad Sci U S A* 2014;111:E5401–5410. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25516983>.

31. Rumi E, Pietra D, Pascutto C, et al. Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood* 2014;124:1062–1069. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24986690>.

32. Guglielmelli P, Rotunno G, Fanelli T, et al. Validation of the differential prognostic impact of type 1/type 1-like versus type 2/type 2-like CALR



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

mutations in myelofibrosis. *Blood Cancer J* 2015;5:e360. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26473532>.

33. Barbui T, Ghirardi A, Carobbio A, et al. Increased risk of thrombosis in JAK2 V617F-positive patients with primary myelofibrosis and interaction of the mutation with the IPSS score. *Blood Cancer J* 2022;12:156. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36385103>.

34. Guglielmelli P, Pancrazzi A, Bergamaschi G, et al. Anaemia characterises patients with myelofibrosis harbouring Mpl mutation. *Br J Haematol* 2007;137:244–247. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17408465>.

35. Tefferi A, Lasho TL, Finke CM, et al. CALR vs JAK2 vs MPL-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia* 2014;28:1472–1477. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24402162>.

36. Milosevic Feenstra JD, Nivarthi H, Gisslinger H, et al. Whole-exome sequencing identifies novel MPL and JAK2 mutations in triple-negative myeloproliferative neoplasms. *Blood* 2016;127:325–332. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26423830>.

37. Guglielmelli P, Biamonte F, Score J, et al. EZH2 mutational status predicts poor survival in myelofibrosis. *Blood* 2011;118:5227–5234. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21921040>.

38. Tefferi A, Jimma T, Sulai NH, et al. IDH mutations in primary myelofibrosis predict leukemic transformation and shortened survival: Clinical evidence for leukemogenic collaboration with JAK2V617F. *Leukemia* 2012;26:475–480. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21912393>.

39. Vannucchi AM, Lasho TL, Guglielmelli P, et al. Mutations and prognosis in primary myelofibrosis. *Leukemia* 2013;27:1861–1869. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23619563>.

40. Guglielmelli P, Lasho TL, Rotunno G, et al. The number of prognostically detrimental mutations and prognosis in primary

myelofibrosis: An international study of 797 patients. *Leukemia* 2014;28:1804–1810. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24549259>.

41. Lundberg P, Karow A, Nienhold R, et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood* 2014;123:2220–2228. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24478400>.

42. Tefferi A, Finke CM, Lasho TL, et al. U2AF1 mutation types in primary myelofibrosis: Phenotypic and prognostic distinctions. *Leukemia* 2018;32:2274–2278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29535431>.

43. Vallapureddy RR, Mudireddy M, Penna D, et al. Leukemic transformation among 1306 patients with primary myelofibrosis: risk factors and development of a predictive model. *Blood Cancer J* 2019;9:12. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30683837>.

44. Santos FPS, Getta B, Masarova L, et al. Prognostic impact of RAS-pathway mutations in patients with myelofibrosis. *Leukemia* 2020;34:799–810. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31628430>.

45. Tefferi A, Abdelmagid M, Loscocco GG, et al. TP53 mutations in myeloproliferative neoplasms: context-dependent evaluation of prognostic relevance. *Am J Hematol* 2025;100:552–560. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39873146>.

46. Rolles B, Filho CMO, Fergusson N, et al. Risk stratification of patients with TP53-mutated myeloproliferative neoplasms. *Leukemia* 2026;40:773–782. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41735484>.

47. Tefferi A, Guglielmelli P, Lasho TL, et al. CALR and ASXL1 mutations-based molecular prognostication in primary myelofibrosis: An international study of 570 patients. *Leukemia* 2014;28:1494–1500. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24496303>.

48. Rotunno G, Pacilli A, Artusi V, et al. Epidemiology and clinical relevance of mutations in postpolycythemia vera and postessential



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

thrombocythemia myelofibrosis: A study on 359 patients of the AGIMM group. *Am J Hematol* 2016;91:681–686. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27037840>.

49. Passamonti F, Elena C, Schnittger S, et al. Molecular and clinical features of the myeloproliferative neoplasm associated with JAK2 exon 12 mutations. *Blood* 2011;117:2813–2816. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21224469>.

50. Tondeur S, Paul F, Riou J, et al. Long-term follow-up of JAK2 exon 12 polycythemia vera: a French Intergroup of Myeloproliferative Neoplasms (FIM) study. *Leukemia* 2021;35:871–875. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32694617>.

51. Tefferi A, Lavu S, Mudireddy M, et al. JAK2 exon 12 mutated polycythemia vera: Mayo-Careggi MPN Alliance study of 33 consecutive cases and comparison with JAK2V617F mutated disease. *Am J Hematol* 2018;93:E93–E96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29274140>.

52. Rotunno G, Mannarelli C, Guglielmelli P, et al. Impact of calreticulin mutations on clinical and hematological phenotype and outcome in essential thrombocythemia. *Blood* 2014;123:1552–1555. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24371211>.

53. Rumi E, Pietra D, Ferretti V, et al. JAK2 or CALR mutation status defines subtypes of essential thrombocythemia with substantially different clinical course and outcomes. *Blood* 2014;123:1544–1551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24366362>.

54. Elala YC, Lasho TL, Gangat N, et al. Calreticulin variant stratified driver mutational status and prognosis in essential thrombocythemia. *Am J Hematol* 2016;91:503–506. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26890983>.

55. Alvarez-Larran A, Martinez D, Arenillas L, et al. Essential thrombocythaemia with mutation in MPL: clinicopathological correlation and comparison with JAK2V617F-mutated and CALR-mutated genotypes.

J Clin Pathol 2018;71:975–980. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29934356>.

56. Perez Encinas MM, Sobas M, Gomez-Casares MT, et al. The risk of thrombosis in essential thrombocythemia is associated with the type of CALR mutation: A multicentre collaborative study. *Eur J Haematol* 2021;106:371–379. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33275803>.

57. Alvarez-Larran A, Senin A, Fernandez-Rodriguez C, et al. Impact of genotype on leukaemic transformation in polycythaemia vera and essential thrombocythaemia. *Br J Haematol* 2017;178:764–771. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28542718>.

58. Finazzi G, Carobbio A, Guglielmelli P, et al. Calreticulin mutation does not modify the IPSET score for predicting the risk of thrombosis among 1150 patients with essential thrombocythemia. *Blood* 2014;124:2611–2612. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25323688>.

59. Tefferi A, Lasho TL, Guglielmelli P, et al. Targeted deep sequencing in polycythemia vera and essential thrombocythemia. *Blood Adv* 2016;1:21–30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29296692>.

60. Senin A, Fernandez-Rodriguez C, Bellosillo B, et al. Non-driver mutations in patients with JAK2V617F-mutated polycythemia vera or essential thrombocythemia with long-term molecular follow-up. *Ann Hematol* 2018;97:443–451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29181548>.

61. Benton CB, Boddu PC, DiNardo CD, et al. Janus kinase 2 variants associated with the transformation of myeloproliferative neoplasms into acute myeloid leukemia. *Cancer* 2019;125:1855–1866. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30811597>.

62. Vardiman JW, Harris NL, Brunning RD. The World Health Organization (WHO) classification of the myeloid neoplasms. *Blood* 2002;100:2292–2302. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12239137>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

63. Vardiman JW, Thiele J, Arber DA, et al. The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: Rationale and important changes. *Blood* 2009;114:937–951. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19357394>.

64. Barbui T, Thiele J, Vannucchi AM, Tefferi A. Rationale for revision and proposed changes of the WHO diagnostic criteria for polycythemia vera, essential thrombocythemia and primary myelofibrosis. *Blood Cancer J* 2015;5:e337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26832847>.

65. Swerdlow SH, Campo E, Harris NL, et al. WHO classification of tumours of haematopoietic and lymphoid tissues (revised 4th edition). International Agency for Research on Cancer; Lyon, France; 2017.

66. Arber DA, Orazi A, Hasserjian RP, et al. International consensus classification of myeloid neoplasms and acute leukemia: Integrating morphological, clinical, and genomic data. *Blood* 2022;140:1200–1228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35767897>.

67. Khoury JD, Solary E, Abla O, et al. The 5th edition of the World Health Organization classification of haematolymphoid tumours: Myeloid and histiocytic/dendritic neoplasms. *Leukemia* 2022;36:1703–1719. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35732831>.

68. WHO Classification of Tumors Editorial Board. Haematolymphoid tumours [Internet]. Lyon (France): International Agency for Research on Cancer; 2024 [cited 2025 April 16]. (WHO classification of tumours series, 5th ed.; vol. 11). Available from: <https://tumourclassification.iarc.who.int/chapters/63>.

69. Mesa RA, Verstovsek S, Cervantes F, et al. Primary myelofibrosis (PMF), post polycythemia vera myelofibrosis (post-PV MF), post essential thrombocythemia myelofibrosis (post-ET MF), blast phase PMF (PMF-BP): Consensus on terminology by the international working group for myelofibrosis research and treatment (IWG-MRT). *Leuk Res* 2007;31:737–740. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17210175>.

70. Barbui T, Thiele J, Passamonti F, et al. Survival and disease progression in essential thrombocythemia are significantly influenced by accurate morphologic diagnosis: an international study. *J Clin Oncol* 2011;29:3179–3184. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21747083>.

71. Guglielmelli P, Pacilli A, Rotunno G, et al. Presentation and outcome of patients with 2016 WHO diagnosis of prefibrotic and overt primary myelofibrosis. *Blood* 2017;129:3227–3236. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28351937>.

72. Jeryczynski G, Thiele J, Gisslinger B, et al. Pre-fibrotic/early primary myelofibrosis vs. WHO-defined essential thrombocythemia: The impact of minor clinical diagnostic criteria on the outcome of the disease. *Am J Hematol* 2017;92:885–891. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28543356>.

73. Mudireddy M, Shah S, Lasho T, et al. Prefibrotic versus overtly fibrotic primary myelofibrosis: clinical, cytogenetic, molecular and prognostic comparisons. *Br J Haematol* 2018;182:594–597. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28677834>.

74. Barbui T, Thiele J, Gisslinger H, et al. Masked polycythemia vera (mPV): Results of an international study. *Am J Hematol* 2014;89:52–54. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23996471>.

75. Alvarez-Larran A, Angona A, Ancochea A, et al. Masked polycythemia vera: presenting features, response to treatment and clinical outcomes. *Eur J Haematol* 2016;96:83–89. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25810304>.

76. Thiele J, Kvasnicka HM, Orazi A, et al. The international consensus classification of myeloid neoplasms and acute leukemias: Myeloproliferative neoplasms. *Am J Hematol* 2023;98:544–545. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36600640>.

77. Delic S, Rose D, Kern W, et al. Application of an NGS-based 28-gene panel in myeloproliferative neoplasms reveals distinct mutation patterns in essential thrombocythemia, primary myelofibrosis and polycythemia



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

vera. Br J Haematol 2016;175:419–426. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27447873>.

78. Thiele J, Kvasnicka HM, Facchetti F, et al. European consensus on grading bone marrow fibrosis and assessment of cellularity. Haematologica 2005;90:1128–1132. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16079113>.

79. McMahon B, Stein BL. Thrombotic and bleeding complications in classical myeloproliferative neoplasms. Semin Thromb Hemost 2013;39:101–111. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23264112>.

80. Kaifia A, Kirschner M, Wolf D, et al. Bleeding, thrombosis, and anticoagulation in myeloproliferative neoplasms (MPN): Analysis from the German SAL-MPN-registry. J Hematol Oncol 2016;9:18. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26944254>.

81. Federici AB, Rand JH, Bucciarelli P, et al. Acquired von Willebrand syndrome: Data from an international registry. Thromb Haemost 2000;84:345–349. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10959711>.

82. Lippi G, Franchini M, Salvagno GL, et al. Correlation between von Willebrand factor antigen, von Willebrand factor ristocetin cofactor activity and factor VIII activity in plasma. J Thromb Thrombolysis 2008;26:150–153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17786534>.

83. Mesa RA, Schwager S, Radia D, et al. The Myelofibrosis Symptom Assessment Form (MFSAF): an evidence-based brief inventory to measure quality of life and symptomatic response to treatment in myelofibrosis. Leuk Res 2009;33:1199–1203. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19250674>.

84. Mesa RA, Kantarjian H, Tefferi A, et al. Evaluating the serial use of the Myelofibrosis Symptom Assessment Form for measuring symptomatic improvement: Performance in 87 myelofibrosis patients on a JAK1 and JAK2 inhibitor (INCB018424) clinical trial. Cancer 2011;117:4869–4877. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21480207>.

85. Scherber R, Dueck AC, Johansson P, et al. The Myeloproliferative Neoplasm Symptom Assessment Form (MPN-SAF): international prospective validation and reliability trial in 402 patients. Blood 2011;118:401–408. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21536863>.

86. Emanuel RM, Dueck AC, Geyer HL, et al. Myeloproliferative neoplasm (MPN) symptom assessment form total symptom score: Prospective international assessment of an abbreviated symptom burden scoring system among patients with MPNs. J Clin Oncol 2012;30:4098–4103. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23071245>.

87. Mesa RA, Gotlib J, Gupta V, et al. Effect of ruxolitinib therapy on myelofibrosis-related symptoms and other patient-reported outcomes in COMFORT-I: A randomized, double-blind, placebo-controlled trial. J Clin Oncol 2013;31:1285–1292. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23423753>.

88. EINahass YH, Mahmoud HK, Mattar MM, et al. MPN10 score and survival of molecularly annotated myeloproliferative neoplasm patients. Leuk Lymphoma 2018;59:844–854. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28828906>.

89. Mesa RA, Niblack J, Wadleigh M, et al. The burden of fatigue and quality of life in myeloproliferative disorders (MPDs): An international Internet-based survey of 1179 MPD patients. Cancer 2007;109:68–76. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17123268>.

90. Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. N Engl J Med 2012;366:799–807. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22375971>.

91. Vannucchi AM, Kiladjian JJ, Griesshammer M, et al. Ruxolitinib versus standard therapy for the treatment of polycythemia vera. N Engl J Med 2015;372:426–435. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25629741>.

92. Harrison CN, Mead AJ, Panchal A, et al. Ruxolitinib vs best available therapy for ET intolerant or resistant to hydroxycarbamide. Blood



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

2017;130:1889–1897. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29074595>.

93. Harrison CN, Schaap N, Vannucchi AM, et al. Janus kinase-2 inhibitor fedratinib in patients with myelofibrosis previously treated with ruxolitinib (JAKARTA-2): A single-arm, open-label, non-randomised, phase 2, multicentre study. *Lancet Haematol* 2017;4:e317–e324. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28602585>.

94. Verstovsek S, Gerds AT, Vannucchi AM, et al. Momelotinib versus danazol in symptomatic patients with anaemia and myelofibrosis (MOMENTUM): Results from an international, double-blind, randomised, controlled, phase 3 study. *Lancet* 2023;401:269–280. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36709073>.

95. Gerds AT, Verstovsek S, Vannucchi AM, et al. Momelotinib versus danazol in symptomatic patients with anaemia and myelofibrosis previously treated with a JAK inhibitor (MOMENTUM): An updated analysis of an international, double-blind, randomised phase 3 study. *Lancet Haematol* 2023;10:e735–e746. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37517413>.

96. Mesa RA, Vannucchi AM, Mead A, et al. Pacritinib versus best available therapy for the treatment of myelofibrosis irrespective of baseline cytopenias (PERSIST-1): An international, randomised, phase 3 trial. *Lancet Haematol* 2017;4:e225–e236. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28336242>.

97. Mascarenhas J, Hoffman R, Talpaz M, et al. Pacritinib vs best available therapy, including ruxolitinib, in patients with myelofibrosis: A randomized clinical trial. *JAMA Oncol* 2018;4:652–659. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29522138>.

98. Pardanani A, Tefferi A, Masszi T, et al. Updated results of the placebo-controlled, phase III JAKARTA trial of fedratinib in patients with intermediate-2 or high-risk myelofibrosis. *Br J Haematol* 2021;195:244–248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34331348>.

99. Cervantes F, Dupriez B, Pereira A, et al. New prognostic scoring system for primary myelofibrosis based on a study of the International Working Group for Myelofibrosis Research and Treatment. *Blood* 2009;113:2895–2901. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18988864>.

100. Passamonti F, Cervantes F, Vannucchi AM, et al. A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). *Blood* 2010;115:1703–1708. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20008785>.

101. Gangat N, Caramazza D, Vaidya R, et al. DIPSS plus: A refined Dynamic International Prognostic Scoring System for primary myelofibrosis that incorporates prognostic information from karyotype, platelet count, and transfusion status. *J Clin Oncol* 2011;29:392–397. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21149668>.

102. Guglielmelli P, Lasho TL, Rotunno G, et al. MIPSS70: Mutation-Enhanced International Prognostic Score System for transplantation-age patients with primary myelofibrosis. *J Clin Oncol* 2018;36:310–318. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29226763>.

103. Tefferi A, Guglielmelli P, Lasho TL, et al. MIPSS70+ Version 2.0: Mutation and Karyotype-Enhanced International Prognostic Scoring System for primary myelofibrosis. *J Clin Oncol* 2018;36:1769–1770. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29708808>.

104. Patnaik MM, Caramazza D, Gangat N, et al. Age and platelet count are IPSS-independent prognostic factors in young patients with primary myelofibrosis and complement IPSS in predicting very long or very short survival. *Eur J Haematol* 2010;84:105–108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19895568>.

105. Tefferi A, Siragusa S, Hussein K, et al. Transfusion-dependency at presentation and its acquisition in the first year of diagnosis are both equally detrimental for survival in primary myelofibrosis--prognostic relevance is independent of IPSS or karyotype. *Am J Hematol*



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

2010;85:14–17. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20029953>.

106. Hussein K, Pardanani AD, Van Dyke DL, et al. International Prognostic Scoring System-independent cytogenetic risk categorization in primary myelofibrosis. *Blood* 2010;115:496–499. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19901264>.

107. Caramazza D, Begna KH, Gangat N, et al. Refined cytogenetic-risk categorization for overall and leukemia-free survival in primary myelofibrosis: A single center study of 433 patients. *Leukemia* 2011;25:82–88. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20944670>.

108. Guerra M, Pasquer H, Daltro de Oliveira R, et al. Comparative clinical and molecular landscape of primary and secondary myelofibrosis: Superior performance of MIPSS70+ v2.0 over MYSEC-PM. *Am J Hematol* 2024;99:741–744. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/38279562>.

109. Shide K, Takenaka K, Kitanaka A, et al. Nationwide prospective survey of secondary myelofibrosis in Japan: Superiority of DIPSS-plus to MYSEC-PM as a survival risk model. *Blood Cancer J* 2023;13:110. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/37463903>.

110. Passamonti F, Giorgino T, Mora B, et al. A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis. *Leukemia* 2017;31:2726–2731. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28561069>.

111. Palandri F, Palumbo GA, Iurlo A, et al. Differences in presenting features, outcome and prognostic models in patients with primary myelofibrosis and post-polycythemia vera and/or post-essential thrombocythemia myelofibrosis treated with ruxolitinib. New perspective of the MYSEC-PM in a large multicenter study. *Semin Hematol* 2018;55:248–255. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30502854>.

112. Mora B, Giorgino T, Guglielmelli P, et al. Value of cytogenetic abnormalities in post-polycythemia vera and post-essential thrombocythemia myelofibrosis: A study of the MYSEC project. *Haematologica* 2018;103:e392–e394. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29622658>.

113. Ianotto JC, Boyer-Perrard F, Gyan E, et al. Efficacy and safety of pegylated-interferon alpha-2a in myelofibrosis: A study by the FIM and GEM French cooperative groups. *Br J Haematol* 2013;162:783–791. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23848933>.

114. Ianotto JC, Chauveau A, Boyer-Perrard F, et al. Benefits and pitfalls of pegylated interferon-alpha2a therapy in patients with myeloproliferative neoplasm-associated myelofibrosis: A French Intergroup of Myeloproliferative neoplasms (FIM) study. *Haematologica* 2018;103:438–446. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29217781>.

115. Silver RT, Barel AC, Lascu E, et al. The effect of initial molecular profile on response to recombinant interferon-alpha (rIFNalpha) treatment in early myelofibrosis. *Cancer* 2017;123:2680–2687. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28518222>.

116. Sorensen AL, Mikkelsen SU, Knudsen TA, et al. Ruxolitinib and interferon-alpha2 combination therapy for patients with polycythemia vera or myelofibrosis: A phase II study. *Haematologica* 2020;105:2262–2272. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33054051>.

117. Kiladjian JJ, Ianotto JC, Soret J, et al. Final results of RUXOPEG, a phase 1/2 adaptive randomized trial of ruxolitinib (rux) and pegylated interferon alpha (IFNa) 2a in patients with myelofibrosis (MF) [abstract]. *Blood* 2022;140:577–578. Available at:

<https://ashpublications.org/blood/article/140/Supplement%201/577/488443/Final-Results-of-Ruxopeg-a-Phase-1-2-Adaptive>.

118. Harrison C, Kiladjian JJ, Verstovsek S, et al. Overall and progression-free survival in patients treated with fedratinib as first-line myelofibrosis (MF) therapy and after prior ruxolitinib (RUX): Results from the JAKARTA and JAKARTA2 trials [abstract]. *EHA Congress 2021:Abstract S203*.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

Available at: <https://library.ehaweb.org/eha/2021/eha2021-virtual-congress/324611/>.

119. Harrison CN, Schaap N, Vannucchi AM, et al. Safety and efficacy of fedratinib, a selective oral inhibitor of Janus kinase-2 (JAK2), in patients with myelofibrosis and low pretreatment platelet counts. *Br J Haematol* 2022;198:317–327. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/35476316>.

120. Harrison CN, Schaap N, Vannucchi AM, et al. Fedratinib in patients with myelofibrosis previously treated with ruxolitinib: An updated analysis of the JAKARTA2 study using stringent criteria for ruxolitinib failure. *Am J Hematol* 2020;95:594–603. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32129512>.

121. Tremblay D, Cavalli L, Sy O, et al. The effect of fedratinib, a selective inhibitor of janus kinase 2, on weight and metabolic parameters in patients with intermediate- or high-risk myelofibrosis. *Clin Lymphoma Myeloma Leuk* 2022;22:e463–e466. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/35120852>.

122. Gupta V, Yacoub A, Mesa RA, et al. Safety and efficacy of fedratinib in patients with myelofibrosis previously treated with ruxolitinib: Primary analysis of FREEDOM trial. *Leuk Lymphoma* 2024;1–11. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/38838026>.

123. Harrison CN, Mesa R, Talpaz M, et al. Efficacy and safety of fedratinib in patients with myelofibrosis previously treated with ruxolitinib (FREEDOM2): Results from a multicentre, open-label, randomised, controlled, phase 3 trial. *Lancet Haematol* 2024;11:e729–e740. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39265613>.

124. Vainchenker W, Leroy E, Gilles L, et al. JAK inhibitors for the treatment of myeloproliferative neoplasms and other disorders. *F1000Res* 2018;7:82. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29399328>.

125. Tefferi A. JAK inhibitors for myeloproliferative neoplasms: Clarifying facts from myths. *Blood* 2012;119:2721–2730. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22279053>.

126. Pardanani A, Harrison C, Cortes JE, et al. Safety and efficacy of fedratinib in patients with primary or secondary myelofibrosis: A randomized clinical trial. *JAMA Oncol* 2015;1:643–651. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26181658>.

127. Harrison CN, Mesa RA, Jamieson C, et al. Case series of potential wernicke's encephalopathy in patients treated with fedratinib [abstract]. *Blood* 2017;130:Abstract 4197. Available at:

http://www.bloodjournal.org/content/130/Suppl_1/4197.abstract.

128. National Organization for Rare Disorders. Wernicke-Korsakoff Syndrome. 2021. Available at: <https://rarediseases.org/rare-diseases/wernicke-korsakoff-syndrome/>. Accessed June 6, 2022.

129. Harrison C, Kiladjian JJ, Al-Ali HK, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med* 2012;366:787–798. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22375970>.

130. Davis KL, Cote I, Kaye JA, et al. Real-world assessment of clinical outcomes in patients with lower-risk myelofibrosis receiving treatment with ruxolitinib. *Adv Hematol* 2015;2015:848473. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26635878>.

131. Mead AJ, Milojkovic D, Knapper S, et al. Response to ruxolitinib in patients with intermediate-1-, intermediate-2-, and high-risk myelofibrosis: results of the UK ROBUST Trial. *Br J Haematol* 2015;170:29–39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25824940>.

132. Al-Ali HK, Griesshammer M, le Coutre P, et al. Safety and efficacy of ruxolitinib in an open-label, multicenter, single-arm phase 3b expanded-access study in patients with myelofibrosis: a snapshot of 1144 patients in the JUMP trial. *Haematologica* 2016;101:1065–1073. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27247324>.

133. Palandri F, Tiribelli M, Benevolo G, et al. Efficacy and safety of ruxolitinib in intermediate-1 IPSS risk myelofibrosis patients: Results from an independent study. *Hematol Oncol* 2018;36:285–290. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28512865>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

134. Al-Ali HK, Griesshammer M, Foltz L, et al. Primary analysis of JUMP, a phase 3b, expanded-access study evaluating the safety and efficacy of ruxolitinib in patients with myelofibrosis, including those with low platelet counts. *Br J Haematol* 2020;189:888–903. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32017044>.

135. Gupta V, Griesshammer M, Martino B, et al. Analysis of predictors of response to ruxolitinib in patients with myelofibrosis in the phase 3b expanded-access JUMP study. *Leuk Lymphoma* 2021;62:918–926. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33210570>.

136. Verstovsek S, Mesa RA, Gotlib J, et al. Efficacy, safety, and survival with ruxolitinib in patients with myelofibrosis: results of a median 3-year follow-up of COMFORT-I. *Haematologica* 2015;100:479–488. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25616577>.

137. Verstovsek S, Mesa RA, Gotlib J, et al. Long-term treatment with ruxolitinib for patients with myelofibrosis: 5-year update from the randomized, double-blind, placebo-controlled, phase 3 COMFORT-I trial. *J Hematol Oncol* 2017;10:55. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28228106>.

138. Cervantes F, Vannucchi AM, Kiladjan JJ, et al. Three-year efficacy, safety, and survival findings from COMFORT-II, a phase 3 study comparing ruxolitinib with best available therapy for myelofibrosis. *Blood* 2013;122:4047–4053. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24174625>.

139. Harrison CN, Vannucchi AM, Kiladjan JJ, et al. Long-term findings from COMFORT-II, a phase 3 study of ruxolitinib vs best available therapy for myelofibrosis. *Leukemia* 2016;30:1701–1707. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27211272>.

140. Verstovsek S, Mesa RA, Gotlib J, et al. The clinical benefit of ruxolitinib across patient subgroups: analysis of a placebo-controlled, Phase III study in patients with myelofibrosis. *Br J Haematol* 2013;161:508–516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23480528>.

141. Vannucchi AM, Kantarjian HM, Kiladjan JJ, et al. A pooled analysis of overall survival in COMFORT-I and COMFORT-II, 2 randomized phase III trials of ruxolitinib for the treatment of myelofibrosis. *Haematologica* 2015;100:1139–1145. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26069290>.

142. Verstovsek S, Kantarjian HM, Estrov Z, et al. Long-term outcomes of 107 patients with myelofibrosis receiving JAK1/JAK2 inhibitor ruxolitinib: Survival advantage in comparison to matched historical controls. *Blood* 2012;120:1202–1209. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22718840>.

143. Guglielmelli P, Gharardi A, Carobbio A, et al. Impact of ruxolitinib on survival of patients with myelofibrosis in the real world: Update of the ERNEST Study. *Blood Adv* 2022;6:373–375. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34753179>.

144. Gupta V, Harrison C, Hexner EO, et al. The impact of anemia on overall survival in patients with myelofibrosis treated with ruxolitinib in the COMFORT studies. *Haematologica* 2016;101:e482–e484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27587385>.

145. Cervantes F, Ross DM, Radinoff A, et al. Efficacy and safety of a novel dosing strategy for ruxolitinib in the treatment of patients with myelofibrosis and anemia: The REALISE phase 2 study. *Leukemia* 2021;35:3455–3465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34017073>.

146. Talpaz M, Prchal J, Afrin L, et al. Safety and efficacy of ruxolitinib in patients with myelofibrosis and low platelet counts (50 - 100 x 10⁹/L): Final analysis of an open-label phase 2 study. *Clin Lymphoma Myeloma Leuk* 2022;22:336–346. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34911667>.

147. Vannucchi AM, Te Boekhorst PAW, Harrison CN, et al. EXPAND, a dose-finding study of ruxolitinib in patients with myelofibrosis and low platelet counts: 48-week follow-up analysis. *Haematologica* 2019;104:947–954. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30442723>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

148. Heine A, Brossart P, Wolf D. Ruxolitinib is a potent immunosuppressive compound: is it time for anti-infective prophylaxis? *Blood* 2013;122:3843–3844. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24288410>.

149. Wysham NG, Sullivan DR, Allada G. An opportunistic infection associated with ruxolitinib, a novel janus kinase 1,2 inhibitor. *Chest* 2013;143:1478–1479. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23648912>.

150. Blechman AB, Cabell CE, Weinberger CH, et al. Aggressive skin cancers occurring in patients treated with the janus kinase inhibitor ruxolitinib. *J Drugs Dermatol* 2017;16:508–511. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28628689>.

151. Palandri F, Derenzini E, Ottaviani E, et al. Association of essential thrombocythemia and non-Hodgkin lymphoma: A single-centre experience. *Leuk Lymphoma* 2009;50:481–484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19241208>.

152. Vannucchi AM, Masala G, Antonioli E, et al. Increased risk of lymphoid neoplasms in patients with Philadelphia chromosome-negative myeloproliferative neoplasms. *Cancer Epidemiol Biomarkers Prev* 2009;18:2068–2073. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19531676>.

153. Rumi E, Passamonti F, Elena C, et al. Increased risk of lymphoid neoplasm in patients with myeloproliferative neoplasm: A study of 1,915 patients. *Haematologica* 2011;96:454–458. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21109692>.

154. Masarova L, Newberry KJ, Pierce SA, et al. Association of lymphoid malignancies and Philadelphia-chromosome negative myeloproliferative neoplasms: Clinical characteristics, therapy and outcome. *Leuk Res* 2015;39:822–827. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26012362>.

155. Porpaczy E, Tripolt S, Hoelbl-Kovacic A, et al. Aggressive B-cell lymphomas in patients with myelofibrosis receiving JAK1/2 inhibitor

therapy. *Blood* 2018;132:694–706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29907599>.

156. Pemmaraju N, Kantarjian H, Nastoupil L, et al. Characteristics of patients with myeloproliferative neoplasms with lymphoma, with or without JAK inhibitor therapy. *Blood* 2019;133:2348–2351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30796023>.

157. Rumi E, Zibellini S, Boveri E, et al. Ruxolitinib treatment and risk of B-cell lymphomas in myeloproliferative neoplasms. *Am J Hematol* 2019;94:E185–E188. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30972817>.

158. Barbui T, Ghirardi A, Masciulli A, et al. Second cancer in Philadelphia negative myeloproliferative neoplasms (MPN-K). A nested case-control study. *Leukemia* 2019;33:1996–2005. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31142846>.

159. Polverelli N, Elli EM, Abruzzese E, et al. Second primary malignancy in myelofibrosis patients treated with ruxolitinib. *Br J Haematol* 2021;193:356–368. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33222197>.

160. Guglielmelli P, Biamonte F, Rotunno G, et al. Impact of mutational status on outcomes in myelofibrosis patients treated with ruxolitinib in the COMFORT-II study. *Blood* 2014;123:2157–2160. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24458439>.

161. Pacilli A, Rotunno G, Mannarelli C, et al. Mutation landscape in patients with myelofibrosis receiving ruxolitinib or hydroxyurea. *Blood Cancer J* 2018;8:122. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30467377>.

162. Patel KP, Newberry KJ, Luthra R, et al. Correlation of mutation profile and response in patients with myelofibrosis treated with ruxolitinib. *Blood* 2015;126:790–797. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26124496>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

163. Maffioli M, Mora B, Ball S, et al. A prognostic model to predict survival after 6 months of ruxolitinib in patients with myelofibrosis. *Blood Adv* 2022;6:1855–1864. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35130339>.

164. Mesa RA, Kiladjan JJ, Catalano JV, et al. SIMPLIFY-1: A phase III randomized trial of momelotinib versus ruxolitinib in janus kinase inhibitor-naive patients with myelofibrosis. *J Clin Oncol* 2017;35:3844–3850. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28930494>.

165. Harrison CN, Vannucchi AM, Platzbecker U, et al. Momelotinib versus best available therapy in patients with myelofibrosis previously treated with ruxolitinib (SIMPLIFY 2): A randomised, open-label, phase 3 trial. *Lancet Haematol* 2018;5:e73–e81. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29275119>.

166. Mesa R, Harrison C, Oh ST, et al. Overall survival in the SIMPLIFY-1 and SIMPLIFY-2 phase 3 trials of momelotinib in patients with myelofibrosis. *Leukemia* 2022;36:2261–2268. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35869266>.

167. Hudgens S, Verstovsek S, Floden L, et al. Meaningful symptomatic change in patients with myelofibrosis from the SIMPLIFY studies. *Value Health* 2024;27:607–613. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38311180>.

168. Gerds AT, Savona MR, Scott BL, et al. Determining the recommended dose of pacritinib: results from the PAC203 dose-finding trial in advanced myelofibrosis. *Blood Adv* 2020;4:5825–5835. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33232476>.

169. Ajufo H, Bewersdorf JP, Harrison C, et al. Pacritinib response is associated with overall survival in myelofibrosis: PERSIST-2 landmark analysis of survival. *Eur J Haematol* 2025;114:238–247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39400386>.

170. Gagelmann N, Bose P, Gupta V, et al. Consistency of spleen and symptom reduction regardless of cytopenia in patients with myelofibrosis

treated with pacritinib. *Clin Lymphoma Myeloma Leuk* 2024;24:796–803. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39034203>.

171. Gagelmann N, Quarder M, Badbaran A, et al. Clearance of driver mutations after transplantation for myelofibrosis. *N Engl J Med* 2025;392:150–160. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39778169>.

172. Salit RB, Hexner EO, Gagelmann N, et al. Defining remission following hematopoietic cell transplant for myelofibrosis: An international expert panel consensus. *Leukemia* 2025;39:2862–2865. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41083679>.

173. McLornan D, Szydlo R, Koster L, et al. Myeloablative and reduced-intensity conditioned allogeneic hematopoietic stem cell transplantation in myelofibrosis: A retrospective study by the Chronic Malignancies Working Party of the European Society for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant* 2019;25:2167–2171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31284069>.

174. Snyder DS, Palmer J, Stein AS, et al. Allogeneic hematopoietic cell transplantation following reduced intensity conditioning for treatment of myelofibrosis. *Biol Blood Marrow Transplant* 2006;12:1161–1168. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17085309>.

175. Kroger N, Holler E, Kobbe G, et al. Allogeneic stem cell transplantation after reduced-intensity conditioning in patients with myelofibrosis: A prospective, multicenter study of the Chronic Leukemia Working Party of the European Group for Blood and Marrow Transplantation. *Blood* 2009;114:5264–5270. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19812383>.

176. Patriarca F, Bacigalupo A, Sperotto A, et al. Outcome of allogeneic stem cell transplantation following reduced-intensity conditioning regimen in patients with idiopathic myelofibrosis: The G.I.T.M.O. Experience. *Mediterr J Hematol Infect Dis* 2010;2:e2010010. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21415963>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

177. Snyder DS, Palmer J, Gaal K, et al. Improved outcomes using tacrolimus/sirolimus for graft-versus-host disease prophylaxis with a reduced-intensity conditioning regimen for allogeneic hematopoietic cell transplant as treatment of myelofibrosis. *Biol Blood Marrow Transplant* 2010;16:281–286. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19786111>.

178. Gupta V, Malone AK, Hari PN, et al. Reduced-intensity hematopoietic cell transplantation for patients with primary myelofibrosis: A cohort analysis from the center for international blood and marrow transplant research. *Biol Blood Marrow Transplant* 2014;20:89–97. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24161923>.

179. Lussana F, Rambaldi A, Finazzi MC, et al. Allogeneic hematopoietic stem cell transplantation in patients with polycythemia vera or essential thrombocythemia transformed to myelofibrosis or acute myeloid leukemia: A report from the MPN Subcommittee of the Chronic Malignancies Working Party of the European Group for Blood and Marrow Transplantation. *Haematologica* 2014;99:916–921. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24389309>.

180. Rondelli D, Goldberg JD, Isola L, et al. MPD-RC 101 prospective study of reduced-intensity allogeneic hematopoietic stem cell transplantation in patients with myelofibrosis. *Blood* 2014;124:1183–1191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24963042>.

181. Robin M, Porcher R, Wolschke C, et al. Outcome after transplantation according to reduced-intensity conditioning regimen in patients undergoing transplantation for myelofibrosis. *Biol Blood Marrow Transplant* 2016;22:1206–1211. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26970380>.

182. Jain T, Kunze KL, Temkit M, et al. Comparison of reduced intensity conditioning regimens used in patients undergoing hematopoietic stem cell transplantation for myelofibrosis. *Bone Marrow Transplant* 2019;54:204–211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29795431>.

183. Rezvani AR, McCune JS, Storer BE, et al. Cyclophosphamide followed by intravenous targeted busulfan for allogeneic hematopoietic cell

transplantation: Pharmacokinetics and clinical outcomes. *Biol Blood Marrow Transplant* 2013;19:1033–1039. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23583825>.

184. Salit RB, Scott BL, Stevens EA, et al. Pre-hematopoietic cell transplant Ruxolitinib in patients with primary and secondary myelofibrosis. *Bone Marrow Transplant* 2020;55:70–76. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30962501>.

185. Deeg HJ, Bredeson C, Farnia S, et al. Hematopoietic cell transplantation as curative therapy for patients with myelofibrosis: Long-term success in all age groups. *Biol Blood Marrow Transplant* 2015;21:1883–1887. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26371371>.

186. Ballen KK, Shrestha S, Sobocinski KA, et al. Outcome of transplantation for myelofibrosis. *Biol Blood Marrow Transplant* 2010;16:358–367. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19879949>.

187. Robin M, de Wreede LC, Wolschke C, et al. Long-term outcome after allogeneic hematopoietic cell transplantation for myelofibrosis. *Haematologica* 2019;104:1782–1788. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30733269>.

188. Hernandez-Boluda JC, Pereira A, Kroger N, et al. Determinants of survival in myelofibrosis patients undergoing allogeneic hematopoietic cell transplantation. *Leukemia* 2021;35:215–224. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32286544>.

189. Kunte S, Rybicki L, Viswabandya A, et al. Allogeneic blood or marrow transplantation with haploidentical donor and post-transplantation cyclophosphamide in patients with myelofibrosis: A multicenter study. *Leukemia* 2022;36:856–864. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/34663912>.

190. Jain T, Tsai HL, DeZern AE, et al. Post-transplantation cyclophosphamide-based graft-versus-host disease prophylaxis with nonmyeloablative conditioning for blood or marrow transplantation for



NCCN Guidelines Version 2.2026 Myeloproliferative Neoplasms

myelofibrosis. Transplant Cell Ther 2022;28:259.e251–259.e211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35158092>.

191. Polverelli N, Mauff K, Kroger N, et al. Impact of spleen size and splenectomy on outcomes of allogeneic hematopoietic cell transplantation for myelofibrosis: A retrospective analysis by the chronic malignancies working party on behalf of European society for blood and marrow transplantation (EBMT). Am J Hematol 2021;96:69–79. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33064301>.

192. Scott BL, Gooley TA, Sorrow ML, et al. The Dynamic International Prognostic Scoring System for myelofibrosis predicts outcomes after hematopoietic cell transplantation. Blood 2012;119:2657–2664. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22234678>.

193. Alchalby H, Yunus DR, Zabelina T, et al. Risk models predicting survival after reduced-intensity transplantation for myelofibrosis. Br J Haematol 2012;157:75–85. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22280409>.

194. Kroger N, Giorgino T, Scott BL, et al. Impact of allogeneic stem cell transplantation on survival of patients less than 65 years of age with primary myelofibrosis. Blood 2015;125:3347–3350. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25784679>.

195. Gowin K, Ballen K, Ahn KW, et al. Survival following allogeneic transplant in patients with myelofibrosis. Blood Adv 2020;4:1965–1973. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32384540>.

196. Hernandez-Boluda JC, Pereira A, Correa JG, et al. Prognostic risk models for transplant decision-making in myelofibrosis. Ann Hematol 2018;97:813–820. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29396714>.

197. Gagelmann N, Ditschkowski M, Bogdanov R, et al. Comprehensive clinical-molecular transplant scoring system for myelofibrosis undergoing stem cell transplantation. Blood 2019;133:2233–2242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30760453>.

198. Hernandez-Boluda JC, Pereira A, Alvarez-Larran A, et al. Predicting survival after allogeneic hematopoietic cell transplantation in myelofibrosis: Performance of the myelofibrosis transplant scoring system (MTSS) and development of a new prognostic model. Biol Blood Marrow Transplant 2020;26:2237–2244. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32717433>.

199. Panagiota V, Thol F, Markus B, et al. Prognostic effect of calreticulin mutations in patients with myelofibrosis after allogeneic hematopoietic stem cell transplantation. Leukemia 2014;28:1552–1555. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24504025>.

200. Kroger N, Panagiota V, Badbaran A, et al. Impact of molecular genetics on outcome in myelofibrosis patients after allogeneic stem cell transplantation. Biol Blood Marrow Transplant 2017;23:1095–1101. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28389256>.

201. Tamari R, Rapaport F, Zhang N, et al. Impact of high-molecular-risk mutations on transplantation outcomes in patients with myelofibrosis. Biol Blood Marrow Transplant 2019;25:1142–1151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30625392>.

202. Ali H, Aldoss I, Yang D, et al. MIPSS70+ v2.0 predicts long-term survival in myelofibrosis after allogeneic HCT with the Flu/Mel conditioning regimen. Blood Adv 2019;3:83–95. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30622146>.

203. Mannina D, Gagelmann N, Badbaran A, et al. Allogeneic stem cell transplantation in patients with myelofibrosis harboring the MPL mutation. Eur J Haematol 2019;103:552–557. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31446640>.

204. Stevens EA, Jenkins IC, Beppu LW, et al. Targeted sequencing improves DIPSS-Plus prognostic scoring in myelofibrosis patients undergoing allogeneic transplantation. Biol Blood Marrow Transplant 2020;26:1371–1374. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32194289>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

205. Martinez-Trillos A, Gaya A, Maffioli M, et al. Efficacy and tolerability of hydroxyurea in the treatment of the hyperproliferative manifestations of myelofibrosis: Results in 40 patients. *Ann Hematol* 2010;89:1233–1237. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20567824>.

206. Gisslinger H, Klade C, Georgiev P, et al. Ropoginterferon alfa-2b versus standard therapy for polycythaemia vera (PROUD-PV and CONTINUATION-PV): A randomised, non-inferiority, phase 3 trial and its extension study. *Lancet Haematol* 2020;7:e196–e208. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32014125>.

207. Mesa R, Gill H, Zhang L, et al. Ropoginterferon alfa-2b in hydroxyurea-intolerant or hydroxyurea-refractory essential thrombocythaemia (SURPASS ET): A multicentre, open-label, randomised, active-controlled, phase 3 study. *Lancet Haematol* 2025;12:e862–e875. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41193116>.

208. Kroger NM, Deeg JH, Olavarria E, et al. Indication and management of allogeneic stem cell transplantation in primary myelofibrosis: A consensus process by an EBMT/ELN international working group. *Leukemia* 2015;29:2126–2133. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26293647>.

209. Masarova L, Verstovsek S, Hidalgo-Lopez JE, et al. A phase 2 study of ruxolitinib in combination with azacitidine in patients with myelofibrosis. *Blood* 2018;132:1664–1674. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30185431>.

210. Chhabra S, Narra RK, Wu R, et al. Fludarabine/busulfan conditioning-based allogeneic hematopoietic cell transplantation for myelofibrosis: Role of ruxolitinib in improving survival outcomes. *Biol Blood Marrow Transplant* 2020;26:893–901. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31982543>.

211. Shanavas M, Popat U, Michaelis LC, et al. Outcomes of allogeneic hematopoietic cell transplantation in patients with myelofibrosis with prior exposure to janus kinase 1/2 inhibitors. *Biol Blood Marrow Transplant*

2016;22:432–440. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26493563>.

212. Gupta V, Kosiorek HE, Mead A, et al. Ruxolitinib therapy followed by reduced-intensity conditioning for hematopoietic cell transplantation for myelofibrosis: Myeloproliferative Disorders Research Consortium 114 Study. *Biol Blood Marrow Transplant* 2019;25:256–264. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30205231>.

213. Kroger N, Bacigalupo A, Barbui T, et al. Indication and management of allogeneic hematopoietic stem-cell transplantation in myelofibrosis: Updated recommendations by the EBMT/ELN International Working Group. *Lancet Haematol* 2024;11:e62–e74. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38061384>.

214. Polverelli N, Hernandez-Boluda JC, Czerw T, et al. Splenomegaly in patients with primary or secondary myelofibrosis who are candidates for allogeneic hematopoietic cell transplantation: A position paper on behalf of the Chronic Malignancies Working Party of the EBMT. *Lancet Haematol* 2023;10:e59–e70. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36493799>.

215. Kroger N, Sbianchi G, Sirait T, et al. Impact of prior JAK-inhibitor therapy with ruxolitinib on outcome after allogeneic hematopoietic stem cell transplantation for myelofibrosis: A study of the CMWP of EBMT. *Leukemia* 2021;35:3551–3560. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34023851>.

216. Hernandez-Boluda JC, Correa JG, Alvarez-Larran A, et al. Clinical characteristics, prognosis and treatment of myelofibrosis patients with severe thrombocytopenia. *Br J Haematol* 2018;181:397–400. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28419426>.

217. Scotch AH, Kosiorek H, Scherber R, et al. Symptom burden profile in myelofibrosis patients with thrombocytopenia: Lessons and unmet needs. *Leuk Res* 2017;63:34–40. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29096334>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

218. Tefferi A, Lasho TL, Jimma T, et al. One thousand patients with primary myelofibrosis: the mayo clinic experience. *Mayo Clin Proc* 2012;87:25–33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22212965>.

219. Jain AG, Gerds AT. How I treat anemia in myelofibrosis. *Blood* 2025;145:1738–1746. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39808793>.

220. Gupta V, Oh S, Devos T, et al. Momelotinib vs. ruxolitinib in myelofibrosis patient subgroups by baseline hemoglobin levels in the SIMPLIFY-1 trial. *Leuk Lymphoma* 2024;65:965–977. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38501751>.

221. Harrison CN, Vannucchi AM, Recher C, et al. Momelotinib versus continued ruxolitinib or best available therapy in JAK inhibitor-experienced patients with myelofibrosis and anemia: Subgroup analysis of SIMPLIFY-2. *Adv Ther* 2024;41:3722–3735. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38990433>.

222. Harrison CN, Mesa R, Talpaz M, et al. Longitudinal assessment of transfusion intensity in patients with JAK inhibitor-naïve or -experienced myelofibrosis treated with momelotinib. *Clin Lymphoma Myeloma Leuk* 2025;25:199–211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39516087>.

223. Oh ST, Mesa RA, Harrison CN, et al. Pacritinib is a potent ACVR1 inhibitor with significant anemia benefit in patients with myelofibrosis. *Blood Adv* 2023;7:5835–5842. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37552106>.

224. McMullin MF, Harrison CN, Niederwieser D, et al. The use of erythropoiesis-stimulating agents with ruxolitinib in patients with myelofibrosis in COMFORT-II: An open-label, phase 3 study assessing efficacy and safety of ruxolitinib versus best available therapy in the treatment of myelofibrosis. *Exp Hematol Oncol* 2015;4:26. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26380150>.

225. Crisa E, Cilloni D, Elli EM, et al. The use of erythropoiesis-stimulating agents is safe and effective in the management of anaemia in myelofibrosis patients treated with ruxolitinib. *Br J Haematol* 2018;182:701–704. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29984826>.

226. Huang J, Tefferi A. Erythropoiesis stimulating agents have limited therapeutic activity in transfusion-dependent patients with primary myelofibrosis regardless of serum erythropoietin level. *Eur J Haematol* 2009;83:154–155. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19366369>.

227. Cervantes F, Isola IM, Alvarez-Larran A, et al. Danazol therapy for the anemia of myelofibrosis: Assessment of efficacy with current criteria of response and long-term results. *Ann Hematol* 2015;94:1791–1796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26122869>.

228. Ashfaq S, Pellegrini MV, Can AS. Ashfaq S, Pellegrini MV, Can AS. Danazol. [Updated 2024 Feb 28]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2026 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK564344/>.

229. Cervantes F, Alvarez-Larran A, Hernandez-Boluda JC, et al. Erythropoietin treatment of the anaemia of myelofibrosis with myeloid metaplasia: results in 20 patients and review of the literature. *Br J Haematol* 2004;127:399–403. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15521916>.

230. Cervantes F, Alvarez-Larran A, Hernandez-Boluda JC, et al. Darbepoetin-alpha for the anaemia of myelofibrosis with myeloid metaplasia. *Br J Haematol* 2006;134:184–186. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16740139>.

231. Tsiara SN, Chaidos A, Bourantas LK, et al. Recombinant human erythropoietin for the treatment of anaemia in patients with chronic idiopathic myelofibrosis. *Acta Haematol* 2007;117:156–161. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17159338>.



NCCN Guidelines Version 2.2026 Myeloproliferative Neoplasms

232. Fenaux P, Kiladjian JJ, Platzbecker U. Luspatercept for the treatment of anemia in myelodysplastic syndromes and primary myelofibrosis. *Blood* 2019;133:790–794. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30602619>.

233. Gerds AT, Harrison CN, Kiladjian JJ, et al. Safety and efficacy of luspatercept for the treatment of anemia in patients with myelofibrosis. *Blood Adv* 2024;8:4511–4522. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38820422>.

234. Kiladjian JJ, Harrison C, Mesa RA, et al. Independence: Enrolling phase III trial to study the efficacy and safety of luspatercept versus placebo in patients with myelofibrosis on JAK2 inhibitor (JAK2i) therapy requiring red blood cell transfusions (RBCTs) [abstract]. *Clin Lymphoma Myeloma Leuk* 2023;23:Abstract MPN–346. Available at: <https://www.sciencedirect.com/science/article/pii/S215226502301234X>.

235. Barosi G, Elliott M, Canepa L, et al. Thalidomide in myelofibrosis with myeloid metaplasia: a pooled-analysis of individual patient data from five studies. *Leuk Lymphoma* 2002;43:2301–2307. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12613516>.

236. Elliott MA, Mesa RA, Li CY, et al. Thalidomide treatment in myelofibrosis with myeloid metaplasia. *Br J Haematol* 2002;117:288–296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11972510>.

237. Merup M, Kutti J, Birgergard G, et al. Negligible clinical effects of thalidomide in patients with myelofibrosis with myeloid metaplasia. *Med Oncol* 2002;19:79–86. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12180484>.

238. Marchetti M, Barosi G, Balestri F, et al. Low-dose thalidomide ameliorates cytopenias and splenomegaly in myelofibrosis with myeloid metaplasia: A phase II trial. *J Clin Oncol* 2004;22:424–431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14752066>.

239. Strupp C, Germing U, Scherer A, et al. Thalidomide for the treatment of idiopathic myelofibrosis. *Eur J Haematol* 2004;72:52–57. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14962263>.

240. Thomas DA, Giles FJ, Albitar M, et al. Thalidomide therapy for myelofibrosis with myeloid metaplasia. *Cancer* 2006;106:1974–1984. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16583431>.

241. Abgrall JF, Guibaud I, Bastie JN, et al. Thalidomide versus placebo in myeloid metaplasia with myelofibrosis: A prospective, randomized, double-blind, multicenter study. *Haematologica* 2006;91:1027–1032. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16885042>.

242. Mesa RA, Steensma DP, Pardanani A, et al. A phase 2 trial of combination low-dose thalidomide and prednisone for the treatment of myelofibrosis with myeloid metaplasia. *Blood* 2003;101:2534–2541. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12517815>.

243. Tefferi A, Cortes J, Verstovsek S, et al. Lenalidomide therapy in myelofibrosis with myeloid metaplasia. *Blood* 2006;108:1158–1164. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16609064>.

244. Quintas-Cardama A, Kantarjian HM, Manshouri T, et al. Lenalidomide plus prednisone results in durable clinical, histopathologic, and molecular responses in patients with myelofibrosis. *J Clin Oncol* 2009;27:4760–4766. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19720904>.

245. Mesa RA, Yao X, Cripe LD, et al. Lenalidomide and prednisone for myelofibrosis: Eastern Cooperative Oncology Group (ECOG) phase 2 trial E4903. *Blood* 2010;116:4436–4438. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20651074>.

246. Chihara D, Masarova L, Newberry KJ, et al. Long-term results of a phase II trial of lenalidomide plus prednisone therapy for patients with myelofibrosis. *Leuk Res* 2016;48:1–5. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27416326>.

247. Santana-Davila R, Tefferi A, Holtan SG, et al. Primary myelofibrosis is the most frequent myeloproliferative neoplasm associated with del(5q): Clinicopathologic comparison of del(5q)-positive and -negative cases. *Leuk Res* 2008;32:1927–1930. Available at: <https://pubmed.ncbi.nlm.nih.gov/18538839/>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

248. Jabbour E, Thomas D, Kantarjian H, et al. Comparison of thalidomide and lenalidomide as therapy for myelofibrosis. *Blood* 2011;118:899–902. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21622644>.

249. Schlenk RF, Stegelmann F, Reiter A, et al. Pomalidomide in myeloproliferative neoplasm-associated myelofibrosis. *Leukemia* 2017;31:889–895. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27774990>.

250. Tefferi A, Al-Ali HK, Barosi G, et al. A randomized study of pomalidomide vs placebo in persons with myeloproliferative neoplasm-associated myelofibrosis and RBC-transfusion dependence. *Leukemia* 2017;31:896–902. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27773929>.

251. Rampal RK, Verstovsek S, Devlin SM, et al. Safety and efficacy of combined ruxolitinib and thalidomide in patients with myelofibrosis: A phase II study [abstract]. *Blood* 2019;134:Abstract 4163. Available at: https://ashpublications.org/blood/article/134/Supplement_1/4163/425729/Safety-and-Efficacy-of-Combined-Ruxolitinib-and.

252. Stegelmann F, Jahn E, Koschmieder S, et al. Clinical and genetic results of the phase IB/II trial MPNSG-0212: Ruxolitinib plus pomalidomide in myelofibrosis with anemia [abstract]. *HemaSphere* 2022;6:945–946. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9429480/>.

253. Daver N, Cortes J, Newberry K, et al. Ruxolitinib in combination with lenalidomide as therapy for patients with myelofibrosis. *Haematologica* 2015;100:1058–1063. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26088933>.

254. Tefferi A, Barosi G, Mesa RA, et al. International Working Group (IWG) consensus criteria for treatment response in myelofibrosis with myeloid metaplasia, for the IWG for Myelofibrosis Research and Treatment (IWG-MRT). *Blood* 2006;108:1497–1503. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16675707>.

255. Tefferi A, Cervantes F, Mesa R, et al. Revised response criteria for myelofibrosis: International Working Group-Myeloproliferative Neoplasms

Research and Treatment (IWG-MRT) and European LeukemiaNet (ELN) consensus report. *Blood* 2013;122:1395–1398. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23838352>.

256. Tefferi A, Barosi G, Passamonti F, et al. Proposals for revised International Working Group-European LeukemiaNet criteria for anemia response in myelofibrosis. *Blood* 2024;144:1813–1820. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39116296>.

257. Tefferi A, Pardanani A. Serious adverse events during ruxolitinib treatment discontinuation in patients with myelofibrosis. *Mayo Clin Proc* 2011;86:1188–1191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22034658>.

258. Newberry KJ, Patel K, Masarova L, et al. Clonal evolution and outcomes in myelofibrosis after ruxolitinib discontinuation. *Blood* 2017;130:1125–1131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28674026>.

259. Mascarenhas J, Mehra M, He J, et al. Patient characteristics and outcomes after ruxolitinib discontinuation in patients with myelofibrosis. *J Med Econ* 2020;23:721–727. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32159402>.

260. Pardanani A, Gotlib JR, Jamieson C, et al. Safety and efficacy of TG101348, a selective JAK2 inhibitor, in myelofibrosis. *J Clin Oncol* 2011;29:789–796. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21220608>.

261. Deininger M, Radich J, Burn TC, et al. The effect of long-term ruxolitinib treatment on JAK2p.V617F allele burden in patients with myelofibrosis. *Blood* 2015;126:1551–1554. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26228487>.

262. Alchalby H, Badbaran A, Zabelina T, et al. Impact of JAK2V617F mutation status, allele burden, and clearance after allogeneic stem cell transplantation for myelofibrosis. *Blood* 2010;116:3572–3581. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20489052>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

263. Lange T, Edelmann A, Siebolts U, et al. JAK2 p.V617F allele burden in myeloproliferative neoplasms one month after allogeneic stem cell transplantation significantly predicts outcome and risk of relapse. *Haematologica* 2013;98:722–728. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23300178>.

264. Cervantes F. How I treat myelofibrosis. *Blood* 2014;124:2635–2642. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25232060>.

265. Kaur A, Wang S, Jayarangaiah A, et al. Real world risk of splenic rupture with G-CSF/GM-CSF therapy: A pharmacovigilance assessment using FDA Adverse Event Reporting System (FAERS) database [abstract]. *Blood* 2020;136:32–33. Available at: <https://ashpublications.org/blood/article/136/Supplement%201/32/473386/Real-World-Risk-of-Splenic-Rupture-with-G-CSF-GM>.

266. Landolfi R, Di Gennaro L, Barbui T, et al. Leukocytosis as a major thrombotic risk factor in patients with polycythemia vera. *Blood* 2007;109:2446–2452. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17105814>.

267. Gangat N, Wolanskyj AP, Schwager SM, et al. Leukocytosis at diagnosis and the risk of subsequent thrombosis in patients with low-risk essential thrombocythemia and polycythemia vera. *Cancer* 2009;115:5740–5745. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19806641>.

268. Carobbio A, Finazzi G, Antonioli E, et al. Thrombocytosis and leukocytosis interaction in vascular complications of essential thrombocythemia. *Blood* 2008;112:3135–3137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18587010>.

269. Carobbio A, Thiele J, Passamonti F, et al. Risk factors for arterial and venous thrombosis in WHO-defined essential thrombocythemia: an international study of 891 patients. *Blood* 2011;117:5857–5859. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21490340>.

270. Campbell PJ, MacLean C, Beer PA, et al. Correlation of blood counts with vascular complications in essential thrombocythemia: analysis of the

prospective PT1 cohort. *Blood* 2012;120:1409–1411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22709688>.

271. Carobbio A, Ferrari A, Masciulli A, et al. Leukocytosis and thrombosis in essential thrombocythemia and polycythemia vera: a systematic review and meta-analysis. *Blood Adv* 2019;3:1729–1737. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31175128>.

272. De Stefano V, Za T, Rossi E, et al. Recurrent thrombosis in patients with polycythemia vera and essential thrombocythemia: incidence, risk factors, and effect of treatments. *Haematologica* 2008;93:372–380. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18268279>.

273. De Stefano V, Za T, Rossi E, et al. Leukocytosis is a risk factor for recurrent arterial thrombosis in young patients with polycythemia vera and essential thrombocythemia. *Am J Hematol* 2010;85:97–100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20052743>.

274. Pasquer H, Daltro de Oliveira R, Vasseur L, et al. Distinct clinico-molecular arterial and venous thrombosis scores for myeloproliferative neoplasms risk stratification. *Leukemia* 2024;38:326–339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38148396>.

275. Marchioli R, Finazzi G, Landolfi R, et al. Vascular and neoplastic risk in a large cohort of patients with polycythemia vera. *J Clin Oncol* 2005;23:2224–2232. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15710945>.

276. Passamonti F, Caramazza D, Mora B, et al. It is time to change thrombosis risk assessment for PV and ET? *Best Pract Res Clin Haematol* 2014;27:121–127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25189723>.

277. Cervantes F, Passamonti F, Barosi G. Life expectancy and prognostic factors in the classic BCR/ABL-negative myeloproliferative disorders. *Leukemia* 2008;22:905–914. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18385755>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

278. Tefferi A, Guglielmelli P, Lasho TL, et al. Mutation-enhanced international prognostic systems for essential thrombocythaemia and polycythaemia vera. *Br J Haematol* 2020;189:291–302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31945802>.

279. Passamonti F, Thiele J, Girodon F, et al. A prognostic model to predict survival in 867 World Health Organization-defined essential thrombocythemia at diagnosis: A study by the International Working Group on Myelofibrosis Research and Treatment. *Blood* 2012;120:1197–1201. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22740446>.

280. Barbui T, Finazzi G, Carobbio A, et al. Development and validation of an International Prognostic Score of thrombosis in World Health Organization-essential thrombocythemia (IPSET-thrombosis). *Blood* 2012;120:5128–5133. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23033268>.

281. Barbui T, Vannucchi AM, Buxhofer-Ausch V, et al. Practice-relevant revision of IPSET-thrombosis based on 1019 patients with WHO-defined essential thrombocythemia. *Blood Cancer J* 2015;5:e369. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26617062>.

282. Haider M, Gangat N, Lasho T, et al. Validation of the revised International Prognostic Score of Thrombosis for Essential Thrombocythemia (IPSET-thrombosis) in 585 Mayo Clinic patients. *Am J Hematol* 2016;91:390–394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26799697>.

283. Landolfi R, Marchioli R, Kutti J, et al. Efficacy and safety of low-dose aspirin in polycythemia vera. *N Engl J Med* 2004;350:114–124. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14711910>.

284. Marchioli R, Finazzi G, Specchia G, et al. Cardiovascular events and intensity of treatment in polycythemia vera. *N Engl J Med* 2013;368:22–33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23216616>.

285. Alvarez-Larran A, Cervantes F, Pereira A, et al. Observation versus antiplatelet therapy as primary prophylaxis for thrombosis in low-risk

essential thrombocythemia. *Blood* 2010;116:1205–1210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20508163>.

286. Alvarez-Larran A, Pereira A, Guglielmelli P, et al. Antiplatelet therapy versus observation in low-risk essential thrombocythemia with a CALR mutation. *Haematologica* 2016;101:926–931. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27175028>.

287. Kaplan ME, Mack K, Goldberg JD, et al. Long-term management of polycythemia vera with hydroxyurea: a progress report. *Semin Hematol* 1986;23:167–171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3749925>.

288. Kiladjian JJ, Chevret S, Dosquet C, et al. Treatment of polycythemia vera with hydroxyurea and pipobroman: Final results of a randomized trial initiated in 1980. *J Clin Oncol* 2011;29:3907–3913. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21911721>.

289. Samuelsson J, Hasselbalch H, Bruserud O, et al. A phase II trial of pegylated interferon alpha-2b therapy for polycythemia vera and essential thrombocythemia: feasibility, clinical and biologic effects, and impact on quality of life. *Cancer* 2006;106:2397–2405. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16639737>.

290. Kiladjian JJ, Cassinat B, Chevret S, et al. Pegylated interferon-alfa-2a induces complete hematologic and molecular responses with low toxicity in polycythemia vera. *Blood* 2008;112:3065–3072. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18650451>.

291. Quintas-Cardama A, Abdel-Wahab O, Manshouri T, et al. Molecular analysis of patients with polycythemia vera or essential thrombocythemia receiving pegylated interferon alpha-2a. *Blood* 2013;122:893–901. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23782935>.

292. Masarova L, Patel KP, Newberry KJ, et al. Pegylated interferon alfa-2a in patients with essential thrombocythaemia or polycythaemia vera: a post-hoc, median 83 month follow-up of an open-label, phase 2 trial. *Lancet Haematol* 2017;4:e165–e175. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28291640>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

293. Barbui T, Vannucchi AM, De Stefano V, et al. Ropeginterferon versus standard therapy for low-risk patients with polycythemia vera. *NEJM Evid* 2023;2:EVIDoa2200335. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38320126>.

294. Fruchtmann SM, Mack K, Kaplan ME, et al. From efficacy to safety: a Polycythemia Vera Study group report on hydroxyurea in patients with polycythemia vera. *Semin Hematol* 1997;34:17–23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9025158>.

295. Finazzi G, Caruso V, Marchioli R, et al. Acute leukemia in polycythemia vera: an analysis of 1638 patients enrolled in a prospective observational study. *Blood* 2005;105:2664–2670. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15585653>.

296. Mascarenhas J, Kosiorek HE, Prchal JT, et al. A randomized phase 3 trial of interferon- α vs hydroxyurea in polycythemia vera and essential thrombocythemia. *Blood* 2022;139:2931–2941. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35007321>.

297. Kiladjian JJ, Klade C, Georgiev P, et al. Long-term outcomes of polycythemia vera patients treated with ropeginterferon Alfa-2b. *Leukemia* 2022;36:1408–1411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35210530>.

298. Gisslinger H, Klade C, Georgiev P, et al. Event-free survival in patients with polycythemia vera treated with ropeginterferon alfa-2b versus best available treatment. *Leukemia* 2023;37:2129–2132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37634011>.

299. Koschmieder S, Isfort S, Wolf D, et al. Efficacy and safety of ruxolitinib in patients with newly-diagnosed polycythemia vera: Futility analysis of the RuxoBEAT clinical trial of the GSG-MPN study group. *Ann Hematol* 2023;102:349–358. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36564535>.

300. Verstovsek S, Vannucchi AM, Griesshammer M, et al. Ruxolitinib versus best available therapy in patients with polycythemia vera: 80-week

follow-up from the RESPONSE trial. *Haematologica* 2016;101:821–829. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27102499>.

301. Kiladjian JJ, Guglielmelli P, Griesshammer M, et al. Efficacy and safety of ruxolitinib after and versus interferon use in the RESPONSE studies. *Ann Hematol* 2018;97:617–627. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29396713>.

302. Kiladjian JJ, Zachee P, Hino M, et al. Long-term efficacy and safety of ruxolitinib versus best available therapy in polycythaemia vera (RESPONSE): 5-year follow up of a phase 3 study. *Lancet Haematol* 2020;7:e226–e237. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31982039>.

303. Passamonti F, Griesshammer M, Palandri F, et al. Ruxolitinib for the treatment of inadequately controlled polycythaemia vera without splenomegaly (RESPONSE-2): a randomised, open-label, phase 3b study. *Lancet Oncol* 2017;18:88–99. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27916398>.

304. Griesshammer M, Saydam G, Palandri F, et al. Ruxolitinib for the treatment of inadequately controlled polycythemia vera without splenomegaly: 80-week follow-up from the RESPONSE-2 trial. *Ann Hematol* 2018;97:1591–1600. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29804268>.

305. Passamonti F, Palandri F, Saydam G, et al. Ruxolitinib versus best available therapy in inadequately controlled polycythaemia vera without splenomegaly (RESPONSE-2): 5-year follow up of a randomised, phase 3b study. *Lancet Haematol* 2022;9:e480–e492. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35597252>.

306. Mesa R, Vannucchi AM, Yacoub A, et al. The efficacy and safety of continued hydroxycarbamide therapy versus switching to ruxolitinib in patients with polycythaemia vera: a randomized, double-blind, double-dummy, symptom study (RELIEF). *Br J Haematol* 2017;176:76–85. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27858987>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

307. Harrison CN, Nangalia J, Boucher R, et al. Ruxolitinib versus best available therapy for polycythemia vera intolerant or resistant to hydroxycarbamide in a randomized trial. *J Clin Oncol* 2023;41:3534–3544. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37126762>.

308. Cortelazzo S, Finazzi G, Ruggeri M, et al. Hydroxyurea for patients with essential thrombocythemia and a high risk of thrombosis. *N Engl J Med* 1995;332:1132–1136. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7700286>.

309. Harrison CN, Campbell PJ, Buck G, et al. Hydroxyurea compared with anagrelide in high-risk essential thrombocythemia. *N Engl J Med* 2005;353:33–45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16000354>.

310. Gisslinger H, Gotic M, Holowiecki J, et al. Anagrelide compared with hydroxyurea in WHO-classified essential thrombocythemia: The ANAHYDRET study, a randomized controlled trial. *Blood* 2013;121:1720–1728. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23315161>.

311. Kanakura Y, Shirasugi Y, Yamaguchi H, et al. A phase 3b, multicenter, open-label extension study of the long-term safety of anagrelide in Japanese adults with essential thrombocythemia. *Int J Hematol* 2018;108:491–498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30121892>.

312. Kellner A, Dombi P, Illes A, et al. Anagrelide influences thrombotic risk, and prolongs progression-free and overall survival in essential thrombocythaemia vs hydroxyurea plus aspirin. *Eur J Haematol* 2020;105:408–418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32557810>.

313. Yacoub A, Mascarenhas J, Kosiorek H, et al. Pegylated interferon alfa-2a for polycythemia vera or essential thrombocythemia resistant or intolerant to hydroxyurea. *Blood* 2019;134:1498–1509. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31515250>.

314. O'Sullivan JM, Hamblin A, Yap C, et al. The poor outcome in high molecular risk, hydroxycarbamide-resistant/intolerant ET is not

ameliorated by ruxolitinib. *Blood* 2019;134:2107–2111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31578205>.

315. Verstovsek S, Passamonti F, Rambaldi A, et al. Ruxolitinib for essential thrombocythemia refractory to or intolerant of hydroxyurea: long-term phase 2 study results. *Blood* 2017;130:1768–1771. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28827411>.

316. Chu DK, Hillis CM, Leong DP, et al. Benefits and risks of antithrombotic therapy in essential thrombocythemia: a systematic review. *Ann Intern Med* 2017;167:170–180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28632284>.

317. Dillinger JG, Sideris G, Henry P, et al. Twice daily aspirin to improve biological aspirin efficacy in patients with essential thrombocythemia. *Thromb Res* 2012;129:91–94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22014557>.

318. Pascale S, Petrucci G, Dragani A, et al. Aspirin-insensitive thromboxane biosynthesis in essential thrombocythemia is explained by accelerated renewal of the drug target. *Blood* 2012;119:3595–3603. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22234683>.

319. Rocca B, Tosetto A, Petrucci G, et al. Long-term pharmacodynamic and clinical effects of twice- versus once-daily low-dose aspirin in essential thrombocythemia: The ARES trial. *Am J Hematol* 2024;99:1462–1474. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38877813>.

320. Rocca B, Tosetto A, Betti S, et al. A randomized double-blind trial of 3 aspirin regimens to optimize antiplatelet therapy in essential thrombocythemia. *Blood* 2020;136:171–182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32266380>.

321. Larsen ML, Pedersen OH, Hvas AM, et al. Once- versus twice-daily aspirin treatment in patients with essential thrombocytosis. *Platelets* 2019;30:322–328. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29442537>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

322. Godfrey AL, Campbell PJ, MacLean C, et al. Hydroxycarbamide plus aspirin versus aspirin alone in patients With essential thrombocythemia age 40 to 59 years without high-risk features. *J Clin Oncol* 2018;36:3361–3369. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30153096>.

323. Barosi G, Mesa R, Finazzi G, et al. Revised response criteria for polycythemia vera and essential thrombocythemia: an ELN and IWG-MRT consensus project. *Blood* 2013;121:4778–4781. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23591792>.

324. Vannucchi AM, Verstovsek S, Guglielmelli P, et al. Ruxolitinib reduces JAK2 p.V617F allele burden in patients with polycythemia vera enrolled in the RESPONSE study. *Ann Hematol* 2017;96:1113–1120. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28456851>.

325. Passamonti F, Rumi E, Pietra D, et al. A prospective study of 338 patients with polycythemia vera: the impact of JAK2 (V617F) allele burden and leukocytosis on fibrotic or leukemic disease transformation and vascular complications. *Leukemia* 2010;24:1574–1579. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20631743>.

326. Alvarez-Larran A, Bellosillo B, Pereira A, et al. JAK2V617F monitoring in polycythemia vera and essential thrombocythemia: clinical usefulness for predicting myelofibrotic transformation and thrombotic events. *Am J Hematol* 2014;89:517–523. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24458835>.

327. Borowczyk M, Wojtaszewska M, Lewandowski K, et al. The JAK2 V617F mutational status and allele burden may be related with the risk of venous thromboembolic events in patients with Philadelphia-negative myeloproliferative neoplasms. *Thromb Res* 2015;135:272–280. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25559461>.

328. Alvarez-Larran A, Perez-Encinas M, Ferrer-Marin F, et al. Risk of thrombosis according to need of phlebotomies in patients with polycythemia vera treated with hydroxyurea. *Haematologica* 2017;102:103–109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27686377>.

329. Barbui T, Carobbio A, Ghirardi A, et al. No correlation of intensity of phlebotomy regimen with risk of thrombosis in polycythemia vera: evidence from ECLAP and CYTO-PV clinical trials. *Haematologica* 2017;102:e219–e221. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28255021>.

330. Enblom-Larsson A, Girodon F, Bak M, et al. A retrospective analysis of the impact of treatments and blood counts on survival and the risk of vascular events during the course of polycythaemia vera. *Br J Haematol* 2017;177:800–805. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28474342>.

331. Hernandez-Boluda JC, Alvarez-Larran A, Gomez M, et al. Clinical evaluation of the European LeukaemiaNet criteria for clinicohaematological response and resistance/intolerance to hydroxycarbamide in essential thrombocythaemia. *Br J Haematol* 2011;152:81–88. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21083657>.

332. Alvarez-Larran A, Kerguelen A, Hernandez-Boluda JC, et al. Frequency and prognostic value of resistance/intolerance to hydroxycarbamide in 890 patients with polycythaemia vera. *Br J Haematol* 2016;172:786–793. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26898196>.

333. Barbui T, Barosi G, Birgegard G, et al. Philadelphia-negative classical myeloproliferative neoplasms: critical concepts and management recommendations from European LeukemiaNet. *J Clin Oncol* 2011;29:761–770. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21205761>.

334. Carobbio A, Finazzi G, Antonioli E, et al. Hydroxyurea in essential thrombocythemia: rate and clinical relevance of responses by European LeukemiaNet criteria. *Blood* 2010;116:1051–1055. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20479281>.

335. Alvarez-Larran A, Pereira A, Cervantes F, et al. Assessment and prognostic value of the European LeukemiaNet criteria for clinicohematologic response, resistance, and intolerance to hydroxyurea in



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

polycythemia vera. Blood 2012;119:1363–1369. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/22160617>.

336. Hernandez-Boluda JC, Pereira A, Cervantes F, et al. Clinical evaluation of the European LeukemiaNet response criteria in patients with essential thrombocythemia treated with anagrelide. Ann Hematol 2013;92:771–775. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23354997>.

337. Barbui T, Tefferi A, Vannucchi AM, et al. Philadelphia chromosome-negative classical myeloproliferative neoplasms: revised management recommendations from European LeukemiaNet. Leukemia 2018;32:1057–1069. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29515238>.

338. Connelly-Smith L, Alquist CR, Aqui NA, et al. Guidelines on the use of therapeutic apheresis in clinical practice - Evidence-based approach from the writing committee of the American Society for Apheresis: The ninth special issue. J Clin Apher 2023;38:77–278. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/37017433>.

339. Guyatt GH, Akl EA, Crowther M, et al. Executive summary: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:7S–47S. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/22315257>.

340. Kreher S, Ochsenreither S, Trappe RU, et al. Prophylaxis and management of venous thromboembolism in patients with myeloproliferative neoplasms: consensus statement of the Haemostasis Working Party of the German Society of Hematology and Oncology (DGHO), the Austrian Society of Hematology and Oncology (OGHO) and Society of Thrombosis and Haemostasis Research (GTH e.V.). Ann Hematol 2014;93:1953–1963. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25307456>.

341. Hernandez-Boluda JC, Arellano-Rodrigo E, Cervantes F, et al. Oral anticoagulation to prevent thrombosis recurrence in polycythemia vera and essential thrombocythemia. Ann Hematol 2015;94:911–918. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25680896>.

342. Enblom Larsson A, Renlund H, Andreasson B, et al. Thrombosis, major bleeding, and mortality in 1079 patients with myelofibrosis: A matched population-based study. Blood Adv 2025;9:2783–2793. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40117492>.

343. Ruggeri M, Rodeghiero F, Tosi A, et al. Postsurgery outcomes in patients with polycythemia vera and essential thrombocythemia: a retrospective survey. Blood 2008;111:666–671. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17909074>.

344. Griesshammer M, Struve S, Barbui T. Management of Philadelphia negative chronic myeloproliferative disorders in pregnancy. Blood Rev 2008;22:235–245. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/18617299>.

345. Passamonti F, Randi ML, Rumi E, et al. Increased risk of pregnancy complications in patients with essential thrombocythemia carrying the JAK2 (617V>F) mutation. Blood 2007;110:485–489. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17426257>.

346. Melillo L, Tieghi A, Candoni A, et al. Outcome of 122 pregnancies in essential thrombocythemia patients: A report from the Italian registry. Am J Hematol 2009;84:636–640. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19705431>.

347. Randi ML, Bertozzi I, Rumi E, et al. Pregnancy complications predict thrombotic events in young women with essential thrombocythemia. Am J Hematol 2014;89:306–309. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/24265194>.

348. Rumi E, Bertozzi I, Casetti IC, et al. Impact of mutational status on pregnancy outcome in patients with essential thrombocythemia. Haematologica 2015;100:e443–445. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26250575>.

349. Gangat N, Wolanskyj AP, Schwager S, Tefferi A. Predictors of pregnancy outcome in essential thrombocythemia: a single institution study of 63 pregnancies. Eur J Haematol 2009;82:350–353. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19243425>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

350. Passamonti F, Rumi E, Randi ML, et al. Aspirin in pregnant patients with essential thrombocythemia: a retrospective analysis of 129 pregnancies. *J Thromb Haemost* 2010;8:411–413. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19912517>.

351. How J, Leiva O, Bogue T, et al. Pregnancy outcomes, risk factors, and cell count trends in pregnant women with essential thrombocythemia. *Leuk Res* 2020;98:106459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33022566>.

352. Alimam S, Bewley S, Chappell LC, et al. Pregnancy outcomes in myeloproliferative neoplasms: UK prospective cohort study. *Br J Haematol* 2016;175:31–36. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27612319>.

353. Maze D, Kazi S, Gupta V, et al. Association of treatments for myeloproliferative neoplasms during pregnancy with birth rates and maternal outcomes: A systematic review and meta-analysis. *JAMA Netw Open* 2019;2:e1912666. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31584685>.

354. Robinson S, Bewley S, Hunt BJ, et al. The management and outcome of 18 pregnancies in women with polycythemia vera. *Haematologica* 2005;90:1477–1483. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16266894>.

355. Beauverd Y, Radia D, Cargo C, et al. Pegylated interferon alpha-2a for essential thrombocythemia during pregnancy: outcome and safety. A case series. *Haematologica* 2016;101:e182–184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26819057>.

356. Okikiolu J, Woodley C, Cadman-Davies L, et al. Real world experience with ropeginterferon alpha-2b (Besremi) in essential thrombocythaemia and polycythaemia vera following exposure to pegylated interferon alfa-2a (Pegasys). *Leuk Res Rep* 2023;19:100360. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36590864>.

357. Bang SY, Lee SE. A case report of ropeginterferon alfa-2b for polycythemia vera during pregnancy. *Hematol Rep* 2023;15:172–179. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36975731>.

358. Daei M, Khalili H, Heidari Z. Direct oral anticoagulant safety during breastfeeding: a narrative review. *Eur J Clin Pharmacol* 2021;77:1465–1471. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33963877>.

359. Bates SM, Rajasekhar A, Middeldorp S, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: venous thromboembolism in the context of pregnancy. *Blood Adv* 2018;2:3317–3359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30482767>.

360. Lapoirie J, Contis A, Guy A, et al. Management and outcomes of 27 pregnancies in women with myeloproliferative neoplasms. *J Matern Fetal Neonatal Med* 2020;33:49–56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29945475>.

361. Bertozzi I, Rumi E, Cavalloni C, et al. Pregnancy outcome and management of 25 pregnancies in women with polycythemia vera. *Am J Hematol* 2018;93:E234–E235. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29989197>.

362. McMullin MFF, Mead AJ, Ali S, et al. A guideline for the management of specific situations in polycythaemia vera and secondary erythrocytosis: A British Society for Haematology Guideline. *Br J Haematol* 2019;184:161–175. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30426472>.

363. Skeith L, Carrier M, Robinson SE, et al. Risk of venous thromboembolism in pregnant women with essential thrombocythemia: a systematic review and meta-analysis. *Blood* 2017;129:934–939. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28049641>.

364. Masarova L, Bose P, Pemmaraju N, et al. Prognostic value of blasts in peripheral blood in myelofibrosis in the ruxolitinib era. *Cancer* 2020;126:4322–4331. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32697338>.



NCCN Guidelines Version 2.2026 Myeloproliferative Neoplasms

365. Masarova L, Bose P, Pemmaraju N, et al. Clinical significance of bone marrow blast percentage in patients with myelofibrosis and the effect of ruxolitinib therapy. *Clin Lymphoma Myeloma Leuk* 2021;21:318–327.e316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33551345>.

366. Tam CS, Kantarjian H, Cortes J, et al. Dynamic model for predicting death within 12 months in patients with primary or post-polycythemia vera/essential thrombocythemia myelofibrosis. *J Clin Oncol* 2009;27:5587–5593. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19786661>.

367. Abdulkarim K, Girodon F, Johansson P, et al. AML transformation in 56 patients with Ph-MPD in two well defined populations. *Eur J Haematol* 2009;82:106–111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19134023>.

368. Nielsen I, Hasselbalch HC. Acute leukemia and myelodysplasia in patients with a Philadelphia chromosome negative chronic myeloproliferative disorder treated with hydroxyurea alone or with hydroxyurea after busulphan. *Am J Hematol* 2003;74:26–31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12949887>.

369. Bjorkholm M, Derolf AR, Hultcrantz M, et al. Treatment-related risk factors for transformation to acute myeloid leukemia and myelodysplastic syndromes in myeloproliferative neoplasms. *J Clin Oncol* 2011;29:2410–2415. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21537037>.

370. Noor SJ, Tan W, Wilding GE, et al. Myeloid blastic transformation of myeloproliferative neoplasms—a review of 112 cases. *Leuk Res* 2011;35:608–613. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20727590>.

371. Quintas-Cardama A, Kantarjian H, Pierce S, et al. Prognostic model to identify patients with myelofibrosis at the highest risk of transformation to acute myeloid leukemia. *Clin Lymphoma Myeloma Leuk* 2013;13:315–318.e312. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23391717>.

372. Klampfl T, Harutyunyan A, Berg T, et al. Genome integrity of myeloproliferative neoplasms in chronic phase and during disease

progression. *Blood* 2011;118:167–176. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21531982>.

373. Mesa RA, Li CY, Ketterling RP, et al. Leukemic transformation in myelofibrosis with myeloid metaplasia: A single-institution experience with 91 cases. *Blood* 2005;105:973–977. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15388582>.

374. Mascarenhas J, Navada S, Malone A, et al. Therapeutic options for patients with myelofibrosis in blast phase. *Leuk Res* 2010;34:1246–1249. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20627294>.

375. Thepot S, Itzykson R, Seegers V, et al. Treatment of progression of Philadelphia-negative myeloproliferative neoplasms to myelodysplastic syndrome or acute myeloid leukemia by azacitidine: A report on 54 cases on the behalf of the Groupe Francophone des Myelodysplasies (GFM). *Blood* 2010;116:3735–3742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20664061>.

376. Badar T, Kantarjian HM, Ravandi F, et al. Therapeutic benefit of decitabine, a hypomethylating agent, in patients with high-risk primary myelofibrosis and myeloproliferative neoplasm in accelerated or blastic/acute myeloid leukemia phase. *Leuk Res* 2015;39:950–956. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26183878>.

377. Rampal RK, Mascarenhas JO, Kosiorek HE, et al. Safety and efficacy of combined ruxolitinib and decitabine in accelerated and blast-phase myeloproliferative neoplasms. *Blood Adv* 2018;2:3572–3580. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30563881>.

378. Mascarenhas JO, Rampal RK, Kosiorek HE, et al. Phase 2 study of ruxolitinib and decitabine in patients with myeloproliferative neoplasm in accelerated and blast phase. *Blood Adv* 2020;4:5246–5256. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33104796>.

379. Bose P, Verstovsek S, Cortes JE, et al. A phase 1/2 study of ruxolitinib and decitabine in patients with post-myeloproliferative neoplasm acute myeloid leukemia. *Leukemia* 2020;34:2489–2492. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32099037>.



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

380. Gangat N, Guglielmelli P, Szuber N, et al. Venetoclax with azacitidine or decitabine in blast-phase myeloproliferative neoplasm: A multicenter series of 32 consecutive cases. *Am J Hematol* 2021;96:781–789. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33844862>.

381. King AC, Weis TM, Derkach A, et al. Multicenter evaluation of efficacy and toxicity of venetoclax-based combinations in patients with accelerated and blast phase myeloproliferative neoplasms. *Am J Hematol* 2022;97:E7–E10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34674293>.

382. Chen J, Wang K, Xiao Z, Xu Z. Efficacy and safety of combination therapies vs monotherapy of hypomethylating agents in accelerated or blast phase of Philadelphia negative myeloproliferative neoplasms: A systematic review and meta-analysis. *Ann Med* 2023;55:348–360. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36644935>.

383. Patel AA, Cahill K, Charnot-Katsikas A, et al. Clinical outcomes of IDH2-mutated advanced-phase Ph-negative myeloproliferative neoplasms treated with enasidenib. *Br J Haematol* 2020;190:e48–e51. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32358888>.

384. Chifotides HT, Masarova L, Alfayez M, et al. Outcome of patients with IDH1/2-mutated post-myeloproliferative neoplasm AML in the era of IDH inhibitors. *Blood Adv* 2020;4:5336–5342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33112940>.

385. Stone RM, Mandrekar SJ, Sanford BL, et al. Midostaurin plus chemotherapy for acute myeloid leukemia with a FLT3 mutation. *N Engl J Med* 2017;377:454–464. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28644114>.

386. Erba HP, Montesinos P, Kim HJ, et al. Quizartinib plus chemotherapy in newly diagnosed patients with FLT3-internal-tandem-duplication-positive acute myeloid leukaemia (QuANTUM-First): A randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2023;401:1571–1583. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37116523>.

387. Ciurea SO, de Lima M, Giral S, et al. Allogeneic stem cell transplantation for myelofibrosis with leukemic transformation. *Biol Blood Marrow Transplant* 2010;16:555–559. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20005966>.

388. Cherington C, Slack JL, Leis J, et al. Allogeneic stem cell transplantation for myeloproliferative neoplasm in blast phase. *Leuk Res* 2012;36:1147–1151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22578777>.

389. Kennedy JA, Atenafu EG, Messner HA, et al. Treatment outcomes following leukemic transformation in Philadelphia-negative myeloproliferative neoplasms. *Blood* 2013;121:2725–2733. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23361905>.

390. Alchalby H, Zabelina T, Stubig T, et al. Allogeneic stem cell transplantation for myelofibrosis with leukemic transformation: A study from the Myeloproliferative Neoplasm Subcommittee of the CMWP of the European Group for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant* 2014;20:279–281. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24201159>.

391. Gagelmann N, Wolschke C, Salit RB, et al. Reduced intensity hematopoietic stem cell transplantation for accelerated-phase myelofibrosis. *Blood Adv* 2022;6:1222–1231. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35051996>.

392. Patel AA, Yoon JJ, Johnston H, et al. Treatment approach and outcomes of patients with accelerated/blast-phase myeloproliferative neoplasms in the current era. *Blood Adv* 2024;8:3468–3477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38739724>.

393. Jeon JY, Zhao Q, Buelow DR, et al. Preclinical activity and a pilot phase I study of pacritinib, an oral JAK2/FLT3 inhibitor, and chemotherapy in FLT3-ITD-positive AML. *Invest New Drugs* 2020;38:340–349. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31102119>.

394. Ma Y, Remsing Rix L, Zhang Q, et al. Pacritinib (PAC) synergistically potentiates azacitidine (5AZA) cytotoxicity in chronic myelomonocytic



NCCN Guidelines Version 2.2026

Myeloproliferative Neoplasms

leukemia (CMML) [abstract]. Blood 2015;126:Abstract 1658. Available at:
<https://ashpublications.org/blood/article/126/23/1658/134818/Pacritinib-PAC-Synergistically-Potentiates>.