



National Comprehensive
Cancer Network®


NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Neuroblastoma

NCCN Evidence Blocks™

Version 2.2026 — April 28, 2026

NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.





NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

***Rochelle Bagatell, MD/Chair €**

Abramson Cancer Center
at the University of Pennsylvania |
Children's Hospital of Philadelphia

Julie R. Park, MD/Vice-Chair €

St. Jude Children's Research Hospital/
The University of Tennessee Health Science Center

Sahaja Acharya, MD §

Johns Hopkins Kimmel Cancer Center |
Johns Hopkins Children's Center

Jennifer Aldrink, MD ¶

The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute | Nationwide
Children's Hospital

Jenna Allison, MD €

Fred & Pamela Buffett Cancer Center |
Children's Hospital & Medical Center

Elizabeth Alva, MD, MSPH €

O'Neal Comprehensive Cancer Center at UAB |
Children's of Alabama

Jennifer Belle, MBA ¥

Children's Oncology Group

Daniel Benedetti, MD, MA €

Vanderbilt-Ingram Cancer Center | Monroe Carell Jr.
Children's Hospital at Vanderbilt

Erin Brown, MD ¶

UC Davis Comprehensive Cancer Center

Alanna Church, MD ≠

Dana-Farber/Brigham and Women's Cancer Center |
Mass General Cancer Center | Dana-Farber/Boston
Children's Cancer and Blood Disorders Center

Ami V. Desai, MD, MSCE ‡ €

The UChicago Medicine
Comprehensive Cancer Center

NCCN

Lisa Hang, PhD

Jamie Nguyen, PharmD

Ryan Schonfeld, BA

Steven DuBois, MD, MS €

Dana-Farber/Brigham and Women's Cancer Center |
Mass General Cancer Center | Dana-Farber/Boston
Children's Cancer and Blood Disorders Center

Joaquim Farinhas, MD, MBA ϕ

Moffitt Cancer Center

Kara Gill, MD ϕ

University of Wisconsin Carbone Cancer Center |
American Family Children's Hospital

Frederick Huang, MD ‡ €

Siteman Cancer Center at Barnes-Jewish Hospital
and WashU Medicine | St. Louis Children's Hospital

Paul Iskander, MD ϕ

UCLA Jonsson Comprehensive Cancer Center |
UCLA Mattel Children's Hospital

Margaret Macy, MD €

University of Colorado Cancer Center |
Children's Hospital Colorado

Alp Oztek, MD, MRCP ϕ ϕ €

Fred Hutchinson Cancer Center |
Seattle Children's Hospital

Farzana Pashankar, MD, MBBS €

Yale Cancer Center/Smilow Cancer Hospital |
Yale New Haven Children's Hospital

Praveen Pendyala, MD §

Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute |
Cleveland Clinic Children's

Stephanie Polites, MD, MPH ¶

Mayo Clinic Comprehensive Cancer Center

Raja Rabah, MD ≠

University of Michigan Rogel Cancer Center |
C.S. Mott Children's Hospital

Hiroyuki Shimada, MD ≠

Stanford Cancer Institute |
Lucile Packard Children's Hospital

Leonora Slatnick, MD ‡

Huntsman Cancer Institute at the University of Utah
| Primary Children's Hospital

Elizabeth Sokol, MD ‡ €

Robert H. Lurie Comprehensive Cancer
Center of Northwestern University |
Ann & Robert H. Lurie Children's Hospital of Chicago

Jessica Sun, MD €

Duke Cancer Institute |
Duke Children's Hospital & Health Center

Mark Tann, MD ϕ

Indiana University Melvin and Bren Simon
Comprehensive Cancer Center | Riley Children's
Health

Clare Twist, MD €

Roswell Park Comprehensive Cancer Center |
Roswell Park Oishei Children's Cancer and Blood
Disorders Program

Kieuhoa Vo, MD €

UCSF Helen Diller Family Comprehensive Cancer
Center | UCSF Benioff Children's Hospital

Tanya Watt, MD €

UT Southwestern Simmons Comprehensive Cancer
Center | Children's Medical Center Dallas

Suzanne Wolden, MD §

Memorial Sloan Kettering Cancer Center

Peter Zage, MD, PhD ‡ €

UC San Diego Moores Cancer Center |
Rady Children's Hospital-San Diego

ϕ Diagnostic radiology
‡ Hematology
‡ Internal medicine
† Medical oncology
ϕ Nuclear medicine
≠ Pathology
¥ Patient advocacy
€ Pediatric oncology
§ Radiotherapy/Radiation oncology
¶ Surgery/Surgical oncology
* Discussion Writing Committee Member

NCCN Guidelines Panel Disclosures



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

[NCCN Neuroblastoma Panel Members](#)
[NCCN Evidence Blocks Definitions \(EB-DEF\)](#)

[Diagnosis/Workup \(NEUROB-1\)](#)
[Risk Classification \(NEUROB-2\)](#)
Primary Treatment
• [Low-Risk Disease \(NEUROB-3\)](#)
• [Intermediate-Risk Disease \(NEUROB-4\)](#)
• [High-Risk Disease \(NEUROB-7\)](#)

[Principles of Pathology \(NEUROB-A\)](#)
[Principles of Imaging \(NEUROB-B\)](#)
[Principles of Risk Classification \(NEUROB-C\)](#)
[Principles of Systemic Therapy \(NEUROB-D\)](#)
[Principles of Surgery \(NEUROB-E\)](#)
[Surveillance/Follow-up \(NEUROB-F\)](#)
[Response Assessment \(NEUROB-G\)](#)
[Principles of Radiation Therapy \(NEUROB-H\)](#)
[Monitoring for Late Effects \(NEUROB-I\)](#)
[Principles of Paraneoplastic Syndromes \(NEUROB-J\)](#)

[International Neuroblastoma Risk Group \(INRG\) Staging \(ST-1\)](#)

[Evidence Blocks](#)

[Abbreviations \(ABBR-1\)](#)

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS

5					
4					
3					
2					
1					

E S Q C A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

Example Evidence Block

5					
4					
3					
2					
1					

E S Q C A

E = 4
S = 4
Q = 3
C = 4
A = 3

Efficacy of Regimen/Agent

5	Highly effective: Cure likely and often provides long-term survival advantage
4	Very effective: Cure unlikely but sometimes provides long-term survival advantage
3	Moderately effective: Modest impact on survival, but often provides control of disease
2	Minimally effective: No, or unknown impact on survival, but sometimes provides control of disease
1	Palliative: Provides symptomatic benefit only

Safety of Regimen/Agent

5	Usually no meaningful toxicity: Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
4	Occasionally toxic: Rare significant toxicities or low-grade toxicities only; little interference with ADLs
3	Mildly toxic: Mild toxicity that interferes with ADLs
2	Moderately toxic: Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
1	Highly toxic: Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

Note: For significant chronic or long-term toxicities, score decreased by 1

Quality of Evidence

5	High quality: Multiple well-designed randomized trials and/or meta-analyses
4	Good quality: One or more well-designed randomized trials
3	Average quality: Low quality randomized trial(s) or well-designed non-randomized trial(s)
2	Low quality: Case reports or extensive clinical experience
1	Poor quality: Little or no evidence

Consistency of Evidence

5	Highly consistent: Multiple trials with similar outcomes
4	Mainly consistent: Multiple trials with some variability in outcome
3	May be consistent: Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
2	Inconsistent: Meaningful differences in direction of outcome between quality trials
1	Anecdotal evidence only: Evidence in humans based upon anecdotal experience

Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)

5	Very inexpensive
4	Inexpensive
3	Moderately expensive
2	Expensive
1	Very expensive



DIAGNOSTIC WORKUP

- **Tissue sampling^a**
 - ▶ Surgical resection should be considered in the setting of localized disease, particularly in the absence of image-defined risk factors (IDRFs). When biopsy rather than upfront resection is indicated, open biopsy may be preferred in some clinical scenarios if adequate tissue is not obtained/obtainable by less invasive means.
 - ◊ See [Principles of Pathology \(NEUROB-A\)](#) for biomarker testing considerations and additional information.
 - ▶ Multiple core biopsies may provide adequate tissue; the need for full histologic and molecular evaluation of specimens must be considered. When sufficient tissue is obtained, the use of additional tissue for research purposes is encouraged.
 - ▶ Fine-needle aspiration is not recommended.
 - ▶ Review of frozen sections by an experienced pathologist is important to determine specimen adequacy, as some samples may be necrotic at the time of initial biopsy.
- **Bilateral bone marrow aspirates and trephine biopsies**
 - ▶ In rare cases, marrow may be the only source of diagnostic material. In those cases, every attempt should be made to obtain material for the full complement of biomarker testing.
 - ◊ See [Principles of Pathology \(NEUROB-A\)](#) for biomarker testing considerations and additional information.

ADDITIONAL WORKUP

- **History and physical (H&P):** attention to abdominal exam for mass and organomegaly; thorough neurologic exam.
- **Family history:** attention to relatives with neuroblastoma or other childhood cancers¹
- **Additional considerations**
 - ▶ **Essential:**
 - ◊ Complete blood count (CBC) with differential
 - ◊ Comprehensive metabolic panel
 - ◊ Homovanillic acid (HVA)/vanillylmandelic acid (VMA): required for diagnosis if only diagnostic tissue is bone marrow or if no biopsy is to be obtained initially
 - ▶ **Useful in selected cases:**
 - ◊ Prothrombin time (PT)/international normalized ratio (INR) if liver is involved or if there is concern for bleeding
 - ◊ Lactate dehydrogenase (LDH), ferritin may be considered
 - ◊ Pregnancy test (for patients of childbearing potential)
 - ◊ **For patients receiving systemic therapy**
 - Audiogram if platinum-containing chemotherapy is indicated
 - Echocardiogram if anthracycline-containing chemotherapy is indicated
 - Electrocardiogram (ECG) if anthracycline-containing chemotherapy is indicated
 - Fertility/fertility preservation: Refer to fertility specialists for discussion of possible options and appropriate timing (for patients likely to receive extensive alkylator therapy)
- **Imaging**
 - ▶ Cross-sectional imaging (MRI with/without contrast or CT with contrast) to evaluate soft tissue disease
 - ▶ MRI of spine with/without contrast in the setting of paraspinal disease or concerns regarding involvement of nerve roots or spinal cord
 - ▶ MRI of brain with/without contrast or CT skull/orbits with contrast if neurologic symptoms are present or if otherwise clinically indicated
 - ▶ Iodine-123 (I-123) meta-iodobenzylguanidine (MIBG) imaging (with SPECT or SPECT/CT, where available) to assess for metastatic disease
 - ▶ Fluorodeoxyglucose (FDG)-PET should be obtained in patients with MIBG non-avid disease or suspected mixed-avidity disease^b
 - ▶ See [Principles of Imaging \(NEUROB-B\)](#) for radiology initial workup and additional information
- **Paraneoplastic syndromes** may occur with neuroblastoma. See [NEUROB-J](#) for additional information

[Risk Classification \(NEUROB-2\)](#)

^a The following patients should have biopsy delayed until after therapy is started and the patient is medically stable:

- ▶ Patients <2 months of age with existing or evolving hepatomegaly (Twist CJ, et al. J Clin Oncol 2019;37:115-124.)
- ▶ Infants in whom safety considerations (coagulopathy, impending organ failure) preclude biopsy
- ▶ In these cases, biopsy should be performed when it is safe to do so to obtain tissue for biomarker testing

Patients <6 months of age with L1 adrenal tumor with maximum diameter ≤3.1 cm if solid or 5 cm if at least 25% cystic component do not require a biopsy (Nuchtern JG, et al. Ann Surg 2012;256:573-580).

^b Not required in patients <6 months of age with L1 adrenal tumor with maximum diameter ≤3.1 cm if solid or 5 cm if at least 25% cystic component. (Nuchtern JG, et al. Ann Surg 2012;256:573-580).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

References

NEUROB-1



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

NEUROBLASTOMA RISK CLASSIFICATION^c

Clinical Stage ^d	MYCN amp	Age	Other Features (Ploidy/International Neuroblastoma Pathology Classification [INPC]/Histology/Extent of Resection/Clinical Symptoms)	Risk Group
L1 ^e	No	Any	Any	Low risk
	Yes	Any	Completely resected	Low risk
	Yes	Any	Incompletely resected	High risk
L2	No	<18 months	Any	Intermediate risk
		18 months to <5 years	FH ^f	Intermediate risk
			UH ^f	High risk ⁱ
		≥5 years	Differentiating per INPC ^f	Intermediate risk
	Undifferentiated, poorly differentiated ^f		High risk ⁱ	
	Yes	Any	Any	High risk
M	No	<12 months	Any	Intermediate risk
	Yes			High risk
	No	≥12 to <18 months	UH, ^f DNA index (DI)=1, ^g or segmental chromosomal aberrations (SCA+) See Principles of Pathology (NEUROB-A)	High risk
			FH, ^f DI >1, ^g and SCA- (category 2B) ²	Intermediate risk ^j
	Yes	≥18 months	Any	High risk
	Any		Any	High risk
MS	Unknown	<12 months	Symptomatic ^h	Intermediate risk
	No		UH, ^f DI = 1, ^g or SCA+	Intermediate risk
			Asymptomatic, FH, ^f DI >1, ^g and SCA-	Low risk
	Yes	12 to <18 months	Any	High risk
	No		UH, ^f DI = 1, ^g or SCA+	High risk
			FH, ^f DI >1, ^g and SCA-	Intermediate risk
Yes	Any	High risk		

FH: Favorable histology; UH: Unfavorable histology

Adapted from Irwin MS, Naranjo A, Zhang FF, et al. Revised Neuroblastoma Risk Classification System: A report from the Children's Oncology Group. J Clin Oncol 2021;39:3229-3241.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

[Footnotes](#)
[References](#)



NEUROBLASTOMA RISK CLASSIFICATION FOOTNOTES

^c See [Principles of Pathology \(NEUROB-A\)](#) and [Principles of Risk Classification \(NEUROB-C\)](#) for additional considerations.

^d See [ST-1](#) for International Neuroblastoma Risk Group (INRG) Staging Criteria.

^e Perinatally diagnosed children with L1 tumors will be discussed in the treatment section. See [Low-Risk Disease \(NEUROB-3\)](#) and [Principles of Systemic Therapy \(NEUROB-D\)](#).

^f Pathology should be reviewed by a pathologist with expertise in classification of neuroblastoma.

^g If information about DI is missing, assign risk based on other available factors. Ploidy may be inferred by next-generation sequencing (NGS).

^h Infants in whom safety considerations (coagulopathy, impending organ failure) preclude biopsy, especially those with hepatomegaly leading to respiratory compromise. Symptomatic patients should start intermediate-risk treatment emergently prior to biopsy. Once symptoms have resolved following initial therapy, patients should undergo biopsy and risk assignment should be revised based on available data. See Hsu LL, et al. Med Pediatr Oncol 1996;27:521-528 for symptoms; a total score ≥ 2 should lead to emergent therapy.

ⁱ Cooperative groups outside North America may treat these patients as intermediate risk.

^j Cooperative groups outside North America may treat all patients with INRG Stage M disease ≥ 12 months as high risk.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

LOW-RISK DISEASE

PRIMARY TREATMENT^{l,m}



^k Obtain MIBG scan to confirm localized disease. If the patient is <6 months of age with adrenal tumor, no MIBG scan is required, unless signs and symptoms are suggestive of metastatic disease.

^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^m [Principles of Surgery \(NEUROB-E\)](#).

ⁿ Use ultrasound (US) for surveillance when clinically indicated and appropriate.

^o Obtain cross-sectional imaging to delineate new baseline disease status at least 1 month postoperatively, then transition to US for surveillance if possible.

^p If resection is incomplete and *MYCN* amplification is detected, see [Risk Classification \(NEUROB-2\)](#).

^q [Disease Surveillance/Follow-up \(NEUROB-F\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026


Neuroblastoma

NCCN Evidence Blocks™

INTERMEDIATE-RISK DISEASE

PRIMARY TREATMENT^m

RESPONSE



^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^m [Principles of Surgery \(NEUROB-E\)](#).

^q [Disease Surveillance/Follow-up \(NEUROB-F\)](#).

^r Favorable biologic features include FH, DI >1, no SCA. If SCA or histology status are not available, consider as unfavorable.

^s Audiologic assessment and echocardiogram should be obtained at diagnosis and again at end of therapy.

^t Historically measured by volume. See [NEUROB-D 1 of 13](#) and [Response Assessment \(NEUROB-G\)](#).

^u Multidisciplinary discussions should be undertaken on an iterative basis. Surveillance should begin once the response or reduction in primary tumor size noted above is achieved, with chemotherapy and/or surgery.

^v If additional chemotherapy is given, consider 2–4 additional cycles with repeat primary tumor imaging after every 2 cycles. Consider surgery if chemotherapy has led to stable disease (SD) or minimal improvement in tumor size. Multidisciplinary discussion is important in decisions surrounding surgery and additional chemotherapy. Consider biopsy as the tumor may have differentiated and additional treatment may not be required. If 8 cycles of chemotherapy have been given and the tumor remains unresectable, consider treatment with second-line chemotherapy as per Twist CJ, et al. J Clin Oncol 2019;37:3243-3255.

^w If the treatment endpoint has been achieved, obtain MIBG scan as part of the end of therapy evaluation.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

INTERMEDIATE-RISK DISEASE PRIMARY TREATMENT^m RESPONSE



^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^m [Principles of Surgery \(NEUROB-E\)](#).

^q [Disease Surveillance/Follow-up \(NEUROB-F\)](#).

^r Favorable biologic features include FH, DI >1, no SCA. If SCA or histology status are not available, consider as unfavorable.

^s Audiologic assessment and echocardiogram should be obtained at diagnosis and again at end of therapy.

^t Historically measured by volume. See [NEUROB-D 1 of 13](#) and [Response Assessment \(NEUROB-G\)](#).

^u Multidisciplinary discussion should be undertaken on an iterative basis. Surveillance should begin once the response or reduction in primary tumor size noted above is achieved, with chemotherapy and/or surgery.

^v If additional chemotherapy is given, consider 2–4 additional cycles with repeat primary tumor imaging after every 2 cycles. Consider surgery if chemotherapy has led to SD or minimal improvement in tumor size. Multidisciplinary discussion is important in decisions surrounding surgery and additional chemotherapy. Consider biopsy as the tumor may have differentiated and additional treatment may not be required. If 8 cycles of chemotherapy have been given and the tumor remains unresectable, consider treatment with second-line chemotherapy as per Twist CJ, et al. J Clin Oncol 2019;37:3243-3255.

^w If the treatment endpoint has been achieved, obtain MIBG scan as part of the end of therapy evaluation.

^x Evaluate all prior sites of disease after 4 cycles if only 4 cycles of chemotherapy are planned. If 8 cycles are planned, evaluate after cycles 4 and 8.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026


Neuroblastoma

NCCN Evidence Blocks™

INTERMEDIATE-RISK DISEASE

PRIMARY TREATMENT^m

RESPONSE



^h Infants in whom safety considerations (coagulopathy, impending organ failure) preclude biopsy, especially those with hepatomegaly leading to respiratory compromise. Symptomatic patients should start intermediate-risk treatment emergently prior to biopsy. Once symptoms have resolved following initial therapy, patients should undergo biopsy and risk assignment should be revised based on available data. See Hsu LL, et al. Med Pediatr Oncol 1996;27:521-528 for symptoms; a total score ≥ 2 should lead to emergent therapy.

^l Principles of Systemic Therapy (NEUROB-D).

^m Principles of Surgery (NEUROB-E).

^q Disease Surveillance/Follow-up (NEUROB-F).

^r Favorable biologic features include FH, DI > 1 , no SCA. If SCA or histology status are not available, consider as unfavorable.

^s Audiologic assessment and echocardiogram should be obtained at diagnosis and again at end of therapy.

^t Historically measured by volume. See [NEUROB-D 1 of 13](#) and [Response Assessment \(NEUROB-G\)](#).

^u Multidisciplinary discussion should be undertaken on an iterative basis. Surveillance should begin once the response or reduction in primary tumor size noted above is achieved, with chemotherapy and/or surgery.

^v If additional chemotherapy is given, consider 2–4 additional cycles with repeat primary tumor imaging after every 2 cycles. Consider surgery if chemotherapy has led to SD or minimal improvement in tumor size. Multidisciplinary discussion is important in decisions surrounding surgery and additional chemotherapy. Consider biopsy as the tumor may have differentiated and additional treatment may not be required. If 8 cycles of chemotherapy have been given and the tumor remains unresectable, consider treatment with second-line chemotherapy as per Twist CJ, et al. J Clin Oncol 2019;37:3243-3255.

^w If the treatment endpoint has been achieved, obtain MIBG scan as part of the end of therapy evaluation.

^y For infants who are too unstable to undergo biopsy before starting treatment, chemotherapy may be initiated and a biopsy obtained when it is safe to do so.

^z Symptomatic patients with ongoing organ dysfunction with inadequate response to initial chemotherapy may benefit from radiation therapy (RT); multidisciplinary discussion is recommended. See [Principles of Radiation Therapy \(NEUROB-H\)](#).

^{aa} Evaluate all prior sites of disease after 2 cycles if only 2 cycles of chemotherapy are planned. If 4 cycles are planned, perform the complete evaluation of all sites of prior disease after 4 cycles.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma


NCCN Evidence Blocks™

OVERVIEW OF TREATMENT FOR HIGH-RISK NEUROBLASTOMA

INDUCTION THERAPY

CONSOLIDATION THERAPY

POST-CONSOLIDATION THERAPY



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

HIGH-RISK DISEASE

INDUCTION THERAPY



^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^m [Principles of Surgery \(NEUROB-E\)](#).

^{bb} Anatomic imaging of primary site prior to surgery.

^{cc} Subtotal resection if gross total will jeopardize vital organs, major nerves, and/or major vessels.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026


Neuroblastoma

NCCN Evidence Blocks™

HIGH-RISK DISEASE

RESPONSE³

TREATMENT AFTER STANDARD INDUCTION



^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^{dd} Bridging therapy may be appropriate depending upon degree of response. See [Principles of Systemic Therapy \(NEUROB-D 4 of 13\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

[References](#)

NEUROB-9




NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

HIGH-RISK DISEASE

CONSOLIDATION THERAPY



^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^{ee} Common high-risk groups include all other categories not included as part of special high-risk groups.

^{ff} Special high-risk groups include:

Stage L2, ≥18 months, *MYCN* non-amplified, and UH

Stage M, 12 to <18 months, *MYCN* non-amplified, and any of the following unfavorable features: UH; diploid DNA content; and/or presence of SCAs.

^{gg} Non-myeloablative approaches have been evaluated in single-institution studies and may be appropriate in select patients (Kushner BH, et al. *Oncotarget* 2016;7:4155-4166; Kushner BH, et al. *Oncotarget* 2017;8:95293-95302). Single transplant is currently standard in Europe (Ladenstein R, et al. *Lancet Oncol* 2017;18:500-514).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

HIGH-RISK DISEASE

POST-CONSOLIDATION THERAPY^l



^l [Principles of Systemic Therapy \(NEUROB-D\)](#).

^{hh} For patients with relapsed/refractory neuroblastoma, treatment decisions are scenario-specific. Repeat tumor biomarker profiling should be performed if feasible. Consultation with a neuroblastoma specialist should be considered (see [NEUROB-D 10 of 13](#)).

ⁱⁱ I-123 MIBG scan (FDG-PET, if MIBG non-avid) recommended after first 3 cycles of post-consolidation therapy. Additional disease evaluations (anatomic imaging of primary site and/or bone marrow evaluation) if residual disease present at start of post-consolidation.

^{jj} Serial monitoring of hearing with audiograms or brainstem auditory evoked response is essential at 6-month intervals or as clinically indicated, as most patients with high-risk neuroblastoma are at a critical age for language development.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

REFERENCES

- ¹ Kamihara J, Bourdeaut F, Foulkes WD, et al. Retinoblastoma and neuroblastoma predisposition and surveillance. Clin Cancer Res 2017;23:e98-e106.
- ² Sokol E, Desai AV, Applebaum MA, et al. Age, diagnostic category, tumor grade, and mitosis-karyorrhexis index are independently prognostic in neuroblastoma: An INRG project. J Clin Oncol 2020;38:1906-1918.
- ³ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute Clinical Trials Planning Meeting. J Clin Oncol 2017;35:2580-2587.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF PATHOLOGY

Sampling

- It is imperative that pathology workflows are designed to accommodate histologic diagnosis with prognostic classification and biomarker profiling as required for treatment.
- Tissue requirements:
 - ▶ Guidance on which evaluations are indicated for each patient based on clinical criteria are outlined below, based on requirements for risk stratification for treatment selection. Enrollment in clinical trials may require additional testing.
 - ▶ Typical tissue considerations:
 - ◊ For primary diagnosis (tumor or metastasis):
 - Surgical resection if clinically indicated
 - Incisional biopsy: >1 cm³
 - Tissue cores: at least 10 cores, ideally 20–30 mm in length obtained using a 16-gauge needle
 - ◊ Bilateral bone marrow biopsies and clot sections alone may not be sufficient to assess characteristics relevant to INPC¹
- These tissue quantity requirements can be challenging to accommodate. In some cases, open rather than minimally invasive procedures may be required to obtain adequate specimens for all required testing.²
- Advanced knowledge of how much tissue is required and the condition of the tissue (formalin-fixed paraffin-embedded [FFPE], fresh, frozen, touch preparations, etc.) will increase the likelihood that required testing can be completed.
- A coordinated diagnostic sampling effort including pathologists, oncologists, surgeons, and radiologists is recommended.^{3,4}

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

NEUROB-A
1 OF 4



PRINCIPLES OF PATHOLOGY

Histologic Classification

- Diagnosis should be determined according to the International Collaboration on Cancer Reporting (ICCR) and INPC.^{5,6,7}
- Determination of INPC should be done using the diagnostic sample obtained prior to the initiation of therapy.
- The histologic diagnosis of neuroblastoma is typically made based on hematoxylin and eosin (H&E) staining. In the setting of small samples, unusual locations, or undifferentiated subtype, immunohistochemical (IHC) stains are helpful.
 - ▶ Neuronal markers: Chromogranin, synaptophysin
 - ▶ Neural crest markers: PHOX2B (strongly recommended), tyrosine hydroxylase
- The INPC classifies peripheral neuroblastic tumors into two prognostic groups (FH and UH),⁵⁻⁸ based on:
 - ◇ Tumor category based on Schwannian stroma development and subtyping based on degree of neuroblastic differentiation:
 - Neuroblastoma (Schwannian stroma-poor)
 - Undifferentiated subtype: This diagnosis relies heavily on ancillary tests, such as IHC and/or molecular analysis
 - Poorly differentiated subtype
 - Differentiating subtype
 - Ganglioneuroblastoma, nodular (composite, Schwannian stroma-rich/stroma-dominant and stroma-poor)
 - Grade of neuroblastic differentiation (undifferentiated subtype, poorly differentiated subtype, or differentiating subtype) is determined for the neuroblastoma component in this category.
 - Ganglioneuroblastoma, intermixed (Schwannian stroma-rich)
 - Ganglioneuroma (Schwannian stroma-dominant)
 - Note that the diagnosis of ganglioneuroblastoma, intermixed (Schwannian stroma-rich) or ganglioneuroma (Schwannian stroma-dominant) from a biopsy sample (rather than resection sample) should include a comment: “Favorable histology based on review of limited material,” indicating the possibility of sampling issue resulting in missing a neuroblastoma nodule.
 - ◇ Mitosis and karyorrhexis index (MKI):
 - Assessed by estimating mitotic and karyorrhectic nuclei per 5000 neuroblastic cells.
 - Based on the MKI, all tumors in the neuroblastoma (Schwannian stroma-poor) category and neuroblastic components of ganglioneuroblastoma, nodular (composite, Schwannian stroma-rich/stroma-dominant) category are further classified into 1 of 3 classes:
 - Low MKI: <100/5000 cells
 - Intermediate MKI: 100–200/5000 cells
 - High MKI: ≥200/5000 cells
 - ◇ Age at diagnosis

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

NEUROB-A
2 OF 4



PRINCIPLES OF PATHOLOGY

Biomarker Testing Considerations:

• Prognosis:

- ▶ Risk stratification for primary treatment depends heavily on tumor biomarker profiling ([Principles of Risk Classification \[NEUROB-C\]](#)).
- ▶ Amplification of *MYCN* is associated with aggressive disease.
 - ◊ Due to the strong prognostic association, it should be assessed on all neuroblastomas and neuroblastomatous nodules of ganglioneuroblastoma nodular tumors.
 - ◊ Can be assessed by fluorescence in situ hybridization (FISH) (using a reference probe), by microarray, or by next-generation sequencing (NGS) (see “assay selection” below)
 - ◊ Status by FISH is defined as follows: not amplified (2 copies per cell), gain (>2–8 copies per cell or <4-fold increase), or amplified (>8 copies per cell or ≥4-fold increase, can be >30-fold increase⁹)
- ▶ SCAs may be associated with aggressive disease, depending on other factors^{10,11}:
 - ◊ The most extensively studied segmental chromosome aberrations include 1p, 11q, 17q, 3p, 4p, 1q, and 2p.
 - ◊ Can be assessed by microarray or by NGS (see “Assay selection” bullet)
- ▶ Ploidy evaluates the amount of DNA within tumor cells.
 - ◊ A DI of 1 may be associated with aggressive disease, depending on other factors including age.
 - ◊ Can be assessed by flow cytometry or estimated by NGS
- Identification of targeted therapy opportunities:
 - ▶ Biomarker-directed therapies are emerging in pediatric solid tumors, including neuroblastoma.^{12,13}
 - ▶ Amplification and sequence variants in *ALK* predict response to targeted agents in patients with relapsed/refractory disease.¹⁴⁻¹⁷
 - ▶ As research and drug discovery continue to evolve, additional testing for therapeutic biomarkers may be indicated.

• Assay selection:

▶ NGS:

- ◊ Has become widely available and is increasingly feasible with FFPE tissue
- ◊ Is recommended to simultaneously evaluate for *MYCN* amplification, SCAs, and *ALK* alterations if the approach:
 - Provides robust assessment of copy number status
 - Provides coverage of the relevant regions of *ALK* and other neuroblastoma-associated genes
- ▶ Assessment of the status of standard prognostic biomarkers can also be done using FISH and chromosomal microarray, but these will not identify sequence variants in *ALK* and other neuroblastoma-associated genes.
- Incidental identification of germline alterations:
 - ▶ Whether using single-gene or multigene panel testing (MGPT) approaches, genomic alterations that may be associated with cancer predisposition or other germline inherited conditions will be identified.^{18,19}
 - ▶ Dedicated germline testing may be indicated based on the family history, clinical presentation, or the presence of alterations identified by tumor biomarker profiling.
 - ▶ Involvement of genetic counselors and molecular laboratory professionals with experience in germline genetics is helpful in resolving questions of etiology of variants (germline vs. somatic) and in supporting the patient and family in navigating this complex issue.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



PRINCIPLES OF PATHOLOGY REFERENCES

- ¹ Burchill SA, Beiske K, Shimada H, et al. Recommendations for the standardization of bone marrow disease assessment and reporting in children with neuroblastoma on behalf of the International Neuroblastoma Response Criteria Bone Marrow Working Group. *Cancer* 2017;123:1095-1105.
- ² Oh C, Youn JK, Han JW, et al. Abdominal tumors in children: Comparison between minimally invasive surgery and traditional open surgery. *Medicine (Baltimore)* 2016;95:e5181.
- ³ Pinches RS, Clinton CM, Ward A, et al. Making the most of small samples: Optimization of tissue allocation of pediatric solid tumors for clinical and research use. *Pediatr Blood Cancer* 2020;67:e28326.
- ⁴ Fisch AS, Church AJ. Special considerations in the molecular diagnostics of pediatric neoplasms. *Clin Lab Med* 2022;42:349-365.
- ⁵ Shimada H, Ambros IM, Dehner LP, et al. Terminology and morphologic criteria of neuroblastic tumors: recommendations by the International Neuroblastoma Pathology Committee. *Cancer* 1999;86:349-363.
- ⁶ Shimada H, Ambros IM, Dehner LP, et al. The International Neuroblastoma Pathology Classification (the Shimada system). *Cancer* 1999;86:364-372.
- ⁷ Jarzembowski JA, Webster F, Beiske K, et al. Data set for reporting of peripheral neuroblastic tumours: recommendations from the international collaboration on cancer reporting (ICCR). *Histopathology* 2025;87:343-356.
- ⁸ Peuchmaur M, d'Amore ES, Joshi VV, et al. Revision of the International Neuroblastoma Pathology Classification: Confirmation of favorable and unfavorable prognostic subsets in ganglioneuroblastoma, nodular. *Cancer* 2003;98:2274-2281.
- ⁹ Ambros PF, Ambros IM, Brodeur GM, et al. International consensus for neuroblastoma molecular diagnostics: report from the International Neuroblastoma Risk Group (INRG) Biology Committee. *Br J Cancer* 2009;100:1471-1482.
- ¹⁰ Irwin MS, Naranjo A, Zhang FF, et al. Revised Neuroblastoma Risk Classification System: A report from the Children's Oncology Group. *J Clin Oncol* 2021;39:3229-3241.
- ¹¹ Ambros IM, Tonini GP, Potschger U, et al. Age dependency of the prognostic impact of tumor genomics in localized resectable MYCN-nonamplified neuroblastomas. Report from the SIOPEN Biology Group on the LNESG Trials and a COG Validation Group. *J Clin Oncol* 2020;38:3685-3697.
- ¹² Church AJ, Corson LB, Kao PC, et al. Molecular profiling identifies targeted therapy opportunities in pediatric solid cancer. *Nat Med* 2022;28:1581-1589.
- ¹³ Zafar A, Wang W, Liu G, et al. Molecular targeting therapies for neuroblastoma: Progress and challenges. *Med Res Rev* 2021;41:961-1021.
- ¹⁴ Bresler SC, Weiser DA, Huwe PJ, et al. ALK mutations confer differential oncogenic activation and sensitivity to ALK inhibition therapy in neuroblastoma. *Cancer Cell* 2014;26:682-694.
- ¹⁵ Bellini A, Potschger U, Bernard V, et al. Frequency and prognostic impact of ALK amplifications and mutations in the European Neuroblastoma Study Group (SIOPEN) High-Risk Neuroblastoma Trial (HR-NBL1). *J Clin Oncol* 2021;39:3377-3390.
- ¹⁶ Qiu B, Matthay KK. Advancing therapy for neuroblastoma. *Nat Rev Clin Oncol* 2022;19:515-533.
- ¹⁷ Berlak M, Tucker E, Dorel M, et al. Mutations in ALK signaling pathways conferring resistance to ALK inhibitor treatment lead to collateral vulnerabilities in neuroblastoma cells. *Mol Cancer* 2022;21:126.
- ¹⁸ Schiend J, Church AJ, Corson LB, et al. Germline sequencing improves tumor-only sequencing interpretation in a precision genomic study of patients with pediatric solid tumor. *JCO Precis Oncol* 2021;5.
- ¹⁹ Parsons DW, Roy A, Yang Y, et al. Diagnostic yield of clinical tumor and germline whole-exome sequencing for children with solid tumors. *JAMA Oncol* 2016;2:616-624.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF IMAGING

General Principles

- Neuroblastoma, ganglioneuroblastoma, and ganglioneuroma are tumors of the sympathetic nervous system that arise from the embryonic neural crest and are referred to collectively as neuroblastic tumors, all with similar imaging appearance.^{1,2}
- A diagnosis of neuroblastic tumor is usually suspected based on the patient's age and the appearance of the tumor on imaging.¹
- The combination of imaging studies to be used during the initial evaluation of a child with a suspected neuroblastic tumor is dependent upon the symptoms that are present and the suspected sites of disease.³
- Local extension of neuroblastoma mainly consists of vascular encasement, infiltration of adjacent soft tissues and organs (mainly the kidneys and liver), and infiltration of the foramina and epidural space of the spinal canal when the primary tumor arises from a paraspinal sympathetic chain.
- Approximately 50% of patients present with localized or regional disease, and approximately 35% of patients have regional lymph node spread at time of diagnosis.⁴
- The International Neuroblastoma Risk Group (INRG) Task Force developed the INRG Staging System (INRGSS) and the INRG Risk Classification System for neuroblastoma, which have profound implications for imaging and an increased role for radiology.¹
- While soft-tissue tumor volume was previously used as measurement of response in the prior International Neuroblastoma Response Criteria (INRC), the revised INRC uses RECIST criteria for measurement of soft-tissue disease, using the single longest dimension.^{5,6}

Goals of Imaging

- Identify features consistent with a neuroblastic tumor.⁷
- Assess for presence of IDRFs to classify tumors as L1 or L2 disease by the INRGSS.^{a,b,2}
- Help estimate potential surgical risk associated with local tumor excision.
- Assess for presence and degree of regional and distant metastatic disease.
- Facilitate post-treatment response assessment and disease surveillance.

Ultrasound

- When an abdominal or pelvic tumor is suspected in a child, ultrasonography (US) is often the first imaging examination performed because it is widely available and noninvasive.
- Although US is not associated with exposure to radiation, interobserver reproducibility is low and there is limited assessment of highly calcified tumors because of acoustic shadowing.
- While US can be helpful as an initial screening imaging study, magnetic resonance or CT images should be obtained at the time of diagnosis for accurate staging and treatment planning.

CT/MRI

- Both MRI and CT are routinely used for imaging of soft-tissue disease, depending on availability and institutional expertise.
- CT is rapidly performed and widely available, with superior detection of calcifications. It allows rapid acquisition without motion artifacts, and reduces the need for sedation.
- MRI is superior for evaluating spinal involvement and does not involve use of ionizing radiation.
- Both CT and MRI may be performed at the initial evaluation, especially if the tumor is paraspinal.
- Both modalities demonstrate typical features of primary tumors and metastasis to the liver, lymph nodes, bone, and skin.^{7,8}

^a [INRG Staging \(ST-1\)](#).

^b [Descriptions of IDRFs \(NEUROB-B 3 of 4\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



PRINCIPLES OF IMAGING

Nuclear Imaging:

- I-123 (I-123 MIBG) is used for imaging of neuroblastic tumors, especially metastatic sites of disease.⁹
 - ▶ Obtain prior to resection of the primary tumor, if possible
 - ▶ Not required in patients <6 months of age with a localized adrenal tumor in longest diameter ≤3.1 cm if solid or 5 cm if >25% cystic component.
 - ▶ Baseline I-123 MIBG imaging may be delayed in patients <2 months of age or those weighing <2.5 kg with suspected primary tumors within the adrenal glands.
 - ▶ Novel radiotracers are in development and may serve a role in select patients, but there are insufficient data currently available to incorporate routinely for diagnostic imaging and response assessment.
- As an analog of norepinephrine, MIBG is taken up by norepinephrine transporters. Uptake is demonstrated in up to 90% of neuroblastoma tumors.
- The high specificity and high sensitivity with improved accuracy for localization make MIBG imaging the test of choice for identification of metastatic disease.¹⁰ I-123 MIBG is the agent of choice for diagnostic imaging. Iodine-131 (I-131) MIBG may be useful in resource-limited settings.⁹
- In addition to MIBG whole-body planar scintigraphy (vertex of head through feet), single-photon emission CT (SPECT) or SPECT/CT should be performed at sites of known or suspected disease, where available, due to improved sensitivity and anatomic localization of disease sites.¹⁰
- Thyroid protection is required as per institutional guidance.
- Interpretation of MIBG tumor uptake is performed by means of semiquantitative scoring of body segments. The modified Curie score¹¹ and the International Society of Pediatric Oncology Europe Neuroblastoma (SIOPEN) score¹² are the two most commonly used systems. In North America, the modified Curie score is used.
- When neuroblastoma tumors are not MIBG avid, or when MIBG imaging and anatomic imaging do not correlate, FDG-PET/CT or PET/MRI are useful alternative or supplemental diagnostic tools.¹³

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

PRINCIPLES OF IMAGING DESCRIPTIONS OF IDRFs²

Anatomic Region	Description
Multiple body compartments	Ipsilateral tumor extension within two body compartments (ie, neck and chest, chest and abdomen, or abdomen and pelvis)
Neck	• Tumor encasing carotid artery, vertebral artery, and/or internal jugular vein • Tumor extending to skull base • Tumor compressing trachea
Cervicothoracic junction	• Tumor encasing brachial plexus roots • Tumor encasing subclavian vessels, vertebral artery, and/or carotid artery • Tumor compressing trachea
Thorax	• Tumor encasing aorta and/or major branches • Tumor compressing trachea and/or principal bronchi • Lower mediastinal tumor infiltrating costovertebral junction between T9 and T12 vertebral levels
Thoracoabdominal junction	Tumor encasing aorta and/or vena cava
Abdomen and pelvis	• Tumor infiltrating porta hepatis and/or hepatoduodenal ligament • Tumor encasing branches of superior mesenteric artery at mesenteric root • Tumor encasing origin of celiac axis and/or origin of superior mesenteric artery • Tumor invading one or both renal pedicles • Tumor encasing aorta and/or vena cava • Tumor encasing iliac vessels • Pelvic tumor crossing sciatic notch
Intraspinal tumor extension	Intraspinal tumor extension (whatever the location) provided that more than one-third of spinal canal in axial plane is invaded, the perimedullary leptomeningeal spaces are not visible, or the spinal cord signal intensity is abnormal
Infiltration of adjacent organs and structures	Pericardium, diaphragm, kidney, liver, duodenopancreatic block, and mesentery

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



PRINCIPLES OF IMAGING REFERENCES

- ¹ Lonergan GJ, Schwab CM, Suarez ES, Carlson CL. Neuroblastoma, ganglioneuroblastoma, and ganglioneuroma: radiologic-pathologic correlation. *RadioGraphics* 2002;22:911-934.
- ² Brisse HJ, McCarville MB, Granata C, et al. Guidelines for imaging and staging of neuroblastic tumors: Consensus report from the International Neuroblastoma Risk Group Project. *Radiology* 2011;261:243-257.
- ³ Sharp SE, Trout AT, Weiss BD, Gelfand MJ. MIBG in Neuroblastoma diagnostic imaging and therapy. *Radiographics* 2016;36:258-278.
- ⁴ Park JR, Eggert A, Caron H. Neuroblastoma: Biology, prognosis, and treatment. *Hematology Oncol Clin North Am* 2010;24:65-86.
- ⁵ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute clinical trials planning meeting. *J Clin Oncol* 2017;35:2580-2587.
- ⁶ Bagatell R, McHugh K, Naranjo A, et al. Assessment of primary site response in children with high-risk neuroblastoma: An international multicenter study. *J Clin Oncol* 2016;34:740-746.
- ⁷ Papaioannou G, McHugh K. Neuroblastoma in childhood: review and radiological findings. *Cancer Imaging* 2005;5:116-127.
- ⁸ Kembhavi SA, Shah S, Rangarajan V, et al. Imaging in neuroblastoma: an update. *Indian J Radiol Imaging* 2015;25:129-136.
- ⁹ Swift CC, Eklund MJ, Kravaka J M, et al. Updates in diagnosis, management, and treatment of neuroblastoma. *Radiographics* 2018;38:566-580.
- ¹⁰ Lai HA, Sharp SE, Bhatia A, et al. Imaging of pediatric neuroblastoma: A COG Diagnostic Imaging Committee/SPR Oncology Committee White Paper. *Pediatr Blood Cancer* 2023;70 Supple 4:e29974.
- ¹¹ Yanik GA, Parisi MT, Naranjo A, et al. Validation of postinduction Curie scores in high-risk neuroblastoma: A Children's Oncology Group and SIOPEN group report on SIOPEN/HR-NBL1. *J Nucl Med* 2018;59:502-508.
- ¹² Ladenstein R, Lambert B, Pütschger U, et al. Validation of the mIBG skeletal SIOPEN scoring method in two independent high-risk neuroblastoma populations: the SIOPEN/HR-NBL1 and COG-A3973 trials. *Eur J Nucl Med Mol Imaging* 2018;45:292-305.
- ¹³ Sharp SE, Shulkin BL, Gelfand MJ, et al. 123I-MIBG scintigraphy and 18F-FDG PET in neuroblastoma. *J Nucl Med* 2009;50:1237-1243.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RISK CLASSIFICATION

Risk classification is assigned¹⁻⁵ based on outcome data generated over recent decades. Event-free survival (EFS) and overall survival (OS) outcomes are influenced by key prognostic risk factors including:

- Age at diagnosis
- INRG Stage ([ST-1](#))
- Tumor *MYCN* status (presence or absence of *MYCN* amplification)
- Histopathology (FH or UH based on International Neuroblastoma Pathology Classification [INPC])
- Presence or absence of SCAs
- DI (diploid or hyperdiploid)

It should be noted that treatment substantially impacts outcome, and outcomes achieved often reflect the impact of different types of therapy within a given risk group. However, risk classification must occur in practice prior to the start of therapy. The risk assignments that follow therefore reflect pre-treatment decision-making, taking into account treatment administered to prior patients with neuroblastoma over time.

Risk	5-year EFS ¹	5-year OS ¹
Low	90.7% ± 1.0%	97.9% ± 0.5%
Intermediate	85.1% ± 1.4%	95.8% ± 0.8%
High	51.2% ± 1.4%	62.5% ± 1.3%

¹ Irwin MS, Naranjo A, Zhang FF, et al. Revised Neuroblastoma Risk Classification System: A report from the Children's Oncology Group. J Clin Oncol 2021;39:3229-3241.

² Sokol E, Desai AV, Applebaum MA, et al. Age, diagnostic category, tumor grade, and mitosis-karyorrhexis index are independently prognostic in neuroblastoma: An INRG Project. J Clin Oncol 2020;38:1906-1918.

³ Shimada H, Ambros IM, Dehner LP, et al. The International Neuroblastoma Pathology Classification (the Shimada system). Cancer 1999;86:364-372.

⁴ Janoueix-Lerosey I, Schleiermacher G, Michels E, et al. Overall genomic pattern is a predictor of outcome in neuroblastoma. J Clin Oncol 2009;27:1026-1033.

⁵ Schleiermacher G, Mosseri V, London WB, et al. Segmental chromosomal alterations have prognostic impact in neuroblastoma: a report from the INRG project. Br Cancer 2012;107:1418-1422.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY LOW- AND INTERMEDIATE-RISK DISEASE

Patients with low- and intermediate-risk (ie, non-high risk) neuroblastoma have excellent survival. Recent clinical trials have focused on reduction of therapy for patients with favorable biology and have been successful in maintaining excellent outcomes with these strategies.¹⁻³

Over the past 2 decades, the staging system, risk classification system, and response criteria definitions for neuroblastoma have evolved. It is important to note that published results from clinical trials for patients with non-high-risk disease utilized a legacy staging system (International Neuroblastoma Staging System [INSS]) and older response criteria⁴ or protocol-specific response criteria.³ These NCCN Guidelines are based on published data from clinical trials of patients with non-high-risk disease, but also bridge the transition to currently used staging, classification, and response criteria systems where applicable.⁵⁻⁷

In all patients with neuroblastoma except those with completely resected INRG Stage L1 disease, detection of *MYCN* amplification in tumor cells leads to assignment to the high-risk group rather than low- or intermediate-risk groups⁷ ([NEUROB-2](#)). For symptomatic infants with INRG stage MS disease who are too ill to undergo biopsy at the time of initial presentation, therapy is started with a presumptive assignment to the intermediate-risk group. However, if a biopsy undertaken when the patient is clinically stable identifies tumor cells with *MYCN* amplification, the patient would then be reassigned to the high-risk group.

Treatment approaches for low risk INRG L1 tumors involve surgical resection, except in patients <6 months of age with adrenal tumor with maximal diameter ≤3.1 cm if solid or 5 cm if at least 25% cystic component. In these infants, observation is recommended. In patients with INRG MS disease who are asymptomatic and have tumors with favorable biology, observation is also the preferred approach.

Intermediate-risk treatment involves a combination of moderate intensity multi-agent chemotherapy and surgical resection. The COG study ANBL0531 aimed to decrease number of chemotherapy cycles given to patients with more favorable tumor biology, by allowing for a larger residual primary tumor following chemotherapy ± surgery, compared to treatment endpoints utilized on legacy clinical trials. Patients were assigned to receive a minimum number of cycles based on age, stage, and tumor biologic features. Using this strategy, after completion of the assigned number of cycles, disease response was assessed and decisions regarding additional systemic therapy, surgery, or surveillance were made. In the ANBL0531 clinical trial, protocol-specific criteria for disease response were used, and primary tumor response was assessed by tumor volume reduction. The goal for groups of patients with localized, favorable biology tumors was to achieve at least 50% reduction in the primary tumor volume, while patients with localized disease with less favorable tumor biologic features were to continue therapy until a goal of 90% reduction in primary tumor volume (very good PR [VGPR]) was achieved. The goal treatment endpoint for metastatic disease response for patients with the equivalent of INRG MS disease and those with intermediate risk INRG Stage M disease were also protocol-specific, and not identical to the legacy INRC.⁴

Since the completion of ANBL0531, the neuroblastoma response criteria have been updated and no longer include VGPR as a response category. Additionally, in the current system of response assessment, tumors are measured with a single dimension as per Response Evaluation Criteria in Solid Tumors (RECIST), rather than by tumor volume ([NEUROB-G 1 of 4](#)). Until additional data become available regarding the use of the revised response criteria in assessment of disease in patients with non-high-risk neuroblastoma, the committee supports utilizing either volume or 1-dimensional assessments of primary tumor response in this group of patients.

For patients who have not achieved the targeted tumor reduction goal with the initial course of chemotherapy prescribed, multidisciplinary discussion should take place. For some patients, surgical resection may be appropriate to reach the targeted treatment endpoint. If surgery cannot be performed safely to achieve the proposed degree of tumor reduction, additional chemotherapy may be given with re-evaluation after every 2 cycles. At these timepoints, the potential risks and benefits of additional chemotherapy or surgery can be discussed by the multidisciplinary treatment team. In some circumstances it may be reasonable to consider biopsy of the residual mass to assess for histologic differentiation, which may support observation of a tumor that does not shrink sufficiently with chemotherapy and for which a surgical debulking is considered unsafe. In COG ANBL0531, cyclophosphamide and topotecan were used as the additional treatment for patients who did not achieve the target response with 8 cycles of the intermediate-risk therapy regimen outlined on [NEUROB-4](#).

SIOPEN treatment regimens can also be considered, as similar outcomes are achieved with the SIOPEN and COG strategies.

[References](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY
Intermediate-Risk Chemotherapy^a

COG ANBL0531 - Carboplatin/Cyclophosphamide/Doxorubicin/Etoposide^{3,8}		Dosing^b
Cycle 1	• Carboplatin/Etoposide	Chemotherapy dosing for young children is evolving toward dose banding.⁹ • Carboplatin 560 mg/m² >12 kg or 18.6 mg/kg ≤12 kg^c D1 • Etoposide 120 mg/m² >12 kg or 4 mg/kg ≤12 kg^c on D1–3 • Cyclophosphamide 1000 mg/m² >12 kg or 33.3 mg/kg ≤12 kg D1, Doxorubicin 30 mg/m² >12 kg or 1 mg/kg ≤12 kg^c D1
Cycle 2	• Carboplatin/Cyclophosphamide/Doxorubicin	
Cycle 3	• Cyclophosphamide/Etoposide	
Cycle 4	• Carboplatin/Doxorubicin/Etoposide	
Cycle 5	• Cyclophosphamide/Etoposide	
Cycle 6	• Carboplatin/Cyclophosphamide/Doxorubicin	
Cycle 7	• Carboplatin/Etoposide	
Cycle 8	• Cyclophosphamide/Doxorubicin	

SIOPEN - Carboplatin/Etoposide alternating with Cyclophosphamide/Doxorubicin/Vincristine¹⁰	Dosing^b
• Carboplatin/Etoposide	• Carboplatin 200 mg/m² or 6.6 mg/kg and Etoposide 150 mg/m² or 5.0 mg/kg on D1–3 • Cyclophosphamide 300 mg/m² or 10 mg/kg D1–5, Doxorubicin 30 mg/m² or 1 mg/kg D4–5, Vincristine D1 and D5 age-, weight-, and body surface area (BSA)-based dosing
• CADO (Cyclophosphamide/Doxorubicin/Vincristine)	
• Number of cycles vary for different subgroups	

- Subsequent-line chemotherapy may be considered if adequate response on imaging has not been achieved, and undifferentiated viable tumor remains.³

[See Evidence Blocks on EB-1](#)

^a Similar outcomes have been seen in Children's Oncology Group (COG) and SIOPEN studies.

^b Mesna and dexrazoxane may be used as clinically indicated.

^c Or per infant dosing strategies.⁹

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

References



PRINCIPLES OF SYSTEMIC THERAPY HIGH-RISK DISEASE

- Patients with newly diagnosed high-risk neuroblastoma have an estimated 5-year EFS rate of 51% from a large analysis reflecting 10 years of data from the Children's Oncology Group (COG).⁷ These outcomes have improved over time as a result of increasingly intensive multimodal therapies divided into induction, consolidation, and post-consolidation phases. Much of the data summarized below is derived from national phase 3 trials and the Panel encourages participation in open clinical trials when available.

Induction Therapy

- The goal of initial induction therapy is to decrease disease burden and to achieve the best possible response prior to subsequent phases of therapy. This goal is achieved through a combination of multiagent chemotherapy and surgical resection of the primary tumor. In addition, autologous peripheral blood stem cells are collected during induction and aliquoted in a way that facilitates subsequent therapy.¹¹
- There are no contemporary data comparing North American induction regimens that have been evaluated in prospective randomized trials. Instead, a number of induction combinations have evolved over the last several decades, mainly based upon cisplatin- and alkylator-intensive regimens developed at Memorial Sloan-Kettering Cancer Center (MSKCC). Several different regimens have been used in North American cooperative group pilot and phase 3 trials, each enrolling >100 patients over the last 2 decades.
- These regimens yield broadly similar end-induction response rates, with approximately 80% of patients having a PR or better and approximately 9% of patients progressing despite these intensive regimens.¹² Given the lack of prospective comparative efficacy data and similar response rates, toxicity considerations have driven the evolution of induction regimens to reduce exposure to nephrotoxic and cardiotoxic agents. The combination of topotecan and cyclophosphamide given as cycles 1 and 2 of induction was studied initially as part of a pilot trial that demonstrated the feasibility and acceptable end-induction response rate (84%) of this approach.¹³ This regimen was adopted for use in ANBL0532, demonstrating that 39.1% of patients had a PR or better after the first 2 cycles with topotecan and cyclophosphamide.¹⁴ A subsequent trial, ANBL12P1, empirically reduced induction to 5 cycles, with an 80% end-induction response rate.¹⁵ Likewise, data from MSKCC demonstrated comparable end-induction response rates with 5 versus 7 induction cycles.¹⁶ Therefore, a 5-cycle induction was included as part of the standard arm of the COG phase 3 trial, ANBL1531. The ANBL1531 induction regimen is similar to the published ANBL12P1 regimen, with minor adjustments to dosing in order to align with updated COG chemotherapy standards. Given extensive contemporary experience with a 5-cycle induction regimen, the Panel recommends either ANBL12P1 or ANBL1531 induction as preferred regimens, with a 6-cycle regimen from ANBL0532 as an additional acceptable regimen. However, the Panel acknowledges the lack of comparative data, and notes that other published chemotherapy regimens that achieve a similar end-induction response rate could be considered as reasonable alternatives for individual patients.

[See Evidence Blocks on EB-2](#)

References

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY HIGH-RISK DISEASE

- A number of strategies to improve induction are under investigation at this time, including early addition of an ALK inhibitor for the subset of patients whose tumors have been found to harbor ALK alterations. In addition, interventions such as the early administration of I-131 MIBG or anti-GD2 monoclonal antibody therapy are being evaluated. As robust safety and efficacy data are not yet available to support these approaches, the Panel does not recommend adoption of these approaches outside the context of clinical trials at this time.
- Surgical resection of the primary tumor and associated locoregional adenopathy is another important goal of induction therapy ([Principles of Surgery \[NEUROB-E\]](#)). Given the aggressive nature of these tumors, upfront resection is rarely feasible and the Panel recommends surgical resection after several cycles of initial cytoreductive chemotherapy. Even after initial chemotherapy, resection with negative margins remains rarely feasible and is not the recommended surgical goal. Instead, two large analyses have demonstrated improved EFS and lower local relapse/progression rates in patients who had >90% resection (North American experience) or a complete macroscopic resection (European experience).^{17,18} The Panel therefore recommends this degree of resection, broadly considered a gross total resection, as the goal of primary site surgery. When vital organs, major nerves, and/or major blood vessels would be threatened or would require resection to achieve this goal, the Panel recommends subtotal resection.
- The Panel recommends full disease reassessment at the end of induction. This end-induction evaluation is a critical decision point in the treatment of patients with high-risk neuroblastoma. Prior analyses have demonstrated that patients with poor end-induction response (less than PR) have inferior outcomes compared to patients with more favorable end-induction response (PR or better).¹² However, analyses of randomized trials evaluating different consolidation strategies support a potential role for modern consolidation therapies even in patients with less than a PR to induction. For example, the CCG-3891 trial reported higher EFS for patients with less favorable end-induction response who were randomized to transplant (vs. continued chemotherapy).¹⁹ Likewise, the ANBL0532 trial (see below) reported a benefit of tandem transplant (vs. single transplant) independent of end-induction response.¹⁴
- The Panel recommends proceeding to consolidation therapy for patients with PR or better to induction, though the Panel acknowledges that bridging therapy to improve response may be appropriate depending upon the nature of the PR.²⁰ Patients with PD during or at the end of induction have not typically been candidates to proceed with consolidation therapy, and the Panel endorses this approach. Instead, the Panel recommends non-myeloablative therapies for these patients, including a chemoimmunotherapy regimen combining anti-GD2 monoclonal antibody with chemotherapy^{21,22} or participation in clinical trials for patients with first relapse. Patients with end-induction MR or SD require individualized decision-making. For patients with end-induction MR or SD not proceeding to consolidation therapy, the Panel recommends a chemoimmunotherapy regimen combining anti-GD2 monoclonal antibody with chemotherapy or participation in clinical trials for patients with refractory disease. Retrospective data suggest that proceeding to consolidation therapy may be appropriate for patients with an inadequate response to standard induction therapy who respond to alternative “bridging” therapies.²⁰ Specifically, patients with a CR to bridging therapy in one series who proceeded to consolidation had favorable outcomes.

References

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY HIGH-RISK DISEASE

Consolidation Therapy

- A standard consolidation phase includes both high-dose chemotherapy with autologous stem cell rescue and consolidative radiotherapy to the primary site. In North America, it is also considered standard to administer radiotherapy to sites of residual metastatic disease remaining at the end-induction disease evaluation.
- High-dose chemotherapy with autologous stem cell rescue has been a hallmark of high-risk neuroblastoma therapy since a series of randomized trials demonstrated improved outcomes with this approach compared with continued conventional chemotherapy.²³ The Panel acknowledges that these randomized trials were conducted in a treatment era that preceded routine use of anti-GD2-directed immunotherapy and that additional work is needed to understand if subgroups of patients might benefit from consolidative approaches that do not rely on high-dose chemotherapy with autologous stem cell rescue. For example, a limited institution retrospective experience suggests that similar OS rates may be achieved with or without high-dose chemotherapy among patients with greater than a PR in the era of anti-GD2 immunotherapy.²⁴
- For patients who are candidates for consolidation therapy, the Panel recommends tandem transplantation with two consecutive rounds of high-dose chemotherapy with autologous stem cell rescue for the majority of patients with high-risk disease (category 1 recommendation). This recommendation is based upon the COG ANBL0532 randomized phase 3 trial.¹⁴ Patients without PD after a 6-cycle induction were eligible for randomization to a single transplant with full-dose CEM (carboplatin/etoposide/melphalan) or to tandem transplant with thiotepa/cyclophosphamide followed 6–10 weeks later with dose-reduced CEM ([NEUROB-10](#)). Patients randomized to the tandem transplant arm had significantly improved EFS (3-year EFS 61.6% vs. 48.4% for single transplant).
- There are two less common subgroups of patients with high-risk disease for whom a single round of high-dose chemotherapy with autologous stem cell rescue may be appropriate: 1) patients with stage L2, ≥18 months at diagnosis, UH, AND *MYCN* non-amplified disease; and 2) patients with stage M, 12 to <18 months at diagnosis, *MYCN* non-amplified, with any of the following other unfavorable features: UH; diploid DNA content; and/or presence of SCAs. Patients in these two groups have historically had more favorable outcomes compared to patients with high-risk disease due to *MYCN* amplification or patients with high-risk disease due to age ≥18 months at diagnosis with stage M disease. For example, in a large series from the COG, patients in these two groups had 5-year EFS rates of approximately 75%–80%.⁷ Patients in these two more favorable subgroups were non-randomly assigned to single transplant with full-dose CEM in ANBL0532 and the Panel endorses this approach. In other contexts, these patients are treated differently.^{25–29}
- Conditioning with BuMel (busulfan and melphalan) is a preferred conditioning regimen in Europe based upon results of a randomized phase 3 trial that showed superior EFS with BuMel compared to CEM following the European rapid COJEC induction regimen.³⁰ In addition, the BuMel regimen was associated with lower rates of most adverse events, though the risk of sinusoidal obstruction syndrome was higher. The COG conducted a pilot trial, ANBL12P1, that demonstrated the feasibility of this approach,¹⁵ but the role of BuMel in the context of North American therapy is not currently defined. Single transplant with BuMel may be an appropriate regimen for patients with a contraindication to tandem transplant or for patients in subgroups for which single transplant is recommended.
- RT to the primary tumor is typically administered upon recovery from high-dose chemotherapy with stem cell rescue ([Principles of Radiation Therapy \[NEUROB-H\]](#)). Neuroblastoma is a radiosensitive tumor and a commonly used dose of 21.6 Gy is recommended. Recent national trials have attempted to improve local control by augmenting radiotherapy. In COG trial A3973, a subset of patients had primary site radiotherapy fields extended to include uninvolved draining nodal stations. These patients had similar local relapse/progression rates compared to patients treated without extending the radiotherapy field and this approach is not recommended.³¹ In COG trial ANBL0532, patients with gross residual tumor following primary site resection received a boost of 14.4 Gy to gross residual tumor.³² This augmented dose did not improve local relapse/progression or EFS rates compared to the historic controls. Based upon these data, the Panel does not recommend either strategy.
- The Panel recommends radiation to sites of residual metastatic disease present at the end-induction disease evaluation, recognizing that not all sites may be feasibly targeted by external beam radiotherapy. While repeat imaging following recovery from transplant and prior to radiation is not routinely performed, it may aid in decision-making in select clinical scenarios. The recommendation to radiate residual metastatic sites of disease is based upon limited institution data supporting a benefit from this approach,^{33,34} though this approach has not been adopted internationally.

References

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY

HIGH-RISK DISEASE

Post-Consolidation Therapy

- Historically, post-consolidation therapy consisted of 6 cycles of isotretinoin administered as a differentiating agent. This approach was adopted based upon the landmark CCG-3891 trial that demonstrated improved outcomes in patients randomized to isotretinoin compared to no further therapy.³⁵ Subsequently, the ANBL0032 trial demonstrated a significant improvement in EFS for patients randomized to receive the anti-GD2 monoclonal antibody dinutuximab + cytokines (sargramostim in cycles 1, 3, and 5; interleukin-2 in cycles 2 and 4) + isotretinoin (2-year EFS from randomization at start of post-consolidation of 66%) compared to patients randomized to receive isotretinoin alone (2-year EFS of 46%).³⁹ Based upon these findings, the ANBL0032 immunotherapy regimen became a standard post-consolidation regimen (category 1 recommendation for use of post-consolidation therapy with anti-GD2 antibody) ([NEUROB-D 9 of 13](#)).³⁶
- More recently, data from the SIOPEN HR-NBL1 trial have called into question the role of interleukin-2 together with anti-GD2 immunotherapy. In this trial, patients were randomized to receive anti-GD2 immunotherapy (dinutuximab beta) with or without subcutaneous interleukin-2.³⁷ Interleukin-2 did not improve outcomes, but was associated with increased toxicity. Based upon these findings, COG high-risk protocols no longer include interleukin-2 and the Panel endorses this approach.
- Other anti-GD2 antibodies may be appropriate as post-consolidation therapy. For example, dinutuximab beta given with isotretinoin but without sargramostim is a commonly used post-consolidation regimen in Europe. A non-randomized comparison showed higher EFS among patients treated with this approach compared to the historic experience with isotretinoin alone.³⁸

Continuation Therapy

- Eflornithine (2,5-diamino-2-[difluoromethyl] pentanoic acid hydrochloride hydrate; DFMO) is an inhibitor of ornithine decarboxylase, a key enzyme required for the synthesis of polyamines that regulate homeostasis and promote survival in cancer cells. This agent was studied as continuation therapy in a multicenter, single-arm, phase 2 trial (Study 3b; NCT02395666) in children with high-risk neuroblastoma that had responded to frontline therapy that included induction, consolidation, and anti-GD2 directed immunotherapy. Patients who had a PR or better following standard frontline therapy were eligible to enroll on study 3b following completion of immunotherapy, and received eflornithine 750 mg/m² ± 250 mg/m² twice daily for up to 2 years. Reported adverse events included transaminitis and hearing loss.³⁹ Data from 92 patients on Study 3b were compared with an external control arm consisting of 852 patients treated with anti-GD2 immunotherapy, cytokines, and isotretinoin on COG ANBL0032 who did not go on to receive eflornithine continuation therapy.⁴⁰ Patients on Study 3b had superior outcomes compared to the external control group [EFS hazard ratio 0.48 (95% CI: 0.27, 0.85); OS hazard ratio 0.32 (95% CI: 0.15, 0.70)]. Further analyses using propensity score matching and sensitivity analyses also demonstrated higher EFS and OS for patients on Study 3b, though the potential for residual confounding remains in the context of a non-randomized comparison. In December 2023, the FDA approved eflornithine for use in the continuation setting for patients with high-risk neuroblastoma who have achieved a PR or better following completion of anti-GD2 immunotherapy. The Panel recommends that clinicians discuss eflornithine as a continuation therapy option with patients who are in remission at the end of immunotherapy.

Disease Evaluations During Frontline Therapy for High-Risk Disease

- Following initial staging evaluations prior to the start of therapy ([ST-1](#)), patients with high-risk disease undergo anatomic imaging (CT or MRI) of the primary site prior to planned surgical resection. Full disease evaluation (anatomic imaging of the primary site, I-123 MIBG scan [or FDG-PET, if MIBG non-avid disease], and bilateral bone marrow aspirates and biopsies) is recommended at the end of induction, start of post-consolidation, and at end of therapy. Patients with >5 residual MIBG-avid sites of disease at the end of induction are encouraged to have a repeat I-123 MIBG scan after recovery from high-dose chemotherapy with stem cell rescue in order to prioritize metastatic sites that might be treated during consolidative radiotherapy. An I-123 MIBG scan (or FDG-PET, if MIBG non-avid disease) is recommended halfway through post-consolidation therapy, with anatomic imaging and bone marrow evaluations reserved for patients with residual disease identified on the disease evaluation at the start of post-consolidation.

Organ Function Evaluations During Frontline Therapy for High-Risk Disease

- Therapy for high-risk neuroblastoma is intensive and associated with both acute and long-term toxicities ([Monitoring for Late Effects \[NEUROB-II\]](#)). During treatment, these patients require frequent laboratory monitoring, including blood counts, chemistry panels, and urinalyses. Detailed evaluation of renal function (often with nuclear medicine measurements of glomerular filtration rate [GFR]) is essential before consolidation high-dose therapy. Serial monitoring of cardiac function with ECGs and echocardiograms is routine. Serial monitoring of hearing with audiograms or brainstem auditory evoked response is essential as most patients with high-risk neuroblastoma are at a critical age for language development.

Therapy for Adolescents and Adults with High-Risk Neuroblastoma

- Neuroblastoma is largely a disease of young children, though adolescents and adults may occasionally present with high-risk disease. The clinical studies that inform these guidelines, including toxicity data, therefore predominantly included patients <5 years of age at initial diagnosis. The general principles of high-risk therapy should be applied to older patients with high-risk disease, though it is acknowledged that these patients may require a more individualized approach to treatment based upon comorbid conditions and tolerance of planned therapy.

References

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

HIGH-RISK DISEASE: CHEMOTHERAPY INDUCTION REGIMENS^{d,e,f,9,14,15}

- Chemotherapy dosing for young children is evolving toward dose banding⁹

Cycle 1	• Topotecan 1.2 mg/m ² /dose Days 1–5
Cycle 2	• Cyclophosphamide 400 mg/m ² /dose Days 1–5
Cycle 3^e	• Cisplatin 50 mg/m ² /dose Days 1–4 • Etoposide 200 mg/m ² /dose Days 1–3
Cycle 4^e	• Vincristine (age-, weight-, and BSA-based dosing) Days 1–3 ⁹ • Doxorubicin 25 mg/m ² /dose Days 1–3 • Cyclophosphamide 2100 mg/m ² /dose Days 1–2
Cycle 5^e	• Cisplatin 50 mg/m ² /dose Days 1–4 • Etoposide 200 mg/m ² /dose Days 1–3
Cycle 6⁹ (Not currently used)	• Vincristine (age-, weight-, and BSA-based dosing) Days 1–3 • Doxorubicin 25 mg/m ² /dose Days 1–3 • Cyclophosphamide 2100 mg/m ² /dose Days 1–2

[See Evidence Blocks on EB-2](#)

^d All cycles are 21-day cycles. Refer to individual literature for dosing strategies for infants/patients with lower weights/BSAs.

^e COG trial ANBL1531 used this same regimen with a slightly different dosing strategy based upon updated COG chemotherapy standards. This induction regimen is another preferred regimen.

^f Dexrazoxane may be used as clinically indicated. Mesna may be used with cycles 1 and 2 as clinically indicated, but must be used with cycles 4 and 6.

⁹ Cycle 6 was used in prior historical trial ANBL0532, but is not used in ongoing COG clinical trials. Park JR, Kreissman SG, London WB, et al. Effect of tandem autologous stem cell transplant vs single transplant on event-free survival in patients with high-risk neuroblastoma: A randomized clinical trial. JAMA 2019;322:746-755.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**

References

NEUROB-D
7 OF 13



PRINCIPLES OF SYSTEMIC THERAPY

HIGH-RISK DISEASE: CHEMOIMMUNOTHERAPY REGIMENS (may be used as bridging therapy)^h

- ANBL1221 - Irinotecan/Sargramostim/Temozolomide + Dinutuximab^{21,41}
 - ▶ Temozolomide 100 mg/m²/dose PO Days 1–5
 - ▶ Irinotecan 50 mg/m²/dose IV over 90 min Days 1–5
 - ▶ Dinutuximab 17.5 mg/m²/day IV Days 2–5
 - ▶ Sargramostim 250 µg/m²/dose SQ Days 6–12
 - ▶ Regimen repeated every 21 days
- Irinotecan/Sargramostim/Temozolomide + Naxitamab-gqgk²²
 - ▶ Temozolomide 150 mg/m²/day PO Days 1–5
 - ▶ Irinotecan 50 mg/m²/day IV Days 1–5
 - ▶ Naxitamab-gqgk 2.25 mg/kg/day IV over 30–60 min on Days 2, 4, 9, and 11
 - ▶ Sargramostim 250 mcg/m²/day SQ on Days 6–10
 - ▶ Regimen repeated every 4 weeks

HIGH-RISK DISEASE: HIGH-DOSE CHEMOTHERAPY REGIMENS FOR CONSOLIDATION THERAPYⁱ

Tandem Transplant ¹⁴	Single Transplant Option 1 ¹⁴	Single Transplant Option 2 ¹⁵
Thiotepa ¹⁴ 300 mg/m ² /dose Days -7 to -5	Carboplatin ^{i,14} (weight-, BSA-, and GFR-based dosing) Days -7 to -4	Busulfan ¹⁵ (age- and weight-based dosing) Days -6 to -3 (dose adjusted based upon real-time pharmacokinetics)
Cyclophosphamide 1500 mg/m ² /dose Days -5 to -2	Etoposide 338 mg/m ² /dose Days -7 to -4	Melphalan 140 mg/m ² /dose Day -1
6–10 Weeks Later and Recovery of Organ Function	Melphalan 70 mg/m ² /dose Days -7 to -5	
Carboplatin ⁱ (age-, BSA-, and GFR-based dosing) Days -7 to -4		
Etoposide 300 mg/m ² /dose (if GFR ≥100 mL/min/1.73 m ²) Days -7 to -4		
Melphalan 60 mg/m ² /dose Days -7 to -5		

[See Evidence Blocks on EB-2](#)

^h Refer to individual protocols for dosing strategies for infants/patients with lower BSAs. COG trial ANBL1531 uses this same regimen with a slightly different dosing strategy based upon updated COG chemotherapy standards.

ⁱ Refer to literature for dosing strategies for infants/patients with lower weights/BSAs and with impaired organ function.

[References](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

HIGH-RISK DISEASE: Post-Consolidation Therapy^{j,35-38}

• Isotretinoin/Sargramostim + Dinutuximab

Cycle 1	• Sargramostim 250 µg/m²/dose SQ Days 1–14 • Dinutuximab 17.5 mg/m²/dose IV Days 4–7 • Isotretinoin 80 mg/m²/dose PO twice daily Days 11–24
Cycle 2	
Cycle 3	
Cycle 4	
Cycle 5	• Isotretinoin 80 mg/m²/dose PO twice daily Days 11–24
Cycle 6	

HIGH-RISK DISEASE: Continuation Therapy⁴⁰

For patients who are in remission, have had at least a PR to prior systemic agents, and have completed post-consolidation immunotherapy with an anti-GD2 antibody	• Consider Eflornithine (category 2B)^k ▶ BSA (m²)–based dosage (recalculate the BSA dosage every 3 months during treatment with Eflornithine) ◇ >1.5: 768 mg (four tablets) PO twice daily ◇ 0.75–≤1.5: 576 mg (three tablets) PO twice daily ◇ 0.5 to <0.75: 384 mg (two tablets) PO twice daily ◇ 0.25 to <0.5: 192 mg (one tablet) PO twice daily
---	---

[See Evidence Blocks on EB-2](#)

^j All cycles are 28-day cycles. Refer to individual protocols for dosing strategies for infants/patients with lower BSAs.

^k Serial monitoring of hearing with audiograms or brainstem auditory evoked response is essential, as most patients with high-risk neuroblastoma are at a critical age for language development.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

NEUROB-D
9 OF 13



PRINCIPLES OF SYSTEMIC THERAPY

Relapsed/Refractory Disease

- Selection of treatment for patients who experience disease progression or relapse during or after completion of high-risk neuroblastoma therapy requires an individualized approach.
- Factors that impact therapy selection include timing of recurrence, extent of disease, MIBG avidity at relapse, prior treatment administered and associated toxicities, clinical condition and organ function, as well as the molecular features of the disease at diagnosis and at relapse.
- Participation in clinical trials is strongly encouraged for this group of patients who have historically had very poor outcomes.
- While other therapies may be selected for specific populations of patients, the combination of an anti-GD2 antibody and cytotoxic chemotherapy is now widely used at the time of first relapse or first declaration of refractory disease. Dinutuximab is often used clinically in North America while dinutuximab beta is widely used in Europe and other areas of the world.
 - ▶ The combination of irinotecan, temozolomide, and GD2-directed antibody therapy has been evaluated in multiple studies and objective response rates of ~40% are among the highest seen in this patient population. It is notable that these response rates are strikingly reproducible.^{21,22,42-46} Other chemotherapy backbones that include camptothecins and alkylators (eg, topotecan and cyclophosphamide or topotecan and temozolomide) in combination with dinutuximab or dinutuximab beta demonstrate anti-tumor activity.^{45,48,49}
- Naxitamab-gqgk is FDA approved for the treatment of children >1 year of age and adults with relapsed/refractory high-risk neuroblastoma specifically in bone or bone marrow (but not soft tissue) who have had a PR, MR, or SD to prior therapy.
- In patients for whom GD2-directed therapy is not appropriate at time of first relapse, the targeted radiopharmaceutical I-131 MIBG may be considered based upon results of clinical trials conducted in the relapsed/refractory population.^{49,50}
- The use of targeted agents for patients with relapsed disease whose tumors harbor actionable aberrations has also been evaluated; repeat biomarker profiling on a current sample at the time of relapse is recommended.⁵¹
 - ▶ ALK inhibitors appear to have activity in patients with tumors that have aberrations in the *ALK* gene.⁵²⁻⁵⁴
 - ▶ Agents targeting other pathways of interest in neuroblastoma are also being studied in ongoing clinical trials.
- In specific contexts, external beam radiation and/or surgery may be indicated to address individual sites of relapsed disease.
- Additional options may be considered at the time of second or greater relapse. Rechallenge with chemotherapy combined with GD2-directed antibody therapy based on a history of prior response can be considered.⁴⁷

[References](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SYSTEMIC THERAPY REFERENCES

- ¹ Baker DL, Schmidt ML, Cohn SL, et al; Children's Oncology Group. Outcome after reduced chemotherapy for intermediate-risk neuroblastoma. *N Engl J Med* 2010;363:1313-1323.
- ² Strother DR, London WB, Schmidt ML, et al. Outcome after surgery alone or with restricted use of chemotherapy for patients with low-risk neuroblastoma: results of Children's Oncology Group study P9641. *J Clin Oncol* 2012;30:1842-1848.
- ³ Twist CJ, Schmidt ML, Naranjo A, et al. Maintaining outstanding outcomes using response- and biology-based therapy for intermediate-risk neuroblastoma: A report from the Children's Oncology Group Study ANBL0531. *J Clin Oncol* 2019;37:3243-3255.
- ⁴ Brodeur GM, Pritchard J, Berthold F, et al. Revisions of the international criteria for neuroblastoma diagnosis, staging, and response to treatment. *J Clin Oncol* 1993;11:1466-1477.
- ⁵ Cohn SL, Pearson AD, London WB, et al; INRG Task Force. The International Neuroblastoma Risk Group (INRG) classification system: an INRG Task Force report. *J Clin Oncol* 2009;27:289-297.
- ⁶ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute Clinical Trials Planning Meeting. *J Clin Oncol* 2017;35:2580-2587.
- ⁷ Irwin MS, Naranjo A, Zhang FF, et al. Revised Neuroblastoma Risk Classification System: A report from the Children's Oncology Group. *J Clin Oncol* 2021;39:3229-3241.
- ⁸ Twist CJ, Naranjo A, Schmidt ML, et al. Defining risk factors for chemotherapeutic intervention in infants with Stage 4S neuroblastoma: A report from Children's Oncology Group Study ANBL0531. *J Clin Oncol* 2019;37:115-124.
- ⁹ Balis FM, Womer RB, Berg S, et al. Dosing anticancer drugs in infants: Current approach and recommendations from the Children's Oncology Group's Chemotherapy Standardization Task Force. *Pediatr Blood Cancer* 2017;64:e26636.
- ¹⁰ Kohler JA, Rubie H, Castel V, et al. Treatment of children over the age of one year with unresectable localised neuroblastoma without MYCN amplification: results of the SIOPEN study. *Eur J Cancer* 2013;49:3671-3679.
- ¹¹ Grupp SA, Cohn SL, Wall D, et al. Collection, storage, and infusion of stem cells in children with high-risk neuroblastoma: saving for a rainy day. *Pediatr Blood Cancer* 2006;46:719-722.
- ¹² Pinto N, Naranjo A, Hibbitts E, et al. Predictors of differential response to induction therapy in high-risk neuroblastoma: A report from the Children's Oncology Group (COG). *Eur J Cancer* 2019;112:66-79.
- ¹³ Park JR, Scott JR, Stewart CF, et al. Pilot induction regimen incorporating pharmacokinetically guided topotecan for treatment of newly diagnosed high-risk neuroblastoma: a Children's Oncology Group study. *J Clin Oncol* 2011;29:4351-4357.
- ¹⁴ Park, JR. Kriessman SG, London WB, et al. Effect of tandem autologous stem cell transplant vs single transplant on event-free survival in patients with high-risk neuroblastoma: A randomized clinical trial. *JAMA* 2019;322:746-755.
- ¹⁵ Granger MM, Naranjo A, Bagatell R, et al. Myeloablative busulfan/melphalan consolidation following induction chemotherapy for patients with newly diagnosed high-risk neuroblastoma: Children's Oncology Group Trial ANBL12P1. *Transplant Cell Ther* 2021;27:490.e1-490.e8.
- ¹⁶ Kushner BH, Kramer K, LaQuaglia MP, et al. Reduction from seven to five cycles of intensive induction chemotherapy in children with high-risk neuroblastoma. *J Clin Oncol* 2004;22:4888-4892.
- ¹⁷ Holmes K, Pötschger U, Pearson ADJ, et al. Influence of surgical excision on the survival of patients with stage 4 high-risk neuroblastoma: A report from the HR-NBL1/SIOPEN study. *J Clin Oncol* 2020;38:2902-2915.
- ¹⁸ von Allmen D, Davidoff AM, London WB, et al. Impact of extent of resection on local control and survival in patients from the COG A3973 study with high-risk neuroblastoma. *J Clin Oncol* 2017;35:208-216.
- ¹⁹ Matthay KK, Villablanca JG, Seeger RC, et al. Treatment of high-risk neuroblastoma with intensive chemotherapy, radiotherapy, autologous bone marrow transplantation, and 13-cis-retinoic acid. Children's Cancer Group. *N Engl J Med* 1999;341:1165-1173.
- ²⁰ Desai AV, Applebaum MA, Karrison TG, et al. Efficacy of post-induction therapy for high-risk neuroblastoma patients with end-induction residual disease. *Cancer* 2022;128:2967-2977.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Continued

NEUROB-D
11 OF 13



PRINCIPLES OF SYSTEMIC THERAPY REFERENCES

- 21 Mody R, Yu AL, Naranjo A, et al. Irinotecan, temozolomide, and dinutuximab with GM-CSF in children with refractory or relapsed neuroblastoma: A report from the Children's Oncology Group. *J Clin Oncol* 2020;38:2160-2169.
- 22 Muñoz, JP, Larrosa C, Chamorro S, et al Early salvage chemo-immunotherapy with irinotecan, temozolomide and naxitamab plus GM-CSF (HITS) for patients with primary refractory high-risk neuroblastoma provide the best chance for long-term outcomes. *Cancers (Basel)* 2023;15:4837.
- 23 Yalçın B, Kremer LCM, van Dalen EC. High-dose chemotherapy and autologous haematopoietic stem cell rescue for children with high-risk neuroblastoma. *Cochrane Database Syst Rev* 2015;2015:CD006301.
- 24 Kushner BH, Ostrovskaya I, Cheung IY, et al. Lack of survival advantage with autologous stem-cell transplantation in high-risk neuroblastoma consolidated by anti-GD2 immunotherapy and isotretinoin. *Oncotarget* 2016;7:4155-4166.
- 25 Meany HJ, London WB, Ambros PF, et al. Significance of clinical and biologic features in stage 3 neuroblastoma: a report from the International Neuroblastoma Risk Group project. *Pediatr Blood Cancer* 2014;61:1932-1939.
- 26 Modak S, Kushner BH, LaQuaglia MP, et al. Management and outcome of stage 3 neuroblastoma. *Eur J Cancer* 2009;45:90-98.
- 27 Pinto N, Naranjo A, Ding X, et al. Impact of genomic and clinical factors on outcome of children ≥18 months of age with stage 3 neuroblastoma with unfavorable histology and without MYCN amplification: A Children's Oncology Group (COG) report. *Clin Cancer Res* 2023;29:1546-1556.
- 28 George RE, London WB, Cohn SL, et al. Hyperdiploidy plus nonamplified MYCN confers a favorable prognosis in children 12 to 18 months old with disseminated neuroblastoma: a Pediatric Oncology Group study. *J Clin Oncol* 2005;23:6466-6473.
- 29 Schmidt ML, Lal A, Seeger RC, et al. Favorable prognosis for patients 12 to 18 months of age with stage 4 nonamplified MYCN neuroblastoma: a Children's Cancer Group Study. *J Clin Oncol* 2005;23:6474-6480.
- 30 Ladenstein R, Pötschger U, Pearson ADJ, et al. Busulfan and melphalan versus carboplatin, etoposide, and melphalan as high-dose chemotherapy for high-risk neuroblastoma (HR-NBL1/SIOPEN): an international, randomised, multi-arm, open-label, phase 3 trial. *Lancet Oncol* 2017;18:500-514.
- 31 Braunstein SE, London WB, Kreissman SG, et al. Role of the extent of prophylactic regional lymph node radiotherapy on survival in high-risk neuroblastoma: A report from the COG A3973 study. *Pediatr Blood Cancer* 2019;66:e27736.
- 32 Liu KX, Naranjo A, Zhang FF, et al. Prospective evaluation of radiation dose escalation in patients with high-risk neuroblastoma and gross residual disease after surgery: A report from the Children's Oncology Group ANBL0532 study. *J Clin Oncol* 2020;38:2741-2752.
- 33 Casey DL, Pitter KL, Kushner BH, et al. Radiation therapy to sites of metastatic disease as part of consolidation in high-risk neuroblastoma: Can long-term control be achieved? *Int J Radiat Oncol Biol Phys* 2018;100:1204-1209.
- 34 Polishchuk AL, Li R, Hill-Kayser C, et al. Likelihood of bone recurrence in prior sites of metastasis in patients with high-risk neuroblastoma. *Int J Radiat Oncol Biol Phys* 2014;89:839-845.
- 35 Yu AL, Gilman AL, Ozkaynak MF, et al. Anti-GD2 antibody with GM-CSF, interleukin-2, and isotretinoin for neuroblastoma. *N Engl J Med* 2010;363:1324-1334.
- 36 Yu AL, Gilman AL, Ozkaynak MF, et al. Long-term follow-up of a phase III study of ch14.18 (dinutuximab) + cytokine immunotherapy in children with high-risk neuroblastoma: COG study ANBL0032. *Clin Cancer Res* 2021;27:2179-2189.
- 37 Ladenstein R, Pötschger U, Valteau-Couanet D, et al. Interleukin 2 with anti-GD2 antibody ch14.18/CHO (dinutuximab beta) in patients with high-risk neuroblastoma (HR-NBL1/SIOPEN): a multicentre, randomised, phase 3 trial. *Lancet Oncol* 2018;19:1617-1629.
- 38 Ladenstein R, Pötschger U, Valteau-Couanet D, et al. Investigation of the role of dinutuximab beta-based immunotherapy in the SIOPEN High-Risk Neuroblastoma 1 Trial (HR-NBL1). *Cancers (Basel)* 2020;12:309.
- 39 Sholler GLS, Ferguson W, Bergendahl G, et al. Maintenance DFMO increases survival in high risk neuroblastoma. *Sci Rep* 2018;8:14445.
- 40 Oesterheld J, Ferguson W, Kravaka JM, et al. Eflornithine as postimmunotherapy maintenance in high-risk neuroblastoma: Externally controlled, propensity score-matched survival outcome comparisons. *J Clin Oncol* 2024;42:90-102.
- 41 Mody R, Naranjo A, Van Ryn C, et al. Irinotecan-temozolomide with temsirolimus or dinutuximab in children with refractory or relapsed neuroblastoma (COG ANBL1221): an open-label, randomised, phase 2 trial. *Lancet Oncol* 2017;18:946-957.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Continued

NEUROB-D
12 OF 13



PRINCIPLES OF SYSTEMIC THERAPY REFERENCES

- ⁴² Lerman BJ, Li Y, Carlowicz C, et al. Progression-free survival and patterns of response in patients with relapsed high-risk neuroblastoma treated with irinotecan/temozolomide/dinutuximab/granulocyte-macrophage colony-stimulating factor. *J Clin Oncol* 2023;41:508-516.
- ⁴³ Olgun N, Cecen E, Ince D, et al. Dinutuximab beta plus conventional chemotherapy for relapsed/refractory high-risk neuroblastoma: A single-center experience. *Front Oncol* 2022;12:1041443.
- ⁴⁴ Wiczorek A, Zaniewska-Tekieli A, Ehlert K, et al. Dinutuximab beta combined with chemotherapy in patients with relapsed or refractory neuroblastoma. *Front Oncol* 2023;13:1082771.
- ⁴⁵ Gray JC, Weston R, Owens C, et al. Dinutuximab beta added to temozolomide-based chemotherapy for children with relapsed and refractory neuroblastoma: Results of the ITCC-SIOPEN BEACON Immuno phase II trial. *J Clin Oncol* 2026;44:176-187.
- ⁴⁶ Mora J, Chan GCF, Morgenstern DA, et al. The anti-GD2 monoclonal antibody naxitamab plus GM-CSF for relapsed or refractory high-risk neuroblastoma: a phase 2 clinical trial. *Nat Commun* 2025;16:1636.
- ⁴⁷ Lerman BJ, King M, Mody R, et al. Efficacy and tolerability of topotecan/cyclophosphamide/dinutuximab in relapsed and refractory high-risk neuroblastoma: A multi-institutional retrospective study. *Pediatr Blood Cancer* 2026;73(5):e70195.
- ⁴⁸ Raiser P, Schleiermacher G, Gambart M, et al. Chemo-immunotherapy with dinutuximab beta in patients with relapsed/progressive high-risk neuroblastoma: Does chemotherapy backbone matter? *Eur J Cancer* 2024;202:114001.
- ⁴⁹ Matthay KK, Yanik G, Messina J, et al. Phase II study on the effect of disease sites, age, and prior therapy on response to iodine-131-metaiodobenzylguanidine therapy in refractory neuroblastoma. *J Clin Oncol* 2007;25:1054-1060.
- ⁵⁰ DuBois SG, Granger MM, Groshen S, et al. Randomized phase II trial of MIBG versus MIBG, vincristine, and irinotecan versus MIBG and vorinostat for patients with relapsed or refractory neuroblastoma: A report from NANT Consortium. *J Clin Oncol* 2021;39:3506-3514.
- ⁵¹ Eleveld TF, Oldridge DA, Bernard V, et al. Relapsed neuroblastomas show frequent RAS-MAPK pathway mutations. *Nat Genet* 2015;47:864-871.
- ⁵² Goldsmith KC, Park JR, Kayser K, et al. Lorlatinib with or without chemotherapy in ALK-driven refractory/relapsed neuroblastoma: phase 1 trial results. *Nat Med* 2023;29:1092-1102.
- ⁵³ Fischer M, Moreno L, Ziegler DS, et al. Ceritinib in paediatric patients with anaplastic lymphoma kinase-positive malignancies: an open-label, multicentre, phase 1, dose-escalation and dose-expansion study. *Lancet Oncol* 2021;22:1764-1776.
- ⁵⁴ Rosswog C, Fassunke J, Ernst A, et al. Genomic ALK alterations in primary and relapsed neuroblastoma. *Br J Cancer* 2023;128:1559-1571.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF SURGERY

- Evaluate primary tumor resectability using cross-sectional imaging; CT or MRI should be performed.
 - ▶ Evaluate for the presence of IDRFs.¹
- Multidisciplinary discussion of risks/benefits and optimal timing of surgery (at diagnosis or following neoadjuvant chemotherapy) should include surgeons, diagnostic and interventional radiologists, medical oncologists, radiation oncologists, pathologists, and dieticians as needed.
- If upfront resection is not appropriate, a biopsy that provides adequate tissue for diagnosis and risk stratification should be performed.

General Principles of Surgery

- Understanding the heterogenous nature of neuroblastoma is critical when determining the optimal timing and extent of surgical resection.
 - ▶ Patients with intermediate-risk disease most often require treatment with a combination of surgery and chemotherapy. If more than minimal risks of surgical morbidity exist, neoadjuvant chemotherapy should be administered initially, as systemic therapy may reduce subsequent surgical risk.²
 - ▶ In the setting of high-risk disease, neoadjuvant chemotherapy is nearly always indicated prior to resection.
- For all risk groups, the surgical approach should limit morbidity and mortality, avoid resection of vital structures, and preserve organ function.

Principles of Tumor Biopsy

- Indicated when resection at diagnosis is not possible or is not recommended due to the clinical context.
 - ▶ Generally indicated for patients with metastatic disease.
 - ▶ Generally indicated for patients with L2 tumors.
- Incisional biopsy via an open or laparoscopic/thoracoscopic approach should result in collection of >1 cm³ of fresh viable tissue.^a
- Core biopsy can be performed by a surgeon or interventional radiologist, and should result in collection of at least 10 cores to ensure that the specimen is adequate for biomarker profiling.^{a,3}
- Pathologist assessment at the time of biopsy is recommended to ensure that the tissue biopsied is viable rather than necrotic, and likely diagnostic.^a

^a See [Principles of Pathology \(NEUROB-A\)](#) for specimen requirements.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



PRINCIPLES OF SURGERY

Surgical Approach by Risk Group

- Low Risk
 - ▶ Patients with INRG Stage L1 tumors are often cured with surgical resection. If upfront surgery will potentially obviate the need for chemotherapy and can be safely performed with minimal morbidity, a resection should be performed. See Low-Risk Disease ([NEUROB-3](#)) regarding infants in whom observation without biopsy or resection is appropriate.
- Intermediate Risk
 - ▶ The decision to proceed with upfront resection without preoperative chemotherapy is based on whether IDRFs are absent or present (L1 vs. L2) and on risk factor analysis (including evaluation of the nature of IDRFs present). When risk of surgical morbidity is deemed significant by the surgeon, neoadjuvant chemotherapy should be considered.
 - ▶ Gross total resection may be attempted following initial systemic therapy. However, preservation of vital structures and of end-organ function is of utmost importance in the intermediate-risk context, as less than CR has been shown to be an acceptable endpoint of therapy for patients with localized intermediate-risk tumors.
 - ▶ Timing of resection will depend upon response to initial therapy and subsequent assessment of surgical risk.
- High Risk
 - ▶ Surgery is generally undertaken prior to completion of induction therapy, as extended preoperative therapy is unlikely to further reduce surgical risk after multiple cycles of chemotherapy have been given.⁴⁻⁷
 - ▶ Gross total resection of the primary tumor is optimal; however, residual disease adherent to critical/vital structures should not be removed if attempts to resect it can be expected to result in surgical morbidity, organ dysfunction, and delays in the initiation of postoperative systemic therapy.
 - ▶ Resection of adjacent organs (kidney, pancreas, intestine, and neurovascular structures) is not recommended. If direct extension/infiltration of the tumor into adjacent organs is identified, tumor may be left behind. Resection of adjacent organs increases the morbidity of the surgical procedure and could impact the dosing/tolerance of adjuvant therapy. Margin-negative resection is rarely achieved in this context and is not the surgical goal.⁸
 - ▶ Nephrectomy should be avoided as neuroblastoma treatment can result in significant renal toxicity, and because postoperative renal dysfunction can limit therapeutic options.⁹
 - ▶ Tumors with vascular encasement require identification, isolation, and clearance of associated major vessels. Exposure should be adequate to safely obtain proximal and distal vascular control. Commonly, the tumor is divided over the involved vascular structure and dissected down to the adventitia to allow for vessel preservation and tumor resection, but dissection should not proceed deeper than the adventitial layer to avoid vascular injury.
 - ◊ Tumors may be resected in multiple segments; en bloc resection is not required.
 - ▶ Resection of nodal disease adherent or adjacent to the primary tumor is recommended as long as resection of this tissue is not expected to increase surgical morbidity. Resection of distant lymph nodes at the time of primary tumor resection is not required.¹⁰
- Special considerations for infants with INRG MS disease
 - ▶ The primary surgical role for children with MS disease is biopsy of the primary or metastatic tumor for histologic and biologic studies when it is safe to do so.
 - ▶ Infants with MS disease who are <3 months of age are at increased risk for surgical complications in the presence of massive hepatomegaly with or without coagulopathy. In some cases, systemic therapy may be required prior to biopsy. In this setting, biopsy (surgical or via percutaneous core needle) can be delayed until it can be undertaken safely.
 - ▶ In extreme situations, for infants with abdominal compartment syndrome, severe respiratory compromise, or life-threatening symptoms secondary to hepatomegaly, laparotomy and silo placement may be considered to alleviate intra-abdominal pressure.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

[References](#)



PRINCIPLES OF SURGERY

Surgical Management: Cervical and Cervicothoracic

- Assess for encasement of the vertebral or carotid arteries, jugular vein, subclavian vessels, brachial plexus roots, or extension across the midline or to the base of the skull and ensure proper identification and preservation.
- Assess for tracheal compression evidenced by patient symptoms or cross-sectional imaging prior to any general anesthesia.
- Assess for extension into the thoracic cavity. If present, consider a cervicothoracic “trap door” or a median sternotomy with cervical extension incision for resection.
- Identification and preservation of vagus nerve, brachial plexus, phrenic nerve, and other major nerves should be ensured. Intraoperative nerve monitoring should be considered. Communication with the anesthesia team regarding the use or withholding of neuromuscular blockade should occur preoperatively.
- Postoperative Horner syndrome should be discussed with the patient/family, as this is not uncommon following resection of neck and apical chest tumors.
- There is no role for complete compartment-oriented cervical lymphadenectomy.

Surgical Management: Thoracic Cavity

- Assess for tumor encasing the thoracic aorta, vena cava, or other great vessels.
- Assess for tracheal compression evidenced by patient symptoms or cross-sectional imaging prior to any general anesthesia.
- A thoracotomy incision is generally indicated for large tumors or those with involvement of critical structures.
- A thoracoscopic approach may be indicated for selective lesions.¹¹
- Avoid aggressive resection, retraction, or use of monopolar cautery near the recurrent laryngeal, phrenic, and vagus nerves.
- Extension into spinal foramina should be assessed preoperatively with a spine MRI if indicated based on primary tumor location or clinical symptoms. Dissection close to the spinal nerve roots should be avoided. Neural foraminal extension may be resected via a thoracic approach, but deep/blind tissue extraction from this region should be avoided. Leaving residual tumor in this location is favored over risking neurologic injury.
- If known intraspinal extension is present, the case should be discussed with the multidisciplinary team.
- Thoracic/spinal imaging of the aortic branches to the spinal cord should be evaluated preoperatively. Arterial branches to the spinal cord, including the artery of Adamkiewicz, should be evaluated preoperatively and preserved whenever possible for multiple level thoracic paraspinal tumors, even if some tumor is left in situ.
- Careful dissection posterior to the carina and near the right hemidiaphragm can minimize injury to the thoracic duct; any visible lymphatics should be sealed or ligated. A thorough visual inspection for lymphatic leak should be performed prior to completion of the procedure.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References](#)



PRINCIPLES OF SURGERY

Surgical Management: Abdominal/Adrenal/Retroperitoneal/Pelvic Cavity

- Assess for tumor infiltrating the porta hepatis and/or the hepatoduodenal ligament or encasing the mesenteric arteries, celiac axis, aorta and/or vena cava, or iliac vessels.
 - ▶ Surgical resection should not compromise major vessels or increase the risk of bowel ischemia.
 - ▶ Proximal and distal vascular control should be obtained when indicated.
- Assess for tumor infiltrating one or both renal pedicles. Nephrectomy should be avoided.⁹
- Ureters may be adherent to the tumor and care should be taken to avoid injury.
- Preoperative MRI to assess nerve root and sacral involvement should be obtained when relevant in light of primary tumor location.
 - ▶ Nerve monitoring should be considered for pelvic tumors due to proximity to the lumbosacral plexus and obturator nerve.
- A laparoscopic approach can be considered depending on the location of the tumor and its relationship to surrounding structures (ie, IDRFs).^{12,13}

Surgical Management: Paraspinal

- A detailed preoperative neurologic examination should be performed and any deficits documented.
- A preoperative spine MRI should be obtained.
- Urgent multidisciplinary evaluation is indicated with consideration of rapid initiation of chemotherapy. Laminectomy is rarely indicated at the time of diagnosis.¹⁴⁻¹⁷
- Progressive neurologic symptoms or acute deterioration following rapid initiation of systemic therapy should prompt urgent multidisciplinary re-evaluation.

Surgical Complications

- Intraoperative complications are site dependent.
- Hemorrhage is a major risk with infiltrative tumors. Appropriate preoperative planning or intraoperative vascular surgical consultation should be sought when necessary. Critical vessels including the aorta, carotid, subclavian, hepatic, mesenteric, or renal arteries should be preserved or repaired if injured, either primarily or with a graft, to restore flow.
- Nerve injuries may also occur, and primary repair should be considered utilizing intraoperative consultation with vascular, neurosurgical, or plastic surgical specialists.
- It is acceptable and recommended to leave a small amount of residual tumor adherent to vital structures if there is significant risk of injury with more aggressive or complete resection. An estimate of the extent of resection should be documented in the operative report.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



PRINCIPLES OF SURGERY REFERENCES

- ¹ Irtan S, Brisse HJ, Minard-Colin V, et al. Image-defined risk factor assessment of neurogenic tumors after neoadjuvant chemotherapy is useful for predicting intra-operative risk factors and the completeness of resection. *Pediatr Blood Cancer* 2015;62:1543-1549.
- ² Medary I, Aronson D, Cheung NK, et al. Kinetics of primary tumor regression with chemotherapy: implications for the timing of surgery. *Ann Surg Oncol* 1996;3:521-525.
- ³ Overman RE, Kartal TT, Cunningham AJ, et al. Optimization of percutaneous biopsy for diagnosis and pretreatment risk assessment of neuroblastoma. *Pediatr Blood Cancer* 2020;67:e28153.
- ⁴ La Quaglia MP, Kushner BH, Su W, et al. The impact of gross total resection on local control and survival in high-risk neuroblastoma. *J Pediatr Surg* 2004;39:412-417.
- ⁵ Yokoyama J, Ikawa H, Endow M, et al. The role of surgery in advanced neuroblastoma. *Eur J Pediatr Surg* 1995;5:23-26.
- ⁶ Von Allmen D, Davidoff AM, London W, et al. Influence of extent of resection on Survival in high risk neuroblastoma patients: a report from the COG A3973 Study. *J Clin Oncol* 2017;35:208-216.
- ⁷ Holmes K, Sarnacki S, Poetschger U, et al. Influence of surgical excision on survival of patients with high risk neuroblastoma. Report from Study 1 of SIOP Europe (SIOPEN). *J Clin Oncol* 2020;38:2902-2915.
- ⁸ Englum BR, Rialon KL, Speicher PJ, et al. Value of surgical resection in children with high-risk neuroblastoma. *Pediatr Blood Cancer* 2015;62:1529-1535.
- ⁹ Shamberger RC, Smith EI, Joshi VV, et al. The risk of nephrectomy during local control in abdominal neuroblastoma. *J Pediatr Surg* 1998;33:161-164.
- ¹⁰ Hayes FA, Green A, Hustu HO, et al. Surgicopathologic staging of neuroblastoma: prognostic significance of regional lymph node metastases. *J Pediatr* 1983;102:59-62.
- ¹¹ Malek MM, Mollen KP, Kane TD, et al. Thoracic neuroblastoma: a retrospective review of our institutional experience with comparison of the thoracoscopic and open approaches to resection. *J Pediatr Surg* 2010;45:1622-1626.
- ¹² Kelleher CM, Smithson L, Nguyen LL, et al. Clinical outcomes in children with adrenal neuroblastoma undergoing open versus laparoscopic adrenalectomy. *J Pediatr Surg* 2013;48:1727-1732.
- ¹³ Gurria JP, Malek MM, Heaton TE, et al. Minimally invasive surgery for neuroblastic tumors: A systematic review by the APSA Cancer Committee. *J Pediatr Surg* 2020;55:2260-2272.
- ¹⁴ De Bernardi B, Pianca C, Pistamiglio P, et al. Neuroblastoma with symptomatic spinal cord compression at diagnosis: treatment and results with 76 cases. *J Clin Oncol* 2001;19:183-190.
- ¹⁵ Katzenstein HM, Kent PM, London WB, et al. Treatment and outcome of 83 children with intraspinal neuroblastoma: the Pediatric Oncology Group experience. *J Clin Oncol* 2001;19:1047-1055.
- ¹⁶ Hoover M, Bowman LC, Crawford SE, et al. Long-term outcome of patients with intraspinal neuroblastoma. *Med Pediatr Oncol* 1999;32:353-359.
- ¹⁷ Trahair T, Sorrentino S, Russell SJ, et al. Spinal canal involvement in neuroblastoma. *J Pediatr* 2017;188:294-298.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

DISEASE SURVEILLANCE/FOLLOW-UP AFTER COMPLETION OF TREATMENT

Risk Category ¹	Workup/Imaging
Low Risk (observation only)	• N/A
Low Risk (patients who underwent surgical resection)	• Interim H&P: ▶ Approximately every 3 months for Year 1, ▶ Then every 6 months for Year 2, ▶ Then every 6–12 months for Year 3, ▶ Then as clinically indicated
	• Laboratory studies: ▶ Urine catecholamine levels are no longer included in the revised INRC due to the lack of standardization in specimen collection and analysis, and due to the influence of diet on results.² However, results of catecholamine testing may be helpful during surveillance. ▶ If elevated at diagnosis, consider obtaining spot catecholamine levels during surveillance.
	• Imaging: ▶ US of primary site^a ◇ Approximately every 3 months for Year 1, ◇ Then every 6–12 months for Years 2–3, ◇ Then as clinically indicated

^a Cross-sectional imaging may be required depending on the location of the primary tumor.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

NEUROB-F
1 OF 4



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

DISEASE SURVEILLANCE/FOLLOW-UP AFTER COMPLETION OF TREATMENT

Risk Category ¹	Workup/Imaging
Intermediate Risk (adapted from ANBL0531 ³ and ANBL1232)	• Interim H&P: ▶ Approximately every 3 months for Year 1, ▶ Then every 6 months for Year 2, ▶ Then annually for Years 3–5 • Consider audiologic assessment depending on degree of exposure to ototoxic agents
	• Laboratory studies: ▶ Urine catecholamine levels are no longer included in the revised INRC due to the lack of standardization in specimen collection and analysis, and due to the influence of diet on results.² However, results of catecholamine testing may be helpful during surveillance. ▶ If elevated at diagnosis, consider obtaining spot catecholamine levels during surveillance. ▶ CBC with differential if bone marrow was involved at diagnosis. Obtain with imaging. ▶ Creatinine ◇ Every 6 months for Year 1, ◇ Then annually for Years 2–3, ◇ Then as clinically indicated ▶ Thyroid studies including thyroid-stimulating hormone (TSH) (and free T4 if TSH is abnormal) – Annually through Year 3 – Then as clinically indicated
	• Imaging: ▶ CT or MRI cross-sectional imaging of primary site ◇ Approximately every 3 months for Year 1, ◇ Then every 6 months for Year 2, ◇ Then annually for Year 3, ◇ Then as clinically indicated ▶ I-123 MIBG scan with SPECT, if available (for MIBG-avid tumors for patients with INRG Stage M disease at diagnosis) ◇ If positive at diagnosis and at completion of therapy, obtain MIBG imaging until a negative scan is achieved or patient is 36 months from completion of therapy – Every 3–6 months in Year 1, – Then annually in Years 2–3, – Then as clinically indicated ▶ FDG-PET scan (for MIBG non-avid tumors) ◇ If positive at diagnosis and at completion of therapy, obtain PET imaging until a negative scan is achieved or patient is 36 months from completion of therapy – Every 3–6 months in Year 1, – Then annually in Years 2–3, – Then as clinically indicated

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

DISEASE SURVEILLANCE/FOLLOW-UP AFTER COMPLETION OF TREATMENT

Risk Category ¹	Workup/Imaging
High Risk (adapted from ANBL0532 ⁴ and ANBL1531)	• Interim H&P: ▶ Every 3 months for Year 1, ▶ Then every 6 months for Years 2–5 ▶ Ongoing evaluation of growth parameters • Laboratory studies: ▶ Urine catecholamine levels are no longer included in the revised INRC due to the lack of standardization in specimen collection and analysis, and due to the influence of diet on results.² However, results of catecholamine testing may be helpful during surveillance. ▶ CBC with differential. Obtain with imaging. ▶ Electrolytes including Ca⁺⁺, PO₄, Mg⁺⁺; and creatinine, alanine aminotransferase (ALT), bilirubin – Every 3 months for Year 1, – Then every 6 months for Years 2–3, – Then annually for Years 4–5 ▶ Thyroid studies including TSH (and free T4 if TSH is abnormal) – Every 6 months for Years 1–2, – Then annually for Years 3–5 ▶ Hemoglobin A1c (HbA1c), ferritin, and reproductive health labs (follicle-stimulating hormone [FSH], luteinizing hormone [LH], anti-Mullerian hormone, inhibin B) – As clinically indicated ▶ Audiologic assessment – Annually for Years 1–5, – Then as clinically indicated – Studies of survivors of high-risk neuroblastoma suggest high rates of severe ototoxicity (requiring hearing aids)⁵ ▶ Echocardiogram – If normal at the end of therapy, obtain every 2–5 years based on total anthracycline dose and radiation dose with potential impact to heart (Monitoring for Late Effects [NEUROB-I]) ▶ Pulmonary function tests – As clinically indicated • Imaging: ▶ CT or MRI cross-sectional imaging of primary site – Every 3–6 months for Year 1, – Then every 6 months for Year 2, – Then annually for Year 3, – Then as clinically indicated ▶ I-123 MIBG scan with SPECT, if available (for MIBG-avid tumors) – Every 3–6 months for Year 1, – Then every 6 months for Year 2, – Then annually for Year 3, – Then as clinically indicated ▶ FDG-PET scan (for MIBG non-avid tumors) – Every 3–6 months for Year 1, – Then every 6 months for Year 2, – Then annually for Year 3, – Then as clinically indicated • Bilateral bone marrow aspirates and biopsies: ▶ If negative at the end of therapy, obtain only as clinically indicated.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

NEUROB-F
3 OF 4



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

DISEASE SURVEILLANCE/FOLLOW-UP AFTER COMPLETION OF TREATMENT REFERENCES

- ¹ Irwin MS, Naranjo A, Zhang FF, et al. Revised Neuroblastoma Risk Classification System: A Report From the Children's Oncology Group. J Clin Oncol 2021;39:3229-3241.
- ² Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute clinical trials planning meeting. J Clin Oncol 2017;35:2580-2587.
- ³ Twist CJ, Schmidt ML, Naranjo A, et al. Maintaining outstanding outcomes using response- and biology-based therapy for intermediate-risk neuroblastoma: A report from the Children's Oncology Group study ANBL0531. J Clin Oncol 2019;37:3243-3255.
- ⁴ Park JR, Kreissman SG, London WB, et al. Effect of tandem autologous stem cell transplant vs single transplant on event-free survival in patients with high-risk neuroblastoma: A randomized clinical trial. JAMA 2019;322:746-755.
- ⁵ Henderson TO, Bardwell JK, Kao PC, et al. Late effects after high-risk neuroblastoma (LEAHRN): a multicentre, cross-sectional cohort study from the Children's Oncology Group. Lancet Child Adolesc Health 2025;9:776-786.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

RESPONSE ASSESSMENT

Component	Method	International Neuroblastoma Response Criteria (INRC) Component Response ¹
Primary Site	RECIST/Functional Imaging ^{1,2}	Complete response (CR): <10 mm residual soft tissue at primary site, AND complete resolution of MIBG or FDG-PET ^a uptake (for MIBG non-avid tumors) at primary site
		Partial response (PR): ≥30% decrease in longest diameter of primary site, AND MIBG or FDG-PET ^a uptake at primary site stable, improved or resolved
		Stable disease (SD): Neither sufficient shrinkage for PR nor sufficient increase for PD at the primary site
		Progressive disease (PD): >20% increase in longest diameter taking as reference the smallest diameter on study (this includes the baseline diameter if that is the smallest on study), AND a minimum absolute increase of 5 mm in longest dimension ^b

Footnotes

^a Used for tumors that do not demonstrate MIBG uptake.

^b Fluctuating MIBG or PET avidity is insufficient for PD in the absence of an increase in measurement that meets criteria for PD.

References

¹ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute clinical trials planning meeting. J Clin Oncol 2017;35:2580-2587.

² Bagatell R, McHugh K, Naranjo A, et al. Assessment of primary site response in children with high-risk neuroblastoma: An international multicenter study. J Clin Oncol 2016;34:740-746.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

RESPONSE ASSESSMENT

Component	Method	INRC Component Response ¹
Metastatic bone/ soft tissue	RECIST/Modified Curie or SIOPEN score ^{1,3,4}	CR: Non-primary target and non-target lesions measure <10 mm, AND Lymph nodes identified as target lesions decrease to a short axis <10 mm, AND MIBG uptake or FDG- PET uptake (for MIBG non-avid tumors) of non-primary lesions resolves completely
		PR: ≥30% decrease in sum of diameters of non-primary target lesions compared to baseline, AND all of the following: • Non-target lesions may be stable or smaller in size AND • No new lesions AND • ≥50% reduction in MIBG absolute bone score (relative MIBG bone score ≥0.1 to ≤0.5) or ≥50% reduction in number of FDG-PET-avid bone lesions^{c,d}
		SD: Neither sufficient shrinkage for PR nor sufficient increase for PD of non-primary lesions
		PD: Any of the following: • Any new soft tissue lesion detected by CT/MRI that is also MIBG avid or FDG-PET avid • Any new soft tissue lesion seen on anatomic imaging that is biopsied and confirmed to be a neuroblastoma or ganglioneuroblastoma • Any new bone site that is MIBG avid • A new bone site that is FDG-PET avid (for MIBG non-avid tumors) AND has CT/MRI findings consistent with tumor OR has been confirmed histologically to be neuroblastoma or ganglioneuroblastoma • >20% increase in the sum of the longest diameters taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study), AND a minimum absolute increase of 5 mm in sum of diameters of target soft tissue lesions • Relative MIBG bone score ≥1.2

Footnotes

^c Complete resolution of MIBG or PET avidity is not required.

^d Relative score is the absolute score for bone lesions at time of response assessment divided by the absolute score of bone lesions at baseline, prior to therapeutic interventions.

References

- ¹ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute clinical trials planning meeting. J Clin Oncol 2017;35:2580-2587.
- ³ Yanik GA, Parisi MT, Shulkin BL, et al. Semiquantitative mIBG scoring as a prognostic indicator in patients with stage 4 neuroblastoma: a report from the Children's Oncology Group. J Nucl Med 2013;54:541-548.
- ⁴ Ladenstein R, Lambert B, Pütschger U, et al. Validation of the mIBG skeletal SIOPEN scoring method in two independent high-risk neuroblastoma populations: the SIOPEN/HR-NBL1 and COG-A3973 trials. Eur J Nucl Med Mol Imaging 2018;45:292-305.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

RESPONSE ASSESSMENT

Component	Method	INRC Component Response ¹
Bone marrow	INRC/Burchill ^{e,5}	CR: Bone marrow with no tumor infiltration noted on reassessment, independent of baseline tumor infiltration
		SD: Bone marrow with tumor infiltration that remains positive with >5% tumor infiltration on reassessment but does not meet criteria for CR, minimal disease (MD), or PD
		MD: Bone marrow with ≤5% tumor infiltration that remains >0 to ≤5% tumor infiltration at subsequent reassessment, OR bone marrow with no tumor infiltration that has ≤5% tumor infiltration on reassessment, OR bone marrow with >20% tumor infiltration that has >0% to ≤5% tumor infiltration on reassessment
		PD: Bone marrow without tumor infiltration that becomes >5% tumor infiltration on reassessment, OR bone marrow with tumor infiltration that increases by >2-fold and is >20% tumor infiltration on reassessment

Footnote

^e Recommend >5% viable tumor as reliable level for detecting response.

References

¹ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute clinical trials planning meeting. J Clin Oncol 2017;35:2580-2587.

⁵ Burchill SA, Beiske K, Shimada H, et al. Recommendations for the standardization of bone marrow disease assessment and reporting in children with neuroblastoma on behalf of the International Neuroblastoma Response Criteria Bone Marrow Working Group. Cancer 2017;123:1095-1105.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

RESPONSE ASSESSMENT

Overall Response ¹	
Complete response (CR)	All components CR
Partial response (PR)	PR in at least one component and all other components CR, MD (bone marrow), PR, or not involved at baseline
Minor response (MR)	PR or CR in at least one component but at least one other component with SD; no component with PD
Stable disease (SD)	SD in one component with no better than SD or not involved at baseline in any other component, no component with PD
Progressive disease (PD)	Any component with PD

Notable features of the INRC

- Changes in urinary catecholamine levels (HVA and VMA) do not impact response assessment.
- Technetium-99m (99mTc) scintigraphy should not be used for response assessment.
- Specific requirements for timing of response are indicated within treatment algorithms and differ across risk groups.
- While novel radiotracers are in development to assess response, insufficient data are currently available to permit incorporation into the INRC.^{6,7}

¹ Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute clinical trials planning meeting. J Clin Oncol 2017;35:2580-2587.

⁶ Fortunati E, Argalia G, Zanoni L, et al. New PET radiotracers for the imaging of neuroendocrine neoplasms. Curr Treat Options Oncol 2022;23:703-720.

⁷ Feng L, Li S, Wang C, Yang J. Current status and future perspective on molecular imaging and treatment of neuroblastoma. Semin Nucl Med 2023;53:517-529.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RADIATION THERAPY

General Principles for RT in Patients with High-Risk Neuroblastoma

- RT is indicated for nearly all cases of high-risk neuroblastoma and in exceedingly rare circumstances for patients with non–high-risk disease. For example, symptomatic patients (stage MS) with severe ongoing organ dysfunction with inadequate response to initial chemotherapy may benefit from RT; multidisciplinary discussion is recommended.
 - ▶ RT will be delivered after recovery from high-dose chemotherapy with stem cell rescue unless there is an indication for emergent RT.
 - ▶ Intensity-modulated RT (IMRT) or proton therapy can be used to reduce the risk and/or magnitude of side effects.
 - ▶ The primary site is always irradiated. Metastatic sites are only irradiated if there is concern for active disease after completion of induction chemotherapy, demonstrated by MIBG/FDG uptake and/or persistent soft-tissue mass $>1 \text{ cm}^3$.¹
 - ▶ If radiation to metastatic sites is indicated, they should be irradiated concurrently with the primary site.
- The Panel recognizes that not all metastatic sites may be feasibly targeted by external beam radiotherapy. While repeat imaging following recovery from transplant and prior to radiation is not routinely performed, it may aid in decision-making in select clinical scenarios. The recommendation to radiate residual metastatic sites of disease is based upon limited institution data supporting a benefit from this approach,^{1,2} though this approach has not been adopted internationally.
- Simulation
 - ▶ Patients should be supine with appropriate immobilization for CT simulation.
 - ▶ Four-dimensional CT (4D-CT) is recommended for targets that are subject to respiratory motion.³
 - ▶ Consider addition of MRI simulation for paraspinal sites.
- Target Volume Definitions
 - ▶ Primary Site:
 - ◊ The gross tumor volume (GTV) will include the postoperative tumor bed, any residual disease identified after surgery, and initially involved regional lymph nodes. The tumor bed will be defined by the post-induction chemotherapy, pre-surgery volume, respecting pushing borders, and operative findings not readily identified on imaging. Relevant CT and/or MRI studies completed after induction chemotherapy and prior to surgery should be fused to the CT simulation for target delineation.
 - Special consideration: If the primary tumor was resected prior to induction chemotherapy, GTV will be based on the tumor volume at diagnosis.
 - ◊ The clinical target volume (CTV) = GTV + 1 cm, confined to anatomic borders. For instance, CTV need not extend into uninvolved kidneys, abdominal organs, or bones.
 - ◊ If the tumor is subject to respiratory motion, an internal target volume (ITV) should be utilized based on the 4D-CT.
 - ◊ The planning target volume (PTV) = CTV + 0.3–0.5 cm, depending on the institution's standard, immobilization method, and modality of in-room verification imaging.
 - ▶ Metastatic Sites:
 - ◊ The metastatic GTV (mGTV) is defined by the post-induction chemotherapy volume.
 - ◊ Metastatic CTV (mCTV) = mGTV + 1 cm, confined to anatomic borders.
 - ◊ Metastatic PTV (mPTV) = mCTV + 0.3–0.5 cm, depending on the institution's standard, immobilization method, and modality of in-room verification imaging.
- Neuroblastoma is a radiosensitive tumor and a commonly used dose of 21.6 Gy is recommended ([NEUROB-H 2 of 3](#)). Augmented RT approaches have not resulted in improved local control based on data from recent trials.^{4,5}

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

References



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY

Dose Definitions	
Site	Dose
Primary site and involved regional lymph nodes ^a	21.6 Gy in 1.8 Gy per fraction
Metastatic sites that are concerning for active disease after induction chemotherapy ^{1,6}	21.6 Gy in 1.8 Gy per fraction

Organs at Risk (OAR) ^b	Dose Constraints
Contralateral kidney ⁷	V18 Gy <25% Mean dose ≤14.4 Gy
Ipsilateral kidney	Mean dose ≤18 Gy V14.4 Gy <100% V18 Gy <75%
Liver ⁸	Mean dose <18 Gy
Vertebrae (applies if PTV extends into vertebral body) ⁹	Minimum 18 Gy to cover vertebrae to prevent asymmetrical growth
Bilateral lungs ¹⁰	V20 Gy <30%
Ipsilateral lung	V20 Gy <30%
Contralateral lung	V20 Gy <10%

- Emergent RT (4.5 Gy in 1.5 Gy/fraction) should be considered in clinical scenarios including the following:
 - ▶ Hepatomegaly leading to respiratory distress
 - ▶ Orbit or optic pathway disease leading to vision loss

^a Boosting residual disease after surgery to 36 Gy did not improve cumulative incidence of local progression.⁴ An ongoing European trial includes a randomization in which patients with residual disease either receive a boost or receive standard dose therapy.

^b Adapted from COG ANBL 1531 (NCT03126916) study protocol.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



PRINCIPLES OF RADIATION THERAPY REFERENCES

- ¹ Casey DL, Pitter KL, Kushner BH, et al. Radiation therapy to sites of metastatic disease as part of consolidation in high-risk neuroblastoma: Can long-term control be achieved? *Int J Radiat Oncol Biol Phys* 2018;100:1204-1209.
- ² Polishchuk AL, Li R, Hill-Kayser C, et al. Likelihood of bone recurrence in prior sites of metastasis in patients with high-risk neuroblastoma. *Int J Radiat Oncol Biol Phys* 2014;89:839-845.
- ³ Kannan S, Teo BK, Solberg T, Hill-Kayser C. Organ motion in pediatric high-risk neuroblastoma patients using four-dimensional computed tomography. *J Appl Clin Med Phys* 2017;18:107-114.
- ⁴ Liu KX, Naranjo A, Zhang FF, et al. Prospective evaluation of radiation dose escalation in patients with high-risk neuroblastoma and gross residual disease after surgery: A report from the Children's Oncology Group ANBL0532 study. *J Clin Oncol* 2020;38:2741-2752.
- ⁵ Braunstein SE, London WB, Kreissman SG, et al. Role of the extent of prophylactic regional lymph node radiotherapy on survival in high-risk neuroblastoma: A report from the COG A3973 study. *Pediatr Blood Cancer* 2019;66:e27736.
- ⁶ Lucas Jr JT, Wakefield DV, Doubrovin M, et al. Risk factors associated with metastatic site failure in patients with high-risk neuroblastoma. *Clin Transl Radiat Oncol* 2022;34:42-50.
- ⁷ Poppe MM, Tai A, Li XA, et al. Kidney disease in childhood cancer survivors treated with radiation therapy: A PENTEC comprehensive review. *Int J Radiat Oncol Biol Phys* 2024;119:560-574.
- ⁸ Hall MD, Howell RM, Jackson A, et al. Liver late effects in childhood cancer survivors treated with radiation therapy: A PENTEC comprehensive review. *Int J Radiat Oncol Biol Phys* 2024;119:575-587.
- ⁹ Nanda RH, Hua CH, Flampouri S, et al. Risks of spinal abnormalities and growth impairment after radiation to the spine in childhood cancer survivors: A PENTEC comprehensive review. *Int J Radiat Oncol Biol Phys* 2024;119:507-521.
- ¹⁰ Briere TM, Agrusa JE, Martel MK, et al. Acute and late pulmonary effects after radiation therapy in childhood cancer survivors: A PENTEC comprehensive review. *Int J Radiat Oncol Biol Phys* 2024;119:533-548.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



MONITORING FOR LATE EFFECTS

Monitoring for Late Effects ≥2 Years After Completion of Systemic Therapy

- Late effects and survivorship care are specific to the individual, the age at which neuroblastoma is diagnosed, and the wide range of therapies used to treat it. Generalized recommendations for screening for late effects would not be appropriate or accurate for many patients. It is strongly recommended to create a survivorship care plan specific to each individual patient.
- Refer to the [COG Survivorship Guidelines](#) for appropriate screening and counseling related to: thyroid, cardiac, pulmonary, renal, bone, reproductive health, subsequent cancers (with special attention to thyroid and kidney), and other treatment-associated late effects.

High-Risk Disease

- Patients with high-risk neuroblastoma (especially those treated with myeloablative therapy) are at particularly high risk of hearing impairment,^a endocrine deficiencies, and growth impairment. Additional late effects common to patients with high-risk disease include:
 - ▶ Chronic kidney disease
 - ▶ Impaired fertility
 - ▶ Cardiotoxicity
 - ▶ Neurocognitive impairment
 - ▶ Chronic lung disease
 - ▶ Second malignant neoplasms
- Screening for sensorineural hearing loss due to platinum chemotherapy and eflornithine is of particular importance. While the [COG Survivorship Guidelines](#) outline the minimum recommended frequency for audiologic assessment, children with impaired hearing should be referred to institutional audiology and/or otolaryngology teams to determine the appropriate schedule for subsequent evaluations.
- Patients treated for high-risk neuroblastoma are at risk for neurocognitive impairment. Risk appears to be impacted by chronic health conditions and specific treatments received.¹
- Fertility preservation is an option for some patients with high-risk disease. When possible, referral to fertility specialists for further discussion prior to initiation of chemotherapy is recommended.

Non-High-Risk Disease

- Patients who receive chemotherapy for non-high-risk disease typically receive lower cumulative doses of chemotherapeutic agents and limited to no external beam radiotherapy, and are therefore at lower risk of long-term toxicities compared to patients with high-risk disease. However, patients with non-high-risk disease constitute a heterogeneous group, and there is a wide range of cumulative chemotherapeutic exposures.²

Footnote

^a Consider audiologic assessment during key periods for speech development, depending on exposure to ototoxic agents.

References

¹ Hesko C, Liu W, Srivastava DK, et al. Neurocognitive outcomes in adult survivors of neuroblastoma: A report from the Childhood Cancer Survivor Study. *Cancer* 2023;129:2904-2914.

² Friedman DN, Goodman PJ, Leisenring WM, et al. Long-term morbidity and mortality among survivors of neuroblastoma diagnosed during infancy: A report from the Childhood Cancer Survivor Study. *J Clin Oncol* 2023;41:1565-1576.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF PARANEOPLASTIC SYNDROMES

Signs and Symptoms of Paraneoplastic Syndromes

- Presence does not imply metastases or incurability

Gastrointestinal

Vasoactive Intestinal Peptide Tumor (VIPoma) Syndrome

- Physiology: Due to tumor secretion of vasoactive intestinal peptide (VIP)
- Incidence: Present in <1% of patients with neuroblastoma
- Diagnosis: Elevated VIP, typically low potassium
- Symptoms:
 - ▶ Chronic watery diarrhea
 - ▶ Weight loss
 - ▶ Electrolyte abnormalities
- Treatment:
 - ▶ Supportive care including fluid and electrolyte replacement
 - ▶ Surgery when feasible
 - ▶ Other tumor-directed therapy per risk classification if resection is not feasible
- Symptoms usually resolve with surgical resection/successful anti-tumor therapy

Neurologic

Opsoclonus-Myoclonus-Ataxia Syndrome (OMAS)

- Physiology: Autoimmunity directed against cerebellum/brainstem
- Incidence: Present in 2% of patients with neuroblastoma
- Diagnosis:
 - ▶ Primarily clinical with supporting evidence of anti-Hu/ANNA1 antibodies¹
 - ▶ Cerebrospinal fluid (CSF) studies may show: Pleocytosis, increased oligoclonal bands, increased neopterin, increased B cells
- Symptoms:
 - ▶ Irregular multidirectional eye movements
 - ▶ Myoclonic jerking of limbs

Neurologic (continued)

- ▶ Ataxia
- ▶ Disordered mood/behavior, including irritability
- Treatment
 - ▶ Urgent neurology referral
 - ◊ Best neurologic outcomes have been observed following addition of immunomodulators to standard anti-tumor treatment or resection
 - ▶ Treatment options²⁻⁴ include: steroids (prednisone/adrenocorticotrophic hormone [ACTH]/dexamethasone), intravenous immunoglobulin (IVIG),³ rituximab
 - ◊ IVIG was studied in a randomized trial³
 - ◊ A stepwise approach may be appropriate
 - ◊ Treat neuroblastoma according to risk classification
 - ◊ Early initiation improves outcomes
 - ▶ Multidisciplinary services
 - ◊ Ongoing physical therapy/occupational therapy
 - ◊ Neuropsychology referral¹
 - ▶ Neurologic symptoms often persist despite anti-tumor treatment
 - ◊ Extended multidisciplinary care is often necessary

Lambert-Eaton Myasthenic Syndrome

- Physiology: Autoimmunity directed against calcium channels at neuromuscular junctions
- Incidence: Rarely associated with neuroblastoma
- Diagnosis: Clinical
- Symptoms:
 - ▶ Proximal lower extremity weakness
 - ▶ Loss of reflexes
 - ▶ Dysautonomia
- Treatment
 - ▶ Supportive care
 - ▶ Neuroblastoma-directed therapy according to risk classification

¹ Rossor T, Yeh EA, Khakoo Y, et al. Diagnosis and management of opsoclonus-myoclonus-ataxia syndrome in children: An international perspective. *Neurol Neuroimmunol Neuroinflamm* 2022;9:e1153.

² Patel A, Fischer C, Lin YC, et al. Treatment and revaccination of children with paraneoplastic opsoclonus-myoclonus-ataxia syndrome and neuroblastoma: The Memorial Sloan Kettering experience. *Pediatr Blood Cancer* 2020;67:e28319.

³ de Alarcon PA, Matthay KK, London WB, et al. Intravenous immunoglobulin with prednisone and risk-adapted chemotherapy for children with opsoclonus myoclonus ataxia syndrome associated with neuroblastoma (ANBL00P3): a randomised, open-label, phase 3 trial. *Lancet Child Adolesc Health* 2018;2:25-34.

⁴ Kumar P, Willard VW, Embry L, et al. Late cognitive and adaptive outcomes of patients with neuroblastoma-associated opsoclonus-myoclonus-ataxia-syndrome: A report from the Children's Oncology Group. *Pediatr Blood Cancer* 2024;71:e31039.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

EVIDENCE BLOCKS FOR SYSTEMIC THERAPY FOR NEUROBLASTOMA

Therapy for Intermediate-Risk Neuroblastoma

Primary Treatment	
COG ANBL0531 - Carboplatin/Cyclophosphamide/Doxorubicin/Etoposide	
SIOPEN - Carboplatin/Etoposide alternating with Cyclophosphamide/Doxorubicin/Vincristine	

Subsequent-Line Therapy	
Cyclophosphamide + topotecan (following COG ANBL0531)	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



EVIDENCE BLOCKS FOR SYSTEMIC THERAPY FOR NEUROBLASTOMA

Therapy for High-Risk Neuroblastoma

Induction Therapy	
COG ANBL12P1	
COG ANBL1531	
COG ANBL0532	

Consolidation Therapy	
Thiotepa, Cyclophosphamide, Carboplatin, Etoposide, Melphalan [tandem high-dose chemotherapy with autologous stem cell rescue]	
CEM (Carboplatin, Etoposide, Melphalan) [single high-dose chemotherapy with autologous stem cell rescue]	
BuMel (Busulfan, Melphalan) [single high-dose chemotherapy with autologous stem cell rescue]	

Chemoimmunotherapy	Post-Induction	Post-Consolidation
ANBL1221 - Irinotecan/Sargramostim/Temozolomide + Dinutuximab		
Irinotecan/Sargramostim/Temozolomide + Naxitamab-gqgk		

Post-Consolidation Therapy	
Isotretinoin/Sargramostim + Dinutuximab	

Continuation Therapy (after Post-Consolidation)	
Eflornithine	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

INTERNATIONAL NEUROBLASTOMA RISK GROUP (INRG) STAGING

Descriptions of INRG Tumor Stages¹

Tumor Stage	Description
L1	Localized tumor not involving vital structures, as defined by the list of IDRFs, and confined to one body compartment
L2	Local-regional tumor with presence of one or more IDRFs
M	Distant metastatic disease (except stage MS tumor)
MS	Metastatic disease in children younger than 18 months, with metastases confined to skin, liver, and/or bone marrow (<10% involvement)

¹ Brisse HJ, McCarville MB, Granata C, et al. Guidelines for Imaging and Staging of Neuroblastic Tumors: Consensus Report from the International Neuroblastoma Risk Group Project. Radiology 2011;261:243-257.



ABBREVIATIONS

4D-CT	four-dimensional computed tomography	ICCR	International Collaboration on Cancer Reporting	NGS	next-generation sequencing
ACTH	adrenocorticotrophic hormone	IDRF	image-defined risk factor	OAR	organ at risk
ALT	alanine aminotransferase	IHC	immunohistochemistry	OMAS	opsoclonus-myoclonus-ataxia syndrome
BSA	body surface area	IMRT	intensity-modulated radiation therapy	OS	overall survival
CBC	complete blood count	INPC	International Neuroblastoma Pathology Classification	PD	progressive disease
COG	Children's Oncology Group	INR	international normalized ratio	PR	partial response
CR	complete response	INRC	International Neuroblastoma Response Criteria	PT	prothrombin time
CSF	cerebrospinal fluid	INRG	International Neuroblastoma Risk Group	PTV	planning target volume
CTV	clinical target volume	INRGSS	INRG Staging System	RECIST	Response Evaluation Criteria in Solid Tumors
DI	DNA index	INSS	International Neuroblastoma Staging System	SCA	segmental chromosomal aberration
ECG	electrocardiogram	ITV	internal target volume	SD	stable disease
EFS	event-free survival	IVIG	intravenous immunoglobulin	SIOPEN	Society of Pediatric Oncology Europe Neuroblastoma
FDG	fluorodeoxyglucose	LDH	lactate dehydrogenase	SPECT	single-photon emission computed tomography
FFPE	formalin-fixed paraffin-embedded	LH	luteinizing hormone	TSH	thyroid-stimulating hormone
FH	favorable histology	mCTV	metastatic clinical target volume	UH	unfavorable histology
FISH	fluorescence in situ hybridization	MD	minimal disease	VGPR	very good partial response
FSH	follicle-stimulating hormone	MGPT	multigene panel testing	VIP	vasoactive intestinal peptide
GFR	glomerular filtration rate	mGTV	metastatic gross tumor volume	VIPoma	vasoactive intestinal peptide tumor
GTV	gross tumor volume	MIBG	meta-iodobenzylguanidine	VMA	vanillylmandelic acid
H&E	hematoxylin and eosin	MKI	mitosis and karyorrhexis index		
H&P	history and physical	mPTV	metastatic planning target volume		
HbA1C	hemoglobin A1c	MR	minor response		
HVA	homovanillic acid				



NCCN Guidelines Version 2.2026

Neuroblastoma

NCCN Evidence Blocks™

NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 2.2026

Neuroblastoma

Discussion

This discussion corresponds to the NCCN Guidelines for Neuroblastoma.
The discussion was last updated on April 28, 2026.

Table of Contents

Overview.....MS-2

Guidelines Development and Update Methodology.....MS-2

Literature Search Criteria.....MS-2

Sensitive/Inclusive Language.....MS-3

Genetic Risk Factors.....MS-3

Clinical Presentation.....MS-3

Workup.....MS-4

Pathology.....MS-6

Imaging.....MS-8

Staging.....MS-9

Risk Classification.....MS-9

Treatment for Patients with Newly Diagnosed Neuroblastoma.....MS-10

Response Assessment.....MS-18

Disease Surveillance.....MS-18

Treatment for Patients with Relapsed/Refractory Neuroblastoma...MS-20

Monitoring for Late Effects Related to Neuroblastoma Treatment...MS-21

Summary.....MS-22

References.....MS-23



NCCN Guidelines Version 2.2026

Neuroblastoma

Overview

Neuroblastoma is a cancer that originates from the developing sympathetic nervous system and is the most common extracranial solid tumor in children.^{1,2} It has been estimated that 600 to 800 patients are diagnosed with neuroblastoma annually in the United States.³ The prevalence is approximately 1 per 7000 live births.² The average age of a patient at the time of a neuroblastoma diagnosis is between 1 and 2 years of age.³ The vast majority of individuals are <5 years of age at diagnosis.^{3,4}

Neuroblastoma is a primitive neoplasm of neuroectodermal origin composed of neuroblasts (or immature nerve cells).⁵ These tumors may occur anywhere in the sympathoadrenal neuroendocrine system including the adrenal gland, connective/soft tissue, retroperitoneum, and mediastinum.⁵

Due to differences in disease severity, symptoms, and clinical behavior of the tumor, neuroblastoma is considered a complex and heterogeneous disease.^{1,6} For example, some patients have tumors that spontaneously regress without any treatment, while others are diagnosed with aggressive metastatic disease that requires multimodal intervention.⁶

In addition to disease burden and treatment-related side effects, some patients with neuroblastoma may face additional challenges. Limited data suggest that health disparities due to race/ethnicity and socioeconomic status exist among patients with neuroblastoma.⁶⁻⁹ For example, a recent study found that the 5-year survival was higher for white (80.7%) or Hispanic (80.8%) patients with neuroblastoma compared with their Black counterparts (72.6%).⁹

These NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Neuroblastoma include up-to-date recommendations for

the treatment of patients with neuroblastoma. These Guidelines were developed by a multidisciplinary Panel of representatives from NCCN Member Institutions with neuroblastoma-focused expertise in the fields of pediatric oncology, surgical oncology, radiation oncology, pathology, and radiology. Treatment recommendations for neuroblastoma in these Guidelines will continue to be updated by the NCCN Neuroblastoma Panel annually based on consensus and emerging clinical evidence.

Guidelines Development and Update Methodology

The complete details of the Development of the NCCN Guidelines® are available at www.NCCN.org.

Literature Search Criteria

Prior to the development of the NCCN Guidelines for Neuroblastoma®, an electronic search of the PubMed database was performed to obtain key literature in neuroblastoma published since 2012, using the search term “neuroblastoma.” The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.¹⁰ The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trials; Meta-Analysis; Systematic Reviews; and Validation Studies. The data from key PubMed articles as well as articles from additional sources deemed as relevant to these Guidelines as discussed by the Panel have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel’s review of lower-level evidence and expert opinion.



NCCN Guidelines Version 2.2026

Neuroblastoma

Sensitive/Inclusive Language

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation.¹¹ NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-weight-biased; and inclusive of individuals of all sexual orientations and gender identities. NCCN Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will continue to use the terms men, women, female, and male when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Genetic Risk Factors

Although cases of familial neuroblastoma have been reported, they are rare (estimated to be about 1%–2% of all cases).^{12,13} Among the cases of familial neuroblastoma that do occur, germline gain-of-function *ALK* mutations and loss-of-function *PHOX2B* mutations have been identified as causative factors.¹⁴⁻¹⁶ Certain genetic syndromes are also associated with neuroblastoma, such as Li-Fraumeni syndrome, and those caused by mutations in RAS pathway genes, including Costello syndrome, Noonan syndrome, and neurofibromatosis.¹³ Other genetic variants likely contribute to familial neuroblastoma or predispose patients to sporadic neuroblastoma; however, additional validation studies are needed.^{12,17,18}

Clinical Presentation

Symptoms associated with neuroblastoma can vary depending on the location of the tumor. Patients with neuroblastoma most commonly present with an abdominal mass or abdominal distension.^{2,19} Other signs and symptoms of the disease can include loss of appetite, weight loss, irritability, constipation, fever, hypertension, anemia, paralysis, bruising or swelling around the eyes, bone pain, and pancytopenia.

Paraneoplastic Syndromes

Paraneoplastic syndromes may also occur in a small subgroup of patients with neuroblastoma.^{5,20-24} Paraneoplastic syndromes associated with neuroblastoma are rare; symptoms generally result from an autoimmune response induced by tumors or from tumor secretion of peptides or hormones.²⁵ The NCCN Panel notes that the presence of a paraneoplastic syndrome does not imply metastases or incurability of the disease.²⁶⁻²⁸

Opsoclonus-myoclonus-ataxia syndrome (OMAS) is the most common paraneoplastic syndrome in children, occurring in approximately 2% of patients.²⁹ Symptoms include irregular eye movements, myoclonic jerking of limbs, ataxia, and disordered mood and behavior.^{30,31}

The optimal treatment strategy for OMAS has not been established, although international consensus guidelines have been developed.²¹ Best neurologic outcomes have been observed following addition of immunomodulators to standard anti-tumor treatment or resection.³² Treatment options include steroids (prednisone/adrenocorticotrophic hormone (ACTH)/dexamethasone), intravenous immunoglobulin (IVIG), and rituximab.^{28,33,34} Of note, the efficacy of IVIG as a treatment option for OMAS has been evaluated in randomized, phase 3 ANBL00P3 clinical trial.²⁸ In this study, the addition of IVIG to prednisone and risk-adapted chemotherapy (IVIG+) was evaluated against no IVIG (NO_IVIG) in 53 patients with neuroblastoma-associated OMAS. Patients randomized to



NCCN Guidelines Version 2.2026

Neuroblastoma

IVIG+ had significantly higher OMA response rates compared to the NO-IVIG group (80.8% vs 40.7%, respectively). The 3-year overall survival rate amongst all patients regardless of treatment type was 98.0%.

Early initiation of neuroblastoma treatment according to risk classification is recommended in patients with neuroblastoma-associated OMAS to improve patient outcomes. Neurological symptoms associated with OMAS may persist despite anti-tumor treatment.^{32,35} A stepwise approach to OMAS treatment may be appropriate.²¹ Following OMAS diagnosis, urgent neurology referral is recommended, as well as coordination of multidisciplinary care (including ongoing occupational and physical therapy, and neuropsychology referral).

Lambert-Eaton myasthenic syndrome, caused by antibodies directed against voltage-gated calcium channels, is another rare neurological paraneoplastic syndrome that may occur with neuroblastoma.³⁶⁻³⁸ Affected patients may present with proximal lower extremity weakness, loss of reflexes, and dysautonomia. Treatment for Lambert-Eaton myasthenic syndrome typically consists of supportive care and neuroblastoma-directed therapy according to risk classification.

Vasoactive intestinal peptide tumor (VIPoma) syndrome is a gastrointestinal paraneoplastic syndrome of neuroblastoma, occurring in <1% of patients.^{24,27,37} It is characterized by chronic watery diarrhea, weight loss, and electrolyte disturbances resulting from tumor secretion of vasoactive intestinal peptide (VIP). Treatment for VIPoma should include supportive care, surgery (when feasible), and other tumor-directed therapy per risk classification (if resection is not feasible).

For additional information on the evaluation and treatment of paraneoplastic syndromes, see the *Principles of Paraneoplastic Syndromes* in the algorithm.

Workup

The workup for suspected neuroblastoma in a patient outside of the perinatal period as recommended by the NCCN Neuroblastoma Panel includes a combination of tissue sampling for diagnostic evaluation and additional evaluations that involve a complete history and physical (H&P), family history assessment, laboratory tests, and imaging.^{1,19} Histologic evaluation and molecular characterization of the neuroblastoma tumor upon diagnosis are also essential for tumor staging, risk classification, and treatment selection in most (but not all) patients.^{1,6,39,40}

Diagnostic Workup

Based on international consensus, one of the following criteria must be met for a definitive diagnosis of neuroblastoma: 1) an unequivocal pathologic diagnosis made from tumor tissue by light microscopy; OR 2) bone marrow aspirate or trephine biopsy containing unequivocal tumor cells (eg, syncytia or immunocytoologically positive clumps of cells) and increased levels of urinary catecholamine metabolites.⁴¹

A neuroblastoma diagnosis should be made primarily based on tissue sampling (see *Diagnostic Workup* in the algorithm). For patients outside of the perinatal period, the NCCN Neuroblastoma Panel recommends that surgical resection should be considered in the setting of localized disease, particularly in the absence of image-defined risk factors (IDRFs; for a description, see the *Principles of Imaging* in the algorithm). When biopsy rather than upfront resection is indicated, minimally invasive biopsy techniques can be used; however, open biopsy may be preferred in some clinical scenarios in which adequate tissue cannot be obtained by less invasive means.

Multiple core biopsies may provide adequate tissue; clinicians should consider the amount of tissue needed for full histologic and molecular evaluation. It is crucial that an adequate amount of tissue be obtained to



NCCN Guidelines Version 2.2026

Neuroblastoma

evaluate the status of key prognostic factors needed for risk classification and treatment selection. Once it has been confirmed that sufficient tissue for clinical purposes has been obtained, preservation of additional tissue for research is encouraged when possible. Fine-needle aspiration (FNA) is not recommended by the Panel. It is important that an experienced pathologist review frozen sections to determine the adequacy of the specimen, as some samples may be necrotic at the time of initial biopsy.

Bilateral bone marrow aspirates and trephine biopsies can be used for diagnostic purposes, particularly in rare cases in which marrow is the only source of tumor material. In these situations, every attempt should be made to obtain adequate material for the full complement of molecular testing.

Patients <2 months of age with existing or evolving hepatomegaly⁴² and infants in whom safety considerations (coagulopathy, impending organ failure) preclude biopsy should not undergo a biopsy until after initiation of therapy and clinical stabilization. In these cases, biopsy should be performed when it is safe to do so to obtain tissue for histologic evaluation and molecular testing. Patients <6 months of age with L1 adrenal tumors with maximum diameter ≤3.1 cm if solid or 5 cm if at least 25% cystic do not require an initial biopsy or resection based on data from a prospective multicenter cooperative group study.⁴³

See the *Pathology* section of this discussion for additional recommendations regarding tissue sampling, histologic classification, and molecular testing considerations.

Additional Workup

The NCCN Neuroblastoma Panel recommends that all patients undergo an H&P, with special attention given to the abdominal exam to evaluate for masses and organomegaly, as well as a thorough neurologic exam. Any

family history of neuroblastoma or other childhood cancers should be noted.

Several laboratory tests are considered essential as part of the workup for neuroblastoma. These include a complete blood count (CBC) with differential and a comprehensive metabolic panel. Urine catecholamine levels (homovanillic acid [HVA] and vanillylmandelic acid [VMA]) are elevated in the majority of patients with neuroblastoma.^{44,45} Analysis of urine HVA and VMA levels is required for diagnosis only if bone marrow is the only diagnostic tissue obtained.

Other tests and assessments have been deemed as useful in selected cases by the NCCN Neuroblastoma Panel. Clinicians may consider obtaining the prothrombin (PT)/international normalized ratio (INR) if the liver is involved or there is a concern for bleeding. Pregnancy tests should be done for all patients of childbearing potential. Analysis of lactate dehydrogenase (LDH) and ferritin levels can be considered in select cases. High levels of LDH and ferritin in patients with neuroblastoma have been associated with worse prognosis; however, neither of these are components of the risk classification criteria recommended by the NCCN Neuroblastoma Panel (see *Neuroblastoma Risk Classification* in the algorithm).^{46,47}

Some assessments are recommended by the Panel only if a certain type of treatment is indicated. For example, audiograms, echocardiograms, or electrocardiograms should be done if certain types of chemotherapy agents are to be administered. A referral to fertility specialists should be made for patients likely to receive extensive alkylator therapy to discuss possible options and timing for fertility preservation.

Imaging studies are also an essential component of the workup for neuroblastoma as imaging results will have a profound impact on tumor staging and risk classification, which will inform the treatment strategy



NCCN Guidelines Version 2.2026

Neuroblastoma

implemented for each patient. See the *Imaging* section below and the *Principles of Imaging* in the algorithm for additional information about imaging for patients with neuroblastoma.

Pathology

Sampling

Pathology workflows should be designed to facilitate histologic diagnosis with prognostic classification and molecular profiling as required for treatment. The NCCN Neuroblastoma Panel recommends a coordinated diagnostic sampling effort that includes pathologists, oncologists, surgeons, and radiologists.^{48,49}

For the pathology diagnosis of neuroblastoma, considerations for acceptable tissue sampling from either the primary site or a site of metastasis are based on the requirements for risk stratification and treatment. These include: surgical resection if clinically indicated, incisional biopsy (>1 cm³), or tissue cores (at least 10 cores, ideally 20–30 mm in length obtained using a 16-gauge needle if possible). Clinicians should be aware that bilateral bone marrow biopsies and clot sections alone may not be sufficient to assess characteristics relevant to the International Neuroblastoma Pathology Classification (INPC) system.⁵⁰

The tissue quantity requirements specified above may be challenging to meet. In some cases, open rather than minimally invasive procedures may be necessary to obtain the adequate specimens for all required testing.⁵¹ Advanced planning for the required amount and condition of the tissue (eg, formalin-fixed paraffin-embedded [FFPE], fresh, frozen, touch preparations) will increase the likelihood of completing the required testing. Additional testing may be required for patients enrolled in clinical trials.

Histologic Classification

Neuroblastoma diagnosis should be determined according to the International Collaboration on Cancer Reporting (ICCR) and INPC.^{39,40} Histologic classification based on the INPC should be done prior to initiation of therapy and is typically made based on hematoxylin and eosin (H&E) staining.

Immunohistochemical stains can be helpful in the setting of small samples, unusual locations, or undifferentiated subtypes. In these situations, the NCCN Panel recommends staining for chromogranin and synaptophysin, which are widely used markers for neuroendocrine cells and tumors.⁵² Staining for the neural crest marker PHOX2B (strongly recommended) and tyrosine hydroxylase are also appropriate for these situations.⁵³

Neuroblastoma is part of a group known as peripheral neuroblastic tumors, which are tumors of the sympathetic nervous system that arise from the embryonic neural crest.^{54,55} The INPC distinguishes peripheral neuroblastic tumors into four categories mainly based on Schwannian stroma development: neuroblastoma (Schwannian stroma-poor); ganglioneuroblastoma, intermixed (Schwannian stroma-rich); ganglioneuroma (Schwannian stroma-dominant); and ganglioneuroblastoma, nodular (composite, Schwannian stroma-rich/stroma-dominant and stroma-poor).^{39,40} Neuroblastoma is the most common type of neuroblastic tumor.

The INPC classifies peripheral neuroblastic tumors into a Favorable Histology group or Unfavorable Histology group. Classification of tumors is based on the grade of neuroblastic differentiation (undifferentiated subtype, poorly differentiated subtype, and differentiating subtype), MKI (Mitosis-Karyorrhexis Index = estimated mitotic and karyorrhectic nuclei per 5000 neuroblastoma cells; low <100, intermediate 100–200, and high ≥200), and age at diagnosis. Favorable Histology tumors include: 1) poorly



NCCN Guidelines Version 2.2026

Neuroblastoma

differentiated or differentiating subtype with low or intermediate MKI, ≤ 547 days of age at diagnosis; and 2) differentiating subtype with low MKI, ≤ 1824 days of age at diagnosis. Unfavorable Histology tumors include: 1) undifferentiated subtype at any age; 2) high MKI at any age; 3) poorly differentiated subtype with any MKI, ≥ 548 days of age at diagnosis; 4) differentiating subtype with intermediate MKI, ≥ 548 days of age at diagnosis; and 5) differentiating subtype with any MKI, ≥ 1825 days of age at diagnosis.

Tumors in the category of ganglioneuroblastoma, intermixed and ganglioneuroma, are usually diagnosed in older children and classified into the Favorable Histology group. Patients in these two categories are expected to have an excellent prognosis. Tumors in the category of ganglioneuroblastoma, nodular, are composed of at least two distinct histologies: one is neuroblastoma while the other is either ganglioneuroblastoma, intermixed or ganglioneuroma. The neuroblastoma component usually makes discrete nodules and dictates clinical behavior of the tumor in this category.⁵⁶ In order to determine the prognostic group (Favorable or Unfavorable Histology group) for tumors categorized as ganglioneuroblastoma, nodular, the same age-dependent criteria of morphologic features (grade of neuroblastic differentiation and MKI) as described in the neuroblastoma category (see above) should be applied to the neuroblastoma component.

See *Principles of Pathology* in the algorithm for more information about the histologic classification of neuroblastoma.

Biomarker Testing Considerations

Risk stratification for the initial treatment of neuroblastoma is heavily dependent on tumor biomarker profiling (see *Neuroblastoma Risk Classification* in the algorithm). The key prognostic biomarkers that should

be assessed include *MYCN* amplification status, segmental chromosomal aberration (SCA) status, and tumor cell ploidy.

Amplification of the *MYCN* oncogene is associated with aggressive disease, and is the strongest independent prognostic risk factor included in the neuroblastoma risk classification system.^{47,57,58} The NCCN Neuroblastoma Panel recommends the assessment of *MYCN* amplification status in all neuroblastomas and neuroblastoma nodules of ganglioneuroblastoma nodular tumors.

SCAs, defined as the loss or gain of a portion of a chromosome arm, may also be associated with inferior outcome in patients with neuroblastoma.^{47,59} The most extensively studied SCAs in neuroblastoma include loss of genetic material from chromosomes 1p and 11q; however, segmental losses involving 3p or 4p and segmental gains involving 17q, 1q, or 2p are also considered when assigning SCA status. The Panel recommends that biomarker testing include analysis of these seven SCAs for appropriate risk stratification.

Ploidy status, a measure of the amount of DNA within cells, is another prognostic factor for a small subset of patients.⁴⁷ In general, a DNA index (DI) = 1 is considered less favorable than DI > 1, and ploidy has historically been used in risk classification in infants.^{6,60} If information about ploidy is not provided, then risk can be assigned based on other available factors.

New biomarker-directed therapies for the treatment of neuroblastoma are emerging.^{61,62} For example, amplification and sequence variants in the *ALK* gene predict response to matched targeted agents in patients with relapsed/refractory disease.^{1,63-65} Biomarker testing requirements will continue to evolve as new targeted therapies become available.



NCCN Guidelines Version 2.2026

Neuroblastoma

Assay Selection

Multiple approaches using different assays can be used for biomarker testing to meet the above recommendations. Next-generation sequencing (NGS) has become widely available and analysis of FFPE tissue with this approach is feasible. Using NGS to simultaneously evaluate for the key neuroblastoma prognostic factors (including the presence of *MYCN* amplification and SCAs) and aberrations that may guide therapy selection (such as amplification or activating mutations in *ALK*) is recommended, as long as the approach enables robust assessment of copy number status and can provide coverage of the relevant regions of neuroblastoma-associated genes. Use of a single assay may be beneficial when tissue is limited.

The status of the standard prognostic biomarkers can also be assessed using fluorescence in situ hybridization (FISH), microarray, and/or flow cytometry; however, clinicians should be aware that these assays will not identify sequence variants in neuroblastoma-related genes such as *ALK*.

See the *Principles of Pathology* section in the algorithm for additional information about biomarker testing considerations.

Imaging

A neuroblastic tumor diagnosis is usually suspected based on the patient's age and the appearance of the tumor on imaging.⁵⁴ The combination of imaging studies to be used during the initial evaluation of a child with a suspected neuroblastic tumor depends on the symptoms and suspected sites of disease.⁶⁶ Local extension of neuroblastoma mainly consists of vascular encasement, infiltration of adjacent soft tissues and organs (most commonly the kidneys and liver), and infiltration of the foramina and epidural space of the spinal canal when the primary tumor arises from a paraspinal sympathetic chain.

Approximately 50% of patients present with localized or regional disease, and approximately 35% of patients have regional lymph node spread at time of diagnosis.⁶⁷ The International Neuroblastoma Risk Group (INRG) Task Force developed the INRG Staging System (INRGSS) and the INRG Risk Classification System for neuroblastoma, which have profound implications for imaging in this disease.^{47,55} Moreover, while soft tissue tumor volume was previously used to measure response in the prior International Neuroblastoma Response Criteria (INRC), the revised INRC use RECIST criteria for the measurement of soft tissue disease, using assessment of the single longest dimension of soft tissue lesions.^{68,69}

The primary goals of imaging in a patient with suspected neuroblastoma are to identify features consistent with a neuroblastic tumor,⁷⁰ assess for the presence of IDRFs to facilitate INRG staging (see the *Staging* section below),⁵⁵ help estimate the potential surgical risk that would be associated with local tumor excision, assess for the presence and degree of regional and distant metastatic disease, and facilitate post-treatment response assessment and disease surveillance.

The NCCN Neuroblastoma Panel recommends cross-sectional imaging (MRI with/without contrast or CT with contrast) to evaluate soft tissue disease at the time of initial evaluation. MRI of the spine with/without contrast is appropriate in the setting of paraspinal disease or if there are concerns regarding involvement of nerve roots or spinal cord. MRI of the brain with/without contrast or CT skull/orbits with contrast should be done if neurological symptoms are present or if otherwise clinically indicated.

Iodine-123 metaiodobenzylguanidine (I-123 MIBG) imaging should be used to assess for metastatic disease due to the high specificity and high sensitivity of this tracer. MIBG, a norepinephrine analog, is taken up by norepinephrine transporters; uptake has been demonstrated in up to 90% of neuroblastoma tumors.⁷¹ Thyroid protection is required per institutional



NCCN Guidelines Version 2.2026

Neuroblastoma

guidance. Novel radiotracers are in development and may serve a role in select patients.⁷²⁻⁷⁴

Interpretation of I-123 MIBG imaging is performed by means of semiquantitative scoring of tracer uptake within body segments. The modified Curie score⁷⁵ and the International Society of Pediatric Oncology Europe Neuroblastoma (SIOPEN) score⁷⁶ are the two most commonly used systems. In North America, the modified Curie score is used. If available, single-photon emission CT (SPECT) or SPECT/CT should be performed at sites of known or suspected disease to improve sensitivity and anatomic localization of disease sites.⁷⁷

Fluorodeoxyglucose (FDG)-PET imaging should be obtained in patients with I-123 MIBG non-avid disease or suspected mixed-avidity disease, with the exception of patients who are <6 months of age with L1 adrenal tumors ≤3.1 cm in diameter if solid or <5 cm if at least 25% cystic component.⁴³ FDG-PET/CT and PET/MRI can also be useful alternative or supplemental diagnostic tools when I-123 MIBG imaging and anatomic imaging do not correlate.⁷⁸ See *Principles of Imaging* in the algorithm for additional information.

Imaging is also an important component of post-treatment response assessment and disease surveillance. See the *Response Assessment* and *Disease Surveillance* sections for more information.

Staging

The NCCN Neuroblastoma Panel recommends using the neuroblastoma staging system developed by the INRG (see *Staging* in the algorithm).⁵⁵ This system was developed to improve the consistency and uniformity of tumor staging for patients with neuroblastoma by ensuring that disease staging could be done before initiation of treatment. Tumor stage defined by the INRGSS is also a key prognostic factor accounted for in the

updated Children's Oncology Group (COG) neuroblastoma risk classification system (see *Neuroblastoma Risk Classification* in the algorithm).⁴⁷ As risk classification is used to guide neuroblastoma treatment, it is essential for disease staging to be completed before initiation of treatment when possible. Initiation of emergent therapy, if needed, should not be delayed for I-123 MIBG or FDG-PET imaging; however, this imaging should be obtained as soon as possible.

In the INRGSS, localized tumors are classified based on the number of IDRFs present. While the currently used list of IDRFs is derived from a consensus list of risk factors initially developed to guide surgical decision-making, IDRFs are thought to be proxies for tumor biologic features that are not yet well understood. See *Principles of Imaging* in the algorithm for a description of IDRFs.

An L1 tumor is defined as a localized tumor that is not involved with vital structures, as defined by the list of IDRFs, and is confined to one body compartment. An L2 tumor is described as a locoregional tumor with the presence of one or more IDRFs. A patient with stage M neuroblastoma has distant metastatic disease, except for the special case of young children with INRG stage MS disease. MS neuroblastoma is diagnosed only in children who are <18 months, with metastases confined to the skin, liver, and/or bone marrow (<10% involvement).

Risk Classification

The risk classification criteria incorporated into the NCCN Guidelines is adapted from an updated risk classifier published by the COG in 2021.⁴⁷ The assigned risk group is based on outcomes data generated over recent decades.^{40,47,79-81} Event-free survival (EFS) and overall survival (OS) outcomes are influenced by key prognostic risk factors, including: age at diagnosis, INRG stage (see the *Staging* section above), tumor *MYCN* status (ie, presence or absence of *MYCN* amplification), histopathology



NCCN Guidelines Version 2.2026

Neuroblastoma

(favorable or unfavorable based on INPC), presence or absence of SCAs, and ploidy status (diploid or hyperdiploid). The Panel notes that if information about ploidy status is missing, risk should be assigned based on the other available factors. Ploidy may be inferred by NGS.

It should be noted that treatment substantially impacts outcome, and outcomes achieved often reflect the effects of different types of therapy within a given risk group. However, risk classification must occur in practice prior to the start of therapy. The risk assignments that follow therefore reflect pre-treatment decision-making, taking into account treatment administered to prior patients with neuroblastoma over time.

Based on this classification system, patients of any age with L1 disease with tumors that are *MYCN* non-amplified are included in the low-risk group. Patients with L1 disease and tumor *MYCN* amplification would only be considered low risk if complete resection is achieved. If residual tumor remains post-resection, patients with L1, *MYCN*-amplified tumors are considered to have high-risk disease. Asymptomatic patients who are <12 months of age with MS disease and Favorable Histology tumor that is demonstrated to be *MYCN* non-amplified, hyperdiploid, and without SCAs are also assigned to the low-risk group.

Patients who are between 12 and 18 months of age with M or MS stage disease with tumors that are found to have unfavorable prognostic factors such as Unfavorable Histology, presence of SCAs, or *MYCN* amplification are assigned to the high-risk group. Patients who are ≥18 months with M stage disease are also assigned to the high-risk group regardless of any other features. Apart from those with L1 disease and completely resected tumors (who are considered to have low-risk disease despite *MYCN* status), detection of *MYCN* amplification leads to assignment to the high-risk group regardless of age or stage.⁴⁷

The intermediate-risk group generally includes patients whose disease characteristics do not meet the criteria for either the low-risk or high-risk groups. See *Neuroblastoma Risk Classification* in the algorithm for the detailed Neuroblastoma Risk Classification table.

For symptomatic infants with INRG stage MS disease who are too ill to undergo biopsy at the time of initial presentation (such as those with coagulopathy or impending organ failure), and especially those with hepatomegaly leading to respiratory compromise,⁸² therapy is started with a presumptive assignment to the intermediate-risk group. However, if a biopsy undertaken when the patient is clinically stable identifies tumor cells with *MYCN* amplification, the patient would then be reassigned to the high-risk group.

Clinicians should be aware that there are differences between the COG risk classifier and criteria used by cooperative groups outside of North America.^{47,83,84} For example, patients with L2, *MYCN* non-amplified disease who are >18 months of age at diagnosis and have Unfavorable Histology or whose tumors are undifferentiated/poorly differentiated are classified as high risk by COG criteria, but may be categorized as intermediate risk by other groups.

For the complete criteria for each risk group, see *Neuroblastoma Risk Classification* and *Principles of Risk Classification* in the algorithm.

Treatment for Patients with Newly Diagnosed Neuroblastoma

Due to the complex nature of the disease, it is recommended that a multidisciplinary team be involved in making treatment decisions for patients with neuroblastoma. The team should include at a minimum: diagnostic radiologists, nuclear medicine physicians, interventional



NCCN Guidelines Version 2.2026

Neuroblastoma

radiologists, surgeons, anatomic pathologists, molecular pathologists, radiation oncologists, and pediatric oncologists.

Non–High-Risk Disease

Over the past two decades, the staging system, risk classification system, and response criteria definitions for neuroblastoma have evolved. It is important to note that published results from clinical trials for patients with non–high-risk disease utilized a legacy staging system (International Neuroblastoma Staging System [INSS]) and older response criteria⁴¹ or protocol-specific response criteria.⁴² These NCCN Guidelines are based on published data from clinical trials of patients with non–high-risk disease, but also bridge the transition to currently used staging, classification, and response criteria systems where applicable.^{47,68,83}

Approximately half of newly diagnosed patients with neuroblastoma have non–high-risk disease.^{85,86} Patients with low- and intermediate-risk (ie, non–high risk) neuroblastoma have excellent survival rates. Five-year survival rates are >95% for patients with low-risk neuroblastoma and approximately 90% to 95% for those with intermediate-risk neuroblastoma.^{3,47} The goal of therapy for these patients is to cure the disease with minimal toxicity. Recent clinical trials have focused on reduction of therapy for patients with favorable biology and have been successful in maintaining excellent outcomes with these strategies.^{42,87,88}

Treatment for Low-Risk Neuroblastoma

Treatment approaches for patients with low-risk INRG L1 tumors involve surgical resection, with the exception of those who are <6 months old with isolated adrenal masses with maximum diameter ≤3.1 cm if solid, or 5 cm if at least 25% of the mass is cystic. For these patients, observation without biopsy is the recommended approach. If upfront surgery will potentially obviate the need for chemotherapy and can be safely performed with minimal morbidity, a resection should be performed (see

Principles of Surgery in the algorithm). If the resection is incomplete, and *MYCN* amplification is detected, the patient should be reassigned to the high-risk group. In patients with INRG MS disease who are asymptomatic and have tumors with favorable biology, observation is also the preferred approach. For complete treatment recommendations for low-risk neuroblastoma, see *Low-Risk Disease* in the algorithm.

Treatment for Intermediate-Risk Neuroblastoma

Intermediate-risk treatment involves a combination of moderate-intensity multiagent chemotherapy and surgical resection (see *Principles of Systemic Therapy* and *Principles of Surgery* in the algorithm).

The North American treatment strategy for patients with intermediate-risk neuroblastoma is based on results of the COG study ANBL0531, which aimed to decrease the number of chemotherapy cycles given to patients with more favorable tumor biology, by allowing for a larger residual primary tumor following chemotherapy with or without surgery, compared to treatment endpoints utilized on legacy clinical trials.⁴² Patients were assigned to receive a minimum number of cycles based on age, stage, and tumor biologic features.

The NCCN Neuroblastoma Panel recommends that patients with intermediate-risk neuroblastoma receive 2 to 8 cycles of chemotherapy, with the number of cycles being dependent on a combination of factors (including disease stage, age, and biologic features; see *Intermediate-Risk Disease* in the algorithm). Favorable biologic features include Favorable Histology, DI >1, and no SCA. If SCA or histology status are unavailable, then clinicians should consider the tumor to have unfavorable biologic features. For infants with stage MS disease who are too unstable to undergo biopsy before starting treatment, the Panel recommends initiating chemotherapy immediately. This approach may be used for patients <2 months of age with existing or evolving hepatomegaly and infants in whom



NCCN Guidelines Version 2.2026

Neuroblastoma

safety considerations (such as coagulopathy, impending organ failure) preclude biopsy. An objective symptom scoring system has been created to aid clinicians in decision-making regarding initiation of therapy in very sick infants.⁸² A score ≥ 2 , indicating impairment of multiple organ systems, is associated with increased mortality, and warrants emergent chemotherapy.⁸² Once symptoms are resolved following initial therapy, biopsy should be performed, and risk assignment reassessed.

RT may be warranted in symptomatic patients with ongoing organ dysfunction whose disease has an inadequate response to initial chemotherapy. Multidisciplinary discussion is recommended. See *Principles of Radiation Therapy* in the algorithm for additional information.

In the ANBL0531 clinical trial, protocol-specific criteria for disease response were used, and primary tumor response was assessed based on tumor volume reduction. The goal for groups of patients with localized, favorable biology tumors was to achieve at least a 50% reduction in the primary tumor volume, while patients with localized disease with less favorable tumor biologic features were to continue therapy until a goal of 90% reduction in primary tumor volume (very good partial response [VGPR]) was achieved. The goal treatment endpoint for metastatic disease response for patients with the equivalent of INRG MS disease and those with intermediate-risk INRG stage M disease were also protocol-specific, and not identical to the legacy INRC.⁴¹

Since the completion of ANBL0531, the neuroblastoma response criteria have been updated and no longer include VGPR as a response category. Additionally, in the current system of response assessment, tumors are measured with a single dimension as per Response Evaluation Criteria in Solid Tumors (RECIST), rather than by tumor volume (see *Response Assessment* in the algorithm).⁶⁸ Until additional data become available regarding the use of the revised response criteria in assessment of disease in patients with non–high-risk neuroblastoma, the committee

supports utilizing either volume or 1-dimensional assessments of primary tumor response in this group of patients.

After completion of the assigned number of chemotherapy cycles, the NCCN Neuroblastoma Panel recommends surveillance if the targeted tumor reduction goal was achieved (see *Intermediate-Risk Disease* in the algorithm). If the targeted tumor reduction goal with the initial course of chemotherapy regimen was not achieved, a multidisciplinary discussion regarding the role of additional therapy, such as surgery or additional chemotherapy, should be undertaken on an iterative basis, and surveillance should only begin once the target response or reduction in primary tumor size, as noted in the Guidelines, is achieved.

For some patients, surgical resection may be appropriate to reach the targeted treatment endpoint. Surgical resection should be considered if chemotherapy has resulted in $<50\%$ reduction in tumor size. The preservation of vital structures and of end-organ function is of utmost importance in the intermediate-risk context, as less than complete response has been shown to be an acceptable endpoint of therapy for patients with localized intermediate-risk tumors. The timing of resection will depend on the response to initial therapy and subsequent assessment of surgical risk. Multidisciplinary discussion regarding the optimal timing of resection should occur.

If surgery cannot be performed safely to achieve the proposed degree of tumor reduction, additional chemotherapy may be given with re-evaluation after every 2 cycles. At these timepoints, the potential risks and benefits of additional chemotherapy or surgery can be further discussed by the multidisciplinary treatment team.

In some circumstances it may be reasonable to consider biopsy of the residual mass to assess for histologic differentiation, which may support observation of a tumor that does not shrink sufficiently with chemotherapy



NCCN Guidelines Version 2.2026

Neuroblastoma

and for which a surgical debulking is considered unsafe. In COG ANBL0531, cyclophosphamide and topotecan were used as additional treatment for patients who did not achieve the target response with 8 cycles of the intermediate-risk therapy regimen outlined in *Intermediate-Risk Disease* in the algorithm.⁴² SIOPEN treatment regimens can also be considered, as similar outcomes are achieved with the SIOPEN and COG strategies.⁸⁹

For complete treatment recommendations for intermediate-risk neuroblastoma, see *Intermediate-Risk Disease* in the algorithm.

High-Risk Disease

Approximately half of newly diagnosed patients with neuroblastoma have high-risk disease.^{85,86} Five-year overall survival rates are approximately 60% for patients with high-risk neuroblastoma.^{47,90} These outcomes have improved over time as a result of increasingly intensive multimodal therapies divided into induction, consolidation, and post-consolidation phases. Much of the data summarized below is derived from cooperative group phase 3 trials and the Panel encourages participation in open clinical trials when available. For complete treatment recommendations for patients with high-risk neuroblastoma, see *High-Risk Disease* in the algorithm.

Induction Therapy

The goal of initial induction therapy is to decrease disease burden and to achieve the best possible response prior to subsequent phases of therapy. This goal is achieved through a combination of multiagent cytoreductive chemotherapy and surgical resection of the primary tumor and locoregional disease (see *High-Risk Disease* in the algorithm). In addition, autologous peripheral blood stem cells are collected during induction and aliquoted in a way that facilitates subsequent therapy.⁹¹

There are no contemporary data comparing North American induction regimens that have been evaluated in prospective randomized trials. Instead, a number of induction combinations have evolved over the last several decades, mainly based upon cisplatin- and alkylator-intensive regimens developed at Memorial Sloan-Kettering Cancer Center (MSKCC).⁹² Several different regimens have been used in North American cooperative group pilot and Phase 3 trials, each enrolling over 100 patients over the last 2 decades.⁹³

These regimens yield broadly similar end-induction response rates, with approximately 80% of patients having a partial response or better and approximately 9% of patients progressing despite these intensive regimens.⁹³ Given the lack of prospective comparative efficacy data and similar response rates, toxicity considerations have driven the evolution of induction regimens to reduce exposure to nephrotoxic and cardiotoxic agents. The combination of topotecan and cyclophosphamide given as cycles 1 and 2 of induction was studied initially as part of a pilot trial that demonstrated the feasibility and acceptable end-induction response rate (84%) of this approach.⁹⁴ This regimen was adopted for use in ANBL0532, demonstrating that 39.1% of patients had a partial response or better after the first 2 cycles with topotecan and cyclophosphamide.⁹⁵

A subsequent trial, ANBL12P1, empirically reduced induction to 5 cycles, with an 80% end-induction response rate.⁹⁶ Likewise, data from MSKCC demonstrated comparable end-induction response rates with 5 versus 7 induction cycles.⁹² Therefore, a 5-cycle induction was included as part of the standard arm of the COG phase 3 trial, ANBL1531. The ANBL1531 induction regimen was similar to the published ANBL12P1 regimen, with minor adjustments to dosing in order to align with updated COG chemotherapy standards. Given extensive contemporary experience with a 5-cycle induction regimen, the NCCN Neuroblastoma Panel recommends either ANBL12P1 or ANBL1531 induction as preferred



NCCN Guidelines Version 2.2026

Neuroblastoma

regimens, with a 6-cycle regimen from ANBL0532 as an additional acceptable regimen (see *Principles of Systemic Therapy* in the algorithm). However, the NCCN Neuroblastoma Panel acknowledges the lack of comparative data, and notes that other published chemotherapy regimens that achieve a similar end-induction response rate could be considered as reasonable alternatives for individual patients.

Several strategies to improve induction are under investigation at this time, including early addition of ALK inhibitors for patients whose tumors harbor *ALK* aberrations.⁹⁷ In addition, interventions such as the early administration of I-131 MIBG or anti-GD2 monoclonal antibody therapy are being evaluated.⁹⁷ As robust safety and efficacy data are not yet available to support these approaches, the Panel does not recommend adoption of these approaches outside the context of clinical trials at this time.

Surgical resection of the primary tumor and associated locoregional adenopathy is another important goal of induction therapy (see *Principles of Surgery* in the algorithm for additional information). Given the aggressive nature of these tumors, upfront resection is rarely feasible and the NCCN Neuroblastoma Panel recommends surgical resection after several cycles of initial cytoreductive chemotherapy. Even after initial chemotherapy, resection with negative margins is rarely feasible and is not the recommended surgical goal. Instead, two large analyses have demonstrated improved EFS and lower local relapse/progression rates in patients who had >90% resection (North American experience) or a complete macroscopic resection (European experience).^{98,99} The NCCN Panel therefore recommends this degree of resection, broadly considered a gross total resection, as the goal of primary site surgery. When vital organs, major nerves, and/or major blood vessels would be threatened or would require resection to achieve this goal, the Panel recommends subtotal resection.

The NCCN Neuroblastoma Panel recommends full disease reassessment at the end of induction. This end-induction evaluation is a critical decision point in the treatment of patients with high-risk neuroblastoma. Prior analyses have demonstrated that patients with poor end-induction response (less than partial response) have inferior outcomes compared to patients with more favorable end-induction response (partial response or better).⁹³ However, analyses of randomized trials evaluating different consolidation strategies support a potential role for modern consolidation therapies even in patients with less than a partial response to induction. For example, the CCG-3891 trial reported higher EFS for patients with less favorable end-induction response who were randomized to transplant (vs continued chemotherapy).¹⁰⁰ Likewise, the ANBL0532 trial (see below) reported a benefit of tandem transplant (vs single transplant) independent of end-induction response.⁹⁵

The NCCN Neuroblastoma Panel recommends proceeding to consolidation therapy for patients with partial response or better to induction, though the Panel acknowledges that bridging therapy to improve response may be appropriate depending upon the nature of the partial response.¹⁰¹ Patients with progressive disease during or at the end of induction have not typically been candidates to proceed with consolidation therapy, and the Panel endorses this approach. Instead, the Panel recommends non-myeloablative therapies for these patients, including a chemoimmunotherapy regimen combining anti-GD2 monoclonal antibody with chemotherapy^{102,103} or participation in clinical trials for patients with first relapse.

Patients with end-induction minor response or stable disease require individualized decision-making. For patients with end-induction minor response or stable disease not proceeding to consolidation therapy, the Panel recommends a chemoimmunotherapy regimen combining anti-GD2 monoclonal antibody with chemotherapy or participation in clinical trials for



NCCN Guidelines Version 2.2026

Neuroblastoma

patients with refractory disease. Retrospective data suggest that proceeding to consolidation therapy may be appropriate for patients with an inadequate response to standard induction therapy whose disease responds to alternative “bridging” therapies.¹⁰¹ Specifically, patients with a complete response to bridging therapy who proceeded to consolidation had favorable outcomes.

Consolidation Therapy

A standard consolidation phase includes both high-dose chemotherapy with autologous stem cell rescue and consolidative radiotherapy to the primary site. In North America, it is also considered standard to administer radiotherapy to sites of residual metastatic disease remaining at the end-induction disease evaluation.

High-dose chemotherapy with autologous stem cell rescue has been a hallmark of high-risk neuroblastoma therapy since a series of randomized trials demonstrated improved outcomes with this approach compared with continued conventional chemotherapy.¹⁰⁴ The NCCN Neuroblastoma Panel acknowledges that these randomized trials were conducted in a treatment era that preceded routine use of anti-GD2-directed immunotherapy and that additional work is needed to understand if subgroups of patients might benefit from consolidative approaches that do not rely on high-dose chemotherapy with autologous stem cell rescue. For example, a limited institution retrospective experience suggests that similar OS rates may be achieved with or without high-dose chemotherapy among patients with greater than a partial response in the era of anti-GD2 immunotherapy.¹⁰⁵

For patients who are candidates for consolidation therapy, the NCCN Neuroblastoma Panel recommends tandem transplantation with two consecutive rounds of high-dose chemotherapy with autologous stem cell rescue for most patients with high-risk disease (a category 1

recommendation). This recommendation is based on data from the COG ANBL0532 randomized phase 3 trial.⁹⁵ Patients without progressive disease after a 6-cycle induction were eligible for randomization to a single transplant with full-dose carboplatin/etoposide/melphalan (CEM) or to tandem transplant with thiotepa/cyclophosphamide followed 6 to 10 weeks later with dose-reduced CEM. Patients randomized to the tandem transplant arm had significantly improved EFS (3-year EFS 61.6% vs 48.4% for single transplant).

There are two less common subgroups of patients with high-risk disease for whom a single round of high-dose chemotherapy with autologous stem cell rescue may be appropriate: 1) patients with stage L2, ≥ 18 months at diagnosis, Unfavorable Histology, AND *MYCN* non-amplified disease; and 2) patients with stage M, 12 to < 18 months at diagnosis, *MYCN* non-amplified, with any of the following other unfavorable features: Unfavorable Histology; diploid DNA content; and/or presence of SCAs. Patients in these two groups have historically had more favorable outcomes compared to patients with high-risk disease due to *MYCN* amplification or patients with high-risk disease due to age ≥ 18 months at time of diagnosis of stage M disease. For example, in a large series from the COG, patients in these two groups had 5-year EFS rates of approximately 75% to 80%.⁴⁷ Patients in these two more favorable subgroups were non-randomly assigned to single transplant with full-dose CEM in ANBL0532, and the Panel endorses this approach. In other contexts, these patients are treated differently.^{60,84,106-108}

The combination of busulfan and melphalan (BuMel) is a preferred conditioning regimen in Europe based upon results of a randomized phase 3 trial that showed superior EFS with BuMel compared to CEM following the European rapid COJEC induction regimen.¹⁰⁹ In addition, the BuMel regimen was associated with lower rates of most adverse events, though the risk of sinusoidal obstruction syndrome was higher. The COG



NCCN Guidelines Version 2.2026

Neuroblastoma

conducted a pilot trial, ANBL12P1, that demonstrated the feasibility of this approach,⁹⁶ but the role of BuMel in the context of North American therapy is not currently defined. Single transplant with BuMel may be an appropriate regimen for patients with a contraindication to tandem transplant or for patients in subgroups for which single transplant is recommended.

Radiation therapy (RT) to the primary tumor is typically administered upon recovery from high-dose chemotherapy with stem cell rescue.

Neuroblastoma is a radiosensitive tumor, and a commonly used dose of 21.6 Gy is recommended. Recent national trials have attempted to improve local control by augmenting radiotherapy.^{110,111} In COG trial A3973, a subset of patients had primary site radiotherapy fields extended to include uninvolved draining nodal stations. As these patients had similar local relapse/progression rates compared to patients treated without extending the radiotherapy field, this approach is not recommended by the Panel.¹¹¹ In COG trial ANBL0532, patients with gross residual tumor following primary site resection received a boost of 14.4 Gy to gross residual tumor.¹¹⁰ This augmented dose did not improve local relapse/progression or EFS rates compared to the historic controls. Based upon these data, the Panel does not recommend either strategy.

The NCCN Neuroblastoma Panel recommends radiation to sites of residual metastatic disease remaining by I-123 MIBG or FDG-PET (if MIBG non-avid) at the end-induction disease evaluation, recognizing that not all sites may be feasibly targeted by external beam RT (EBRT). While repeat imaging following recovery from transplant and prior to radiation is not routinely performed, it may aid in decision making in select clinical scenarios. This recommendation is based upon limited institution data supporting a benefit from this approach,^{112,113} though this approach has not been adopted internationally. See *Principles of Radiation Therapy* in the algorithm for additional information.

Post-Consolidation Therapy

The NCCN Neuroblastoma Panel recommends post-consolidation therapy containing an anti-GD2 antibody with sargramostim and isotretinoin for those who do not experience disease progression after consolidation therapy, while chemoimmunotherapy or participation in a clinical trial is recommended for those with progressive disease (see the section on *Treatment for Patients with Relapsed/Refractory Neuroblastoma* below).

Historically, post-consolidation therapy consisted of 6 cycles of isotretinoin administered as a differentiating agent. This approach was adopted based upon the landmark CCG-3891 trial that demonstrated improved outcomes in patients randomized to isotretinoin compared to no further therapy.¹⁰⁰ Subsequently, the ANBL0032 trial demonstrated a significant improvement in EFS for patients randomized to receive the anti-GD2 monoclonal antibody dinutuximab + cytokines (sargramostim in cycles 1, 3, and 5; interleukin-2 in cycles 2 and 4) + isotretinoin (2-year EFS from randomization at start of post-consolidation therapy of 66%) compared to patients randomized to isotretinoin alone (2-year EFS of 46%).¹¹⁴ Based upon these findings, the ANBL0032 immunotherapy regimen became a standard post-consolidation regimen (category 1 recommendation for use of post-consolidation therapy with anti-GD2 antibody) (see *Principles of Systemic Therapy* in the algorithm).¹¹⁵

Data from the SIOPEN HR-NBL1 trial called into question the role of interleukin-2 together with anti-GD2 immunotherapy. In this trial, patients were randomized to receive anti-GD2 immunotherapy (dinutuximab beta) with or without subcutaneous interleukin-2.¹¹⁶ Interleukin-2 did not improve outcomes and was associated with increased toxicity. Based upon these findings, COG high-risk protocols no longer include interleukin-2, and the Panel endorses this approach.



NCCN Guidelines Version 2.2026

Neuroblastoma

Other anti-GD2 antibodies may be appropriate as post-consolidation therapy. For example, dinutuximab beta given with isotretinoin but without sargramostim is a commonly used post-consolidation regimen in Europe. A non-randomized comparison showed higher EFS among patients treated with this approach compared to the historic experience with isotretinoin alone.¹¹⁷

Continuation Therapy

Eflornithine [2,5-diamino-2-(difluoromethyl) pentanoic acid hydrochloride hydrate; DFMO] is an inhibitor of ornithine decarboxylase, a key enzyme required for the synthesis of polyamines that regulate homeostasis and promote survival in cancer cells. This agent was studied as continuation therapy in a multicenter, single-arm, phase 2 trial (Study 3b; NCT02395666) in children with high-risk neuroblastoma that had responded to frontline therapy that included induction, consolidation, and anti-GD2 directed immunotherapy. Patients who had a partial response or better following standard frontline therapy were eligible to enroll in Study 3b following completion of immunotherapy and received eflornithine 750 mg/m² ± 250 mg/m² twice daily for up to 2 years. Reported adverse events included transaminitis and hearing loss.¹¹⁸

Data from 92 patients on Study 3b were compared with an external control arm consisting of 852 patients treated with anti-GD2 immunotherapy, cytokines, and isotretinoin on COG ANBL0032 who did not go on to receive eflornithine continuation therapy.¹¹⁹ Patients on Study 3b had superior outcomes compared to the external control group (EFS hazard ratio [HR], 0.48 [95% CI, 0.27–0.85]; OS HR, 0.32 [95% CI, 0.15–0.70]). Further analyses using propensity score matching and sensitivity analyses also demonstrated higher EFS and OS for patients on Study 3b, though the potential for residual confounding remains in the context of a non-randomized comparison.

Eflornithine is approved by the U.S. Food and Drug Administration (FDA) for use in the continuation setting for patients with high-risk neuroblastoma who have achieved a partial response or better following completion of anti-GD2 immunotherapy.¹²⁰ The NCCN Neuroblastoma Panel suggests that clinicians discuss eflornithine as a continuation therapy option with patients who are in remission at the end of immunotherapy; this is a category 2B recommendation. Serial monitoring of hearing with audiograms or brainstem auditory evoked response is essential at 6-month intervals or as clinically indicated, as most patients with high-risk neuroblastoma are at a critical age for language development.

Disease Evaluations During Frontline Therapy for High-Risk Disease

Following initial staging evaluations prior to the start of therapy (see the *Staging* section above), patients with high-risk disease undergo anatomic imaging (CT or MRI) of the primary site prior to planned surgical resection. Full disease evaluation (anatomic imaging of the primary site, I-123 MIBG scan [or FDG-PET, if MIBG non-avid disease], and bilateral bone marrow aspirates and biopsies) is recommended at the end of induction, start of post-consolidation, and end of therapy. Patients with more than five residual MIBG-avid sites of disease at the end of induction are encouraged to have a repeat I-123 MIBG scan after recovery from high-dose chemotherapy with stem cell rescue to prioritize metastatic sites that might be treated during consolidative radiotherapy. An I-123 MIBG scan (or FDG-PET, if MIBG non-avid disease) is recommended halfway through post-consolidation therapy, with anatomic imaging and bone marrow evaluations reserved for patients with residual disease identified on the disease evaluation at the start of post-consolidation therapy.

Organ Function Evaluations During Frontline Therapy for High-Risk Disease

Therapy for high-risk neuroblastoma is intensive and associated with both acute and long-term toxicities (see the *Monitoring for Late Effects Related*



NCCN Guidelines Version 2.2026

Neuroblastoma

to *Neuroblastoma Treatment* section below). During treatment, these patients require frequent laboratory monitoring, including blood counts, chemistry panels, and urinalyses. Detailed evaluation of renal function (often with nuclear medicine measurements of glomerular filtration rate) is essential before consolidation high-dose therapy. Serial monitoring of cardiac function with electrocardiograms and echocardiograms is routine. Serial monitoring of hearing with audiograms or brainstem auditory evoked response is essential, as most patients with high-risk neuroblastoma are at a critical age for language development.

Therapy for Adolescents and Adults with High-Risk Neuroblastoma

Neuroblastoma is largely a disease of young children, though adolescents and adults may occasionally present with high-risk disease. The clinical studies that inform these guidelines, including toxicity data, therefore predominantly included patients <5 years of age at initial diagnosis. The general principles of high-risk therapy should be applied to older patients with high-risk disease, though it is acknowledged that these patients may require a more individualized approach to treatment based upon comorbid conditions and tolerance of planned therapy.

Response Assessment

The disease response criteria recommended by the NCCN Neuroblastoma Panel are based on the INRC, which were revised in 2017 based on the availability of modern imaging modalities and new methods for bone marrow disease assessment.⁶⁸ Changes in urinary catecholamine levels (HVA and VMA) are no longer taken into account for response assessment due to the lack of standardization and the influence of diet. See the treatment algorithm for the specific requirements regarding timing of response, as these differ across risk groups.

Response assessment in the primary tumor site or metastatic bone or soft tissue should be based on RECIST criteria and functional imaging

(anatomical and MIBG/FDG-PET).^{68,69} If either primary neuroblastoma tumors or metastatic bone or soft tissue lesions are not MIBG avid, or when MIBG imaging and anatomic imaging do not correlate, FDG-PET imaging is recommended.

Interpretation of MIBG tumor uptake in metastatic bone or soft tissue is performed by means of semiquantitative scoring of body segments. The modified Curie score¹²¹ and the SIOPEN score⁷⁶ are the two most commonly used systems. In North America, the modified Curie score is used.

MIBG is considered more sensitive and specific than technetium-99m (99mTc)-based bone scintigraphy, which is no longer recommended for use in response assessment in bone. Novel radiotracers for assessing response are in development and may serve a role in select patients.⁷²⁻⁷⁴

For bone marrow response assessment, the Panel recommends using the approach described by Burchill et al and included in the INRC (based on immunocytology and/or immunohistochemistry).⁵⁰

The definition of overall response is based on the combination of primary tumor, metastatic bone or soft tissue, and bone marrow responses.

For the complete neuroblastoma response assessment criteria recommended by the NCCN Neuroblastoma Panel, see *Response Assessment* in the algorithm.

Disease Surveillance

Although outcomes have improved for patients with neuroblastoma, approximately half of patients with high-risk disease will still develop relapsed or refractory disease.⁸⁶



NCCN Guidelines Version 2.2026

Neuroblastoma

Close surveillance of patients treated for neuroblastoma is essential to detect disease progression and other side effects related to prior treatment. Surveillance recommendations stratified by risk group are detailed below; see *Disease Surveillance/Follow-up After Completion of Treatment* in the algorithm for complete surveillance recommendations. Refer to the *Treatment for Relapsed/Refractory Neuroblastoma* section below for patients with relapsed or refractory disease.

Low-Risk Disease

For patients with low-risk neuroblastoma who received no treatment (observation only), the use of ultrasound for surveillance is recommended when clinically indicated and appropriate (see *Low-Risk Disease* in the algorithm).

For those with low-risk disease whose tumors were surgically resected, the NCCN Neuroblastoma Panel recommends cross-sectional imaging to delineate new baseline disease status at least 1 month postoperatively (see *Low-Risk Disease* in the algorithm). This is followed by a transition to ultrasound for surveillance if possible (~every 3 months for year 1, every 6 to 12 months for years 2–3, and then as clinically indicated). Cross-sectional imaging for surveillance may be required depending on the location of the primary tumor. Additionally, an interim H&P is recommended every 3 months in year 1, then every 6 months in year 2, followed by every 6 to 12 months in year 3, and then as clinically indicated. In terms of laboratory studies, the Panel notes that urine catecholamine levels are no longer included in the revised INRC.⁶⁸ However, obtaining spot catecholamine testing can be considered during surveillance, if levels were elevated at diagnosis.

Intermediate-Risk Disease

Surveillance recommendations for patients with intermediate-risk disease were adapted from the ANBL0531 and ANBL1232 studies.^{42,97} For

patients who were treated for intermediate-risk disease, the NCCN Panel recommends an interim H&P approximately every 3 months for year 1, every 6 months for year 2, and then annually for years 3 through 5. Audiologic assessment should be considered depending on the degree of exposure to ototoxic agents. CBC with differential is recommended at the same frequency as imaging if bone marrow was involved at diagnosis. Creatinine should be assessed every 6 months for year 1, then annually for years 2 and 3, and then as clinically indicated. Thyroid function can also be impacted by neuroblastoma treatment. Thyroid studies including thyroid-stimulating hormone (TSH) are recommended annually through year 3, and then as clinically indicated. The Panel recommends including free T4 analysis if TSH is abnormal. Like surveillance in patients with low-risk disease, urine catecholamine levels are no longer recommended as part of INRC.⁶⁸ However, obtaining spot catecholamine levels can be considered during surveillance if elevated at diagnosis.

If the treatment endpoint has been achieved in patients with intermediate-risk disease, the NCCN Neuroblastoma Panel recommends an MIBG scan as part of the end-of-therapy evaluation. For patients with MIBG-avid tumors and INRG stage M disease at diagnosis, an I-123 MIBG scan with SPECT is recommended, if available, while imaging by FDG-PET is recommended for patients with MIBG non-avid tumors. If either MIBG or FDG-PET scan was positive at diagnosis and at completion of therapy, these studies should be obtained until a negative scan is achieved or the patient is 36 months from completion of therapy. This should be done every 3 to 6 months in year 1, annually in years 2 and 3, and then as clinically indicated.

The Panel also recommends CT or MRI cross-sectional imaging of the primary site approximately every 3 months for year 1, then every 6 months for year 2, followed by annually for year 3, and then as clinically indicated.



NCCN Guidelines Version 2.2026

Neuroblastoma

High-Risk Disease

Surveillance recommendations for high-risk disease were adapted from the ANBL0532 and ANBL1531 studies.^{95,122} For patients who were treated for high-risk disease, the NCCN Neuroblastoma Panel recommends that H&P be performed approximately every 3 months for year 1, then every 6 months for years 2 through 5, and ongoing evaluation of growth parameters. The NCCN Panel strongly recommends audiologic assessment annually for 5 years, and then as clinically indicated, as studies have suggested that high rates of severe ototoxicity (requiring hearing aids) are seen in survivors of high-risk neuroblastoma.¹²³ CBC with differential is recommended at the same frequency as imaging. Assessment of electrolytes (including Ca⁺⁺, PO₄, and Mg⁺⁺), as well as creatinine, alanine aminotransferase (ALT), and bilirubin, are recommended every 3 months for year 1, followed by every 6 months for years 2 and 3, and then annually for years 4 and 5. Thyroid studies including TSH should be done every 6 months for years 1 and 2, and then annually for years 3 through 5. The Panel recommends including free T4 analysis if TSH is abnormal.

Depending on total anthracycline dose and radiation dose administered with potential impact to the heart (and if normal at the end of therapy), an echocardiogram should be obtained every 2 to 5 years if normal at the end of therapy, as cardiotoxicity is a known late effect in patients with high-risk neuroblastoma.^{86,124} Hemoglobin A1c, ferritin, reproductive health laboratory tests (follicle-stimulating hormone [FSH], luteinizing hormone [LH], anti-Müllerian hormone, inhibin B), and pulmonary function tests should be evaluated if clinically indicated. Like surveillance in patients with low-risk and intermediate-risk disease, urine catecholamine levels are no longer recommended as part of INRC.⁶⁸ However, spot catecholamine levels can be considered during surveillance if elevated at diagnosis.

For patients with MIBG-avid tumors, an I-123 MIBG scan with SPECT is recommended, if available, while imaging by FDG-PET is recommended for those with MIBG non-avid tumors. Imaging should be done every 3 to 6 months in year 1, every 6 months in year 2, followed by annually in year 3, and then as clinically indicated.

The Panel also recommends CT or MRI cross-sectional imaging of the primary site approximately every 3 to 6 months for year 1, then every 6 months for year 2, followed by annually for year 3, and then as clinically indicated.

If the bone marrow is no longer involved at the end of therapy, bilateral bone marrow aspirates and biopsies should be obtained only if clinically indicated.

Treatment for Patients with Relapsed/Refractory Neuroblastoma

For patients with relapsed or refractory neuroblastoma during or after completion of high-risk neuroblastoma therapy, treatment decisions are scenario-specific and require an individualized approach. Consultation with a neuroblastoma specialist should be considered. Factors that impact therapy selection include recurrence timing, extent of disease, MIBG avidity at relapse, prior treatment administered and associated toxicities, clinical condition and organ function, as well as the molecular features of the disease. Participation in a clinical trial for this group of patients is strongly encouraged by the NCCN Panel.

While other therapies may be selected for specific populations of patients, the combination of an anti-GD2 antibody and cytotoxic chemotherapy is now widely used at the time of first relapse or first declaration of refractory disease. Dinutuximab is approved by the FDA for the treatment of pediatric patients with high-risk neuroblastoma who achieve at least a



NCCN Guidelines Version 2.2026

Neuroblastoma

partial response to prior first-line multiagent, multimodality therapy¹²⁰ and often used clinically in North America, while dinutuximab beta is widely used in Europe¹²⁵ and other areas of the world. The combination of irinotecan, temozolomide, and GD2-directed antibody therapy has been evaluated in multiple studies, and objective response rates are strikingly reproducible.^{102,103,126-130} Other chemotherapy backbones that include camptothecins and alkylators (eg, topotecan and cyclophosphamide or topotecan and temozolomide) in combination with dinutuximab or dinutuximab beta demonstrate anti-tumor activity.^{129,131,132}

Naxitamab-gqgk is FDA-approved for treatment of children >1 year of age and adults with relapsed/refractory high-risk neuroblastoma specifically in bone or bone marrow (but not soft tissue) who have had a partial response, minor response, or stable disease to prior therapy.¹²⁰

In patients for whom GD2-directed therapy is not appropriate at time of first relapse, the targeted radiopharmaceutical I-131 MIBG may be considered based upon results of clinical trials conducted in the relapsed/refractory population.^{133,134} One phase 2 study that evaluated I-131 MIBG in 164 patients with progressive, refractory, or relapsed high-risk neuroblastoma reported that the overall response rate was 36%, with a 2-year OS rate of 29%.¹³³

The use of targeted agents for patients with relapsed disease whose tumors harbor actionable aberrations has also been evaluated. Repeat genomic profiling on a current sample at the time of relapse is recommended. A study that evaluated 23 paired diagnostic and relapsed neuroblastoma samples using NGS observed that more mutations and alterations were identified in relapsed tumors than primary tumors.¹³⁵

Tumors from patients with relapsed neuroblastoma appear to have a higher frequency of *ALK* mutations than those from patients with newly diagnosed neuroblastoma.¹³⁶ Several studies have indicated that *ALK*

inhibitors have activity in patients with *ALK*-aberrant tumors.¹³⁶⁻¹³⁸ Agents targeting other pathways of interest in neuroblastoma are also being studied in ongoing clinical trials.^{86,97} In specific contexts, external beam radiation and/or surgery may be indicated to address individual sites of relapsed disease.

Additional options may be considered at the time of second or greater relapse. Rechallenge with chemotherapy combined with GD2-directed antibody therapy based on a history of prior response can be considered. For example, one retrospective study that evaluated 54 patients with relapsed/refractory high-risk neuroblastoma found that patients who previously experienced a disease response with chemoimmunotherapy that included dinutuximab had a higher objective response rate after receiving topotecan, cyclophosphamide, and dinutuximab than those without a prior response to chemoimmunotherapy (54% vs 23%).¹³²

Monitoring for Late Effects Related to Neuroblastoma Treatment

The therapies used to treat neuroblastoma may result in a wide range of late effects, which are treatment-related complications that can occur months or years after completion of treatment.¹³⁹ The late effects that may develop in survivors of neuroblastoma are specific to each individual, the age at which neuroblastoma is diagnosed, and the wide range of therapies used to treat it. Monitoring for late effects is becoming increasingly important, especially as survival outcomes for patients with high-risk neuroblastoma who receive multimodality therapy continue to improve.⁶ Monitoring should be done at each follow-up visit. Typically, monitoring for late effects as a component of survivorship follow-up starts ≥2 years after completion of systemic therapy. The NCCN Neuroblastoma Panel strongly recommends that a personalized survivorship care plan be developed for each patient. Generalized recommendations for screening for late effects in those with neuroblastoma would not be appropriate for many patients.



NCCN Guidelines Version 2.2026

Neuroblastoma

The NCCN Neuroblastoma Panel recommends consulting the COG Survivorship Guidelines for appropriate screening and counseling related to thyroid, cardiac, pulmonary, renal, bone, reproductive health, SMNs (with special attention to thyroid and kidney), and other treatment-associated late effects for patients who received treatment for neuroblastoma.¹⁴⁰

Patients with high-risk neuroblastoma (especially those treated with myeloablative therapy) are at particularly high risk of hearing impairment, endocrine deficiencies, and growth impairment.^{86,124,141,142} Additional late effects common to those with high-risk disease include chronic kidney disease, impaired fertility, cardiotoxicity, neurocognitive impairment, chronic lung disease, and second malignant neoplasms (SMNs).^{139,141,143}

An analysis of the retrospective Childhood Cancer Survivor Study (CCSS) found that survivors of high-risk neuroblastoma were at higher risk of late morbidity and mortality compared with those treated for intermediate- or low-risk disease.¹⁴⁴ All-cause mortality was higher across all risk groups, and the risk for SMNs was higher among survivors of high- and intermediate-risk disease. Individuals treated for high-risk neuroblastoma were also at higher risk of grade 3–5 chronic health conditions compared to their siblings.

Patients who receive chemotherapy for non–high-risk disease typically receive lower cumulative doses of chemotherapeutic agents and limited or no EBRT; therefore, these patients are at lower risk of long-term toxicities compared to those with high-risk disease. However, patients with non–high-risk disease constitute a heterogeneous group, and there is a wide range of cumulative chemotherapeutic exposures.¹⁴⁵

It has been reported that many patients with neuroblastoma who receive platinum-based chemotherapy experience ototoxicity.¹²⁴ Hearing loss is also a reported side effect of eflornithine treatment, with 13% of patients

experiencing new or worsening hearing loss after initiation of eflornithine.¹²⁰ Screening for sensorineural hearing loss in patients who are treated with platinum chemotherapy and eflornithine is of particular importance to minimize delays in speech and language development, which in turn can impact academic performance and social development. While the COG Survivorship Guidelines outline the minimum recommended frequency for audiologic assessment, the NCCN Neuroblastoma Panel recommends a referral to institutional audiology and/or otolaryngology teams for children with impaired hearing to determine the appropriate schedule for subsequent evaluations.¹⁴⁰

Fertility preservation is an option for some patients with high-risk disease. When possible, referral to fertility specialists for further discussion is recommended prior to initiation of chemotherapy.

See *Monitoring for Late Effects* in the algorithm for complete recommendations.

Summary

The NCCN Guidelines for Neuroblastoma were developed by a multidisciplinary Panel and provide recommendations for the diagnosis, risk stratification, and treatment of neuroblastoma based on clinical evidence and consensus. The recommended strategies for treating neuroblastoma will continue to evolve based on emerging data from clinical trials. The NCCN Guidelines for Neuroblastoma will continue to be updated annually.



NCCN Guidelines Version 2.2026

Neuroblastoma

References

1. Qiu B, Matthay KK. Advancing therapy for neuroblastoma. *Nat Rev Clin Oncol* 2022;19:515–533. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35614230>.
2. PDQ Pediatric Treatment Editorial Board. PDQ® Pediatric Treatment Editorial Board. PDQ Neuroblastoma Treatment. Bethesda, MD: National Cancer Institute.; 2025.
3. Key Statistics for Neuroblastoma. Available at: <https://www.cancer.org/cancer/types/neuroblastoma/key-statistics.html>. Accessed April 24, 2026.
4. Siegel DA, King JB, Lupo PJ, et al. Counts, incidence rates, and trends of pediatric cancer in the United States, 2003-2019. *J Natl Cancer Inst* 2023;115:1337–1354. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37433078>.
5. Neuroblastoma. PathologyOutlines.com website. 2026. Available at: <https://www.pathologyoutlines.com/topic/adrenalneuroblastoma.html>. Accessed April 24, 2026.
6. Bagatell R, DuBois SG, Naranjo A, et al. Children's Oncology Group's 2023 blueprint for research: Neuroblastoma. *Pediatr Blood Cancer* 2023;70 Suppl 6:e30572. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37458162>.
7. Henderson TO, Bhatia S, Pinto N, et al. Racial and ethnic disparities in risk and survival in children with neuroblastoma: a Children's Oncology Group study. *J Clin Oncol* 2011;29:76–82. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21098321>.
8. Bona K, Li Y, Winestone LE, et al. Poverty and targeted immunotherapy: survival in Children's Oncology Group clinical trials for high-risk neuroblastoma. *J Natl Cancer Inst* 2021;113:282–291. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33227816>.
9. Campbell K, Siegel DA, Umaretiya PJ, et al. A comprehensive analysis of neuroblastoma incidence, survival, and racial and ethnic disparities from 2001 to 2019. *Pediatr Blood Cancer* 2024;71:e30732. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37867409>.
10. National Institutes of Health. PubMed Overview. Available at: <https://pubmed.ncbi.nlm.nih.gov/about/>.
11. Freedman-Cass DA, Fischer T, Alpert AB, et al. The value and process of inclusion: Using sensitive, respectful, and inclusive language and images in NCCN content. *J Natl Compr Canc Netw* 2023;21:434–441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37156485>.
12. Matthay KK, Maris JM, Schleiermacher G, et al. Neuroblastoma. *Nat Rev Dis Primers* 2016;2:16078. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27830764>.
13. Kamihara J, Bourdeaut F, Foulkes WD, et al. Retinoblastoma and neuroblastoma predisposition and surveillance. *Clin Cancer Res* 2017;23:e98–e106. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28674118>.
14. Mosse YP, Laudenslager M, Longo L, et al. Identification of ALK as a major familial neuroblastoma predisposition gene. *Nature* 2008;455:930–935. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18724359>.
15. Janoueix-Lerosey I, Lequin D, Brugieres L, et al. Somatic and germline activating mutations of the ALK kinase receptor in neuroblastoma. *Nature* 2008;455:967–970. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18923523>.
16. Trochet D, Bourdeaut F, Janoueix-Lerosey I, et al. Germline mutations of the paired-like homeobox 2B (PHOX2B) gene in neuroblastoma. *Am J Hum Genet* 2004;74:761–764. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15024693>.
17. Witkowski L, Nichols KE, Jongmans M, et al. Germline pathogenic SMARCA4 variants in neuroblastoma. *J Med Genet* 2023;60:987–992. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36813544>.



NCCN Guidelines Version 2.2026 Neuroblastoma

18. Kim J, Vaksman Z, Egolf LE, et al. Germline pathogenic variants in neuroblastoma patients are enriched in BARD1 and predict worse survival. *J Natl Cancer Inst* 2024;116:149–159. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37688579>.
19. Early Detection, Diagnosis, and Staging of Neuroblastoma. Available at: <https://www.cancer.org/cancer/types/neuroblastoma/detection-diagnosis-staging.html>. Accessed April 24, 2026.
20. Pang KK, de Sousa C, Lang B, Pike MG. A prospective study of the presentation and management of dancing eye syndrome/opsoclonus-myoclonus syndrome in the United Kingdom. *Eur J Paediatr Neurol* 2010;14:156–161. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19423368>.
21. Rossor T, Yeh EA, Khakoo Y, et al. Diagnosis and management of opsoclonus-myoclonus-ataxia syndrome in children: An international perspective. *Neurol Neuroimmunol Neuroinflamm* 2022;9:e1153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35260471>.
22. Hasegawa S, Matsushige T, Kajimoto M, et al. A nationwide survey of opsoclonus-myoclonus syndrome in Japanese children. *Brain Dev* 2015;37:656–660. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25454391>.
23. Tate ED, Allison TJ, Pranzatelli MR, Verhulst SJ. Neuroepidemiologic trends in 105 US cases of pediatric opsoclonus-myoclonus syndrome. *J Pediatr Oncol Nurs* 2005;22:8–19. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15574722>.
24. Bourdeaut F, de Carli E, Timsit S, et al. VIP hypersecretion as primary or secondary syndrome in neuroblastoma: A retrospective study by the Societe Francaise des Cancers de l'Enfant (SFCE). *Pediatr Blood Cancer* 2009;52:585–590. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19143025>.
25. Mitur-Lesiuk M, Dubaj M, Raniewicz M, et al. Review of paraneoplastic syndromes in children with malignancy. *Med Sci Monit* 2025;31:e947393. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40159669>.
26. Rudnick E, Khakoo Y, Antunes NL, et al. Opsoclonus-myoclonus-ataxia syndrome in neuroblastoma: clinical outcome and antineuronal antibodies—a report from the Children's Cancer Group Study. *Med Pediatr Oncol* 2001;36:612–622. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11344492>.
27. Dornelles Penteado Pacheco ESL, Monteiro Caran EM. Vasoactive Intestinal Peptide-Producing Neuroblastic Tumors: A Rare Cause of Refractory Diarrhea. *Cureus* 2024;16:e67861. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39328672>.
28. de Alarcon PA, Matthay KK, London WB, et al. Intravenous immunoglobulin with prednisone and risk-adapted chemotherapy for children with opsoclonus myoclonus ataxia syndrome associated with neuroblastoma (ANBL00P3): a randomised, open-label, phase 3 trial. *Lancet Child Adolesc Health* 2018;2:25–34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29376112>.
29. Meena JP, Seth R, Chakrabarty B, et al. Neuroblastoma presenting as opsoclonus-myoclonus: a series of six cases and review of literature. *J Pediatr Neurosci* 2016;11:373–377. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28217170>.
30. Pranzatelli MR, Tate ED, McGee NR. Demographic, Clinical, and Immunologic Features of 389 Children with Opsoclonus-Myoclonus Syndrome: A Cross-sectional Study. *Front Neurol* 2017;8:468. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28959231>.
31. Graus F, Vogrig A, Muniz-Castrillo S, et al. Updated Diagnostic Criteria for Paraneoplastic Neurologic Syndromes. *Neurol Neuroimmunol Neuroinflamm* 2021;8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34006622>.
32. Russo C, Cohn SL, Petrucci MJ, de Alarcon PA. Long-term neurologic outcome in children with opsoclonus-myoclonus associated with neuroblastoma: a report from the Pediatric Oncology Group. *Med Pediatr Oncol* 1997;28:284–288. Available at:



NCCN Guidelines Version 2.2026 Neuroblastoma

33. Patel A, Fischer C, Lin YC, et al. Treatment and revaccination of children with paraneoplastic opsoclonus-myoclonus-ataxia syndrome and neuroblastoma: The Memorial Sloan Kettering experience. *Pediatr Blood Cancer* 2020;67:e28319. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32543116>.
34. Kumar P, Willard VW, Embry L, et al. Late cognitive and adaptive outcomes of patients with neuroblastoma-associated opsoclonus-myoclonus-ataxia-syndrome: a report from the Children's Oncology Group. *Pediatr Blood Cancer* 2024;71:e31039. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38689540>.
35. De Grandis E, Parodi S, Conte M, et al. Long-term follow-up of neuroblastoma-associated opsoclonus-myoclonus-ataxia syndrome. *Neuropediatrics* 2009;40:103–111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20020394>.
36. de Buys Roessingh AS, Lorient MH, Wiesenauer C, Lallier M. Lambert-Eaton myasthenic syndrome revealing an abdominal neuroblastoma. *J Pediatr Surg* 2009;44:E5–7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19635290>.
37. Zhang YT, Feng LH, Zhang Z, et al. Different kinds of paraneoplastic syndromes in childhood neuroblastoma. *Iran J Pediatr* 2015;25:e266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26199695>.
38. Jayarangaiah A, Lui F, Theetha Kariyanna P. Lambert-Eaton Myasthenic Syndrome. StatPearls. Treasure Island (FL): StatPearls Publishing. Copyright © 2026, StatPearls Publishing LLC.; 2026.
39. Shimada H, Ambros IM, Dehner LP, et al. Terminology and morphologic criteria of neuroblastic tumors: recommendations by the International Neuroblastoma Pathology Committee. *Cancer* 1999;86:349–363. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10421272>.
40. Shimada H, Ambros IM, Dehner LP, et al. The International Neuroblastoma Pathology Classification (the Shimada system). *Cancer* 1999;86:364–372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10421273>.
41. Brodeur GM, Pritchard J, Berthold F, et al. Revisions of the international criteria for neuroblastoma diagnosis, staging, and response to treatment. *J Clin Oncol* 1993;11:1466–1477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8336186>.
42. Twist CJ, Schmidt ML, Naranjo A, et al. Maintaining outstanding outcomes using response- and biology-based therapy for intermediate-risk neuroblastoma: A report from the Children's Oncology Group study ANBL0531. *J Clin Oncol* 2019;37:3243–3255. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31386611>.
43. Nuchtern JG, London WB, Barnewolt CE, et al. A prospective study of expectant observation as primary therapy for neuroblastoma in young infants: a Children's Oncology Group study. *Ann Surg* 2012;256:573–580. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22964741>.
44. Williams CM, Greer M. Homovanillic acid and vanilmandelic acid in diagnosis of neuroblastoma. *JAMA* 1963;183:836–840. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14000837>.
45. Newly diagnosed with neuroblastoma. 2011. Available at: <https://www.childrensoncologygroup.org/newly-diagnosed-with-neuroblastoma>. Accessed April 30, 2024.
46. Moroz V, Machin D, Hero B, et al. The prognostic strength of serum LDH and serum ferritin in children with neuroblastoma: A report from the International Neuroblastoma Risk Group (INRG) project. *Pediatr Blood Cancer* 2020;67:e28359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32472746>.
47. Irwin MS, Naranjo A, Zhang FF, et al. Revised Neuroblastoma Risk Classification System: A report from the Children's Oncology Group. *J Clin Oncol* 2021;39:3229–3241. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34319759>.
48. Pinches RS, Clinton CM, Ward A, et al. Making the most of small samples: Optimization of tissue allocation of pediatric solid tumors for clinical and research use. *Pediatr Blood Cancer* 2020;67:e28326. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32667141>.



NCCN Guidelines Version 2.2026 Neuroblastoma

49. Fisch AS, Church AJ. Special considerations in the molecular diagnostics of pediatric neoplasms. Clin Lab Med 2022;42:349–365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36150816>.

50. Burchill SA, Beiske K, Shimada H, et al. Recommendations for the standardization of bone marrow disease assessment and reporting in children with neuroblastoma on behalf of the International Neuroblastoma Response Criteria Bone Marrow Working Group. Cancer 2017;123:1095–1105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27984660>.

51. Oh C, Youn JK, Han JW, et al. Abdominal tumors in children: Comparison between minimally invasive surgery and traditional open surgery. Medicine (Baltimore) 2016;95:e5181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27861341>.

52. Franquemont DW, Mills SE, Lack EE. Immunohistochemical detection of neuroblastomatous foci in composite adrenal pheochromocytoma-neuroblastoma. Am J Clin Pathol 1994;102:163–170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8042583>.

53. Warren M, Matsuno R, Tran H, Shimada H. Utility of Phox2b immunohistochemical stain in neural crest tumours and non-neural crest tumours in paediatric patients. Histopathology 2018;72:685–696. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28986989>.

54. Lonergan GJ, Schwab CM, Suarez ES, Carlson CL. Neuroblastoma, ganglioneuroblastoma, and ganglioneuroma: radiologic-pathologic correlation. Radiographics 2002;22:911–934. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12110723>.

55. Brisse HJ, McCarville MB, Granata C, et al. Guidelines for imaging and staging of neuroblastic tumors: consensus report from the International Neuroblastoma Risk Group Project. Radiology 2011;261:243–257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21586679>.

56. Peuchmaur M, d'Amore ES, Joshi VV, et al. Revision of the International Neuroblastoma Pathology Classification: confirmation of favorable and unfavorable prognostic subsets in ganglioneuroblastoma,

nodular. Cancer 2003;98:2274–2281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14601099>.

57. Brodeur GM, Seeger RC, Schwab M, et al. Amplification of N-myc in untreated human neuroblastomas correlates with advanced disease stage. Science 1984;224:1121–1124. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6719137>.

58. Seeger RC, Brodeur GM, Sather H, et al. Association of multiple copies of the N-myc oncogene with rapid progression of neuroblastomas. N Engl J Med 1985;313:1111–1116. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/4047115>.

59. Ambros IM, Tonini GP, Potschger U, et al. Age dependency of the prognostic impact of tumor genomics in localized resectable MYCN-nonamplified neuroblastomas. Report from the SIOPEN biology group on the LNESG trials and a COG validation group. J Clin Oncol 2020;38:3685–3697. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32903140>.

60. George RE, London WB, Cohn SL, et al. Hyperdiploidy plus nonamplified MYCN confers a favorable prognosis in children 12 to 18 months old with disseminated neuroblastoma: a Pediatric Oncology Group study. J Clin Oncol 2005;23:6466–6473. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16116152>.

61. Zafar A, Wang W, Liu G, et al. Molecular targeting therapies for neuroblastoma: Progress and challenges. Med Res Rev 2021;41:961–1021. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33155698>.

62. Church AJ, Corson LB, Kao PC, et al. Molecular profiling identifies targeted therapy opportunities in pediatric solid cancer. Nat Med 2022;28:1581–1589. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35739269>.

63. Bresler SC, Weiser DA, Huwe PJ, et al. ALK mutations confer differential oncogenic activation and sensitivity to ALK inhibition therapy in neuroblastoma. Cancer Cell 2014;26:682–694. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25517749>.



NCCN Guidelines Version 2.2026 Neuroblastoma

64. Bellini A, Potschger U, Bernard V, et al. Frequency and prognostic impact of ALK amplifications and mutations in the European Neuroblastoma Study Group (SIOPEN) High-Risk Neuroblastoma Trial (HR-NBL1). *J Clin Oncol* 2021;39:3377–3390. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34115544>.

65. Berlak M, Tucker E, Dorel M, et al. Mutations in ALK signaling pathways conferring resistance to ALK inhibitor treatment lead to collateral vulnerabilities in neuroblastoma cells. *Mol Cancer* 2022;21:126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35689207>.

66. Sharp SE, Trout AT, Weiss BD, Gelfand MJ. MIBG in neuroblastoma diagnostic imaging and therapy. *Radiographics* 2016;36:258–278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26761540>.

67. Park JR, Eggert A, Caron H. Neuroblastoma: Biology, prognosis, and treatment. *Hematol Oncol Clin North Am* 2010;24:65–86. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20113896>.

68. Park JR, Bagatell R, Cohn SL, et al. Revisions to the International Neuroblastoma Response Criteria: A consensus statement from the National Cancer Institute Clinical Trials Planning Meeting. *J Clin Oncol* 2017;35:2580–2587. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28471719>.

69. Bagatell R, McHugh K, Naranjo A, et al. Assessment of primary site response in children with high-risk neuroblastoma: An international multicenter study. *J Clin Oncol* 2016;34:740–746. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26755515>.

70. Papaioannou G, McHugh K. Neuroblastoma in childhood: review and radiological findings. *Cancer Imaging* 2005;5:116–127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16305949>.

71. Vik TA, Pfluger T, Kadota R, et al. (123)I-mIBG scintigraphy in patients with known or suspected neuroblastoma: Results from a prospective multicenter trial. *Pediatr Blood Cancer* 2009;52:784–790. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19185008>.

72. Bar-Sever Z, Blassoni L, Shulkin B, et al. Guidelines on nuclear medicine imaging in neuroblastoma. *Eur J Nucl Med Mol Imaging* 2018;45:2009–2024. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29938300>.

73. Li S, Zhou Z, Wang X, et al. A head-to-head prospective comparative analysis of (68)Ga-DOTATATE PET/CT and (123)I-MIBG SPECT/CT in central nervous system metastases of neuroblastoma and ganglioneuroblastoma. *Eur J Nucl Med Mol Imaging* 2025;53:452–466. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40637785>.

74. Piccardo A, Treglia G, Fiz F, et al. The evidence-based role of catecholaminergic PET tracers in Neuroblastoma. A systematic review and a head-to-head comparison with mIBG scintigraphy. *Eur J Nucl Med Mol Imaging* 2024;51:756–767. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37962616>.

75. Yanik GA, Parisi MT, Naranjo A, et al. Validation of postinduction Curie scores in high-risk neuroblastoma: A Children's Oncology Group and SIOPEN Group report on SIOPEN/HR-NBL1. *J Nucl Med* 2018;59:502–508. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28887399>.

76. Ladenstein R, Lambert B, Potschger U, et al. Validation of the mIBG skeletal SIOPEN scoring method in two independent high-risk neuroblastoma populations: the SIOPEN/HR-NBL1 and COG-A3973 trials. *Eur J Nucl Med Mol Imaging* 2018;45:292–305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28940046>.

77. Lai HA, Sharp SE, Bhatia A, et al. Imaging of pediatric neuroblastoma: A COG Diagnostic Imaging Committee/SPR Oncology Committee white paper. *Pediatr Blood Cancer* 2023;70 Suppl 4:e29974. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36184716>.

78. Sharp SE, Shulkin BL, Gelfand MJ, et al. 123I-MIBG scintigraphy and 18F-FDG PET in neuroblastoma. *J Nucl Med* 2009;50:1237–1243. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19617326>.

79. Sokol E, Desai AV, Applebaum MA, et al. Age, diagnostic category, tumor grade, and mitosis-karyorrhexis index are independently prognostic



NCCN Guidelines Version 2.2026 Neuroblastoma

in neuroblastoma: An INRG project. J Clin Oncol 2020;38:1906–1918. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32315273>.

80. Janoueix-Lerosey I, Schleiermacher G, Michels E, et al. Overall genomic pattern is a predictor of outcome in neuroblastoma. J Clin Oncol 2009;27:1026–1033. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19171713>.

81. Schleiermacher G, Mosseri V, London WB, et al. Segmental chromosomal alterations have prognostic impact in neuroblastoma: a report from the INRG project. Br J Cancer 2012;107:1418–1422. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22976801>.

82. Hsu LL, Evans AE, D'Angio GJ. Hepatomegaly in neuroblastoma stage 4s: Criteria for treatment of the vulnerable neonate. Med Pediatr Oncol 1996;27:521–528. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8888811>.

83. Cohn SL, Pearson AD, London WB, et al. The International Neuroblastoma Risk Group (INRG) classification system: an INRG Task Force report. J Clin Oncol 2009;27:289–297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19047291>.

84. Meany HJ, London WB, Ambros PF, et al. Significance of clinical and biologic features in Stage 3 neuroblastoma: a report from the International Neuroblastoma Risk Group project. Pediatr Blood Cancer 2014;61:1932–1939. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25044743>.

85. Meany HJ. Non-high-risk neuroblastoma: Classification and achievements in therapy. Children (Basel) 2019;6. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30626019>.

86. DuBois SG, Macy ME, Henderson TO. High-risk and relapsed neuroblastoma: Toward more cures and better outcomes. Am Soc Clin Oncol Educ Book 2022;42:1–13. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35522915>.

87. Baker DL, Schmidt ML, Cohn SL, et al. Outcome after reduced chemotherapy for intermediate-risk neuroblastoma. N Engl J Med

2010;363:1313–1323. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20879880>.

88. Strother DR, London WB, Schmidt ML, et al. Outcome after surgery alone or with restricted use of chemotherapy for patients with low-risk neuroblastoma: results of Children's Oncology Group study P9641. J Clin Oncol 2012;30:1842–1848. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22529259>.

89. Kohler JA, Rubie H, Castel V, et al. Treatment of children over the age of one year with unresectable localised neuroblastoma without MYCN amplification: results of the SIOPEN study. Eur J Cancer 2013;49:3671–3679. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23907002>.

90. Neuroblastoma Survival Rates by Risk Group. Available at: <https://www.cancer.org/cancer/types/neuroblastoma/detection-diagnosis-staging/survival-rates.html>. Accessed April 24, 2026.

91. Grupp SA, Cohn SL, Wall D, et al. Collection, storage, and infusion of stem cells in children with high-risk neuroblastoma: saving for a rainy day. Pediatr Blood Cancer 2006;46:719–722. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16429413>.

92. Kushner BH, Kramer K, LaQuaglia MP, et al. Reduction from seven to five cycles of intensive induction chemotherapy in children with high-risk neuroblastoma. J Clin Oncol 2004;22:4888–4892. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15611504>.

93. Pinto N, Naranjo A, Hibbitts E, et al. Predictors of differential response to induction therapy in high-risk neuroblastoma: A report from the Children's Oncology Group (COG). Eur J Cancer 2019;112:66–79. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30947024>.

94. Park JR, Scott JR, Stewart CF, et al. Pilot induction regimen incorporating pharmacokinetically guided topotecan for treatment of newly diagnosed high-risk neuroblastoma: a Children's Oncology Group study. J Clin Oncol 2011;29:4351–4357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22010014>.



NCCN Guidelines Version 2.2026

Neuroblastoma

95. Park JR, Kreissman SG, London WB, et al. Effect of tandem autologous stem cell transplant vs single transplant on event-free survival in patients with high-risk neuroblastoma: A randomized clinical trial. *JAMA* 2019;322:746–755. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31454045>.

96. Granger MM, Naranjo A, Bagatell R, et al. Myeloablative busulfan/melphalan consolidation following induction chemotherapy for patients with newly diagnosed high-risk neuroblastoma: Children's Oncology Group trial ANBL12P1. *Transplant Cell Ther* 2021;27:490.e491–490.e498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33823167>.

97. Clinicaltrials.gov. Available at: <https://www.clinicaltrials.gov/>. Accessed April 24, 2026.

98. Holmes K, Potschger U, Pearson ADJ, et al. Influence of surgical excision on the survival of patients with stage 4 high-risk neuroblastoma: A report from the HR-NBL1/SIOOPEN study. *J Clin Oncol* 2020;38:2902–2915. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32639845>.

99. von Allmen D, Davidoff AM, London WB, et al. Impact of extent of resection on local control and survival in patients from the COG A3973 study with high-risk neuroblastoma. *J Clin Oncol* 2017;35:208–216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27870572>.

100. Matthay KK, Villablanca JG, Seeger RC, et al. Treatment of high-risk neuroblastoma with intensive chemotherapy, radiotherapy, autologous bone marrow transplantation, and 13-cis-retinoic acid. Children's Cancer Group. *N Engl J Med* 1999;341:1165–1173. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10519894>.

101. Desai AV, Applebaum MA, Karrison TG, et al. Efficacy of post-induction therapy for high-risk neuroblastoma patients with end-induction residual disease. *Cancer* 2022;128:2967–2977. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35665495>.

102. Mody R, Yu AL, Naranjo A, et al. Irinotecan, temozolomide, and dinutuximab with GM-CSF in children with refractory or relapsed neuroblastoma: A report from the Children's Oncology Group. *J Clin Oncol*

2020;38:2160–2169. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32343642>.

103. Munoz JP, Larrosa C, Chamorro S, et al. Early salvage chemo-immunotherapy with irinotecan, temozolomide and naxitamab plus GM-CSF (HITS) for patients with primary refractory high-risk neuroblastoma provide the best chance for long-term outcomes. *Cancers (Basel)* 2023;15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37835531>.

104. Yalcin B, Kremer LC, van Dalen EC. High-dose chemotherapy and autologous haematopoietic stem cell rescue for children with high-risk neuroblastoma. *Cochrane Database Syst Rev* 2015;2015:CD006301. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26436598>.

105. Kushner BH, Ostrovnaya I, Cheung IY, et al. Lack of survival advantage with autologous stem-cell transplantation in high-risk neuroblastoma consolidated by anti-GD2 immunotherapy and isotretinoin. *Oncotarget* 2016;7:4155–4166. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26623730>.

106. Modak S, Kushner BH, LaQuaglia MP, et al. Management and outcome of stage 3 neuroblastoma. *Eur J Cancer* 2009;45:90–98. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC3727624/>.

107. Pinto N, Naranjo A, Ding X, et al. Impact of genomic and clinical factors on outcome of children ≥18 months of age with stage 3 neuroblastoma with unfavorable histology and without MYCN amplification: A Children's Oncology Group (COG) report. *Clin Cancer Res* 2023;29:1546–1556. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36749880>.

108. Schmidt ML, Lal A, Seeger RC, et al. Favorable prognosis for patients 12 to 18 months of age with stage 4 nonamplified MYCN neuroblastoma: a Children's Cancer Group study. *J Clin Oncol* 2005;23:6474–6480. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16116154>.

109. Ladenstein R, Potschger U, Pearson ADJ, et al. Busulfan and melphalan versus carboplatin, etoposide, and melphalan as high-dose



NCCN Guidelines Version 2.2026

Neuroblastoma

chemotherapy for high-risk neuroblastoma (HR-NBL1/SIOPEN): an international, randomised, multi-arm, open-label, phase 3 trial. *Lancet Oncol* 2017;18:500–514. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28259608>.

110. Liu KX, Naranjo A, Zhang FF, et al. Prospective evaluation of radiation dose escalation in patients with high-risk neuroblastoma and gross residual disease after surgery: A report from the Children's Oncology Group ANBL0532 study. *J Clin Oncol* 2020;38:2741–2752. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32530765>.

111. Braunstein SE, London WB, Kreissman SG, et al. Role of the extent of prophylactic regional lymph node radiotherapy on survival in high-risk neuroblastoma: A report from the COG A3973 study. *Pediatr Blood Cancer* 2019;66:e27736. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30968542>.

112. Casey DL, Pitter KL, Kushner BH, et al. Radiation therapy to sites of metastatic disease as part of consolidation in high-risk neuroblastoma: Can long-term control be achieved? *Int J Radiat Oncol Biol Phys* 2018;100:1204–1209. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29439882>.

113. Polishchuk AL, Li R, Hill-Kayser C, et al. Likelihood of bone recurrence in prior sites of metastasis in patients with high-risk neuroblastoma. *Int J Radiat Oncol Biol Phys* 2014;89:839–845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24867534>.

114. Yu AL, Gilman AL, Ozkaynak MF, et al. Anti-GD2 antibody with GM-CSF, interleukin-2, and isotretinoin for neuroblastoma. *N Engl J Med* 2010;363:1324–1334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20879881>.

115. Yu AL, Gilman AL, Ozkaynak MF, et al. Long-term follow-up of a phase III study of ch14.18 (dinutuximab) + cytokine immunotherapy in children with high-risk neuroblastoma: COG study ANBL0032. *Clin Cancer Res* 2021;27:2179–2189. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33504555>.

116. Ladenstein R, Potschger U, Valteau-Couanet D, et al. Interleukin 2 with anti-GD2 antibody ch14.18/CHO (dinutuximab beta) in patients with high-risk neuroblastoma (HR-NBL1/SIOPEN): a multicentre, randomised, phase 3 trial. *Lancet Oncol* 2018;19:1617–1629. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30442501>.

117. Ladenstein R, Potschger U, Valteau-Couanet D, et al. Investigation of the role of dinutuximab beta-based immunotherapy in the SIOPEN high-risk neuroblastoma 1 trial (HR-NBL1). *Cancers (Basel)* 2020;12:309. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32013055>.

118. Sholler GLS, Ferguson W, Bergendahl G, et al. Maintenance DFMO increases survival in high risk neuroblastoma. *Sci Rep* 2018;8:14445. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30262852>.

119. Oesterheld J, Ferguson W, Kravaka JM, et al. Eflornithine as postimmunotherapy maintenance in high-risk neuroblastoma: Externally controlled, propensity score-matched survival outcome comparisons. *J Clin Oncol* 2024;42:90–102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37883734>.

120. US Food & Drug Administration. Available at: <https://www.fda.gov/>. Accessed April 24, 2026.

121. Yanik GA, Parisi MT, Shulkin BL, et al. Semiquantitative mIBG scoring as a prognostic indicator in patients with stage 4 neuroblastoma: a report from the Children's Oncology Group. *J Nucl Med* 2013;54:541–548. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23440556>.

122. Weiss BD, Yanik G, Naranjo A, et al. A safety and feasibility trial of (131) I-MIBG in newly diagnosed high-risk neuroblastoma: A Children's Oncology Group study. *Pediatr Blood Cancer* 2021;68:e29117. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34028986>.

123. Diller L, London W, Bardwell J, et al. Surviving high risk neuroblastoma: A preliminary descriptive report from Project LEAHRN (Late Effects After High-Risk Neuroblastoma). [abstract]. Presented at the Advances in Neuroblastoma Research (ANR) Conference, January 25-27, 2021.



NCCN Guidelines Version 2.2026 Neuroblastoma

124. Friedman DN, Henderson TO. Late effects and survivorship issues in patients with neuroblastoma. *Children (Basel)* 2018;5:107. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30082653>.
125. European Medicines Agency. Available at: <https://www.ema.europa.eu/>. Accessed April 24, 2026.
126. Lerman BJ, Li Y, Carlowicz C, et al. Progression-Free Survival and Patterns of Response in Patients With Relapsed High-Risk Neuroblastoma Treated With Irinotecan/Temozolomide/Dinutuximab/Granulocyte-Macrophage Colony-Stimulating Factor. *J Clin Oncol* 2023;41:508–516. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36206505>.
127. Olgun N, Cecen E, Ince D, et al. Dinutuximab beta plus conventional chemotherapy for relapsed/refractory high-risk neuroblastoma: A single-center experience. *Front Oncol* 2022;12:1041443. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36620564>.
128. Wieczorek A, Zaniewska-Tekieli A, Ehlert K, et al. Dinutuximab beta combined with chemotherapy in patients with relapsed or refractory neuroblastoma. *Front Oncol* 2023;13:1082771. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36816982>.
129. Gray JC, Weston R, Owens C, et al. Dinutuximab Beta Added to Temozolomide-Based Chemotherapy for Children With Relapsed and Refractory Neuroblastoma: Results of the ITCC-SIOPE BEACON Immuno Phase II Trial. *J Clin Oncol* 2026;44:176–187. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41385757>.
130. Mora J, Chan GCF, Morgenstern DA, et al. The anti-GD2 monoclonal antibody naxitamab plus GM-CSF for relapsed or refractory high-risk neuroblastoma: a phase 2 clinical trial. *Nat Commun* 2025;16:1636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39952926>.
131. Raiser P, Schleiermacher G, Gambart M, et al. Chemo-immunotherapy with dinutuximab beta in patients with relapsed/progressive high-risk neuroblastoma: does chemotherapy backbone matter? *Eur J Cancer* 2024;202:114001. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38489858>.
132. Lerman BJ, King M, Mody R, et al. Efficacy and Tolerability of Topotecan/Cyclophosphamide/Dinutuximab in Relapsed and Refractory High-Risk Neuroblastoma: A Multi-Institutional Retrospective Study. *Pediatr Blood Cancer* 2026:e70195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41823156>.
133. Matthay KK, Yanik G, Messina J, et al. Phase II study on the effect of disease sites, age, and prior therapy on response to iodine-131-metaiodobenzylguanidine therapy in refractory neuroblastoma. *J Clin Oncol* 2007;25:1054–1060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17369569>.
134. DuBois SG, Granger MM, Groshen S, et al. Randomized Phase II Trial of MIBG Versus MIBG, Vincristine, and Irinotecan Versus MIBG and Vorinostat for Patients With Relapsed or Refractory Neuroblastoma: A Report From NANT Consortium. *J Clin Oncol* 2021;39:3506–3514. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34270348>.
135. Eleveld TF, Oldridge DA, Bernard V, et al. Relapsed neuroblastomas show frequent RAS-MAPK pathway mutations. *Nat Genet* 2015;47:864–871. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26121087>.
136. Rosswog C, Fassunke J, Ernst A, et al. Genomic ALK alterations in primary and relapsed neuroblastoma. *Br J Cancer* 2023;128:1559–1571. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36807339>.
137. Goldsmith KC, Park JR, Kayser K, et al. Lorlatinib with or without chemotherapy in ALK-driven refractory/relapsed neuroblastoma: phase 1 trial results. *Nat Med* 2023;29:1092–1102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37012551>.
138. Fischer M, Moreno L, Ziegler DS, et al. Ceritinib in paediatric patients with anaplastic lymphoma kinase-positive malignancies: an open-label, multicentre, phase 1, dose-escalation and dose-expansion study. *Lancet Oncol* 2021;22:1764–1776. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34780709>.
139. Laverdiere C, Liu Q, Yasui Y, et al. Long-term outcomes in survivors of neuroblastoma: a report from the Childhood Cancer Survivor Study. *J*



NCCN Guidelines Version 2.2026

Neuroblastoma

Natl Cancer Inst 2009;101:1131–1140. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19648511>.

140. Survivorship Guidelines. Available at:
<https://childrensoncologygroup.org/survivorshipguidelines>. Accessed April 24, 2026.

141. Hesko C, Liu W, Srivastava DK, et al. Neurocognitive outcomes in adult survivors of neuroblastoma: A report from the Childhood Cancer Survivor Study. Cancer 2023;129:2904–2914. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/37199722>.

142. Landier W, Knight K, Wong FL, et al. Ototoxicity in children with high-risk neuroblastoma: prevalence, risk factors, and concordance of grading scales--a report from the Children's Oncology Group. J Clin Oncol 2014;32:527–534. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/24419114>.

143. Henderson TO, Bardwell JK, Kao PC, et al. Late effects after high-risk neuroblastoma (LEAHRN): a multicentre, cross-sectional cohort study from the Children's Oncology Group. Lancet Child Adolesc Health 2025;9:776–786. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/41016399>.

144. Friedman DN, Goodman PJ, Leisenring WM, et al. Impact of Risk-based therapy on late morbidity and mortality in neuroblastoma survivors: a report from the Childhood Cancer Survivor Study. J Natl Cancer Inst 2024. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38460547>.

145. Friedman DN, Goodman PJ, Leisenring WM, et al. Long-term morbidity and mortality among survivors of neuroblastoma diagnosed during infancy: A report from the Childhood Cancer Survivor study. J Clin Oncol 2023;41:1565–1576. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/36525618>.