



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)


Pediatric

Hodgkin Lymphoma

NCCN Evidence Blocks™

Version 1.2026 — May 13, 2026

NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.





NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

***Kara Kelly, MD/Chair €**

Roswell Park Comprehensive Cancer Center
| Roswell Park Oishei Children's Cancer and
Blood Disorders Program

***Susan M. Hiniker, MD/Vice Chair §**

Stanford Cancer Institute | Lucile Packard
Children's Hospital

Saro Armenian, DO, MPH € ‡

City of Hope National Medical Center

Jennifer Belsky, DO, MS €

Indiana University Melvin and Bren Simon
Comprehensive Cancer Center | Riley
Children's Health

Adam J. Bobbey, MD ϕ

The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute | Nationwide
Children's Hospital

Vivian Chang, MD €

UCLA Jonsson Comprehensive Cancer Center |
UCLA Mattel Children's Hospital

Stacy Cooper, MD ‡

Johns Hopkins Kimmel Cancer Center | Johns
Hopkins Children's Center

Don W. Coulter, MD €

Fred & Pamela Buffett Cancer Center |
Children's Hospital & Medical Center

Branko Cuglievan, MD €

The University of Texas
MD Anderson Cancer Center

Adam DuVall, MD, MPH ‡ †

The UChicago Medicine
Comprehensive Cancer Center

Matthew J. Ehrhardt, MD, MS €

St. Jude Children's Research Hospital

Abigail Foederer, DNP, FNP-BC ¥

Patient Advocate

Anusha Preethi Ganesan, MD, PhD ‡

UC San Diego Moores Cancer Center |
Rady Children's Hospital-San Diego

Mallorie Heneghan, MD, MS €

Huntsman Cancer Institute
at the University of Utah |
Primary Children's Hospital

Bradford S. Hoppe, MD, MPH §

Mayo Clinic Comprehensive Cancer Center

Leidy Isenalumhe, MD, MS € ‡ †

Moffitt Cancer Center

Leslie Kersun, MD, MSCE, MSeD € ‡

Abramson Cancer Center at the University
of Pennsylvania | Children's Hospital of
Philadelphia

Adam J. Lamble, MD €

Fred Hutchinson Cancer Center | Seattle
Children's Hospital

Sandra Luna-Fineman, MD €

University of Colorado Cancer Center |
Children's Hospital Colorado

Margaret Ferris, MD, PhD € ‡

Siteman Cancer Center at Barnes-
Jewish Hospital and WashU Medicine |
St. Louis Children's Hospital

Kwadwo Oduro, MD, PhD ≠ ‡

University of Wisconsin Carbone Cancer Center
| American Family Children's Hospital

Arun Ranjan Panigrahi, MD € ‡

UC Davis Comprehensive Cancer Center

Anita P. Price, MD ϕ

Memorial Sloan Kettering Cancer Center

Kenneth B. Roberts, MD §

Yale Cancer Center/Smilow Cancer Hospital |
Yale New Haven Children's Hospital

Seth Rotz, MD, ≠ €

Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute |
University Hospitals Rainbow Babies &
Children's Hospital

Ksenya Shliakhtsitsava, MD €

UT Southwestern Simmons
Comprehensive Cancer Center | Children's
Medical Center Dallas

Christine M. Smith, MD € ‡

Vanderbilt-Ingram Cancer Center | Monroe
Carell Jr. Children's Hospital at Vanderbilt

Aliyah R. Sohani, MD ≠

Mass General Brigham Cancer Institute

Emily Walling, MD, MPHS €

University of Michigan Rogel Cancer Center |
C.S. Mott Children's Hospital

Jane N. Winter, MD ‡

Robert H. Lurie Comprehensive Cancer
Center of Northwestern University | Ann &
Robert H. Lurie Children's Hospital of Chicago

Ana C. Xavier, MD, PhD €

O'Neal Comprehensive Cancer Center at UAB |
Children's of Alabama

Ken Young, MD, PhD ≠

Duke Cancer Institute | Duke Children's Hospital
& Health Center

NCCN

Sarah Montgomery, BS
Katie Stehman, PA-C, MMS

ϕ Diagnostic radiology
‡ Hematology/Hematology oncology
† Medical oncology
≠ Pathology
¥ Patient advocacy
€ Pediatric oncology
§ Radiotherapy/Radiation oncology
* Discussion Section Writing Committee



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

[NCCN Pediatric Hodgkin Lymphoma Panel Members](#) [NCCN Evidence Blocks Definitions \(EB-DEF\)](#)

[Diagnosis and Workup \(PHL-1\)](#)

[Summary of Clinical Trial Risk Groups \(PHL-2\)](#)

Primary Treatment of CHL

- [Stage I–II CHL Without Risk Factors \(PHL-3\)](#)
- [Stage I–II CHL With Risk Factors \(PHL-4\)](#)
- [Stage III–IV CHL \(PHL-5\)](#)

[Primary Treatment of Nodular Lymphocyte-Predominant Hodgkin Lymphoma \(PHL-6\)](#)

[Follow-up After Completion of Treatment and Monitoring for Late Effects \(PHL-8\)](#)

[Suspected Relapsed/Refractory CHL \(PHL-9\)](#)

[Principles of Criteria for Response-Adapted Radiation Therapy \(PHL-A\)](#)

[Principles of Pathology \(PHL-B\)](#)

[Principles of Imaging \(PHL-C\)](#)

[Principles of Staging \(PHL-D\)](#)

[Principles of Systemic Therapy \(PHL-E\)](#)

[Principles of Radiation Therapy \(PHL-F\)](#)

[Select Principles of Supportive Care \(PHL-G\)](#)

[Evidence Blocks](#)

[Staging \(ST-1\)](#)

[Abbreviations \(ABBR-1\)](#)

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS

5					
4					
3					
2					
1					

E S Q C A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

Example Evidence Block

5					
4					
3					
2					
1					

E S Q C A

E = 4
S = 4
Q = 3
C = 4
A = 3

Efficacy of Regimen/Agent

5	Highly effective: Cure likely and often provides long-term survival advantage
4	Very effective: Cure unlikely but sometimes provides long-term survival advantage
3	Moderately effective: Modest impact on survival, but often provides control of disease
2	Minimally effective: No, or unknown impact on survival, but sometimes provides control of disease
1	Palliative: Provides symptomatic benefit only

Safety of Regimen/Agent

5	Usually no meaningful toxicity: Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
4	Occasionally toxic: Rare significant toxicities or low-grade toxicities only; little interference with ADLs
3	Mildly toxic: Mild toxicity that interferes with ADLs
2	Moderately toxic: Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
1	Highly toxic: Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

Note: For significant chronic or long-term toxicities, score decreased by 1

Quality of Evidence

5	High quality: Multiple well-designed randomized trials and/or meta-analyses
4	Good quality: One or more well-designed randomized trials
3	Average quality: Low quality randomized trial(s) or well-designed non-randomized trial(s)
2	Low quality: Case reports or extensive clinical experience
1	Poor quality: Little or no evidence

Consistency of Evidence

5	Highly consistent: Multiple trials with similar outcomes
4	Mainly consistent: Multiple trials with some variability in outcome
3	May be consistent: Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
2	Inconsistent: Meaningful differences in direction of outcome between quality trials
1	Anecdotal evidence only: Evidence in humans based upon anecdotal experience

Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)

5	Very inexpensive
4	Inexpensive
3	Moderately expensive
2	Expensive
1	Very expensive



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

INTRODUCTION

- Consultation with centers participating in pediatric cooperative group trials is encouraged. The recommendations in these Guidelines are from the previous and most recently published trials.
- Referral to current clinical trials is encouraged where available.
- Treatment approaches arising from more recent clinical trials have been collaborative efforts including adult and pediatric patients.
- The Pediatric Hodgkin Lymphoma Panel considers “pediatric” to include any patient aged ≤18 years, and may be applicable to adolescent and young adult (AYA) patients aged ≤39 years. Therefore, these Guidelines are intended to include AYA patients and may apply to patients treated in adult oncology settings. For general oncologic care of AYA patients, see the [NCCN Guidelines for Adolescent and Young Adult \(AYA\) Oncology](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

DIAGNOSTIC WORKUP

- Excisional or incisional biopsy^a
- Immunohistochemistry evaluation^b

ADDITIONAL WORKUP

Essential:

- History and physical (H&P) including:
 - B symptoms (unexplained recurrent fever >38°C within the last month; drenching night sweats; or weight loss >10% of body weight within 6 months of diagnosis)
 - Examination of lymphoid regions, spleen
 - Screening for personal history of autoimmune disorder(s)^c
- Complete blood count (CBC) with differential
- Erythrocyte sedimentation rate (ESR) ± C-reactive protein (CRP)
- Comprehensive metabolic panel (CMP)
- Echocardiogram (ECHO) (especially if anthracycline-based chemotherapy is indicated)
- Chest x-ray posteroanterior (PA) and lateral views (if cross-sectional imaging not available or necessitated to determine bulk of disease for a clinical trial)^d
- CT neck/chest/abdomen/pelvis with (IV ± oral) contrast or CT chest and MRI neck/abdomen/pelvis^d
- FDG-PET/CT^e or FDG-PET/MRI^e (whole-body)^d
- Pregnancy test for patients of childbearing potential
- Counseling on infertility risk^{f,g} (see [NCCN Guidelines for AYA Oncology](#))
- Psychosocial assessment (for AYA, see [NCCN Guidelines for AYA Oncology](#))
- Counseling on cessation of smoking, drugs/illicit substances, vaping, and alcohol (see [NCCN Guidelines for Smoking Cessation](#))

Useful in selected cases:

- Pulmonary function tests (PFTs) (including diffusing capacity [DLCO] if bleomycin indicated)^h
- Electrocardiogram (ECG)
- HIV and hepatitis B/C testing (encouraged)
- Consider immunodeficiency workup (if young age [<5 years], recurrent infections, atypical presentation, personal or family history of immunodeficiency)
- Only consider bilateral bone marrow biopsy if there are cytopenias and negative FDG-PETⁱ
- If spleen irradiation is warranted, vaccines should be given prior to or after RT (ie, pneumococcal, *Haemophilus influenzae* type b, meningococcal). See [Principles of Radiation Therapy \(PHL-F\)](#).
- Assess social determinants of health^j

CLINICAL PRESENTATION

Classic
Hodgkin
lymphoma
(CHL)^k

[Summary
of Clinical
Trial Risk
Groups
\(PHL-2\)](#)

Nodular
lymphocyte-
predominant
Hodgkin
lymphoma
(NLPHL)^l

[PHL-6](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Footnotes



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

DIAGNOSTIC WORKUP FOOTNOTES

- ^a Core needle biopsy may be adequate if it is diagnostic. Fine-needle aspiration (FNA) is not an appropriate diagnostic test. See [Principles of Pathology \(PHL-B\)](#).
- ^b For typical immunophenotype, see [Principles of Pathology \(PHL-B\)](#).
- ^c CPIs should be used with caution in those with active or previously documented autoimmune disease and/or current use of immunosuppressive agents.
- ^d Diagnostic imaging should be done prior to initiating chemotherapy to allow for staging and risk stratification. Consultation with radiation oncologist when considering treatment options and adequacy of imaging for potential future radiation therapy (RT) is strongly recommended. See [Principles of Imaging \(PHL-C\)](#) and [Principles of Staging \(PHL-D\)](#).
- ^e In cases of FDG-PET positivity where sites of disease are inconsistent with usual presentation of Hodgkin lymphoma (HL) or if there is an unusual disease presentation [ie, HIV], additional clinical evaluation may be required for staging. See [Principles of Staging \(PHL-D\)](#). If FDG-PET is negative for anatomic lesions of concern, biopsy should be considered.
- ^f Fertility preservation is an option for some patients. Refer to fertility clinic for further discussion when able, prior to initiation of chemotherapy.
- ^g Close A, et al. *Pediatr Blood Cancer* 2023;70:e30407.
- ^h In general, forced expiratory volume in the first second (FEV1)/forced vital capacity (FVC) >60% by PFT is acceptable for use of bleomycin, unless due to large mediastinal mass from HL. For children who are unable to cooperate for PFTs, the criteria are: no evidence of dyspnea at rest, no exercise intolerance, and a pulse oximetry reading of >92% on room air.
- ⁱ In most instances, if the FDG-PET/CT displays a homogeneous pattern of marrow uptake (thought to be secondary to cytokine release) bone marrow involvement is *not* assumed. If there are multifocal (≥3) skeletal FDG-PET lesions without cortical destruction on CT, marrow involvement may be assumed and a bone marrow biopsy is not needed to confirm involvement (Purz S, et al. *J Clin Oncol* 2011;29:3523-3528).
- ^j Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health. See [NCCN Guidelines for Distress Management \(DIS-A\)](#).
- ^k CHL includes nodular sclerosis, mixed cellularity, lymphocyte-depleted subtypes, and lymphocyte-rich subtypes. If grey-zone, see [NCCN Guidelines for B-Cell Lymphomas](#).
- ^l Per WHO 2022, NLPHL remains under the family of HL, while in the ICC 2022 update, the term NLPHL was replaced with new terminology, nodular lymphocyte-predominant B-cell lymphoma (NLPBL) (Alaggio R, et al. *Leukemia* 2022;36:1720-1748; Campo E, et al. *Blood* 2022;140:1229-1253). Recommendations for management of NLPHL remain in the pediatric HL guidelines as treatment regimens are typically guided by HL experience in clinical practice.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

SUMMARY OF CLINICAL TRIAL RISK GROUPS^m

	Early Stage								Advanced Stage							
Ann Arbor Stage	I				II				III				IV			
B Symptoms	A		B		A		B		A		B		A		B	
Risk Factors (see Table 1)	-	+	-	+	-	+	-	+	-	+						
Clinical Trial																
EuroNet-PHL-C1 ⁿ																
OEPA ± ISRT (Treatment Group [TG] 1)	L		L		L											
OEPA/COPDAC ± ISRT (TG 2)		I		I		I	I		I							
OEPA/COPDAC ± ISRT (TG 3)								H		H	H	H	H	H	H	H
AHOD0031 ^o (ABVE-PC ± ISRT ^p)		I	I	I		I	I		I	I						
AHOD1331 ^q (Bv-AVE-PC ± ISRT)								H			H	H	H	H	H	H
S1826 ^r (Nivolumab-AVD ± RSRT)									H	H	H	H	H	H	H	H
HLHR13 ^s (AEPA/CAPDAC ± RNRT)							H	H			H	H	H	H	H	H

HISTORICAL RISK STRATIFICATION^m

L	Low Risk
I	Intermediate Risk
H	High Risk

- Stage I–II CHL without risk factors ([PHL-3](#))
- Stage I–II CHL with risk factors ([PHL-4](#))
- Stage III–IV CHL ([PHL-5](#))

Table 1				
Risk Factors	EuroNET-PHL		COG	
	C1	C2	AHOD0031	AHOD1331
ESR and B Symptoms		>30 mm/h		
Mediastinal Mass			MMR >0.33	MMR >0.33
E Lesions ^t	Any	Any	Any	
Bulky		Contiguous tumor volume >200 mL	>6 cm extra-mediastinal nodal conglomerate	>6 cm extra-mediastinal nodal conglomerate

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Footnotes



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

SUMMARY OF CLINICAL TRIAL RISK GROUPS^j FOOTNOTES

- ^m Pediatric HL risk stratification is evolving from traditional pediatric risk group definitions to a stage-based system aligned with adult staging. See [NCCN Guidelines for Hodgkin Lymphoma \(Adult\)](#). Clinicians should be aware that historical pediatric risk classifications and contemporary stage-based approaches may overlap during this transition.
- ⁿ Mauz-Korholz C, et al. J Clin Oncol 2010;28:3680-3686; Mauz-Korholz C, et al. Lancet Oncol 2022;23:125-137; Mauz-Korholz C, et al. Lancet Oncol 2023;24:252-261.
- ^o Friedman DL, et al. J Clin Oncol 2014;32:3651-3658.
- ^p Involved-site RT (ISRT) can safely replace involved-field RT (IFRT) ([PHL-F](#)).
- ^q Kelly KM, et al. Br J Haematol 2019;187:39-48; Castellino SM, et al. N Engl J Med 2022;387:1649-1660.
- ^r Herrera AF, et al. N Engl J Med 2024;391:1379-1389.
- ^s Metzger ML, et al. J Clin Oncol 2021;39:2276-2283.
- ^t An E-lesion is defined as a contiguous infiltration of a lymph node mass into extralymphatic structures or organs (eg, lung, bone). Pleural and pericardial involvement should be considered E-lesions, but a pleural or pericardial effusion alone is not considered an E-lesion. Disease that extends beyond the lymphatic system without adjacent lymphatic involvement is considered stage IV; liver or bone marrow involvement is always considered stage IV disease (Zijtregtop EAM, et al. Blood Adv 2023;7:6303-6319). CNS disease is considered extra-axial.


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



CLINICAL PRESENTATION: CHL

PRIMARY TREATMENT^{u,v,w}

RESPONSE^x



^u Regimens are based off of studies with pediatric data.

^v [Principles of Systemic Therapy \(PHL-E\)](#).

^w For growth factor recommendations, see [Select Principles of Supportive Care \(PHL-G\)](#).

^x [Principles of Criteria for Response-Adapted Radiation Therapy \(PHL-A\)](#).

^y See [PHL-2](#) for risk factors as defined by EuroNET-PHL and COG.

^z FDG-PET/CT or FDG-PET/MRI and contrast-enhanced diagnostic CT or MRI of original sites of disease if not included with FDG-PET and treatment approach includes need for assessment of anatomic sites of disease for evaluation of response. See [Principles of Imaging \(PHL-C\)](#).

^{aa} Omission of ISRT should be more strongly considered for patients who meet the GPOH-2002 response criteria (Mauz-Korholz C, et al. J Clin Oncol 2010;28:3680-3686).

^{bb} [Principles of Radiation Therapy \(PHL-F\)](#).

[See Evidence Blocks on EB-1](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: CHL

PRIMARY TREATMENT^{u,v,w}

RESPONSE^x

ADDITIONAL TREATMENT^{v,x}

Preferred

Ongoing clinical trial

or

ABVE-PC
x 2 cycles
(category 1)
(AHOD0031)
(not for stage
IIBX)

Response
assessment^z

Adequate
response (RER)^{dd}

ABVE-PC
x 2 cycles

End of therapy
assessment^z

See Follow-up ([PHL-8](#))
or
If < complete
response (CR),
ISRT^{p,bb} to all sites

Inadequate
response (SER)^{dd}

ABVE-PC
x 2 cycles

ISRT^{p,bb}

End of
therapy
assessment^z

See Follow-up
([PHL-8](#))
or
If concern for
persistent
disease, see
Therapy for
relapsed or
refractory
disease ([PHL-9](#))

or

Bv-AVE-PC
x 2 cycles^{cc}
(category 1)
(AHOD1331)
(stage IIBX
only)

Response
assessment^z

Adequate
response (RRL)^{dd}

Bv-AVE-PC
x 3 cycles

ISRT^{bb} to sites
of LMA^{dd}

Inadequate
response (SRL)^{dd}

Bv-AVE-PC
x 3 cycles

ISRT^{bb} to sites of LMA^{dd} and
SRL^{dd} and boost to sites of
inadequate response at end of
systemic therapy (EOST)

or

Other Recommended

[PHL-4A](#)

Stage
I–II CHL
with risk
factors^y



Footnotes on [PHL-4B](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: CHL

**PRIMARY
TREATMENT^{u,v,w}**
(continued from
[PHL-4](#))

RESPONSE^x

ADDITIONAL TREATMENT^{v,x}

Other Recommended

OEPA x 2 cycles
(category 1)
(EuroNet-PHL-C1)

Response
assessment^z

Adequate
response

COPDAC x 2 cycles
(not for stage IIBE)
COPDAC x 4 cycles
(category 2B) (stage IIBE
and IIBX only)

Inadequate
response

COPDAC x 2 cycles
(not for stage IIBE)
COPDAC x 4 cycles
(category 2B) (stage
IIBE and IIBX only)

ISRT^{bb} to
all sites and
boost to sites
of inadequate
response
at end of
chemotherapy

End of
therapy
assessment^z

See Follow-up
([PHL-8](#))
or
If concern for
persistent
disease, see
Therapy for
relapsed or
refractory
disease ([PHL-9](#))

or

AEPA x 2 cycles^{cc}
(HLHR13) (stage
IIB and IIBX only)

Response
assessment^z

Adequate
response

CAPDAC x
4 cycles

Inadequate
response

CAPDAC
x 4 cycles

Residual-node RT
(RNRT)^{bb} to sites
of inadequate
response

End of
therapy
assessment^z

ABVD

or

Consider additional adult regimens if aged >18 y →

[NCCN Guidelines for Hodgkin Lymphoma \(Adult\)](#)

[See Evidence Blocks on EB-2](#)

Footnotes on [PHL-4B](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRIMARY TREATMENT OF STAGE I-II CHL WITH RISK FACTORS FOOTNOTES

^p ISRT can safely replace IFRT ([PHL-F](#)).

^u Regimens are based off of studies with pediatric data.

^v [Principles of Systemic Therapy \(PHL-E\)](#).

^w For growth factor recommendations, see [Select Principles of Supportive Care \(PHL-G\)](#).

^x [Principles of Criteria for Response-Adapted Radiation Therapy \(PHL-A\)](#).

^y See [PHL-2](#) for risk factors as defined by EuroNET-PHL and COG.

^z FDG-PET/CT or FDG-PET/MRI and contrast-enhanced diagnostic CT or MRI of original sites of disease if not included with FDG-PET and treatment approach includes need for assessment of anatomic sites of disease for evaluation of response. See [Principles of Imaging \(PHL-C\)](#).

^{bb} [Principles of Radiation Therapy \(PHL-F\)](#).

^{cc} *Pneumocystis jiroveci* pneumonia (PJP) prophylaxis is recommended for all Bv-containing regimens.

^{dd} RER = rapid early response; SER = slow early response; LMA = large mediastinal adenopathy; RRL = rapidly responding lesion; SRL = slow responding lesion.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: CHL



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Footnotes on [PHL-5B](#)



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: CHL

PRIMARY TREATMENT^{u,v,w}
(continued from
[PHL-5](#))

RESPONSE^x

ADDITIONAL TREATMENT^{v,x}

Useful in Certain Circumstances

Bv-AVD x 6 cycles (S1826)^{cc,ee}
(if unable to receive
or tolerate a checkpoint
inhibitor [CPI])

Response
assessment^z

Adequate
response

Inadequate
response

See Follow-up
([PHL-8](#))

Consider RSRT^{bb} to
sites of inadequate
response
or
see Therapy for
relapsed or refractory
disease ([PHL-9](#))

or

ABVE-PC
x 2 cycles
(category 2B)
(AHOD0031)
(stage IIIA
only)

Response
assessment^z

Adequate
response (RER)^{dd}

Inadequate
response (SER)^{dd}

ABVE-PC
x 2 cycles

End of therapy
assessment^z

See Follow-up ([PHL-8](#))
or
If <CR, ISRT^{p,bb} to all
sites

See Follow-up ([PHL-8](#))
or
If concern for
persistent disease, see
Therapy for relapsed
or refractory disease
([PHL-9](#))

ABVE-PC
x 2 cycles

ISRT^{p,bb}

End of therapy
assessment^z

or

Consider BrECADD^{hh} or
additional adult regimens if
aged >18 y

[NCCN Guidelines for Hodgkin Lymphoma \(Adult\)](#)

[See Evidence Blocks on EB-3](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Footnotes on [PHL-5B](#)



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRIMARY TREATMENT OF STAGE III–IV CHL FOOTNOTES

- ^c CPIs should be used with caution in those with active or previously documented autoimmune disease and/or current use of immunosuppressive agents.
- ^p ISRT can safely replace IFRT ([PHL-F](#)).
- ^u Regimens are based off of studies with pediatric data.
- ^v [Principles of Systemic Therapy \(PHL-E\)](#).
- ^w For growth factor recommendations, see [Select Principles of Supportive Care \(PHL-G\)](#).
- ^x [Principles of Criteria for Response-Adapted Radiation Therapy \(PHL-A\)](#).
- ^z FDG-PET/CT or FDG-PET/MRI and contrast-enhanced diagnostic CT or MRI of original sites of disease if not included with FDG-PET and treatment approach includes need for assessment of anatomic sites of disease for evaluation of response. See [Principles of Imaging \(PHL-C\)](#).
- ^{bb} [Principles of Radiation Therapy \(PHL-F\)](#).
- ^{cc} PJP prophylaxis is recommended for all Bv-containing regimens.
- ^{dd} RER = rapid early response; SER = slow early response; LMA = large mediastinal adenopathy; RRL = rapidly responding lesion; SRL = slow responding lesion.
- ^{ee} This regimen has only been studied in patients ≥12 years of age with stage III–IV disease.
- ^{ff} Nivolumab is dosed per kg in patients <18 years of age. See [Principles of Systemic Therapy \(PHL-E\)](#).
- ^{gg} False-positive findings may be observed on FDG-PET/CT or FDG-PET/MRI after treatment with a regimen that includes a CPI given their potential for delayed response or pseudo-progression. Clinical judgment is often necessary in such settings. Repeat imaging at short intervals (per LYRIC recommendations) may be considered prior to obtaining a biopsy. Cheson BD, et al. Blood 2016;128:2489-2496. See [Principles of Imaging \(PHL-C\)](#).
- ^{hh} BrECADD regimen: brentuximab vedotin, etoposide, cyclophosphamide, doxorubicin, dacarbazine, and dexamethasone.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: NLPHL

PRIMARY TREATMENT^{u,v,jj}



[See Evidence Blocks on EB-4](#)

^u Regimens are based off of studies with pediatric data.

^v [Principles of Systemic Therapy \(PHL-E\)](#).

^{bb} [Principles of Radiation Therapy \(PHL-F\)](#).

ⁱⁱ Although AHOD03P1 only included completely resected single-node disease for observation, additional retrospective data support observation after complete resection of stage I, multiple nodes (Appel BE, et al. J Clin Oncol 2016;34:2372-2379; Mauz-Korholz C, et al. Cancer 2007;110:179-185; Punnett A, et al. Br J Haematol 2023;201:1081-1087; Shankar A, et al. Br J Haematol 2022;197:679-690). However, extensive node or repeat node dissection is discouraged.

^{jj} The following risk factors may also inform decisions about primary treatment of NLPHL: age ≥45 years, hemoglobin <10.5 g/dL, and splenic involvement (Binkley MS, et al. J Clin Oncol 2024;42:2271-2280).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: NLPHL



^u Regimens are based off of studies with pediatric data.

^v [Principles of Systemic Therapy \(PHL-E\)](#).

^x [Principles of Criteria for Response-Adapted Radiation Therapy \(PHL-A\)](#).

^z FDG-PET/CT or FDG-PET/MRI and contrast-enhanced diagnostic CT or MRI of original sites of disease if not included with FDG-PET and treatment approach includes need for assessment of anatomic sites of disease for evaluation of response. See [Principles of Imaging \(PHL-C\)](#).

^{bb} [Principles of Radiation Therapy \(PHL-F\)](#).

ⁱⁱ Although AHOD03P1 only included completely resected single-node disease for observation, additional retrospective data support observation after complete resection of stage I, multiple nodes (Appel BE, et al. J Clin Oncol 2016;34:2372-2379; Mauz-Korholz C, et al. Cancer 2007;110:179-185; Punnett A, et al. Br J Haematol 2023;201:1081-1087; Shankar A, et al. Br J Haematol 2022;197:679-690). However, extensive node or repeat node dissection is discouraged.

^{jj} The following risk factors may also inform decisions about primary treatment of NLPHL: age ≥45 years, hemoglobin <10.5 g/dL, and splenic involvement (Binkley MS, et al. J Clin Oncol 2024;42:2271-2280).

^{kk} Data are limited on the use of rituximab for early-stage NLPHL.

^{ll} Advanced-stage NLPHL is rare in pediatric patients. Confirm pathologic diagnosis prior to treatment. See [Principles of Pathology \(PHL-B\)](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

FOLLOW-UP AFTER COMPLETION OF TREATMENT AND MONITORING FOR LATE EFFECTS

Pediatric CHL		
Disease Surveillance/ Follow-up After Completion of Treatment	• Interim H&P: ▶ Every 3–4 mo for 1–2 y, then every 6–12 mo until year 3, then annually until 5 y • Laboratory studies: ▶ CBC with differential, ESR or CRP, chemistry profile as clinically indicated. ▶ Thyroid-stimulating hormone (TSH) at least annually if radiation therapy (RT) to neck. • Consider PFTs (if bleomycin, pulmonary RT, significant pulmonary involvement, or other clinical concerns). • Immunizations: ▶ Annual influenza vaccine is recommended, even during therapy. ▶ Other vaccines as per CDC Guidelines, typically starting 6 mo after completion of therapy (see Children's Oncology Group Survivorship Guidelines). • Counsel regarding reproduction, health habits, and psychosocial issues (for AYA, see NCCN Guidelines for AYA Oncology). • Screen for delayed immune-mediated adverse events (imAEs), such as hypothyroidism, which may present within the first 2 years following completion of treatment, and may be needed for those who received a CPI (Durbin SM, et al. JAMA Netw Open 2025;8:e252668). • It is recommended that the patient be provided with a treatment summary at the completion of therapy, including details of RT and chemotherapy, including cumulative anthracycline and alkylator dosages given.	• Imaging: ▶ Consider end-of-therapy ECHO. ▶ Imaging studies are only recommended when relapse is suspected, because most patients will clinically declare themselves and there is no survival advantage in pre-emptive imaging. ▶ If clinical concern, chest x-ray PA and lateral views, CT with contrast, or MRI of original sites of disease may be performed and followed at 3- to 6-mo intervals up to 2 y following completion of therapy. ▶ MRI is acceptable in place of CT scan for neck/abdomen/pelvis, but not for chest; diagnostic CT of chest is needed to evaluate lung parenchyma. ▶ FDG-PET/CT or FDG-PET/MRI if previous FDG-PET was positive (Deauville 4–5), to confirm CR at the end of all prescribed therapy including RT. Once negative, repeat FDG-PET should not be done unless evaluating suspicious findings on H&P or CT or MRI. ◊ Wait at least 8–12 wk after end of RT to perform FDG-PET to minimize false-positive results. ◊ Surveillance FDG-PET is discouraged due to risk for false positives. ▶ If concern for relapse, management decisions should not be based on FDG-PET scan alone; clinical and pathologic correlation is needed. See Principles of Pathology (PHL-B) and Therapy for Relapsed or Refractory Disease (PHL-9).
Monitoring for Late Effects (≥2 years after completion of systemic therapy)	• Appropriate screening and counseling related to: thyroid, cardiac, pulmonary, bone, fertility, and reproductive health; subsequent cancers (with special attention to thyroid and breast cancer); and other treatment-associated late effects (see Children's Oncology Group Survivorship Guidelines)	



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

NCCN Guidelines Version 1.2026


Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

CLINICAL PRESENTATION: CHL

SUSPECTED RELAPSED/ REFRACTORY DISEASE

RE-INDUCTION THERAPYⁿⁿ



^{bb} [Principles of Radiation Therapy \(PHL-F\)](#).

^{gg} False-positive findings may be observed on FDG-PET/CT or FDG-PET/MRI after treatment with a regimen that includes a CPI given their potential for delayed response or pseudo-progression. Clinical judgment is often necessary in such settings. Repeat imaging at short intervals (per LYRIC recommendations) may be considered prior to obtaining a biopsy. Cheson BD, et al. Blood 2016;128:2489-2496. See [Principles of Imaging \(PHL-C\)](#).

^{mm} A biopsy must be obtained to confirm relapse and pathology. See [Principles of Pathology \(PHL-B\)](#).

ⁿⁿ There are no data to support a superior outcome with any of the treatment modalities. Individualized treatment is recommended.

^{oo} Consider consultation with radiation oncology depending on level of clinical suspicion.

^{pp} [Principles of Systemic Therapy for Relapsed or Refractory Disease \(PHL-E, 3 of 6\)](#).

^{qq} It is reasonable to try multiple different re-induction regimens as needed prior to ASCR to minimize disease burden with a goal of achieving a metabolic CR prior to transplant. If less than a metabolic CR, proceed to subsequent therapy.

^{rr} Recommendations for those who may avoid ASCR: initial stage other than IIIB or IVB, no prior exposure to RT, duration of first CR >1 year, absence of extranodal disease or B symptoms at relapse, and adequate response to <4 cycles of chemotherapy in those with relapse between 3 and 12 months (Hoppe BS, et al. JAMA Oncol 2025;11:340-342; Daw S, et al. JAMA Oncol 2025;11:249-257; Daw S, et al. JAMA Oncol 2025;11:258-267).

^{ss} Strongly consider RT for selected sites that have not been previously irradiated.

^{tt} Therapy following a Deauville 5 response to reinduction for relapsed/refractory disease is not well established. Individualized treatment is necessary. Referral to or consultation with a center of expertise is strongly encouraged.

^{uu} Allogeneic transplant is an option in select patients who experience relapse post-ASCT as a category 3 recommendation.

^{vv} RT is usually performed as consolidation after transplant. If unable to get to a metabolic CR, then RT can be used prior to transplant.

^{ww} A higher risk of engraftment syndrome may be observed in patients who have been treated with CPIs (Park JJ, et al. Clin Lymphoma Myeloma Leuk 2021;21:e726-e730).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF CRITERIA FOR RESPONSE-ADAPTED RADIATION THERAPY

Regimen	Criteria for RT
OEPA/OEPA-COPDAC (EuroNet-PHL-C1)^{1,2,3}	• < CR on imaging after 2 cycles of OEPA (applies to patients in TG1 [stage IA, IB, and IIA without E] on GPOH HD-2002) ▶ CR: FDG-PET negative (Deauville 1–2) and volume reduction >95% and ≤2 mL ▶ CR, unconfirmed (CRu): Volume reduction >75% or ≤2 mL • All patients in TG2 (stage IA/B + E, IIA + E, IIB, and IIIA) and TG3 (stage IIB + E, IIIA + E, IIIB, and IVA/B ± E) on HD-2002 received RT • Patients in all TGs on C1 received RT only if FDG-PET positive (Deauville 3–5) or not in at least partial response (PR) after 2 cycles of OEPA ▶ PR: No CR or CRu and >50% volume reduction or residual tumor volume <5 mL ▶ Consider boost for residual mass >100 mL or <75% volume reduction and minimum volume >5 mL at early response assessment Note: Volume = (a x b x c)/2 where a, b, and c are three dimensions of a node or conglomerate
ABVE-PC (AHOD0031)⁴	• SER on imaging after 2 cycles if <60% reduction in product of perpendicular diameters (PPD)^a for all target lesions or • RER on imaging after 2 cycles if not in CR on imaging after 4 cycles ▶ CR: ≥80% reduction in PPD^a with negative FDG-PET at end of therapy (comparable to Deauville 1–2) • Consider boost for persistent FDG-PET–positive (Deauville 3–5) lesions at end of chemotherapy

^a PPD = Transverse x axial plane.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**

[Continued](#)
[References](#)

PHL-A
1 OF 3



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF CRITERIA FOR RESPONSE-ADAPTED RADIATION THERAPY

Regimen	Criteria for RT
Bv-AVE-PC (AHOD1331)^{5,6}	• SRL on imaging after 2 cycles⁶ ▶ Inadequate or SRL: Deauville 4–5 ▶ Adequate or RRL: Deauville ≤3 • All LMA • Boost for persistent FDG-PET–positive lesions (Deauville 3–5) at end of chemotherapy • Consider treating without a boost for LMA that was Deauville 3 after 2 cycles and continues to be Deauville 3 after 5 cycles
AEPA-CAPDAC (HLHR13)⁷	• Residual FDG-PET–positive lesions (Deauville 4–5) after 2 cycles of therapy, or • FDG-PET–negative (Deauville 1–3) lesions with <75% anatomic response (as measured by the product of 2 perpendicular diameters of lesions by CT or MRI)
AVPC - mixed cellularity CHL only (AHOD0431)⁸	• <CR on imaging after 3 cycles ▶ CR: ≥80% reduction in PPD^a and FDG-PET negative; only mediastinal nodes >2 cm ▶ FDG-PET positive (Deauville 3–5): Uptake greater than mediastinal blood pool
AVPC - NLPHL (AHOD03P1)⁹	• Residual FDG-PET–positive lesions after 3 cycles of AVPC, or • FDG-PET–negative lesions with <80% anatomic response (as measured by the product of 2 perpendicular diameters of lesions by CT or MRI), or • Not returned to normal size with residual nodal max dimension >2.0 cm
Nivolumab-AVD (S1826)¹⁰	• Residual FDG-PET–positive lesions after 6 cycles of therapy ▶ Deauville 4–5, only 1–2 lesions, ≥30% reduction in transverse diameter, residual nodal mass ≥2.5 cm in axial diameter, or residual extranodal lesion >1 cm in axial diameter^b
Bv-AVD (S1826)¹⁰	• Residual FDG-PET–positive lesions after 6 cycles of therapy ▶ Deauville 4–5, only 1–2 lesions, ≥30% reduction in transverse diameter, residual nodal mass ≥2.5 cm in axial diameter, or residual extranodal lesion >1 cm in axial diameter^b

^a PPD = Transverse x axial plane.

^b Patients with ≥3 Deauville 4-5 lesions and/or a lesion that that did not achieve ≥30% reduction in transverse diameter were considered to have refractory disease.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**

References

PHL-A
2 OF 3



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF CRITERIA FOR RESPONSE-ADAPTED RADIATION THERAPY REFERENCES

- ¹ Mauz-Körholz C, Hasenclever D, Dörffel W, et al. Procarbazine-free OEPA-COPDAC chemotherapy in boys and standard OPPA-COPP in girls have comparable effectiveness in pediatric Hodgkin's lymphoma: the GPOH-HD-2002 study. *J Clin Oncol* 2010;28:3680-3686.
- ² Mauz-Körholz C, Landman-Parker J, Balwierz W, et al. Response-adapted omission of radiotherapy and comparison of consolidation chemotherapy in children and adolescents with intermediate-stage and advanced-stage classical Hodgkin lymphoma (EuroNet-PHL-C1): a titration study with an open-label, embedded, multinational, non-inferiority, randomised controlled trial. *Lancet Oncol* 2022;23:125-137. Erratum in *Lancet Oncol* 2022;23:125-137.
- ³ Mauz-Körholz C, Landman-Parker J, Fernández-Teijeiro A, et al. Response-adapted omission of radiotherapy in children and adolescents with early-stage classical Hodgkin lymphoma and an adequate response to vincristine, etoposide, prednisone, and doxorubicin (EuroNet-PHL-C1): a titration study. *Lancet Oncol* 2023;24:252-261.
- ⁴ Friedman DL, Chen L, Wolden S, et al. Dose-intensive response-based chemotherapy and radiation therapy for children and adolescents with newly diagnosed intermediate-risk hodgkin lymphoma: a report from the Children's Oncology Group Study AHOD0031. *J Clin Oncol* 2014;32:3651-3658.
- ⁵ Castellino SM, Pei Q, Parsons SK, et al. Brentuximab vedotin with chemotherapy in pediatric high-risk Hodgkin's lymphoma. *N Engl J Med* 2022;387:1649-1660.
- ⁶ Flerlage JE, Kelly KM, Beishuizen A, et al. Staging Evaluation and Response Criteria Harmonization (SEARCH) for Childhood, Adolescent and Young Adult Hodgkin Lymphoma (CAYAHL): Methodology statement. *Pediatr Blood Cancer* 2017;64:e26421.
- ⁷ Metzger ML, Link MP, Billett AL, et al. Excellent outcome for pediatric patients with high-risk Hodgkin lymphoma treated with brentuximab vedotin and risk-adapted residual node radiation. *J Clin Oncol* 2021;39:2276-2283.
- ⁸ Keller FG, Castellino SM, Chen L, et al. Results of the AHOD0431 trial of response adapted therapy and a salvage strategy for limited stage, classical Hodgkin lymphoma: A report from the Children's Oncology Group. *Cancer* 2018;124:3210-3219.
- ⁹ Appel BE, Chen L, Buxton AB, et al. Minimal treatment of low-risk, pediatric lymphocyte-predominant Hodgkin lymphoma: A report from the Children's Oncology Group. *J Clin Oncol* 2016;34:2372-2379.
- ¹⁰ Herrera AF, LeBlanc M, Castellino SM, et al. Nivolumab+AVD in advanced-stage classic Hodgkin's lymphoma. *N Engl J Med* 2024;391:1379-1389.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGY

Histologic Classification

- **Diagnosis should be established according to guidelines in the 2022 World Health Organization (WHO) Classification of Haematolymphoid Tumours, 5th edition or the 2022 International Consensus Classification (ICC) of Mature Lymphoid Neoplasms.^{1,2}**
- **There are two types of Hodgkin lymphoma (HL): CHL and NLPHL. Distinction between these types is important for therapy and prognosis.**
 - ▶ **Per WHO 2022, NLPHL remains under the family of HL, while ICC 2022 replaces the term NLPHL with nodular lymphocyte-predominant B-cell lymphoma (NLPBL).^{1,2}**
 - ▶ **Both classifications recognize the clinical and biologic similarities of NLPHL to an indolent B-cell lymphoma and NLPBL is an acceptable term per WHO 2022.¹**

CHL

- **There are four morphologic variants of CHL^a:**
 - ▶ **Nodular sclerosis**
 - ▶ **Mixed cellularity**
 - ▶ **Lymphocyte-rich**
 - ▶ **Lymphocyte-depleted**
- **CHL subtyping is not necessary for treatment in the vast majority of cases and may not be possible in all cases.^a If considering treatment based on the AHOD0431 trial,³ discussion with a pediatric pathologist or hematopathologist is recommended to determine if tissue is sufficient to establish a diagnosis of mixed-cellularity subtype.**
- **CHL can occur in patients with immunodeficiency (primary immunodeficiency, HIV infection, post-transplant immunodeficiency, and iatrogenic immunodeficiency). Other polymorphic lymphoproliferative disorders and Hodgkin-like lesions are also associated with immunodeficiency and should be distinguished from CHL since management and treatment recommendations differ. These are challenging cases and expert hematopathology evaluation is suggested. Referral to a center of expertise may be necessary.**

NLPHL

- **There are six immunoarchitectural patterns of NLPHL and a mixture of patterns is commonly seen histologically^{b,4}:**
 - ▶ **B-cell-rich nodular (pattern A)**
 - ▶ **Serpiginous/interconnected nodular (pattern B)**
 - ▶ **Nodular with prominent extranodular lymphocyte-predominant (LP) cells (pattern C)**
 - ▶ **T-cell-rich nodular (pattern D)**
 - ▶ **T-cell-rich diffuse or T-cell/histiocyte-rich large B-cell lymphoma (THRLBCL)-like (pattern E)**
 - ▶ **Diffuse B-cell-rich (pattern F)**
- **Including variant histologic patterns in NLPHL in the pathology report may be beneficial, where possible, as T-cell-rich and/or diffuse patterns may be associated with higher risk of disease progression, transformation, and relapse and shorter time to relapse; however, variant histologic patterns are not currently utilized to guide treatment selection.^{5,6,7}**

^a The different morphologic variants of CHL have different clinicopathologic associations and differential diagnoses. Refer to the 2022 WHO Classification or ICC 2022 for more details.

^b The different immunoarchitectural patterns of NLPHL have different prognostic implications and differential diagnoses. In particular, pattern E (T-cell-rich diffuse or THRLBCL-like) is similar to de novo THRLBCL at the molecular level and behaves more aggressively than other NLPHL variants (Hartmann S, et al. PLoS One 2013;8:e78812). There are insufficient data directly comparing the clinical behavior of NLPHL with predominant T-cell-rich diffuse histology with that of THRLBCL to determine whether such cases are best managed like NLPHL or a large B-cell lymphoma.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

[Continued](#)
[References](#)

PHL-B
1 OF 5



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGY

Tissue Adequacy for Diagnosis

- An excisional or incisional biopsy where possible is recommended. Fine-needle aspiration (FNA) is not an appropriate diagnostic test.^c
- For CHL, ample tissue may be necessary to exclude other entities in the differential diagnosis^d and for specific morphologic subtyping.^e A core biopsy may be appropriate in some settings.^f
- For NLPHL, ample tissue may be necessary to assess for the presence of variant histologic patterns and, in cases with predominantly T-cell-rich or diffuse patterns (patterns D–F), to exclude other entities in the differential diagnosis, such as THRLBCL, peripheral T-cell lymphoma, or another non-Hodgkin lymphoma (NHL). For cases with predominantly T-cell-rich histology, at least one nodule with LP cells in a B-cell-rich background is needed to distinguish a variant NLPHL pattern from THRLBCL. Examination of additional sections of submitted tissue and/or immunohistochemical staining of additional slides may be necessary in such cases.
- NLPHL may be associated with concomitant diffuse large B-cell lymphoma (DLBCL) at the same or different site.⁸⁻¹⁰ Excision specimens should be adequately sampled. A re-biopsy may be considered in situations where imaging findings are discordant with the rendered histopathologic diagnosis or discordant between different sites. Splenic involvement has been associated with increased risk of transformation.^{11,12}

^c Sparse neoplastic cells, extensive fibrosis, and presence of Reed Sternberg-like cells in some conditions other than in HL are some reasons a limited biopsy may not be diagnostic.

^d For example, mediastinal gray zone lymphoma or rare composite tumors of CHL and primary mediastinal large B-cell lymphoma may not be demonstrable in limited biopsies.

^e Fibrotic bands completely surrounding nodules are important in distinguishing nodular sclerosis CHL from mixed cellularity CHL but may not be demonstrable in small biopsies.

^f For example, less accessible anatomic sites such as mediastinal mass with sedation risks or retroperitoneum.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**

[Continued](#)
[References](#)

PHL-B
2 OF 5



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGY

Immunohistochemical Considerations and Ancillary Testing

- Consider clinical differential diagnoses (eg, T-cell lymphoblastic lymphoma) and pathologic differential diagnoses: HL versus NHL,⁹ CHL versus NLPHL versus progressive transformation of germinal centers, HL versus infection (eg, cytomegalovirus [CMV], Epstein-Barr virus [EBV]), and HL versus reactive proliferations.^h
- Diagnosis is based on morphologic AND immunohistochemical findings.
- Typical immunophenotype of HL:
 - ▶ CHL: Neoplastic Hodgkin-Reed-Sternberg (HRS) cells are PAX5+ (weaker expression than background B lymphocytes), CD30+, CD15+, CD45-, or CD3-. CD20 may be variably expressed in 20%–40% of cases and, when present, is typically expressed by a subset of HRS cells. This serves as an essential panel of markers for immunohistochemical evaluation of CHL. Evaluation of an expanded panel of markers (ie, CD79a-, ALK-, MUM1+, OCT2-/weak, BOB1-/weak) and assessing for EBV with EBER-ISHⁱ should be considered in cases with equivocal or imperfect morphologic or immunophenotypic features or to exclude entities in the differential diagnosis.
 - ▶ NLPHL: Neoplastic LP cells are PAX5+, CD20+, OCT2+ (strong), CD30-, CD15-, or CD3-. They are also CD45+, CD79a+, BCL6+, EMA+, or MUM1-/weak. Characteristics of the immune microenvironment (identification of small B cells, T-follicular helper cells, and follicular dendritic cells) are also helpful in the diagnostic workup.
- EBV+ CHL cases may benefit from additional studies, such as EBV serology and evaluation for underlying immunodeficiency. EBV+ LP cells have been reported only rarely in NLPHL and this finding should prompt consideration of other entities in the differential diagnosis.¹³
- Flow cytometry is not helpful in diagnosing HL.^j However, it may be helpful in the evaluation of other entities in the clinical or pathologic differential diagnosis.
- While a few papers have reported on molecular/genetic findings in HLs, molecular/genetic testing does not currently have established diagnostic or therapeutic utility in these diseases and is not recommended as part of standard routine practice. For challenging cases where the differential diagnosis between CHL and a T-cell lymphoma remains after extensive morphologic and immunophenotypic assessment, B-cell and T-cell molecular clonality testing may be helpful. Note that HRS cells in bonafide CHL can occasionally show aberrant expression of T-cell antigens, but the majority of CHL is negative for T-cell clonality.¹⁴ B-cell clonality studies are often negative in CHL due to the sparse tumor cells. These factors need to be taken into consideration in the evaluation of these infrequent challenging cases.

⁹ NHL examples, primary mediastinal large B-cell lymphoma, ALK+ anaplastic large cell lymphoma, THRLBCL, EBV+ DLBCL, and peripheral T-cell lymphoma.

^h For example, reactive lymph node with CD30+ immunoblasts (vs. CHL) and progressive transformation of germinal centers (vs. NLPHL) and granulomatous lymphadenitis (vs. CHL or NLPHL).

ⁱ EBER-ISH = EBV-encoded RNA (EBER) in situ hybridization (ISH).

^j Identification of CD4+ CD8dim+ T cells can support a diagnosis of NLPHL, but this population may also be seen in progressive transformation of germinal centers. Neoplastic cells in CHL may also be identified using sophisticated flow cytometry techniques, which are not readily available.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

[Continued](#)
[References](#)

PHL-B
3 OF 5



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGY

Pathology Considerations in the Relapse/Refractory Disease Setting

- Pathologic confirmation is necessary to confirm relapse or refractory disease given the risk for transformation and high false-positive rate of FDG-PET/CT. Re-biopsy is also recommended for residual FDG-PET-avid disease at the end of therapy.¹⁵ In the case of NLPHL, such residual FDG-PET-avid disease could represent foci of concurrent or transformed DLBCL.
- If original diagnosis slides are available, limited immunohistochemical evaluation may be performed on the relapse/refractory specimen.
- For CHL cases, consider the possibility of misdiagnosis at original presentation (eg, primary mediastinal large B-cell lymphoma, mediastinal gray zone lymphoma,^{16,17} or other lymphoma subtypes).
- For NLPHL cases, consider the possibility of DLBCL (either conventional with sheets of large cells or THRLBCL-like) transformation from NLPHL¹⁸ or reactive lymph node with progressive transformation of germinal centers.¹⁹ Although advanced-stage NLPHL can occur, particularly in cases with variant morphology, this is uncommon. Therefore, expert hematopathology re-evaluation of original pathology slides is suggested for patients with advanced disease. Misdiagnosed CHL or THRLBCL are diagnostic considerations in this setting and should be excluded. Referral to a center of expertise may be necessary.
- Prior monoclonal antibody therapy targeting CD30 (for CHL) or CD20 (for NLPHL) may result in weak or negative staining for these antigens by immunohistochemistry.
- There are insufficient data to recommend programmed death ligand 1 (PD-L1) testing by immunohistochemistry as a prerequisite for immune CPI therapy. Robust cut-offs for optimally predicting response to immune CPI therapy have not been established.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGY REFERENCES

- ¹ Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Lymphoid Neoplasms. *Leukemia* 2022;36:1720-1748.
- ² Campo E, Jaffe ES, Cook JR, et al. The International Consensus Classification of Mature Lymphoid Neoplasms: A report from the Clinical Advisory Committee. *Blood* 2022;140:1229-1253.
- ³ Keller FG, Castellino SM, Chen L, et al. Results of the AHOD0431 trial of response adapted therapy and a salvage strategy for limited stage, classical Hodgkin lymphoma: A report from the Children's Oncology Group. *Cancer* 2018;124:3210-3219.
- ⁴ Fan Z, Natkunam Y, Bair E, et al. Characterization of variant patterns of nodular lymphocyte predominant hodgkin lymphoma with immunohistologic and clinical correlation. *Am J Surg Pathol* 2003;27:1346-1356.
- ⁵ Hartmann S, Eichenauer DA, Plütschow A, et al. The prognostic impact of variant histology in nodular lymphocyte-predominant Hodgkin lymphoma: a report from the German Hodgkin Study Group (GHSG). *Blood* 2013;122:4246-4252; quiz 4292.
- ⁶ Hartmann S, Plütschow A, Mottok A, et al. The time to relapse correlates with the histopathological growth pattern in nodular lymphocyte predominant Hodgkin lymphoma. *Am J Hematol* 2019;94:1208-1213.
- ⁷ Binkley MS, Flerlage JE, Savage KJ, et al. International prognostic score for nodular lymphocyte-predominant Hodgkin lymphoma. *J Clin Oncol* 2024;42:2271-2280.
- ⁸ Cotta CV, Coleman JF, Li S, Hsi ED. Nodular lymphocyte predominant Hodgkin lymphoma and diffuse large B-cell lymphoma: a study of six cases concurrently involving the same site. *Histopathology* 2011;59:1194-1203.
- ⁹ Hartmann S, Eray M, Döring C, et al. Diffuse large B cell lymphoma derived from nodular lymphocyte predominant Hodgkin lymphoma presents with variable histopathology. *BMC Cancer* 2014;14:332.
- ¹⁰ Song JY, Egan C, Bouska AC, et al. Genomic characterization of diffuse large B-cell lymphoma transformation of nodular lymphocyte-predominant Hodgkin lymphoma. *Leukemia* 2020;34:2238-2242.
- ¹¹ Al-Mansour M, Connors JM, Gascoyne RD, et al. Transformation to aggressive lymphoma in nodular lymphocyte-predominant Hodgkin's lymphoma. *J Clin Oncol* 2010;28:793-799.
- ¹² Kenderian SS, Habermann TM, Macon WR, et al. Large B-cell transformation in nodular lymphocyte-predominant Hodgkin lymphoma: 40-year experience from a single institution. *Blood* 2016;127:1960-1966.
- ¹³ Huppmann AR, Nicolae A, Slack GW, et al. EBV may be expressed in the LP cells of nodular lymphocyte-predominant Hodgkin lymphoma (NLPHL) in both children and adults. *Am J Surg Pathol* 2014;38:316-324.
- ¹⁴ Venkataraman G, Song JY, Tzankov A, et al. Aberrant T-cell antigen expression in classical Hodgkin lymphoma is associated with decreased event-free survival and overall survival. *Blood* 2013;121:1795-1804.
- ¹⁵ El-Galaly TC, Gormsen LG, Hutchings M. PET/CT for staging; past, present, and future. *Semin Nucl Med* 2018;48:4-16.
- ¹⁶ Pilichowska M, Pittaluga S, Ferry J, et al. Clinicopathologic consensus study of gray zone lymphoma with features intermediate between DLBCL and classical HL. *Blood Adv* 2017;1:2600-2609.
- ¹⁷ Traverse-Glehen A, Pittaluga S, Gaulard P, et al. Mediastinal gray zone lymphoma: the missing link between classic Hodgkin's lymphoma and mediastinal large B-cell lymphoma. *Am J Surg Pathol* 2005;29:1411-1421.
- ¹⁸ Al-Mansour M, Connors J, Gascoyne R, et al. Transformation to aggressive lymphoma in nodular lymphocyte-predominant Hodgkin's lymphoma. *J Clin Oncol* 2010;28:793-799.
- ¹⁹ Shaikh F, Ngan B, Alexander S, Grant R. Progressive transformation of germinal centers in children and adolescents: an intriguing cause of lymphadenopathy. *Pediatr Blood Cancer* 2013;60:26-30.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF IMAGING¹⁻⁸

General Considerations

- The final report for any FDG-PET/CT examination to define response should include the Deauville 5-point scale score, which is a visual score.
- A second opinion of the FDG-PET/CT studies may be helpful if the report differs from the clinical presentation, especially if interpretation is by nuclear medicine and/or a radiologist with less experience in lymphoma imaging and/or if no Deauville scores are reported.
- A comprehensive imaging atlas that aids in the identification of affected nodal or extranodal regions and organs is available.⁹

Staging or Initial Workup (should be performed within 2 to 4 weeks prior to initiation of therapy)

- CT neck/chest/abdomen/pelvis with contrast (IV ± oral) or CT chest and MRI neck/abdomen/pelvis
- Chest x-ray PA and lateral views (if cross-sectional imaging not available or necessitated to determine bulk of disease for a clinical trial)
- FDG-PET/CT^{a,b,c} or FDG-PET/MRI^d
 - ▶ Whole-body is recommended.
 - ▶ Diagnostic-quality CT or MRI is still needed for initial staging; when available FDG-PET/CT and diagnostic CT should be performed as a combined examination to limit radiation exposure.
- MRI may be helpful in distinguishing E-lesions and bone marrow, liver, or spleen involvement in equivocal cases.

Interim and End of Therapy

- FDG-PET/CT^{a,b,c} or FDG-PET/MRI^d
 - ▶ Wait at least 8 to 12 weeks after end of RT to perform FDG-PET to minimize false-positive results.
- Special attention should be given to end-of-therapy scans when utilizing CPIs given their potential for delayed response or pseudo-progression.¹⁰ Repeat short-interval FDG-PET imaging may be necessary to differentiate between progression and pseudoprogression. Biopsy may be warranted for persistent PET avidity on end-of-therapy imaging.
- Diagnostic-quality CT with contrast or MRI should only be performed for original sites of disease if the treatment approach includes need for assessment of anatomic sites of disease for evaluation of response.

Follow-up/Surveillance

- Imaging should only be obtained if significant clinical concern for relapse or as mandated if enrolled in an active protocol.
 - ▶ If imaging is necessary, it may include diagnostic-quality chest x-ray PA and lateral views, and CT or MRI at 3- to 6-month intervals for up to 2 years.
- FDG-PET/CT or FDG-PET/MRI is not advised due to risk of false positives.
 - ▶ Repeat FDG-PET may be considered for persistent positive disease or equivocal finding on post-therapy FDG-PET.^{a,b,c,d}

Relapsed or Refractory (confirmed or highly suspected)

- CT neck/chest/abdomen/pelvis with contrast (IV ± oral) or CT chest and MR neck/abdomen/pelvis
- FDG-PET/CT^{a,b,c} or FDG-PET/MRI^d

^a FDG-PET should be read by an experienced nuclear diagnostic radiologist experienced in reading Deauville scores for FDG-PET–adapted therapy. FDG-PET/CT should be obtained in accordance with American College of Radiology (ACR) practice guidelines.

^b In cases of FDG-PET positivity where sites of disease are inconsistent with usual presentation of HL or if there is an unusual disease presentation (ie, HIV), additional clinical evaluation may be required for staging. See [Principles of Staging \(PHL-D\)](#). If FDG-PET negative at anatomic lesion of concern, biopsy should be considered. In most instances, if the FDG-PET/CT displays a homogeneous pattern of marrow uptake (thought to be secondary to cytokine release) bone marrow involvement is not assumed. If there are multifocal (≥3) skeletal FDG-PET lesions without cortical destruction on CT, marrow involvement may be assumed.

^c Measures to reduce brown fat activation, such as warming or pharmacologic suppression, may be considered to minimize false-positive findings.

^d If FDG-PET/MRI is obtained, diagnostic CT of chest is needed to evaluate lung parenchyma.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF IMAGING¹⁻⁸

Interpretation

- The Panel supports the American College of Radiology (ACR)¹¹ and Society of Nuclear Medicine and Molecular Imaging (SNMMI)¹² recommendation for FDG-PET/CT interpretation, including the requirement that FDG-PET/CT examinations should be performed under the supervision of and interpreted by a physician with the following qualifications:
 - ▶ Board certification in radiology or diagnostic radiology, nuclear radiology, or nuclear medicine
OR
 - ▶ Completion of a formal Accreditation Council for Graduate Medical Education (ACGME)-approved general nuclear medicine program in addition to 1000 hours of clinical training in general nuclear medicine, 20 hours of continuing medical education (CME) in FDG-PET, and at least 150 oncologic FDG-PET/CT examinations interpreted or multi-read during the previous 3 years.¹¹
- Continuing experience/education should include interpretation of a minimum of 150 FDG-PET/CT scans in 3 years (multi-read is acceptable) and completion of 150 hours (including 75 hours of Category 1 CME) during the preceding 3 years pertinent to the physician's practice patterns, including FDG-PET imaging.¹¹
- The interpreting radiology or nuclear medicine physician should have adequate training and CME/experience in interpreting FDG-PET/CT for patients with lymphoma including the use of the Deauville 5-point scoring system.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

References

PHL-C
2 OF 3



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF IMAGING REFERENCES

- ¹ Cerci JJ, Etchebehere EC, Nadel H, et al. Is true whole-body 18F-FDG-PET/CT required in pediatric lymphoma? An IAEA multicentre prospective study. J Nucl Med 2019;60:1087-1093.
- ² Kluge R, Kurch L, Georgi T, Metzger M. Current role of FDG-PET in pediatric Hodgkin's lymphoma. Semin Nucl Med 2017;47:242-257.
- ³ Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. J Clin Oncol 2014;32:3059-3068.
- ⁴ McCarten KM, Nadel HR, Shulkin BL, Cho SY. Imaging for diagnosis, staging, and response assessment of Hodgkin lymphoma and non-Hodgkin lymphoma. Pediatr Radiol 2019;49:1545-1564.
- ⁵ Voss SD, Cairo MS. Surveillance imaging in pediatric lymphoma. Pediatr Radiol 2019;49:1565-1573.
- ⁶ Verhagen MV, Menezes LJ, Neriman D, et al. 18F-FDG PET/MRI for staging and interim response assessment in pediatric and adolescent Hodgkin lymphoma: a prospective study with 18F-FDG PET/CT as the reference standard. J Nucl Med 2021;62:1524-1530.
- ⁷ Rathore N, Eissa HM, Margolin JF, et al. Pediatric Hodgkin lymphoma: Are we over-scanning our patients? Pediatr Hematol Oncol 2012;29:415-423.
- ⁸ Spijkers S, Littooij AS, Humphries PD, et al. Imaging features of extranodal involvement in paediatric Hodgkin lymphoma. Pediatr Radiol 2019;49:266-276.
- ⁹ Stoevesandt D, Steglich J, Bartelt JM, et al. CT, MRI, and FDG PET/CT in the assessment of lymph node involvement in pediatric Hodgkin lymphoma: An expert consensus definition by an international collaboration on Staging Evaluation and Response Criteria Harmonization for Children, Adolescent, and Young Adult Hodgkin Lymphoma (SEARCH for CAYAH). Radiology 2025;314:e232650.
- ¹⁰ Cheson BD, Ansell S, Schwartz L, et al. Refinement of the Lugano Classification lymphoma response criteria in the era of immunomodulatory therapy. Blood 2016;128:2489-2496.
- ¹¹ American College of Radiology. ACR-ACNM-SNMMI-SPR Practice Parameters for Performing FDG-PET/CT in Oncology. 2021. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/FDG-PET-CT.pdf?la=en>. Accessed September 14, 2021.
- ¹² Boellaard R, Delgado-Bolton R, Oyen WJG, et al. FDG PET/CT: EANM procedure guidelines for tumour imaging: version 2.0. Eur J Nucl Med Mol Imaging 2015;42:328-354.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF STAGING^a

- These are only guiding principles of initial staging adapted from criteria of various protocols. This table is not intended to replace protocol-specific staging. Refer to applicable study protocol for complete staging details.
- While these principles are based on Panel consensus, this remains an area of ongoing research.
- The Lugano criteria are not included in this table, because there is inadequate harmonization between adult and pediatric staging criteria. For example, size cut-offs in adults are larger than they are for children.

Site Involvement	Imaging Modality ^{b,c,d}	Protocols
Lymph nodes	Diagnostic CT/MRI and FDG-PET/CT	• Long axis ≥ 2 cm is considered involved on diagnostic CT/MRI¹ • Long axis 1–2 cm, if FDG-PET positive
Splenic	Ultrasound	• Any lesion large enough to characterize unless imaging characteristics indicate an alternative etiology irrespective of the FDG-PET result
	FDG-PET/CT or FDG-PET/MRI	• Focal FDG-PET–positive lesions that are confirmed by CT with IV contrast or MRI or ultrasound • Splenic involvement must include focal imaging abnormality. Splenomegaly and diffuse uptake greater than liver alone are not considered involvement.
Lung ^e	FDG-PET/CT	• E-lesions^f: Extralymphatic structures (lung lesions) contiguous with nodal masses are considered to be E-lesions. • At least 1–2 small foci (between 5–10 mm) within whole lung if no other etiology is suspected • At least 1 intrapulmonary focus >1 cm on CT if no other etiology is suspected • FDG-PET–positive lesions <1 cm if no other etiology is suspected Note: If all lesions are exclusively in 1 lung, then only this particular lung is considered as involved. However, even if there is just one additional smaller focus found within the other lung, then both lungs are considered involved.
Liver	Ultrasound	• Any focal mass lesion or lesions on diagnostic imaging large enough to characterize in a visceral organ are considered lymphomatous involvement unless the imaging characteristics indicate an alternative etiology.
	FDG-PET/CT or FDG-PET/MRI	• Liver involvement is defined as any hepatic lesion on CT or MRI that correlates with 18F-FDG uptake greater than background liver.²
Bone marrow	Bilateral biopsy	• Positive by histopathology on previous high-risk trials; current trial recommendations are based on FDG-PET alone • European-based GPOH-HD-2002 staging: Not recommended
	FDG-PET/CT or FDG-PET/MRI	• ≥ 3 FDG-PET–positive lesions in bone marrow without cortical bone destruction
Bone	FDG-PET/CT or FDG-PET/MRI	• FDG-PET–positive lesion with cortical bone destruction on CT or MRI³ • If no cortical bone destruction, a bone scan may be helpful. • Extralymphatic structures (bone lesions) contiguous with nodal masses are considered to be E-lesions.^f
Waldeyer's Ring	Diagnostic CT/MRI and FDG-PET/CT	Must meet all of the following criteria:^{4,5} • Abnormal asymmetric findings on both FDG-PET/CT and CT/MRI • No other explanation on imaging (eg, adjacent nodal group or mass, infection) • Interim response equal to adjacent sites of disease

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Footnotes and References



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF STAGING^a

ASSESSMENT OF BULK DISEASE

Site Involvement	U.S.-Based Protocols	European-Based Protocols ⁹
Peripheral nodes	Contiguous extramediastinal nodal aggregate >6 cm in the longest diameter (LDi) measured in axial, coronal, or sagittal dimension (including oblique measurement)	Volume of the largest contiguous lymph node mass ≥200 mL
Mediastinal mass	Tumor diameter >1/3 of the maximal thoracic diameter of an upright PA chest radiograph	Tumor volume ≥200 mL

PET 5-POINT SCALE (DEAUVILLE CRITERIA)^b

Score	PET/CT Scan Result
1	No uptake
2	Uptake ≤ mediastinum
3	Uptake > mediastinum but ≤ liver
4	Uptake moderately higher than liver
5	Uptake markedly higher than liver and/or new lesions
X	New areas of uptake unlikely to be related to lymphoma

With kind permission from Springer International Publishing: Barrington SF, Mikhaeel NG, Kostakoglu L, et al. Role of imaging in the staging and response assessment of lymphoma: consensus of the International Conference on Malignant Lymphomas Imaging Working Group. J Clin Oncol 2014;32:3048-3058.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

Footnotes and References



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF STAGING FOOTNOTES

- ^a Clinical interpretation of staging at diagnosis should not be based on reports alone. Treating clinician notes should summarize interpretation of sites of involvement prior to initiation of treatment.
- ^b FDG-PET should be read by an experienced nuclear diagnostic radiologist experienced in reading Deauville scores for FDG-PET–adapted therapy. This is a visual analysis and does not include standardized uptake value (SUV).
- ^c There may be FDG-PET–avid lesions that need clinical correlation to determine if it is related to lymphoma.
- ^d [Principles of Imaging \(PHL-C\)](#).
- ^e There are inconsistencies in staging between protocols and providers. Careful attention to staging of lung involvement is important as it may change the risk group of the patient.
- ^f An E-lesion is defined as a contiguous infiltration of a lymph node mass into extralymphatic structures or organs (eg, lung or bone). Pleural and pericardial involvement should be considered E-lesions, but a pleural or pericardial effusion alone is not considered an E-lesion. Disease that extends beyond the lymphatic system without adjacent lymphatic involvement is considered stage IV; liver or bone marrow involvement is always considered stage IV disease (Zijtregtop EAM, et al. Blood Adv 2023;7:6303-6319). CNS disease is considered extra-axial.
- ^g Volume = $(a \times b \times c)/2$ where a, b, and c are three dimensions of a node or conglomerate (in cm). Mauz-Körholz C, et al. Lancet Oncol 2022;23:125-137.

REFERENCES

- Stoevesandt D, Steglich J, Bartelt JM, et al. CT, MRI, and FDG PET/CT in the assessment of lymph node involvement in pediatric Hodgkin lymphoma: An expert consensus definition by an international collaboration on staging evaluation and response criteria harmonization for children, adolescent, and young adult Hodgkin Lymphoma (SEARCH for CAYAHL). Radiology 2025;314:e232650.
- Hagleitner MM, Metzger ML, Flerlage JE, et al. Liver involvement in pediatric Hodgkin lymphoma: A systematic review by an international collaboration on Staging Evaluation and Response Criteria Harmonization (SEARCH) for Children, Adolescent, and Young Adult Hodgkin Lymphoma (CAYAHL). Pediatr Blood Cancer 2020;67:e28365.
- Lewis J, McCarten K, Kurch L, et al. Definition of cortical bone involvement in the staging of newly diagnosed pediatric Hodgkin lymphoma: A report from the International Working Group on Staging Evaluation and Response Criteria Harmonization (SEARCH). Pediatr Blood Cancer 2020;67:e28142.
- Kurch L, Mauz-Körholz C, Fossa A, et al. Assessment of Waldeyer's ring in pediatric and adolescent Hodgkin lymphoma patients-Importance of multimodality imaging: Results from the EuroNet-PHL-C1 trial. Pediatr Blood Cancer 2021;68:e28903.
- Seelisch J, De Alarcon PA, Flerlage JE, et al. Expert consensus statements for Waldeyer's ring involvement in pediatric Hodgkin lymphoma: The staging, evaluation, and response criteria harmonization (SEARCH) for childhood, adolescent, and young adult Hodgkin lymphoma (CAYAHL) group. Pediatr Blood Cancer 2020;67:e28361.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY^{a,b} Primary Systemic Therapy

Primary Systemic Therapy - Recommended Dosing

AVPC¹

Note: Cyclophosphamide and doxorubicin dosing in AHOD0431 differs from AHOD03P1.

• CHL (AHOD0431)¹

- ▶ Doxorubicin^c 25 mg/m² IV Days 1 and 2
- ▶ Vincristine 1.4 mg/m² IV Days 1 and 8; 2.8 mg/dose maximum
- ▶ Prednisone 20 mg/m² PO twice daily on Days 1–7
- ▶ Cyclophosphamide 600 mg/m² IV Days 1 and 2
- ▶ Regimen repeated every 21 days for 3 cycles

• NLPHL (AHOD03P1)²

- ▶ Doxorubicin^c 50 mg/m² IV Day 1
- ▶ Vincristine 1.4 mg/m² IV Days 1 and 8; 2.8 mg/dose maximum
- ▶ Prednisone 20 mg/m² PO twice daily on Days 1–7
- ▶ Cyclophosphamide 800 mg/m² IV Day 1
- ▶ Regimen repeated every 21 days for 3 cycles

Bv-AVE-PC (AHOD1331)^{d,3}

- Brentuximab vedotin 1.8 mg/kg IV Day 1 (prior to other chemotherapy); 180 mg/dose maximum per dose
- Doxorubicin^c 25 mg/m² IV Days 1 and 2
- Vincristine 1.4 mg/m² IV Day 8; 2.8 mg/dose maximum per dose
- Etoposide 125 mg/m² IV daily on Days 1–3
- Prednisone 40 mg/m² PO divided into two doses daily on Days 1–7
- Cyclophosphamide 600 mg/m² IV Days 1 and 2
- Regimen repeated every 21 days for 5 cycles

ABVE-PC (AHOD0031)⁴

- Doxorubicin^c 25 mg/m² IV Days 1 and 2
- Bleomycin 5 U/m² IV Day 1, 10 U/m² IV Day 8
- Vincristine 1.4 mg/m² IV Days 1 and 8; 2.8 mg/dose maximum per dose
- Etoposide 125 mg/m² IV daily on Days 1–3
- Prednisone 40 mg/m² PO divided into two doses daily on Days 1–7
- Cyclophosphamide 800 mg/m² IV Day 1
- Regimen repeated every 21 days for 4 cycles

Bv-AVD (S1826)^{d,5}

- Brentuximab vedotin 1.2 mg/kg (up to 100 mg max) IV Days 1 and 15
- Doxorubicin^c 25 mg/m² IV Days 1 and 15
- Vinblastine 6 mg/m² IV Days 1 and 15
- Dacarbazine 375 mg/m² IV Days 1 and 15
- Regimen repeated every 28 days for up to 6 cycles

Nivolumab-AVD (S1826)^{e,5}

- Nivolumab 240 mg in patients ≥18 years or 3 mg/kg for patients <18 years (up to 240 mg max) IV Days 1 and 15
- Doxorubicin^c 25 mg/m² IV Days 1 and 15
- Vinblastine 6 mg/m² IV Days 1 and 15
- Dacarbazine 375 mg/m² IV Days 1 and 15
- Regimen repeated every 28 days for 6 cycles

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY^{a,b} Primary Systemic Therapy

Primary Systemic Therapy - Recommended Dosing

OEPA (GPOH-HD-2002)⁶

- Vincristine 1.5 mg/m² IV Days 1, 8, and 15;
2 mg/dose maximum
- Etoposide 125 mg/m² IV daily on Days 1–5
- Prednisone 60 mg/m² PO daily on Days 1–15
- Doxorubicin^c 40 mg/m² IV Days 1 and 15
- Regimen repeated every 28 days for 2 cycles

OEPA-COPDAC (GPOH-HD-2002; EuroNET-PHL-C1)^{6,7}

- **OEPA:**
 - ▶ Vincristine 1.5 mg/m² IV Days 1, 8, and 15;
2 mg/dose maximum
 - ▶ Etoposide 125 mg/m² IV daily on Days 1–5
 - ▶ Prednisone 60 mg/m² PO daily on Days 1–15
 - ▶ Doxorubicin^c 40 mg/m² IV on Days 1 and 15
 - ▶ Regimen repeated every 28 days for 2 cycles
- **COPDAC:**
 - ▶ Cyclophosphamide 500 mg/m² IV Days 1 and 8
 - ▶ Vincristine 1.5 mg/m² IV Days 1 and 8; 2 mg/dose maximum
 - ▶ Prednisone 40 mg/m² PO daily on Days 1–15; 80 mg maximum per day
 - ▶ Dacarbazine 250 mg/m² IV daily on Days 1–3
 - ▶ Regimen repeated every 28 days for 2 cycles for stages IAE, IBE, IIAE, IIB, or IIIA, or 4 cycles for stages IIBE, IIIAE, IIIBE, IIIB, IVA, or IVB

AEPA-CAPDAC (HLHR13)^{d,8}

- **AEPA**
 - ▶ Brentuximab vedotin 1.2 mg/kg IV Days 1, 8, and 15; 120 mg/dose maximum per dose
 - ▶ Etoposide 125 mg/m² IV on Days 1–5
 - ▶ Prednisone 60 mg/m²/day PO divided TID on Days 1–15; 30 mg/dose TID maximum
 - ▶ Doxorubicin^c 40 mg/m² IV Days 1 and 15
 - ◊ Regimen repeated every 28 days for 2 cycles
- **CAPDAC**
 - ▶ Cyclophosphamide 500 mg/m² IV Days 1 and 8
 - ▶ Brentuximab vedotin 1.2 mg/kg IV Days 1 and 8; 120 mg/dose maximum per dose
 - ▶ Prednisone 40 mg/m²/day PO divided TID on Days 1–15; 20 mg/dose TID maximum
 - ▶ Dacarbazine 250 mg/m² IV on Days 1–3
 - ◊ Regimen repeated every 21 days for 4 cycles

CVbP ± Rituximab^{f,9,10}

- Cyclophosphamide 500 mg/m² IV on Day 1
- Vinblastine 6 mg/m² IV Days 1 and 8
- Prednisolone 40 mg/m² PO on Days 1–8
 - ▶ Regimen repeated at 2–3 week intervals for 3 cycles
- Rituximab 375 mg/m² IV Day 1

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY^{a,b} Treatment for Relapsed or Refractory Disease

- Consider the following when selecting re-induction or subsequent therapy:
 - ▶ Referral to a center with expertise
 - ▶ Clinical trial enrollment
 - ▶ Primary therapy⁹ and prior RT exposure
 - ▶ Cumulative short- and long-term toxicity
 - ▶ Opportunity to harvest stem cells
 - ▶ Counseling on infertility risk (for AYA, [NCCN Guidelines for AYA Oncology](#))
 - ▶ Psychosocial assessment (for AYA, see [NCCN Guidelines for AYA Oncology](#))
- Consider use of RT, in consultation with a center with expertise, as part of therapy for relapsed/refractory disease.^h
- Recommend inclusion of a CPI, particularly if not included as part of front-line therapy.

Relapsed/Refractory Disease			
	Re-Induction and Subsequent Therapy Options With Pediatric Data ^{i,j}	Useful in Certain Circumstances Options with or without Pediatric Data	Maintenance (post-transplant)
CHL	CPI-Based Regimens (Preferred):^e • Brentuximab vedotin (Bv) + nivolumab ± bendamustine (AHOD1721)^{d,k,11,12,13} <u>Other Recommended: Non-CPI-Containing Regimens (in alphabetical order)</u> • Bortezomib, ifosfamide, + vinorelbine¹⁴ • Bv + gemcitabine^{d,k,15} • EPIC (etoposide, prednisolone, ifosfamide, cisplatin)¹⁶ • GDP (gemcitabine, dexamethasone, cisplatin)¹⁷ (category 2B) • GV (gemcitabine, vinorelbine)^{k,18,19,20}	CPI-Based Regimens (in alphabetical order)^e • GVD + pembrolizumab (gemcitabine, vinorelbine, doxorubicin liposome)^{l,21} • ICE + nivolumab^{l,22} • ICE + pembrolizumab^{l,23} • Nivolumab^{k,l,24,25} • Pembrolizumab^{k,m,26,27} Non-CPI-Containing Regimens (in alphabetical order) • Brentuximab vedotin + bendamustine^{d,k,28} • DHAP (dexamethasone, cytarabine, cisplatin) (category 2B) • ICE (ifosfamide, carboplatin, etoposide)²⁹ Additional options may be considered for patients >18 years; see NCCN Guidelines for Hodgkin Lymphoma (Adult). Consider referral for allogeneic HCT in appropriate patients.	<u>Useful in Certain Circumstances</u> • Bv (for select patients with high-risk disease^{d,n,o})^{30,31}
NLPHL	Refer to a center of expertise. See NCCN Guidelines for Hodgkin Lymphoma (Adult) .		

[See Evidence Block on EB-5](#)

[See Evidence Blocks on EB-6](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Footnotes](#)
[References](#)

PHL-E
3 OF 6



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY FOOTNOTES

- ^a An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.
- ^b For growth factor recommendations, see [Select Principles of Supportive Care \(PHL-G\)](#).
- ^c Strongly consider administering dexrazoxane to children and adolescents expected to receive ≥ 250 mg/m² of cumulative doxorubicin, or an equivalent amount of an alternative anthracycline (Chow EJ, et al. J Clin Oncol 2023;41:2248-2257; de Baat EC, et al. Lancet Child Adolesc Health 2022;6:885-894).
- ^d PJP prophylaxis is recommended for all Bv-containing regimens.
- ^e CPIs should be used with caution in those with active or previously documented autoimmune disease and/or current use of immunosuppressive agents.
- ^f Data are limited on the use of rituximab for early-stage NLPHL.
- ^g Consider cross-toxicity with prior lines of therapy (including but not limited to cumulative dose of anthracycline) for both acute and late toxicities.
- ^h Considerations for RT use should include site of relapse, history of prior RT, and eligibility for HDT/ASCT. RT alone may also be considered as consolidation following second-line therapy. Mailhot Vega RB, et al. Int J Radiat Oncol Biol Phys 2023;117:400-403.
- ⁱ It is reasonable to try multiple different re-induction regimens as needed prior to ASCR to minimize disease burden with a goal of achieving a metabolic CR prior to transplant.
- ^j Subsequent therapy options include re-induction options that were not previously used.
- ^k Should be considered in patients with heavily pretreated (with platinum or anthracycline-based chemotherapy) disease or if a decrease in cardiac function is observed.
- ^l There are no pediatric data for this regimen.
- ^m Pembrolizumab is indicated for the treatment of pediatric patients with refractory CHL, or who have relapsed after two or more prior lines of therapy.
- ⁿ High-risk disease: progressive disease, refractory disease, or relapse within 1 year of original diagnosis (Moskowitz CH, et al. Blood 2018;132:2639-2642).
- ^o The role of maintenance Bv has not been well-defined in patients who received Bv or a CPI prior to maintenance therapy.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY REFERENCES

- 1 Keller FG, Castellino SM, Chen L, et al. Results of the AHOD0431 trial of response adapted therapy and a salvage strategy for limited stage, classical Hodgkin lymphoma: a report from the Children's Oncology Group. *Cancer* 2018;124:3210-3219.
- 2 Appel BE, Chen L, Buxton AB, et al. Minimal treatment of low-risk, pediatric lymphocyte-predominant hodgkin lymphoma: a report from the Children's Oncology Group. *J Clin Oncol* 2016;34:2372-2379.
- 3 Castellino SM, Pei Q, Parsons SK, et al. Brentuximab vedotin with chemotherapy in pediatric high-risk Hodgkin's lymphoma. *N Engl J Med* 2022;387:1649-1660.
- 4 Friedman DL, Chen L, Wolden S, et al. Dose-intensive response-based chemotherapy and radiation therapy for children and adolescents with newly diagnosed intermediate-risk hodgkin lymphoma: a report from the Children's Oncology Group Study AHOD0031. *J Clin Oncol* 2014;32:3651-3658.
- 5 Herrera AF, LeBlanc M, Castellino SM, et al. Nivolumab+AVD in advanced-stage classic Hodgkin's lymphoma. *N Engl J Med* 2024;391:1379-1389.
- 6 Mauz-Korholz C, Hasenclever D, Dorffel W, et al. Procarbazine-free OEPA-COPDAC chemotherapy in boys and standard OPBA-COPP in girls have comparable effectiveness in pediatric Hodgkin's lymphoma: the GPOH-HD-2002 study. *J Clin Oncol* 2010;28:3680-3686.
- 7 Mauz-Korholz C, Landman-Parker J, Balwierz W, et al. Response-adapted omission of radiotherapy and comparison of consolidation chemotherapy in children and adolescents with intermediate-stage and advanced-stage classical Hodgkin lymphoma (EuroNet-PHL-C1): a titration study with an open-label, embedded, multinational, non-inferiority, randomised controlled trial. *Lancet Oncol* 2022;23:125-137.
- 8 Metzger ML, Link MP, Billett AL, et al. Excellent outcome for pediatric patients with high-risk Hodgkin lymphoma treated with brentuximab vedotin and risk-adapted residual node radiation. *J Clin Oncol* 2021;39:2276-2283.
- 9 Shankar A, Hall GW, Gorde-Grosjean S, et al. Treatment outcome after low intensity chemotherapy [CVP] in children and adolescents with early stage nodular lymphocyte predominant Hodgkin's lymphoma - an Anglo-French collaborative report. *Eur J Cancer* 2012;48:1700-1706.
- 10 Shankar A, Hall GW, McKay P, et al. Management of children and adults with all stages of nodular lymphocyte predominant Hodgkin lymphoma - All STAGES: A consensus-based position paper from the Hodgkin lymphoma subgroup of the UK National Cancer Research Institute. *Br J Haematol* 2022;197:679-690.
- 11 Cole PD, Mauz-Körholz C, Mascarini M, et al. Nivolumab and brentuximab vedotin (BV)-based, response-adapted treatment in children, adolescents, and young adults (CAYA) with standard-risk relapsed/refractory classical Hodgkin lymphoma (R/R cHL): Primary analysis [abstract]. *J Clin Oncol* 2020;38(Suppl): Abstract 8013.
- 12 Harker-Murray P, Mauz-Korholz C, Leblanc T, et al. Nivolumab and brentuximab vedotin with or without bendamustine for R/R Hodgkin lymphoma in children, adolescents, and young adults. *Blood* 2023;141:2075-2084.
- 13 Greve P, Beishuizen A, Hagleitner M, et al. Nivolumab plus brentuximab vedotin +/- bendamustine combination therapy: A safe and effective treatment in pediatric recurrent and refractory classical Hodgkin lymphoma. *Front Immunol* 2023;14:1229558.
- 14 Horton TM, Drachtman R, Chen L, et al. A phase 2 study of bortezomib in combination with ifosfamide/vinorelbine in paediatric patients and young adults with refractory/recurrent Hodgkin lymphoma: a Children's Oncology Group study. *Br J Haematol* 2015;170:118-122.
- 15 Cole PD, McCarten KM, Pei Q, et al. Brentuximab vedotin with gemcitabine for paediatric and young adult patients with relapsed or refractory Hodgkin's lymphoma (AHOD1221): a Children's Oncology Group, multicentre single-arm, phase 1-2 trial. *Lancet Oncol* 2018;19:1229-1238.
- 16 Shankar A, Hayward J, Kirkwood A, et al. Treatment outcome in children and adolescents with relapsed Hodgkin lymphoma--results of the UK HD3 relapse treatment strategy. *Br J Haematol* 2014;165:534-544.
- 17 Baetz T, Belch A, Couban S, et al. Gemcitabine, dexamethasone and cisplatin is an active and non-toxic chemotherapy regimen in relapsed or refractory Hodgkin's disease: a phase II study by the National Cancer Institute of Canada Clinical Trials Group. *Ann Oncol* 2003;14:1762-1767.
- 18 Jan MRU, Saeed H, Abubakar M, Wali RM. Response rates, long term outcomes and toxicity profile of gemcitabine and vinorelbine based outpatient chemotherapy regimen in primary progressive and relapsed childhood Hodgkin lymphoma. *Leuk Lymphoma* 2020;61:3369-3377.
- 19 Cole PD, Schwartz CL, Drachtman RA, et al. Phase II study of weekly gemcitabine and vinorelbine for children with recurrent or refractory Hodgkin's disease: A children's oncology group report. *J Clin Oncol* 2009;27:1456-1461.
- 20 Mahdy A, Hamoda A, Zaher A, et al. Outcome and toxicity of ifosfamide, carboplatin, and etoposide versus gemcitabine and vinorelbine regimen for pediatric patients with relapsed or refractory Hodgkin's lymphoma. *Front Oncol* 2023;13:1153128.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Continued](#)

PHL-E
5 OF 6



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY REFERENCES

- 21 Moskowitz AJ, Shah G, Schoder H, et al. Phase II trial of pembrolizumab plus gemcitabine, vinorelbine, and liposomal doxorubicin as second-line therapy for relapsed or refractory classical Hodgkin lymphoma. *J Clin Oncol* 2021;39:3109-3117.
- 22 Mei MG, Lee HJ, Palmer JM, et al. Response-adapted anti-PD-1-based salvage therapy for Hodgkin lymphoma with nivolumab alone or in combination with ICE. *Blood* 2022;139:3605-3616.
- 23 Bryan LJ, Casulo C, Allen PB, et al. Pembrolizumab added to ifosfamide, carboplatin, and etoposide chemotherapy for relapsed or refractory classic Hodgkin lymphoma: A multi-institutional phase 2 investigator-initiated nonrandomized clinical trial. *JAMA Oncol* 2023;9:683-691.
- 24 Younes A, Santoro A, Shipp M, et al. Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort single-arm phase 2 trial. *Lancet Oncol* 2016;17:1283-1294.
- 25 Armand P, Engert A, Younes A, et al. Nivolumab for relapsed/refractory classic Hodgkin lymphoma after failure of autologous hematopoietic cell transplantation: Extended follow-up of the multicohort single-arm phase II CheckMate 205 trial. *J Clin Oncol* 2018;36:1428-1439.
- 26 Chen R, Zinzani PL, Fanale MA, et al. Phase II study of the efficacy and safety of pembrolizumab for relapsed/refractory classic Hodgkin lymphoma. *J Clin Oncol* 2017;35:2125-2132.
- 27 Georger B, Kang HJ, Yalon-Oren M, et al. Pembrolizumab in paediatric patients with advanced melanoma or a PD-L1-positive, advanced, relapsed, or refractory solid tumour or lymphoma (KEYNOTE-051): interim analysis of an open-label, single-arm, phase 1–2 trial. *Lancet Oncol* 2020;21:121-133.
- 28 O'Connor OA, Lue JK, Sawas A, et al. Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin's lymphoma: an international, multicenter, single-arm, phase 1-2 trial. *Lancet Oncol* 2018;19:257-266.
- 29 Moskowitz CH, Nimer SD, Zelenetz AD, et al. A 2-step comprehensive high-dose chemoradiotherapy second-line program for relapsed and refractory Hodgkin disease: analysis by intent to treat and development of a prognostic model. *Blood* 2001;97:616-623.
- 30 Moskowitz CH, Walewski J, Nademanee A, et al. Five-year PFS from the AETHERA trial of brentuximab vedotin for Hodgkin lymphoma at high risk of progression or relapse. *Blood* 2018;132:2639-2642.
- 31 Forlenza CJ, Rosenzweig J, Mauguén A, et al. Brentuximab vedotin after autologous transplantation in pediatric patients with relapsed/refractory Hodgkin lymphoma. *Blood Adv* 2023;7:3225-3231.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY¹⁻⁶

General Principles

- Treatment with photons, electrons, or protons may all be appropriate, depending on clinical circumstances.
- In specific instances, advanced RT technologies may be used to spare important organs at risk (OARs) and decrease the risk for late normal tissue damage while still achieving the primary goal of local tumor control.
 - ▶ Advanced technologies including intensity-modulated RT (IMRT)/volumetric modulated arc therapy (VMAT), breath hold or respiratory gating and/or image-guided RT (IGRT), or proton therapy may offer significant and clinically relevant advantages.⁷
 - ▶ OARs: heart (including coronary arteries, valves, and left ventricle), lungs, kidneys, spinal cord, esophagus, carotid arteries, bone marrow, breasts, stomach, muscle/soft tissue, and salivary glands.
- Dose-sparing for OARs reflects best clinical practice, as it reduces the risk of late complications from normal tissue damage. Achieving highly conformal dose distributions is especially important for patients who are being treated with curative intent or who have long life expectancies following therapy.
- Breath hold techniques have been shown to decrease incidental dose to the heart and lungs in many disease presentations, including mediastinal HL. Strategies include:
 - ▶ Four-dimensional CT (4D-CT) for simulation or deep inspiration breath hold (DIBH)
 - ▶ Respiratory gating
 - ▶ IGRT during treatment delivery
- Since the advantages of these techniques include tightly conformal doses and steep gradients next to normal tissues, target definition and delineation and treatment delivery verification require careful monitoring to avoid the risk of tumor geographic miss and subsequent decrease in tumor control.
 - ▶ Initial diagnostic imaging with contrast-enhanced CT, MRI, FDG-PET/CT or FDG-PET/MRI, ultrasound, and other imaging modalities facilitate target definition.
 - ▶ Image guidance may be required to provide assurance of accurate daily delivery.
- Randomized studies to test these concepts are unlikely to be done since these techniques are designed to decrease late effects, which take ≥10 years to develop. In light of that, the modalities and techniques (including proton therapy) that are found to best reduce the doses to the OARs for a given patient in a clinically meaningful way without compromising target coverage should be used.
- Historic RT approaches that resulted in high rates of second malignancies and cardiac toxicity observed among long-term survivors of HL were based on outdated RT techniques, including total nodal irradiation, subtotal nodal irradiation, and mantle field irradiation, typically delivered to prescription doses of 40–45 Gy. These strategies exposed substantial volumes of normal tissues—particularly the breast, heart, and lungs—to doses in the range of 30–40 Gy. RT for HL has since evolved to reduce the risk of late effects through the use of significantly smaller treatment volumes—such as involved-field RT (IFRT), ISRT, and RSRT—as well as lower prescription doses and more conformal delivery techniques, which have resulted in fewer late effects.^{8,9,10}

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY¹⁻⁶

Volume

- ISRT is recommended as the appropriate field for HL. If the protocol used IFRT then it should be replaced by ISRT.
- Planning for ISRT requires CT-based simulation and treatment planning capabilities. Incorporating other modern imaging such as FDG-PET/CT or FDG-PET/MRI often enhances treatment volume determination.
- ISRT targets the site of the originally involved lymph node(s). The volume encompasses the original or suspected extent of disease prior to chemotherapy or surgery.^a However, it spares adjacent uninvolved organs (eg, lungs, bone, muscle, kidney) when lymphadenopathy regresses following chemotherapy.
 - Pre-chemotherapy or pre-biopsy gross tumor volume (GTV) provides the basis for determining the clinical target volume (CTV). Concerns for questionable subclinical disease and uncertainties in original imaging accuracy or localization may lead to expansion of the CTV and are determined individually using clinical judgment.
 - Movement of the CTV by respiration as determined by 4D-CT or fluoroscopy should be used to create an internal target volume (ITV).
- The planning target volume (PTV) is an additional expansion of the CTV that accounts only for setup variations and may differ by site and immobilization technique. Daily image guidance is recommended to minimize the PTV expansion.
- Outline OARs for optimizing treatment plan decisions.
 - These should include contouring of breast tissue (conventional breast tissue and glandular breast tissue) and cardiac substructures (left ventricle and coronary vessels), especially when contemporary RT techniques are being used (IMRT, VMAT, and proton therapy).
- The treatment plan can be designed using conventional, three-dimensional conformal RT (3D-CRT), IMRT, or proton therapy techniques using clinical treatment planning considerations of coverage and normal tissue avoidance.
- The treatment of extranodal disease is individualized, but similar principles of GTV/CTV/PTV definition should be applied as for nodal disease.
- Chest wall extension: Effort should be made to include regions of initial chest wall extension to definitive doses.
- Lung involvement:
 - Areas of extension into the lung from mediastinal or hilar disease may be treated with lower doses (15 Gy) unless the relative volume is small, in which case higher doses may be used.
 - Careful consideration of partial lung tolerance is essential.
 - Pulmonary nodular disease is usually not treated following chemotherapy unless residual disease is present.
- Pleural or pericardial effusions are not included in the GTV. Nodular pericardial involvement may be included with consideration of cardiac tolerance.
- Bone: Areas of osseous disease may be treated with a CTV expansion beyond the GTV defined by imaging. In the presence of vertebral body disease, the entire vertebra is generally treated.
- If spleen is irradiated, vaccines should be given prior to or after RT (ie, pneumococcal, *Haemophilus influenzae* type b, meningococcal).

^a In HLHR13, post-chemotherapy volumes were irradiated.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY¹⁻⁶

- In general, RT fields and doses should be delivered per protocol guidelines used for systemic therapy.

RT Fields

- ISRT can safely replace IFRT or modified IFRT.
- RSRT should be used only when dictated by the protocol or as a “boost” following standard ISRT.
- RT should be given according to the protocol being followed. For patients with stage III/IV disease it is preferable to avoid a protocol that calls for IFRT/ISRT to all sites of disease and instead use a protocol that only irradiates sites that are bulky or if inadequate response.¹¹
- Some protocols, such as HLHR13, incorporate RNRT, in which individually involved nodes are assessed for response to determine need for RT to each node. If this approach is selected, particular attention to detail is required in response assessment and radiation contouring.

Stage I–II With or Without Risk Factors

- ISRT, consider
 - ▶ All sites of disease - 21 Gy
 - ▶ Sites of slow response could receive a boost of up to 9 Gy (total dose 21–30 Gy)
 - ▶ Sites of PR should receive a boost of 9–19 Gy (total dose 30–40 Gy)

Stage III–IV

- Avoid regimens that require ISRT to all sites of disease.
- ISRT, consider:
 - ▶ Bulky disease - 21 Gy
 - ▶ Slow responding sites could receive a boost of up to 9 Gy (total dose 21–30 Gy)
 - ▶ Partial responding sites should receive a boost of 9–19 Gy (total dose 30–40 Gy)

Relapsed/Refractory Disease

- If no HDT/ASCR planned: ISRT 30 Gy
- In conjunction with HDT/ASCR
 - ▶ ISRT, 30 Gy to relapsed/refractory sites, and consider 21 Gy to initial sites that are no longer present (depending on the size of the field)
 - ▶ If Deauville 4–5 after several lines of therapy consider RT to achieve metabolic CR prior to transplant. Boost to FDG-PET-positive sites, 10–15 Gy (total dose 40–45 Gy).

RT Dose Constraints

- See RT Dose Constraint Guidelines for Lymphoma in the [NCCN Guidelines for Hodgkin Lymphoma \(Adult\)](#).

Common Contouring Pitfalls¹²

- Care should be taken to ensure that the initially involved nodes are contoured correctly, recognizing that image fusions may be imperfect due to differences in position.
- Attention should be given to ensuring that all intended target volumes are contoured for a given patient.
- Normal tissues that are adjacent to site of disease but not involved do not need to be included in target volume.

[References](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY REFERENCES

- ¹ Bates J, Howell RM, Liu Q, et al. Therapy-related cardiac risk in childhood cancer survivors: An analysis of the childhood cancer survivor study. *J Clin Oncol* 2019;37:1090-1101.
- ² van Nimwegen FA, Ntetas G, Darby SC, et al. Risk of heart failure in survivors of Hodgkin lymphoma: effects of cardiac exposure to radiation and anthracyclines. *Blood* 2017;129:2257-2265.
- ³ Dietz AC, Chen Y, Yasui Y, et al. Risk and impact of pulmonary complications in survivors of childhood cancer: A report from the Childhood Cancer Survivor Study. *Cancer* 2016;122:3687-3696.
- ⁴ Cella L, Conson M, Caterino M, et al. Thyroid V30 predicts radiation-induced hypothyroidism in patients treated with sequential chemo-radiotherapy for Hodgkin's lymphoma. *Int J Radiat Oncol Biol Phys* 2012;82:1802-1808.
- ⁵ Eisbruch A, Ten Haken RK, Kim HM, et al. Dose, volume, and function relationships in parotid salivary glands following conformal and intensity-modulated irradiation of head and neck cancer. *Int J Radiat Oncol Biol Phys* 1999;45:577-587.
- ⁶ Travis LB, Hill DA, Dores GM, et al. Breast cancer following radiotherapy and chemotherapy among young women with Hodgkin disease. *JAMA* 2003;290:465-475.
- ⁷ Hoppe BS, Mailhot-Vega RB, Renfro LA, et al. Comparison of contemporary radiation therapy approaches in combined modality treatment on pediatric high-risk classic Hodgkin lymphoma study: AHOD 1331. *Int J Radiat Oncol Biol Phys* 2025;123:972-979.
- ⁸ Lagerlöf I, Fohlin H, Enblad G, et al. Limited, but not eliminated, excess long-term morbidity in stage I-IIA Hodgkin lymphoma treated with doxorubicin, bleomycin, vinblastine, and dacarbazine and limited-field radiotherapy. *J Clin Oncol* 2022;40:1487-1496.
- ⁹ Lagerlöf I, Holte H, Glimelius I, et al. No excess long-term mortality in stage I-IIA Hodgkin lymphoma patients treated with ABVD and limited field radiotherapy. *Br J Haematol* 2020;188:685-691.
- ¹⁰ Xiang M, Chang DT, Pollom EL. Second cancer risk after primary cancer treatment with three-dimensional conformal, intensity-modulated, or proton beam radiation therapy. *Cancer* 2020;126:3560-3568.
- ¹¹ Hall MD, Terezakis SA, Lucas JT, et al. Radiation therapy across Pediatric Hodgkin Lymphoma Research Group Protocols: A report from the Staging, Evaluation, and Response Criteria Harmonization (SEARCH) for Childhood, Adolescent, and Young Adult Hodgkin Lymphoma (CAYAHL) Group. *Int J Radiat Oncol Biol Phys* 2022;112:317-334.
- ¹² Lo AC, Milgrom SA, Vega RM, et al. Enhancing radiation therapy quality assurance in lymphoma: A rigorous real-time central review process in AHOD2131 [published online ahead of print December 24, 2025]. *Int J Radiat Oncol Biol Phys*, doi: 10.1016/j.ijrobp.2025.12.011.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

SELECT PRINCIPLES OF SUPPORTIVE CARE

Immunotherapy

• Prior to initiating therapy:

- ▶ Patients and caregivers should receive counseling on the warning signs and symptoms of immune-related adverse events (irAEs).
- ▶ A thorough clinical assessment is also recommended, including physical examination alongside patient and relevant family history of any of the following:

- ◊ Autoimmune/organ-specific disease
- ◊ Endocrinopathy
- ◊ Infectious disease

▶ Blood work:

- ◊ CBC, CMP
- ◊ TSH, free thyroxine (FT4)
- ◊ Consider serum cortisol (morning preferred)

• During therapy:

▶ Blood work:

- ◊ CBC, CMP prior to each treatment or every 4 weeks
- ◊ TSH, T4 every 4–6 weeks
- ◊ Consider serum cortisol every 4–6 weeks

- ▶ A clinical exam with irAE symptom assessment should be performed at each visit, assessing for any new signs or symptoms, including severe fatigue, headache, rash, cough, shortness of breath, chest pain, abdominal bloating, change in bowel pattern, weight loss, vision changes or eye pain, severe muscle weakness, severe muscle or joint pains, and/or mood changes.

- For other pre-assessment, monitoring, and irAE management recommendations, see [NCCN Guidelines for Immune Checkpoint Inhibitor-Related Toxicities](#).

Peripheral Neuropathy

- For patients receiving neurotoxic chemotherapy (eg, Bv, vincristine, vinblastine), consider the following grading and/or screening methods^{1,2,3}:

- ▶ Common Terminology Criteria for Adverse Events (CTCAE) v5.0 or Balis score
- ▶ Patient-reported outcome measures
- ▶ Pediatric-modified Total Neuropathy Score (peds-mTNS) and/or Total Neuropathy Score-Pediatric Version (TNS-PV)

- For management of chemotherapy-induced peripheral neuropathy, consider the following:

- ▶ Dose reduction in or hold of neurotoxic chemotherapy based on protocol recommendations^{4,5}
- ▶ Early referral to physical therapy
- ▶ To date, no pharmacologic intervention has shown benefit in the pediatric and adolescent cancer population. Consideration of duloxetine is recommended for patients ≥18 years of age (see [NCCN Guidelines for Adult Cancer Pain](#)).

Growth Factors (filgrastim or pegfilgrastim)^a

- Routine use of granulocyte colony-stimulating factor (G-CSF) is not recommended with ABVD.^{6,7}
- G-CSF support is recommended with use of Bv-AVD.⁸
- G-CSF support is optional with nivolumab-AVD.⁹
- For additional information, see [NCCN Guidelines for Hematopoietic Growth Factors](#).

^a An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Continued](#)

[References](#)

PHL-G
1 OF 3



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

SELECT PRINCIPLES OF SUPPORTIVE CARE

Management of Brentuximab Vedotin-Related Infusion Reactions

- In the absence of corticosteroid therapy, anti-allergy premedications, such as an H1-antagonist (eg, diphenhydramine, cetirizine), an H2-antagonist (eg, famotidine), and acetaminophen are recommended at least 30 minutes prior to all cycles of Bv-containing regimens as per institutional policy. Repeat dosing at appropriate intervals may be required for prolonged infusions.
 - ▶ Systemic corticosteroids (eg, hydrocortisone or methylprednisolone) and/or montelukast may also be considered.
- Management of Grade 1–2 reactions:^b
 - ▶ Hold Bv infusion for 30–60 minutes.
 - ▶ For patients not already receiving premedications, administer an H1-antagonist as per institutional policy. Acetaminophen may also be considered. Depending on reaction severity, methylprednisolone 1 mg/kg/dose or equivalent corticosteroid may be administered.
 - ▶ Following resolution of symptoms, consider resuming Bv infusion at a reduced rate with administration of an H2-antagonist and/or montelukast administered 30 minutes prior (if not previously given for symptom management).
 - ◊ Infusion rate can be slowed by 50% to start and slowed further if needed. If the infusion is tolerated at a decreased rate, the infusion can be titrated back to the original rate as tolerated.
 - ◊ Premedications can be repeated to allow for safe delivery of infusion.
 - ▶ For all subsequent doses, premedication with an H1-antagonist with or without acetaminophen, an H2-antagonist, and/or montelukast is recommended.
 - ◊ For any recurrent grade 2 reactions, methylprednisolone 1 mg/kg/

dose or IV corticosteroid equivalent should be added to premedications.

- Management of Grade 3 reactions:^b
 - ▶ Hold Bv infusion and do not resume during current cycle.
 - ▶ Administer an H1-antagonist and methylprednisolone 1 mg/kg/dose or equivalent corticosteroid +/- other supportive care medications as per institutional policy.
 - ▶ Consult with allergy/immunology specialist for guidance of re-challenge.
 - ▶ For all subsequent cycles, slow infusion rate to 50% or slower as tolerated and premedicate with an H1-antagonist, methylprednisolone 1 mg/kg/dose or equivalent corticosteroid, and acetaminophen. Montelukast and an H2-antagonist may also be considered.
 - ▶ For any recurrent grade 3 reactions, permanently discontinue regimen.
- Management of Grade 4 reactions:^b
 - ▶ Hold Bv infusion permanently.
 - ▶ Administer an H1-antagonist and methylprednisolone 1 mg/kg/dose or equivalent corticosteroid, and acetaminophen.
 - ▶ Epinephrine, bronchodilators, and/or repeated or higher doses of corticosteroids may be utilized as indicated.
 - ▶ Hospital admission for observation should be considered based on severity.

Risk of Cardiac Toxicity

- Strongly consider administering dexrazoxane to children and adolescents expected to receive ≥ 250 mg/m² of cumulative doxorubicin, or an equivalent amount of an alternative anthracycline.^{10,11}

^b All toxicity grades refer to CTCAE V5.0. National Institutes of Health; National Cancer Institute. Common Terminology Criteria for Adverse Events (CTCAE) version 5.0 2017. Available at: https://ctep.cancer.gov/protocoldevelopment/electronic_applications/ctc.htm. Accessed February 1, 2024.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.

References



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

SELECT PRINCIPLES OF SUPPORTIVE CARE REFERENCES

- ¹ Common Terminology Criteria for Adverse Events (CTCAE) V5.0. 2017. Available at: https://ctep.cancer.gov/protocoldevelopment/electronic_applications/ctc.htm#ctc_50. Accessed February 1, 2024.
- ² Smith EML, Kuisell C, Kanzawa-Lee GA, et al. Approaches to measure paediatric chemotherapy-induced peripheral neurotoxicity: a systematic review. *Lancet Haematol* 2020;7:e408-e417.
- ³ Parsons SK, Rodday AM, Pei Q, et al. Performance of the FACT-GOG-Ntx to assess chemotherapy-induced peripheral neuropathy (CIPN) in pediatric high risk Hodgkin lymphoma: Report from the Children's Oncology Group AHOD 1331 study. *J Patient Rep Outcomes* 2023;7:113.
- ⁴ Loprinzi CL, Lacchetti C, Bleeker J, et al. Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: ASCO Guideline Update. *J Clin Oncol* 2020;38:3325-3348.
- ⁵ Heneghan MB, Parsons SK, Keller FG, et al. Protocol-stipulated dose modification to manage chemotherapy-induced peripheral neuropathy in children, adolescents, and young adults with high-risk Hodgkin lymphoma. *JCO Oncol Pract* 2024;OP2400089.
- ⁶ Evens AM, Cilley J, Ortiz T, et al. G-CSF is not necessary to maintain over 99% dose-intensity with ABVD in the treatment of Hodgkin lymphoma: Low toxicity and excellent outcomes in a 10-year analysis. *Br J Haematol* 2007;137:545-552.
- ⁷ Ansell SM, Ristow KM, Colgan JP, et al. G-CSF administration increases pulmonary toxicity for Hodgkin's disease patients treated with bleomycin-containing chemotherapy. *Blood* 2004;104:1317-1317.
- ⁸ Ansell SM, Radford J, Connors JM, et al. Overall survival with brentuximab vedotin in stage III or IV Hodgkin's lymphoma. *N Engl J Med* 2022;387:310-320.
- ⁹ Herrera AF, LeBlanc M, Castellino SM, et al. Nivolumab+AVD in advanced-stage classic Hodgkin's lymphoma. *N Engl J Med* 2024;391:1379-1389.
- ¹⁰ Chow EJ, Aggarwal S, Doody DR, et al. Dexrazoxane and long-term heart function in survivors of childhood cancer. *J Clin Oncol* 2023;41:2248-2257.
- ¹¹ de Baat EC, van Dalen EC, Mulder RL, et al. Primary cardioprotection with dexrazoxane in patients with childhood cancer who are expected to receive anthracyclines: recommendations from the International Late Effects of Childhood Cancer Guideline Harmonization Group. *Lancet Child Adolesc Health* 2022;6:885-894.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

EVIDENCE BLOCKS FOR PRIMARY TREATMENT OF STAGE I-II CHL WITHOUT RISK FACTORS

Preferred Regimens	
OEPA x 2 cycles (if adequate response)	
OEPA x 2 cycles + ISRT (if inadequate response)	
Useful in Certain Circumstances	
AVPC x 3 cycles (for mixed cellularity only, if adequate response)	
AVPC x 3 cycles + ISRT (for mixed cellularity only, if inadequate response)	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

EVIDENCE BLOCKS FOR PRIMARY TREATMENT OF STAGE I-II CHL WITH RISK FACTORS

Preferred Regimens	
ABVE-PC x 4 cycles (if adequate response [RER])	
ABVE-PC x 4 cycles + ISRT (if inadequate response [SER], or if initially RER but <CR at end of therapy)	
Bv-AVE-PC x 5 cycles + ISRT to sites of LMA (if adequate response [RRL])	
Bv-AVE-PC x 5 cycles + ISRT to sites of LMA and SRL and boost to sites of inadequate response at EOST (if inadequate response [SRL])	
Other Recommended Regimens	
OEPA x 2 cycles + COPDAC x 2 cycles (if adequate response)	
OEPA x 2 cycles + COPDAC x 2 cycles + ISRT (if inadequate response)	
OEPA x 2 cycles + COPDAC x 4 cycles (if adequate response)	
OEPA x 2 cycles + COPDAC x 4 cycles + ISRT (if inadequate response)	
AEPA x 2 cycles + CAPDAC x 4 cycles (if adequate response)	
AEPA x 2 cycles + CAPDAC x 4 cycles + RNRT (if inadequate response)	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)

EVIDENCE BLOCKS FOR PRIMARY TREATMENT OF STAGE III-IV CHL

Preferred Regimens	
Nivolumab-AVD x 6 cycles (if adequate response)	
Nivolumab-AVD x 6 cycles + RSRT (if inadequate response)	
Other Recommended Regimens	
Bv-AVE-PC x 5 cycles + ISRT (if adequate response [RRL])	
Bv-AVE-PC x 5 cycles + ISRT (if inadequate response [SRL])	
OEPA x 2 cycles + COPDAC x 4 cycles (if adequate response)	
OEPA x 2 cycles + COPDAC x 4 cycles + ISRT (if inadequate response)	
AEPA x 2 cycles + CAPDAC x 4 cycles (if adequate response)	
AEPA x 2 cycles + CAPDAC x 4 cycles + RNRT (if inadequate response)	
Useful in Certain Circumstances	
Bv-AVD x 6 cycles (if adequate response)	
Bv-AVD x 6 cycles + RSRT (if inadequate response)	
ABVE-PC x 4 cycles + ISRT (if adequate response [RER])	
ABVE-PC x 4 cycles + ISRT (if inadequate response [SER], or if initially RER but <CR at end of therapy)	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-DEF.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

5	Cancer Network, Inc.				
4					
3					
2					
1					
	E	S	Q	C	A

Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

EVIDENCE BLOCKS FOR PRIMARY TREATMENT OF NLPHL

Stage IA (complete resection; single node) confirmed by FDG-PET/CT	
Other Recommended Regimen	
CVbP x 3 cycles	

Stage IA or IIA (incomplete resection and non-bulky; multiple nodes)	
Preferred Regimens	
AVPC x 3 cycles (if adequate response)	
AVPC x 3 cycles + ISRT (if inadequate response)	
CVbP x 3 cycles (if adequate response)	
CVbP x 3 cycles + ISRT (if inadequate response)	
CVbP + rituximab x 3 cycles (if adequate response)	
CVbP + rituximab x 3 cycles + ISRT (if inadequate response)	
Other Recommended Regimens	
OEPA x 2 cycles (if adequate response)	
OEPA x 2 cycles + ISRT (if inadequate response)	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)

EVIDENCE BLOCKS FOR RELAPSED/REFRACTORY PEDIATRIC CHL: RE-INDUCTION AND SUBSEQUENT THERAPY OPTIONS WITH PEDIATRIC DATA

CPI-Based Regimens (Preferred)	
Bv + nivolumab	
Bv + nivolumab + bendamustine	
Other Recommended: Non-CPI-Containing Regimens	
Bortezomib + ifosfamide + vinorelbine	
Bv + gemcitabine	
EPIC (etoposide, prednisolone, ifosfamide, cisplatin)	
GDP (gemcitabine, dexamethasone, cisplatin)	
GV (gemcitabine, vinorelbine)	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

EVIDENCE BLOCKS FOR RELAPSED/REFRACTORY PEDIATRIC CHL: USEFUL IN CERTAIN CIRCUMSTANCES OPTIONS WITH OR WITHOUT PEDIATRIC DATA

CPI-Based Regimens	
GVD (gemcitabine, vinorelbine, doxorubicin liposome) + pembrolizumab	
ICE + nivolumab	
ICE + pembrolizumab	
Nivolumab	
Pembrolizumab	

Non-CPI-Containing Regimens	
Bv + bendamustine	
DHAP (dexamethasone, cytarabine, cisplatin)	
ICE (ifosfamide, carboplatin, etoposide)	

EVIDENCE BLOCKS FOR RELAPSED/REFRACTORY PEDIATRIC CHL: MAINTENANCE THERAPY (POST-TRANSPLANT)

Brentuximab vedotin	
---------------------	--

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

COTSWOLDS-MODIFIED ANN ARBOR STAGING SYSTEM

Stage	Definition
I	One nodal group or lymphoid organ (eg, spleen or thymus)
IE	Local extension from one nodal group to another site ^a
II	Two or more nodal groups, same side of the diaphragm
IIE	Localized extension from one nodal group to an extranodal site with stage II criteria, both on the same side of the diaphragm ^a
III	Nodal groups on both sides of the diaphragm
IIIS ₁	With splenic involvement
IIIE ₂	With localized extension from one nodal group to an extranodal site ^a
IIISE	Both IIIS1 and IIIE2
IV	Disseminated involvement of one or more extralymphatic organs (eg, lung, bone, bone marrow, liver) with or without any nodal involvement

Additional Sub-staging Variables	
A	Asymptomatic
B	Presence of B symptoms (unexplained recurrent fever >38°C within last month; drenching night sweats; or weight loss >10% of body weight within 6 months of diagnosis)
X	Bulky nodal disease: nodal mass >1/3 of intrathoracic diameter or 6 cm ^b in dimension

Lister TA, Crowther D, Sutcliffe SB, et al. Report of a committee convened to discuss the evaluation and staging of patients with Hodgkin's disease: Cotswolds meeting. J Clin Oncol 1989;7:1630-1636.

^a Based on Panel consensus.

^b In adults, 10 cm dimension is used.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

ABBREVIATIONS

3D-CRT	three-dimensional conformal radiation therapy	FEV1	forced expiratory volume in the first second	NLPHL	nodular lymphocyte-predominant Hodgkin lymphoma
4D-CT	four-dimensional computed tomography	FNA	fine-needle aspiration	OAR	organ at risk
ASCR	autologous stem cell rescue	FT4	free thyroxine	PA	posteroanterior
AYA	adolescent and young adult	FVC	forced vital capacity	PD-L1	programmed death ligand 1
CBC	complete blood count	G-CSF	granulocyte colony-stimulating factor	peds-mTNS	Pediatric-modified Total Neuropathy Score
CHL	classic Hodgkin lymphoma	GTV	gross tumor volume	PJP	<i>Pneumocystis jiroveci</i> pneumonia
CME	continuing medical education	H&P	history and physical	PR	partial response
CMP	comprehensive metabolic panel	HDT	high-dose therapy	PFT	pulmonary function test
CMV	cytomegalovirus	HIV	human immunodeficiency virus	PPD	product of perpendicular diameters
CPI	checkpoint inhibitor	HL	Hodgkin lymphoma	PTV	planning target volume
CR	complete response	HRS	Hodgkin-Reed-Sternberg	RER	rapid early response
CRP	C-reactive protein	ICC	International Consensus Classification	RNRT	residual-node radiation therapy
CRu	complete response, unconfirmed	IFRT	involved-field radiation therapy	RRL	rapidly responding lesion
CTCAE	Common Terminology Criteria for Adverse Events	IGRT	image-guided radiation therapy	RSRT	residual-site radiation therapy
CTV	clinical target volume	imAE	immune-mediated adverse event	SER	slow early response
DIBH	deep-inspiratory breath hold	IMRT	intensity-modulated radiation therapy	SRL	slow responding lesion
DLBCL	diffuse large B-cell lymphoma	irAE	immune-related adverse event	SUV	standardized uptake value
DLCO	diffusing capacity of the lung for carbon monoxide	ISRT	involved-site radiation therapy	TG	treatment group
EBER-ISH	Epstein-Barr virus-encoded RNA in situ hybridization	ITV	internal target volume	THRLBCL	T-cell/histiocyte-rich large B-cell lymphoma
EBV	Epstein-Barr virus	LDi	longest diameter	TNS-PV	Total Neuropathy Score-Pediatric Version
ECG	electrocardiogram	LMA	large mediastinal adenopathy	TSH	thyroid-stimulating hormone
ECHO	echocardiogram	LP	lymphocyte predominant	VMAT	volumetric modulated arc therapy
EOST	end of systemic therapy	MMR	mediastinal mass ratio		
ESR	erythrocyte sedimentation rate	NHL	non-Hodgkin lymphoma		
FDG	fluorodeoxyglucose	NLPBL	nodular lymphocyte-predominant B-cell lymphoma		



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

NCCN Evidence Blocks™

NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Discussion

This discussion corresponds to the NCCN Guidelines for Pediatric Hodgkin Lymphoma. Last updated: May 13, 2026.

Table of Contents

Overview	MS-2
Guidelines Update Methodology	MS-3
Literature Search Criteria	MS-3
Sensitive/Inclusive Language Usage	MS-3
Diagnosis and Workup for Hodgkin Lymphoma	MS-3
Clinical Staging and Risk Stratification	MS-5
Principles of Imaging	MS-6
Principles of Radiation Therapy	MS-7
Management of CHL	MS-8
Stage I–II CHL without Risk Factors	MS-8
Stage I–II CHL with Risk Factors and Stage III–IV CHL	MS-9
Relapsed or Refractory CHL	MS-13
Management of NLPHL	MS-17
Stage IA NLPHL with Complete Resection confirmed by FDG-PET/CTMS-17	
Stage IA or IIA NLPHL with Incomplete Resection and Non-Bulky ...	MS-18

Stage IA–IIA with Bulk, IB–IIB, III, IV NLPHL	MS-19
Relapsed or Refractory NLPHL	MS-19
Select Principles of Supportive Care	MS-20
Immunotherapy	MS-20
Peripheral Neuropathy	MS-20
Risk of Cardiac Toxicity	MS-20
Follow-Up After Completion of Treatment	MS-21
Summary	MS-22
References	MS-23



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Overview

Hodgkin lymphoma (HL) is an uncommon malignancy involving lymph nodes and the lymphatic system.¹ The WHO classification divides HL into two main types: classic Hodgkin lymphoma (CHL) and nodular lymphocyte-predominant HL (NLPHL).² CHL accounts for the majority of childhood HL, with NLPHL making up 5% to 10% of cases.¹

CHL is characterized by the presence of large binucleate or multinucleated neoplastic cells or mononuclear variants (collectively termed Hodgkin Reed-Sternberg [HRS] cells) in an inflammatory background, whereas NLPHL lacks HRS cells but is characterized by the presence of lymphocyte-predominant (LP) cells, sometimes termed *popcorn cells*.¹ In CHL, HRS cells nearly always express CD30 and often express CD15 (in ~70%), while CD20 is expressed in only ~6% to 10%. In contrast, in NLPHL, the characteristic LP cells express CD20 and rarely express CD30 and CD15.¹

CHL is divided into four subtypes: nodular sclerosis CHL; mixed cellularity CHL; lymphocyte-depleted CHL; and lymphocyte-rich CHL.² There are six immunoarchitectural patterns of NLPHL: B-cell-rich nodular (pattern A), serpiginous/interconnected nodular (pattern B), nodular with prominent extranodular LP cells (pattern C), T-cell-rich nodular (pattern D), T-cell-rich diffuse or T-cell/histiocyte-rich large B-cell lymphoma (THRLBCL)-like (pattern E), and diffuse B-cell-rich (pattern F).³ Most patients are diagnosed with HL between 15 and 30 years of age, followed by another peak in adults aged ≥ 55 years. Although the exact etiology is unknown, some risk factors for HL include prior infection with Epstein-Barr virus (EBV) and immunocompromising conditions, including immunosuppression after organ transplantation or infection with human immunodeficiency virus (HIV).⁴⁻⁶

In 2026, an estimated 8920 people will be diagnosed with HL in the United States and 1100 people will die from the disease.⁷ In adolescents (aged

15–19 years), HL is the most commonly diagnosed cancer⁸; it is estimated that 4200 adolescents and young adults (AYAs) 15 to 39 years of age were diagnosed with HL in 2020, with 800 in patients 15 to 19 years of age.⁹ The incidence is less common in children (aged 5–14 years); in 2014, a report estimated that 380 children would be diagnosed with HL that year.⁸

The past few decades have seen significant progress in the management of pediatric HL, with estimated 5-year survival rates of >98% after treatment with chemotherapy alone or combined with radiation therapy (RT).^{10,11} However, the potential long-term effects of treatment remain an important consideration.^{10,11}

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Pediatric Hodgkin Lymphoma were developed as a result of meetings convened by a multidisciplinary Panel of pediatric HL experts, with the goal of providing recommendations on standard treatment approaches based on current evidence. The NCCN Guidelines currently focus on clinical staging of HL, and treatment strategies are adapted according to risk. Given the complexity of HL treatment regimens and the required supportive care measures, the NCCN Pediatric Hodgkin Lymphoma Panel recommends a consultation with centers participating in pediatric cooperative group trials. Consistent with NCCN philosophy, participation in clinical trials is always encouraged.

The Panel also considers the term “pediatric” to include any patient ≤ 18 years of age, and recommendations in the Guidelines may extend to AYA patients up to 39 years of age. Across treatment centers, practice patterns vary with regard to AYA patients in terms of whether patients with HL are treated primarily by pediatric or adult oncologists. This Guideline is intended to apply to pediatric patients and may also apply to AYA patients treated in an adult oncology setting.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Guidelines Update Methodology

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Literature Search Criteria

Prior to the update of the NCCN Guidelines® for Pediatric Hodgkin Lymphoma, an electronic search of the PubMed database was performed to obtain key literature published in pediatric HL, using the following search terms: Hodgkin lymphoma and childhood or pediatric or adolescent or young adult. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.¹² Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Meta-Analysis; Randomized Controlled Trial; Meta-Analysis; Multicenter Study; Systematic Reviews; and Validation Studies.

The data from key PubMed articles as well as articles from additional sources deemed as relevant to these Guidelines as discussed by the Panel have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel's review of lower-level evidence and expert opinion.

Sensitive/Inclusive Language Usage

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation. NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-weight-biased; and inclusive of individuals of all sexual orientations and gender identities. NCCN Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will

continue to use the terms men, women, female, and male when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Diagnosis and Workup for Hodgkin Lymphoma

For evaluation and initial workup of HL, the Panel recommends that an excisional or incisional lymph node biopsy generally be performed. A core needle biopsy may be adequate if diagnostic for CHL, but is not adequate for the diagnosis of NLPHL. Fine-needle aspiration (FNA) is not an appropriate diagnostic test. Diagnosis should be established according to guidelines in the 2022 World Health Organization (WHO) Classification Haematolymphoid Tumours, 5th edition² or 2022 International Consensus Classification (ICC) of Mature Lymphoid Neoplasms.¹³

Immunostaining for CD30, CD15, CD20, CD3, and PAX5 is recommended. Evaluation of an expanded panel of markers (ie, CD45, CD79a, ALK, MUM1, OCT2, BOB1) and assessing for EBV with EBER-ISH should be considered in cases with equivocal or imperfect morphologic features, or to exclude entities in the differential diagnosis. The HRS cells of CHL express CD30 in all patients, express CD15 in the majority of patients, and are usually negative for CD45 and CD3. CD20 may be detectable in a minority of cases. The LP cells of NLPHL express CD20 and also express CD45. Characteristics of the immune microenvironment (identification of small B cells, T-follicular helper cells, and follicular dendritic cells) are also helpful in the diagnostic workup of NLPHL. Cases of EBV+ CHL may benefit from additional studies such as



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

EBV serology and evaluation for underlying immunodeficiency. EBV+ LP cells have been reported only rarely in NLPHL and this finding should prompt consideration of other entities in the differential diagnosis.¹⁴ Per WHO 2022, NLPHL remains under the family of HL, while ICC 2022 replaces the term NLPHL with nodular lymphocyte predominant B-cell lymphoma (NLPBL).^{2,13} Both classifications recognize the clinical and biologic similarities of NLPHL to an indolent B-cell lymphoma and NLPBL is an acceptable term per WHO 2022. While a few papers have reported on molecular/genetic findings in HL, molecular/genetic testing does not currently have established diagnostic or therapeutic utility in these diseases and is not recommended as part of standard routine practice. For additional information, see *Principles of Pathology* in the algorithm.

The workup should include a thorough history and physical examination, including determination of one or more B symptoms (unexplained recurrent fevers >38°C within the last month; drenching night sweats within the last month; weight loss of >10% of body weight within 6 months of diagnosis) and examination of lymphoid regions and spleen. Screening for a personal history of autoimmune disorder(s) is also recommended, as checkpoint inhibitors (CPIs) should be used with caution in those with active or previously documented autoimmune disease and/or current use of immunosuppressive agents. Other essential workup components include standard laboratory tests (complete blood count [CBC] with differential; erythrocyte sedimentation rate [ESR] with or without C-reactive protein [CRP]; and a comprehensive metabolic panel). A pregnancy test should be performed before individuals of childbearing potential undergo treatment. HIV and hepatitis B and C testing are encouraged for patients with risk factors for HIV or unusual disease presentations. An immunodeficiency workup should also be considered for patients <5 years of age, those with recurrent infections, those with an atypical presentation, and those with a personal or family history of immunodeficiency.

¹⁸F-fluorodeoxyglucose (FDG)-PET scans are essential for initial staging and for evaluating residual masses at the end of treatment¹⁵ (see *Principles of Imaging* and *Principles of Staging* in the algorithm). For staging and risk assessment, diagnostic imaging should be done before initiating chemotherapy, including: FDG-PET/CT or FDG-PET/MRI scans (whole-body); diagnostic contrast-enhanced (IV ± oral) CT (neck, chest, abdomen, and pelvis); or CT of chest and MRI (neck, abdomen, and pelvis). When available, FDG-PET/CT and diagnostic CT should be performed as a combined examination to limit radiation exposure. FDG-PET scans should be assessed by a nuclear diagnostic radiologist experienced in reading Deauville scores for FDG-PET–adapted therapy. The final report for any FDG-PET/CT examination to define response should include the Deauville 5-point scale score, which is a visual score. If cross-sectional imaging is not available or is needed for a clinical trial, posterior-anterior and lateral chest x-rays are recommended to determine bulk of disease (mediastinal mass). Consultation with a radiation oncologist is strongly recommended when considering treatment options and to determine the adequacy of imaging for potential future RT.

In cases of FDG-PET positivity where sites of disease are not consistent with usual presentation of HL, or if there are unusual disease presentations (ie, HIV), additional clinical evaluation may be needed for staging. If FDG-PET is negative for anatomic lesions of concern, a biopsy should be considered.

In most cases, if the FDG-PET/CT displays a homogeneous pattern of marrow uptake, which is thought to be secondary to cytokine release,^{16,17} bone marrow involvement is not assumed. If there are multifocal (≥3) skeletal FDG-PET lesions without cortical destruction on CT, marrow involvement may be assumed and a bone marrow biopsy is not needed to confirm involvement.¹⁸ In select cases, if there are cytopenias and the FDG-PET scan is negative, a bilateral bone marrow biopsy may be considered.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

If anthracycline-based chemotherapy is indicated, an echocardiogram is recommended. Pulmonary function tests (PFTs), including diffusing capacity of the lungs for carbon monoxide (DLCO), are recommended for patients receiving bleomycin-based chemotherapy. In general, a forced expiratory volume (FEV1)/forced vital capacity (FVC) of at least 60% is acceptable for bleomycin use, unless this is due to large mediastinal mass from HL. For children who are unable to cooperate for PFTs, the criteria for bleomycin use are: no evidence of dyspnea at rest, no exercise intolerance, and a pulse oximetry reading of >92% on room air.

The Panel recommends counseling on infertility risk.¹⁹ In select cases and if the patients are interested, the Panel recommends consideration of fertility preservation (eg, semen cryopreservation, ovarian tissue, oocyte cryopreservation) prior to the initiation of therapy. In general, the Panel also recommends providing referrals for counseling as needed that address smoking cessation, substance use disorders, and psychosocial concerns. For additional recommendations, see the NCCN Guidelines for Supportive Care (available at www.NCCN.org). Assessing social determinants of health may also be beneficial in selected cases (see the NCCN Guidelines for Distress Management [DIS-A], available at www.NCCN.org).

Clinical Staging and Risk Stratification

Physical examination and diagnostic imaging evaluations are used to designate the clinical stage.¹ The most widely used staging scheme for both pediatric and adult HL is the Ann Arbor Staging System, which may include the Cotswolds modification—which includes the prognostic significance of bulky disease.^{1,20,21} Staging is generally defined as follows²¹:

- Stage I: One nodal group or lymphoid organ (eg, spleen, thymus, Waldeyer's ring)

- Stage II: Two or more nodal groups on the same side of the diaphragm
- Stage III: Nodal groups on both sides of the diaphragm
- Stage IV: Disseminated involvement of one or more extralymphatic organs (eg, lung, bone, bone marrow, liver) with or without any nodal involvement

Additional substaging variables include these terms:

- A: Asymptomatic
- B: Presence of B symptoms
- X: Bulky nodal disease, which is nodal mass greater than one-third of intrathoracic diameter on a chest x-ray or as defined by the protocol. Note: Pediatric protocols have also defined bulk disease as contiguous extramediastinal nodal mass >6 cm in the longest diameter, measured in axial, coronal, or sagittal dimension (including oblique measurement), and EuroNet defines bulk as volume of the largest contiguous lymph node mass ≥ 200 mL.
- E: Involvement of extralymphatic by direct extension from an adjacent nodal site

Note: The Ann Arbor Staging System is in need of revision, as it does not fully represent the current practice in staging pediatric HL. Refer to the original protocol for appropriate staging of “E-lesions.” An E-lesion is defined as a contiguous infiltration of a lymph node mass into extralymphatic structures or organs (eg, lung, bone). Pleural and pericardial involvement should be considered E-lesions, but a pleural or pericardial effusion alone is not considered an E-lesion. Disease that extends beyond the lymphatic system without adjacent lymphatic involvement is considered stage IV; liver or bone marrow involvement is always considered stage IV disease.²² A comprehensive imaging atlas that aids in the identification of



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

affected nodal or extranodal regions and organs is available.²³

Currently, there is no uniform risk stratification for pediatric HL, although several factors are considered to confer poor prognosis, including B symptoms, mediastinal and peripheral lymph node bulk, extranodal disease, number of nodal sites, Ann Arbor stage, serum markers for inflammation, gender, and response to initial chemotherapy.¹ To facilitate the interpretation and comparison of global clinical trials, an international collaborative effort was developed: the Staging Evaluation and Response Criteria Harmonization (SEARCH) for Childhood, Adolescent, and Young Adult Hodgkin Lymphoma (CAYAHL) working group.²⁴ As the SEARCH effort for CAYAHL develops, so will the evolution of harmonized risk stratification for pediatric HL.¹¹

There are several cooperative groups, including the Children's Oncology Group (COG) and the European Network for Pediatric Hodgkin Lymphoma (EuroNET-PHL).¹¹ In the Guidelines, the Panel has summarized clinical stage and associated clinical trial risk groups (see *Summary of Clinical Trial Risk Groups* in the algorithm) but notes pediatric HL risk stratification is evolving from traditional pediatric risk group definitions to a stage-based system aligned with adult staging (see the NCCN Guidelines for Hodgkin Lymphoma [Adult], available at www.NCCN.org). Clinicians should be aware that historical pediatric risk classifications and contemporary stage-based approaches may overlap during this transition. Due to the evolving nature of risk stratification, enrollment in a clinical trial is preferred. In addition, for patient treatment, the Panel recommends considering consultation with a center of expertise.

Principles of Imaging

Clinical management of pediatric CHL involves initial treatment with chemotherapy and assessment of treatment response with FDG-PET to determine the need for additional treatment.²⁵⁻²⁷

Given the avidity of pediatric lymphomas for FDG,²⁶ the Deauville criteria were defined for the interpretation of interim and end-of-treatment FDG-PET scans based on the visual assessment of FDG uptake in the involved sites (See Deauville Criteria Table in *Principles of Staging* in the algorithm). These criteria use a 5-point scale (5-PS) to determine the FDG uptake in the involved sites relative to that of the mediastinum and the liver.²⁸⁻³⁰ In the 5-PS (Deauville criteria), scores of 1 to 4 refer to initially involved sites and a score of 5 refers to an initially involved site and/or new lesions related to lymphoma.^{29,30} These criteria vary across different protocols as they have yet to be validated in a large pediatric trial. However, interim or end-of-treatment FDG-PET scans with a score of 1, 2, or 3 are generally considered “negative” and FDG-PET scans with a score of 4 and 5 are considered “positive.”³¹ A score of 4 can be difficult to assess when FDG uptake in mediastinal masses cannot clearly be differentiated from thymic uptake or inflammatory reactions,^{28,32,33} and treatment decisions in these cases will require clinical judgment. In addition, Deauville 4 may represent just a single area of persistent disease or failure to respond in any site. The 5-PS (Deauville criteria) has been validated in international multicenter trials for FDG-PET–guided interim response assessment and risk-adapted therapy in adult patients with HL.

FDG-PET imaging is important as a baseline measurement before therapy to determine the initial sites of involvement and to perform an early response assessment (ERA) (often after the initial 2 cycles of chemotherapy).¹⁵ Although the optimal time point for assessment and criteria for response are not uniform, interim assessment of response by FDG-PET is incorporated into pediatric HL treatment.¹ The Panel recommends diagnostic contrast-enhanced CT or MRI to adequately evaluate all sites of involvement, and FDG-PET/CT or FDG-PET/MRI for interim and end-of-therapy assessments. End-of-therapy assessments may also include contrast-enhanced diagnostic CT or MRI of original sites of disease if not included with FDG-PET and if the treatment approach



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

includes the need for assessment of anatomic sites of disease for evaluation of response. Special attention should be given to end-of-therapy scans when utilizing CPIs given their potential for delayed response or pseudo-progression.³⁴ Repeat short-interval FDG-PET imaging may be necessary to differentiate between progression and pseudoprogression. Biopsy may be warranted for persistent PET avidity on end of therapy imaging. In addition, the Panel recommends waiting for at least 8 to 12 weeks after the end of RT to perform FDG-PET to minimize false-positive results. The Panel also recommends consideration of measures such as warming or pharmacologic suppression to reduce brown fat activation to minimize false-positive FDG-PET findings.

In some cases, routine surveillance scans in the first year following completion of therapy may have utility; however, they are recommended to be limited thereafter.³⁵ During follow-up, scans should only be obtained if there is significant concern for relapse or for up to 2 years.

Principles of Radiation Therapy

RT can be delivered with photons, electrons, or protons, depending upon clinical circumstances. Although advanced RT techniques emphasize tightly conformal doses and steep gradients adjacent to normal tissues, the “low-dose bath” to normal structures such as the breasts must be considered in choosing the final RT technique. Therefore, target definition, delineation, and treatment delivery verification require careful monitoring to avoid the risk of tumor geographic miss and subsequent decrease in tumor control. Initial diagnostic imaging with contrast-enhanced CT, MRI, FDG-PET/CT or FDG-PET/MRI, ultrasound (US), and other imaging modalities facilitate target definition.

Data have shown that significant dose reduction to organs at risk (OARs; eg, lungs, heart, breasts, kidneys, spinal cord, esophagus, carotid arteries, bone marrow, stomach, muscle, soft tissue, salivary glands) can be achieved with advanced RT planning and delivery techniques such as

four-dimensional CT (4D-CT) simulation, intensity-modulated RT (IMRT)/volumetric modulated arc therapy (VMAT), image-guided RT (IGRT), respiratory gating, deep inspiration breath hold, or proton therapy. These techniques offer significant and clinically relevant advantages in specific instances to spare OARs and decrease the risk for normal tissue damage and late effects without compromising the primary goal of local tumor control.³⁶⁻⁴² However, the Panel notes that randomized prospective studies to test these concepts are unlikely to be done since these techniques are designed to decrease late effects, which usually develop ≥ 10 years after completion of treatment. Therefore, the guidelines recommend that RT delivery techniques that are found to best reduce the doses to the OARs in a clinically meaningful manner without compromising target coverage should be considered in these patients, who are likely to enjoy long life expectancies following treatment. Care should be taken to ensure that involved areas are contoured appropriately.⁴³

Involved-site RT (ISRT) is recommended as the appropriate field for HL and can safely replace involved-field RT (IFRT) or modified IFRT from earlier trials, though in HLHR13 involved-node RT was used to treat post-chemotherapy volumes. ISRT targets the originally involved nodal sites and possible extranodal extensions (which generally defines a smaller field than the classical IFRT),⁴⁴ and is intended to spare the adjacent uninvolved organs (such as lungs, bone, muscle, or kidney) when lymphadenopathy regresses following chemotherapy. Treatment planning for ISRT requires the use of CT-based simulation, and additional imaging techniques such as FDG-PET and MRI often enhance the treatment planning.

For patients with stage I–II disease with or without risk factors, the Panel recommends an RT dose of 21 Gy to all sites of disease. Sites of slow response (usually defined with specific anatomic and/or FDG-PET criteria) can receive a boost of up to 9 Gy (total dose of 21–30 Gy). Sites of partial



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

response (PR) should receive a boost of 9–19 Gy (total dose of 30–40 Gy). For patients with stage III–IV disease, the Panel discourages using regimens that require ISRT to all sites of disease. Instead, for bulky disease, a dose of 21 Gy may be considered. The RT doses recommended for sites of slow response and sites of PRs in the setting of stage I–II disease with or without risk factors are the same in this context. The Panel notes that residual site RT should only be used when dictated by the protocol or as a boost following standard ISRT.

For patients with relapsed or refractory disease, if no high-dose therapy (HDT) or autologous stem cell rescue (ASCR) is planned, an RT dose of 30 Gy is recommended. If HDT/ASCR is planned, an RT dose of 30 Gy to relapsed or refractory sites may be used, with a consideration of 21 Gy to initial sites that are no longer present with active disease. If FDG-PET positive (Deauville 4 to 5) after several lines of therapy, RT may be considered to achieve metabolic complete response (CR) before transplant. Boost RT doses of 10 to 15 Gy (total dose of 40–45 Gy) to FDG-PET–positive sites may also be considered.

Management of CHL

In this section, data from select clinical trials that are recommended in the Guidelines are reviewed to provide a rationale for their inclusion in the Guidelines. Treatment approaches arising from more recent clinical trials have been collaborative efforts including adult and pediatric patients.

Stage I–II CHL without Risk Factors

Approximately 26% to 34% of children and adolescents with HL present with low-risk disease.⁴⁵ Outcomes for children and adolescents with low-risk HL are high, so recent trials are focused on modifying treatment (ie, reduction or elimination of specific chemotherapeutic agents or RT).⁴⁵ For instance, the German Society of Pediatric Oncology and Hematology Hodgkin's Disease (GPOH-HD) study series has demonstrated that RT

can be eliminated from a combined modality treatment scheme for patients in treatment group (TG)-1 (early stages: I and IIA) who achieve a CR after chemotherapy (GPOH-HD-95 trial).⁴⁶

In the GPOH-HD-2002 study, the main goal was to replace a component of chemotherapy (ie, procarbazine with etoposide and dacarbazine) to decrease gonad toxicity.⁴⁷ In this trial, all patients were aged <18 years (n = 573); for induction, males (n = 287) received 2 courses of OEPA (vincristine, etoposide, prednisone, and doxorubicin), and females (n = 286) received 2 courses of OPPA (vincristine, procarbazine, prednisone, and doxorubicin).⁴⁷ After chemotherapy, all patients received IFRT at 19 Gy except patients in TG-1 (early stages: IA, IB, and IIA) who were in CR (residual tumor volume ≤95% and ≤2 mL of the initial volume). In TG-1, overall event-free survival (EFS) was 92% ± 2.0%, with no significant impact of RT on EFS.⁴⁷ In TG-2 (intermediate stages: IE, IIB, IIAE, and IIIA) and TG-3 (advanced stages: IIBE, IIIAE, IIIB, IVA, IVB, and IVE), there was no significant difference in EFS between males and females (90.2% ± 2.3% vs 84.7% ± 2.7%, respectively; *P* = .12).⁴⁷ This trial suggested that both regimens could be used in intermediate and advanced stages, but also confirmed findings from GPOH-HD-95 that RT could be eliminated in patients in TG-1 who experience CR after chemotherapy.^{46,47}

Building on the GPOH-HD studies, an international intergroup phase III titration study for CHL in children and adolescents (EuroNET-PHL-C1) aimed to investigate whether RT can be omitted in patients with an adequate morphological and metabolic response to OEPA.⁴⁸ Among 714 patients assigned to and treated on TG-1, an adequate response, defined as ≥50% reduction in tumor volume and PET activity ≤ mediastinal blood pool or background activity, was achieved in 62%. In this group that did not receive RT, the 5-year EFS was 86.5% (95% CI, 83.3%–89.8%), below the target of 90%. Post hoc analyses demonstrated that the target



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

EFS was achieved in patients without risk factors of bulky involvement or an elevated ESR.

In the COG AHOD0431 trial, the goal was to evaluate the efficacy of a lower-intensity regimen, AVPC (doxorubicin, vincristine, prednisone, and cyclophosphamide), in pediatric and AYA patients (≤ 21 years of age) with non-bulky, stage IA and IIA CHL.⁴⁹ All patients ($n = 278$) were treated with 3 cycles of AVPC, and patients who did not experience CR after 3 cycles received 21 Gy of IFRT. Patients who experienced a protocol-defined, low-risk relapse after chemotherapy alone were eligible for an integrated salvage regimen composed of vinorelbine, ifosfamide, dexamethasone, etoposide, cisplatin, and cytarabine, with growth factor support for 2 cycles, and IFRT.⁴⁹ At 4 years, 49.0% of patients had been treated with 3 cycles of AVPC without RT and 88% achieved CR without receiving HDT/ASCR or >21 Gy of IFRT. The OS rate was 99.6%. Patients with mixed cellularity histology had a 4-year EFS of 95.2% compared to an EFS of 75.8% in patients with nodular sclerosis histology ($P = .008$).⁴⁹ In this study, a negative FDG-PET scan after 1 cycle of chemotherapy (PET1) and an ESR rate ≤ 20 mm/h were associated with a favorable EFS outcome.⁴⁹ Overall there were 37 relapses, including 32 classified as low risk.⁵⁰ Twenty patients received protocol specified therapy for R/R disease, 15 of whom were able to avoid HDT/ASCR. The 5-year and 8-year second EFS rates were 80.5% (95% CI, 61.6%–90.8%) and 76.3% (95% CI, 56.3%–88.0%), respectively, providing additional support for consideration of a transplant-free approach for select patients with relapsed disease low-risk features.

NCCN Recommendations for Stage I–II CHL without Risk Factors

For patients with stage I–II CHL without risk factors (see *Clinical Staging of CHL* in the algorithm for risks factors as defined by clinical trials), the Panel recommends enrollment in an ongoing clinical trial or treatment with

OEPA according to GPOH-2002 or EuroNet-PHL-C1 (a category 1 recommendation) as the preferred strategies. In certain circumstances, for patients with mixed cellularity histology only, 3 cycles of AVPC may be considered per the AHOD0431 trial.

After initial cycles of chemotherapy, patients who experience adequate response may be followed. Omission of ISRT should be more strongly considered for patients that meet the GPOH-2002 response criteria, in recognition of the lower EFS observed among patients with risk factors treated on the PHL-C1 trial.⁴⁷ Patients who experience inadequate response receive ISRT (to all sites and boost to sites of inadequate response at end of chemotherapy per EuroNet-PHL-C1). Based on an end-of-therapy FDG-PET assessment, patients may be followed or may consider re-induction therapies if there is a concern for persistent disease.

In some pediatric patients with CHL, the ABVD regimen (doxorubicin, bleomycin, vinblastine, and dacarbazine) may be considered.^{51–55} The Panel recommends referring to the adult NCCN Guidelines for Hodgkin Lymphoma (available at www.NCCN.org) to review relevant data and context.

Stage I–II CHL with Risk Factors and Stage III–IV CHL

The phase III COG AHOD0031 study evaluated the role of early chemotherapy response in tailoring subsequent therapy in pediatric intermediate-risk HL (stages IB, IAE, IIB, IIAE, IIIA, IVA with or without bulk, and IA or IIA with bulk).⁵⁶ Patients with newly diagnosed intermediate-risk HL ($n = 1712$; aged <22 years) received 2 cycles of ABVE-PC (doxorubicin, bleomycin, vincristine, etoposide, cyclophosphamide, and prednisone) followed by ERA with PET/CT. For patients who experienced adequate response (rapid early response [RER], based on anatomic criteria), 2 additional cycles of ABVE-PC were given followed by an evaluation for CR. Those who experienced RER with CR ($\geq 80\%$ reduction in the product of perpendicular parameters [PPD] or a



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

return to normal size for all target lesions, plus no residual extramediastinal nodal mass >2.0 cm, no residual disease in non-measurable sites, and a negative gallium or FDG-PET scan) were randomly assigned to IFRT (21 Gy) or observation, and those who experienced RER with less than CR were nonrandomly assigned to receive IFRT. Patients who experienced inadequate response (slow early response [SER]) after 2 cycles of ABVE-PC were randomly assigned to receive or not receive 2 cycles of chemo-intensification with DECA (dexamethasone, etoposide, cisplatin, and cytarabine) followed by 2 additional cycles of ABVE-PC. All patients in the SER group were randomized to receive IFRT.

The overall 4-year EFS was 85% (86.9% in the setting of RER and 77.4% in the setting of SER; $P < .001$), and the 4-year OS was 97.8% (98.5% in the setting of RER and 95.3% in the setting of SER; $P < .001$).⁵⁶ In those who experienced RER with CR at the end of chemotherapy, there was no significant difference in the 4-year EFS rate between patients who received IFRT versus those who did not receive IFRT (87.9% vs 84.3%, respectively). For those who experienced SER who received either DECA or no DECA, the 4-year EFS was 79.3% versus 75.2%, respectively ($P = .11$). PET response imaging was not required but was obtained for the majority of patients as part of clinical care. Analysis of these data demonstrated that those who experienced SER with PET-positive lesions after 2 cycles had a marginal improvement in EFS on the DECA arm (70.7% vs 54.6%; $P = .05$). Overall, this study showed that RT can be omitted in those who experience RER with CR at the end of chemotherapy, and that augmenting chemotherapy in those who experience SER with PET-positive disease may be beneficial.⁵⁶

In the phase III, randomized COG AHOD1331 trial, the anti-CD30 antibody-drug conjugate brentuximab vedotin (Bv) with doxorubicin, vincristine, etoposide, prednisone, and cyclophosphamide (Bv-AVE-PC) was compared to ABVE-PC in pediatric patients with previously untreated

stage IIB with bulk tumor, stage IIIB, stage IVA, or stage IVB disease ($n = 587$; aged 2–21 years).⁵⁷ ISRT (21 Gy) was administered following cycle 5 to slow-responding lesions (defined by a Deauville score of 4 or 5 on FDG-PET following cycle 2) and to LMA present at diagnosis. With a median follow-up of 42.1 months, 3-year EFS was 92.1% (95% CI, 88.4%–94.7%) in the Bv-AVE-PC group compared to 82.5% (95% CI, 77.4%–86.5%) in the ABVE-PC group (hazard ratio [HR], 0.41; 95% CI, 0.25–0.67; $P < .001$). The percentage of patients requiring ISRT did not differ significantly between the two groups (53.4% in the Bv-AVE-PC group vs 56.8% in the ABVE-PC group, respectively). OS rates at 3 years were 99.3% (95% CI, 97.3%–99.8%) in the Bv-AVE-PC group compared to 98.5% (95% CI, 96.0%–99.4%) in the ABVE-PC group. Toxicity was similar in the two groups, with an overall incidence of clinically significant adverse events (AEs) in 73.5% of patients in the Bv-AVE-PC group versus 68.2% in the ABVE-PC group. There was no significant difference in rates of grade ≥ 2 peripheral neuropathy between the two groups (25.5% in the Bv-AVE-PC group vs 24.9% in the ABVE-PC group).

Building on TG-3 from the GPOH-HD2002 trial, the HLHR13 multicenter trial studied the efficacy of Bv in frontline treatment of pediatric patients ≤ 18 years with stage IIB, IIIB, or IV CHL.⁵⁸ In this study, Bv replaced vincristine in the OEPA/COPDAC regimen. Patients were treated with 2 cycles of AEPA (A = Bv) followed by 4 cycles of CAPDAC. ERA was obtained following AEPA cycles and lymph nodes not achieving CR were targeted with response-based residual node radiation (RNRT). Compared to a historical control group treated with the Stanford V regimen (prednisone, nitrogen mustard, doxorubicin, vincristine, etoposide, bleomycin, and vinblastine) from the HOD99 study, more patients achieved CR at ERA with AEPA than with Stanford V (35.1% vs 17%; $P = .004$) and did not require RT. Three-year EFS and OS rates with AEPA were 97.4% and 98.7% respectively, compared to 80.8% ($P = .0008$) and 96.5% ($P = .311$) with Stanford V.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Other clinical studies evaluating the intermediate-risk group (TG-2) and high-risk group (TG-3) include the GPOH-HD-2002 and EuroNET-PHL-C1 trials^{47,59} as described under *Stage I–II CHL without Risk Factors*.

In a follow-up titration study of EuroNET-PHL-C1, patients were randomly assigned to 2 (TG-2) or 4 (TG-3) cycles of COPP or COPDAC after an initial 2 cycles of OEPA.⁶⁰ RT was omitted in patients who experienced adequate response (partial remission or greater and FDG-PET negativity [Deauville score 1–2]) following 2 cycles of OEPA. Those who experienced inadequate response received ISRT and an additional boost to sites that responded slowly to treatment. Five-year EFS in those who experienced adequate responses and for whom RT was omitted was 90.1% compared to 87.1% in those who experienced inadequate response and went on to RT. Five-year EFS in all patients treated with COPP was 89.9% versus 86.1% in all those treated with COPDAC, with a difference of -3.7%. A subgroup analysis further investigated responses to COPP versus COPDAC in those who experienced adequate versus inadequate response and found a treatment difference of -9.1% (91.9% with COPP and 82.9% with COPDAC). There was no difference in OS between COPP and COPDAC. Results of this trial indicate that RT can be safely omitted in patients with intermediate- and advanced-stage disease who experience adequate response to OEPA when treated with either COPP or COPDAC. Results also show noninferiority of the less gonadotoxic COPDAC compared to COPP in patients who received RT; however, COPDAC was inferior to COPP in those for whom RT was omitted.

The randomized phase II SWOG S1826 trial compared the safety and efficacy of 6 cycles of nivolumab-AVD (doxorubicin, vinblastine, and dacarbazine) to brentuximab vedotin (Bv)-AVD in patients ≥12 years of age (n = 970 in the intention-to-treat cohort; median age 27.6 years; n = 236 within the age range of 12–17 years) with stage III–IV HL.⁶¹ Use of growth factor was optional with N-AVD but mandatory with Bv-AVD during the trial. Two-year progression-free survival (PFS) was significantly

improved with nivolumab-AVD (92%) compared to Bv-AVD (83%) (HR for disease progression or death, 0.45; 95% CI, 0.3–0.65) in the overall cohort. Specifically within the adolescent cohort, 3-year PFS was 93% in the nivolumab-AVD arm compared to 82% in the Bv-AVD arm (HR, 0.37; 95% CI, 0.17–0.80) and 3-year EFS was 89% with nivolumab-AVD versus 80% with Bv-AVD (HR, 0.55; 95% CI, 0.29–1.06).⁶² EFS events were from progression/relapse (N-AVD n = 9; Bv-AVD n = 22); nonprotocol therapy (N-AVD n = 5; Bv-AVD n = 2); and death (n = 1 on Bv-AVD arm before therapy initiation).. RT was administered sparingly in both arms (1 patient with nivolumab-AVD compared to 2 patients with Bv-AVD). Despite a higher use of growth factor in the Bv-AVD arm, only 3% of patients had grade ≥3 febrile neutropenia and 1% with sepsis after either regimen. Overall rates of immune-related AEs of any grade were low. As expected, thyroid dysfunction was more frequent after N-AVD, whereas sensory peripheral neuropathy was more frequent after Bv-AVD. The majority of adolescent patients received dexrazoxane for cardiac protection.

Data for Bv-AVD otherwise comes from adult trials, including the randomized phase III ECHELON-1 trial, which compared the safety and efficacy of ABVD to Bv-AVD in patients ≥18 years with previously untreated stage III or IV CHL.⁶³ Five-year follow-up data confirmed a PFS benefit for Bv-AVD compared to ABVD that was consistent among all patient subgroups independent of disease stage, age, and IPS. A more recent interim analysis revealed a significant OS benefit with Bv-AVD compared to ABVD ($P = .009$).

NCCN Recommendations for Stage I–II CHL with Risk Factors

For patients with stage I–II CHL with risk factors, the Panel recommends enrollment in an ongoing clinical trial or treatment according to AHOD0031 (not for stage 2BX) or AHOD1331 (for stage 2BX only) as the category 1, preferred strategies.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

For the AHOD0031 regimen, after 2 initial cycles of ABVE-PC, patients who experience adequate response are treated with 2 additional cycles of ABVE-PC. Based on an end-of-therapy FDG-PET assessment and CR achievement with CT criteria, patients may either be followed or treated with ISRT to all sites if less than CR. Patients who experience inadequate response receive 2 additional cycles of ABVE-PC and ISRT.

For the Bv-AVE-PC regimen based on AHOD1331, after 2 initial cycles of Bv-AVE-PC, patients who experience adequate response (rapidly responding lesions [RRLs]) are treated with 3 additional cycles of Bv-AVE-PC and ISRT to sites of LMA. Patients who experience inadequate response (slow responding lesions [SRLs]) receive 3 additional cycles of Bv-AVE-PC and ISRT to sites of LMA and SRL. The addition of boost is dependent on FDG-PET–positive lesions at the end of systemic therapy. Pneumocystis jiroveci pneumonia (PJP) prophylaxis is recommended for BV-AVE-PC and for all other BV containing regimens.

Treatment according to HLHR13 or EuroNet PHL-C1 may be useful in certain circumstances. In the regimen based on HLHR13, after 2 initial cycles of AEPA, patients who experience adequate response are treated with 4 cycles of CAPDAC. Patients who experience inadequate response are treated with 4 cycles of CAPDAC and RNRT to sites of inadequate response. In a similar fashion, as per EuroNet PHL-C1, following 2 initial cycles of OEPA, patients who experience adequate response are treated with 2 cycles (as per TG-2; not for stage IIBE; category 1 recommendation) or 4 cycles (as per TG-3; for stage IIBE and IIBX only; category 2B recommendation) of COPDAC and those who experience inadequate response are treated with the same chemotherapy regimens but with the addition of ISRT to all sites and boost to sites of inadequate response at end of chemotherapy.

In all cases, based on an end-of-therapy FDG-PET assessment and CT scan, patients may either be followed or considered for biopsy to confirm persistent active disease.

As noted for stage I–II CHL without risk factors, the ABVD regimen, as well as other adult regimens may be considered for pediatric patients >18 years of age.⁵¹⁻⁵⁵ The Panel recommends referring to the adult NCCN Guidelines for Hodgkin Lymphoma (available at www.NCCN.org) to review relevant data and context for these regimens.

NCCN Recommendations for Stage III–IV CHL

For patients with stage III–V CHL, the Panel recommends enrollment in an ongoing clinical trial or treatment with nivolumab-AVD according to SWOG S1826 as the preferred strategies. Nivolumab-AVD has only been studied in patients ≥12 years of age and 6 cycles are recommended prior to response assessment. Patients who experience adequate response can be followed. For those experiencing inadequate response, regimens for relapsed/refractory disease or residual-site RT (RSRT) to sites of inadequate response can be considered. Of note, false-positive findings may be observed on FDG-PET/CT or FDG-PET/MRI after treatment with nivolumab-AVD, given the potential for delayed response or pseudo-progression with CPIs. Clinical judgment is often necessary in such settings. Repeat imaging at short intervals (per LYRIC recommendations) may be considered prior to obtaining a biopsy.³⁴

Other recommended regimens include the Bv-AVE-PC regimen based on AHOD1331 (not for stage 3A), OEPA followed by 4 cycles of COPDAC based on EuroNet-PHL-C1 (not for stage 3A, other than stage 3AE), and AEPA followed by 4 cycles of CAPDAC based on HLHR13 as described under *NCCN Recommendations for Stage I–II CHL with Risk Factors*.

Bv-AVD as per SWOG S1826 may be useful in certain circumstances for patients that are unable to receive or tolerate a CPI. Mirroring



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

recommendations for nivolumab-AVD, Bv-AVD is only recommended for patients ≥ 12 years of age, with 6 cycles recommended prior to response assessment. Therapy for relapsed/refractory disease or RSRT to sites of inadequate response can be considered for those experiencing inadequate response to Bv-AVD.

ABVE-PC per AHOD0031 as described under *NCCN Recommendations for Stage I–II CHL with Risk Factors* may be useful in certain circumstances for patients with stage 3A disease only.

For patients > 18 years of age, BrECADD (Bv, etoposide, cyclophosphamide, doxorubicin, dacarbazine, dexamethasone) or other adult regimens may be considered. The Panel recommends referring to the adult NCCN Guidelines for Hodgkin Lymphoma (available at www.NCCN.org) to review relevant data and context for these regimens.

In all cases, based on an end-of-therapy FDG-PET assessment, patients may either be followed or considered for biopsy to confirm persistent active disease.

Relapsed or Refractory CHL

Although the outcomes for pediatric HL are excellent, approximately 10% of patients with early-stage disease and up to 25% of patients with advanced-stage disease experience relapse.^{10,64,65} For patients with relapsed or refractory disease, treatment options include standard-dose chemotherapy (re-induction therapy), HDT with ASCR, or novel approaches.⁶⁴ Re-induction regimens can be divided into four major categories.⁶⁴

1) **Targeted therapy and immunotherapy-based regimens** include Bv combined with nivolumab with or without bendamustine,⁶⁶⁻⁶⁸ Bv combined with bendamustine^{69,70} or gemcitabine,⁷¹ or nivolumab, nivolumab with or without ICE,⁷²⁻⁷⁴ or pembrolizumab with or without ICE or GVD.⁷⁵⁻⁷⁸

The combination of Bv and nivolumab, a human monoclonal programmed cell death protein 1 (PD-1)–directed antibody, has been evaluated in adult patients with relapsed/refractory (R/R) HL prior to ASCT.⁷⁹ In this phase 1/2 study, patients received up to 4 cycles of Bv and nivolumab, with staggered dosing in cycle 1 followed by same-day dosing in cycles 2–4. CR and objective response rates were 61% and 82%, respectively. The most common AEs were nausea in 49% of patients, fatigue in 41%, and infusion reactions in 44%. Grade ≥ 3 AEs occurred in 31% of patients. Peripheral neuropathy was reported in 20% of patients.⁷⁹ Long-term follow-up data revealed an estimated 3-year PFS rate of 77% (91% for those who proceeded to ASCT following study treatment) and 3-year OS of 93%.⁸⁰ At a median of 34.3 months, no new AEs were observed.⁸⁰

In the phase II Checkmate 744 trial, children and AYA patients with standard-risk R/R HL ($n = 44$; median age range, 9–30 years) were treated with 4 cycles of Bv-nivolumab and experienced complete metabolic response (CMR) and overall response rate (ORR) rates of 59% and 82%, respectively (according to blinded independent central review [BICR]).⁶⁶ Those who experienced no metabolic response or partial metabolic response went on to receive intensification with Bv and bendamustine for 2 or 4 cycles, with CMR and ORR rates of 82% and 100% following 2 cycles and 67% and 100% following 4 cycles according to BICR. Combined CMR rate (with and without intensification) was 94% and 1-year PFS was 91%. The majority of immune-mediated AEs were grade 1 or grade 2. In the low-risk cohort, ($n = 28$; median age range 6–27 years) the goal was to replace HDT/ASCR with ISRT.⁶⁸ CMR per BICR after 4 cycles of Bv-nivolumab was 82%, with an ORR of 96.4%. Treatment was intensified to Bv and bendamustine for those experiencing less than a CMR. CMR per BICR at any time point before ISRT was 93%, with an ORR rate of 100%. Three-year EFS and PFS rates were 87% and 95%, respectively. Treatment was discontinued due to serious AEs in only 2 patients and immune-mediated adverse events were infrequent.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

In a group of heavily pretreated patients ≥ 18 years of age with R/R HL (n = 64) and anaplastic large cell lymphoma (n = 1), the safety and clinical activity of Bv and bendamustine was evaluated.⁶⁹ An overall response was achieved in 29 of 37 patients (78%).⁶⁹ The combination of Bv and bendamustine as part of salvage therapy was also assessed in a multicenter retrospective analysis of pediatric patients with R/R HL (n = 29; median age, 16 years).⁷⁰ CMR rate and objective response rates were 66% and 79%, respectively. Seventeen patients were able to undergo successful stem cell mobilization and 16 went on to consolidative transplant (13 autologous, 3 allogeneic). Three-year EFS and OS rates were 65% and 89%, respectively. Treatment was well tolerated overall, with the most common grade 3 or 4 AEs being hematologic. There were two cases of grade 3 hyperbilirubinemia and three episodes of grade 3 infusion reactions. There were no reports of significant neuropathy.

In a COG study (AHOD1221), the safety and efficacy of Bv and gemcitabine was evaluated in children and young adults with primary refractory or early relapsed HL (n = 46; aged < 30 years).⁷¹ Of 42 evaluable patients, 24 (57%) achieved a CR within the first 4 cycles of treatment; 4 of 13 patients (31%) who experienced a PR or who had stable disease had all target lesions with Deauville scores of ≤ 2 after 4 cycles of treatment.⁷¹ Using a Deauville score threshold of ≤ 3 , 28 of 42 (67%) achieved a CR.

Multiple studies have also demonstrated efficacy of nivolumab and pembrolizumab, another human monoclonal PD-1-directed antibody, in adult patients with R/R HL, as single agents or in combinations.⁷²⁻⁷⁸

In a study evaluating the efficacy of pembrolizumab in pediatric patients with R/R PD-L1-positive solid tumors or lymphomas (n = 154 evaluable patients; median age, 13 years; interquartile range [IQR], 8–15 years), 9 of 15 patients with R/R HL achieved an objective response (60%).⁷⁶ In the phase III KEYNOTE-204 study, heavily pretreated adult patients with R/R

CHL were randomized to receive either pembrolizumab or Bv (n = 300 evaluable patients; pembrolizumab arm, n = 148; Bv arm, n = 152; aged ≥ 18 years).⁸¹ The median PFS in the pembrolizumab treatment arm was statistically longer than the Bv treatment arm (13.2 months vs 8.3 months, respectively; HR, 0.65; 95% CI, 0.48–0.88; $P = .00271$).⁸¹

In a phase II trial, the combination of pembrolizumab with ICE chemotherapy was investigated in adults with R/R HL (n = 37 with evaluable data; age range, 19–70 years).⁷⁷ Patients received 2 cycles of pembrolizumab with ICE, following which stem cells were mobilized and collected, then pembrolizumab monotherapy for 1 cycle prior to FDG-PET/CT. CR and ORR rates were 86.5% and 97.3%, respectively. Two-year PFS and OS were estimated at 87.2% and 95.1%, respectively. Grade 3–4 adverse events, mostly cytopenias, were reported in 52.4% of patients.

The combination of pembrolizumab with GVD chemotherapy for adults with R/R HL was investigated in another phase II trial.⁷⁸ Patients (n = 38 with evaluable data, ages 21–71 years) received 2 or 4 cycles of pembrolizumab with GVD, with CR rates of 100% and 95%, respectively. Ninety-five percent of patients proceeded to HDT/ASCR. The majority of adverse events were grade 1–2.

The safety and efficacy of nivolumab monotherapy for adults with R/R HL who had previously undergone HDT/ASCR was investigated in the multi-cohort phase II CheckMate 205 trial.^{73,82} Patients (n = 243; median age, 34 years) were enrolled into cohorts based on prior treatment: cohort A had never received Bv; cohort B received Bv following HDT/ASCR, and cohort C received Bv before and/or after HDT/ASCR.⁸² Nivolumab was administered every 2 weeks until progression or unacceptable toxicity. With a median follow-up of 18 months, ORR, as assessed by an independent radiologic review committee, was 69%. Median duration of response and PFS were 16.6 months and 14.7 months, respectively.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Median PFS duration of response and PFS were highest in cohort A, though cohort B had the highest 1-year OS rate of 95%, compared to 93% in cohort A and 90% in cohort C. Elevated lipase, elevated ALT, and neutropenia were the most common grade 3–4 drug related adverse events, though most drug related adverse events were grade 1–2.

The efficacy of PET-adapted nivolumab for R/R HL, with or without ICE chemotherapy, as a bridge to HDT/ASCR was investigated in a phase II trial.⁷⁴ Patients (n = 43; age range, 18–70 years) received 3 cycles of nivolumab monotherapy prior to FDG-PET/CT. Patients achieving CR or PR received an additional 3 cycles of nivolumab monotherapy followed by repeat FDG-PET/CT. Patients experiencing progressive disease following either 3 or 6 cycles of nivolumab monotherapy received 2 cycles of nivolumab-ICE. Patients achieving CR after 6 cycles of nivolumab monotherapy or CR or PR following nivolumab-ICE could proceed to HDT/ASCR. Among the 42 patients with evaluable response data, ORR and CR rates following nivolumab monotherapy and nivolumab-ICE were 81% and 71% and 100% and 89%, respectively. Seventy-nine percent of patients were successfully bridged to HDT/ASCR, the majority of which had received nivolumab monotherapy. Among patients who were bridged to HDT/ASCR, 2-year PFS was 94% compared to 72% in the entire cohort. OS in the entire cohort was 95%. Three patients experienced grade ≥3 adverse events related to nivolumab.

2) **Gemcitabine-based regimens** such as GV⁸³⁻⁸⁵ (gemcitabine and vinorelbine).

The GV regimen was evaluated in heavily pretreated pediatric patients with R/R HL (n = 30; median age, 17.7 years; range, 10.7–29.4 years).⁸³ All patients had received at least two prior chemotherapy regimens and 17 patients had undergone prior HDT/ASCR. Overall, 19 of 25 patients had measurable disease responses for an observed response rate of 76%.⁸³ Patients who had received transplant before GV tended to be more likely

to have disease response to therapy compared to patients who had not received transplant.⁸³ In retrospective studies, the 3-year OS rate among pediatric patients with R/R HL who received GV has been reported at 74% to 80%.^{84,85}

3) **Ifosfamide-/etoposide-based regimens** including: ICE⁸⁶ (ifosfamide, carboplatin, and etoposide); IV⁸⁷ (ifosfamide, and vinorelbine); and a regimen composed of bortezomib and IV.⁸⁸ The ICE regimen was developed to decrease non-hematologic toxicities observed with cisplatin-containing regimens.^{65,86} In a study of patients with primary refractory or relapsed HL (n = 65; median age, 27 years; range, 12–59 years), after treatment with 2 biweekly cycles of ICE, patients whose disease responded to therapy received HDT/ASCR and IFRT. In this study, the response rate to ICE was 88% and the EFS for patients who underwent transplantation was 68%.⁸⁶

In a study evaluating the efficacy of the IV regimen, 66 patients <30 years of age with R/R HL were treated with 2 cycles of IV.⁸⁷ The ORR of 72% allowed most of the patients to undergo subsequent HDT/ASCR.⁸⁷ It is worth noting that this regimen eliminates etoposide, a chemotherapeutic agent associated with secondary malignancy after transplantation.^{87,89} Addition of bortezomib to the IV regimen does not improve anatomic CR after 2 cycles, but may improve the ORR at the completion of therapy.⁸⁸

4) **Platinum-based regimens** including: DHAP⁹⁰ (dexamethasone, cytarabine, and cisplatin); EPIC⁹¹ (etoposide, prednisolone, ifosfamide, and cisplatin); and GDP⁹² (gemcitabine, dexamethasone, and cisplatin). In a study of patients with relapsed/refractory (R/R) HL (n = 102; median age, 34 years; range, 21–64 years), the response rate after 2 cycles of DHAP was 89%.⁹⁰ In a retrospective study of 80 children with relapsed or primary refractory HL, treatment with the EPIC regimen (55% of patients received HDT/ASCR after first relapse following EPIC regimen) resulted in a 5-year OS and PFS from relapse of 75.8% and 59.9%, respectively.⁹¹ In



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

a study of patients with R/R HL (n = 23; median age, 36 years; range, 19–57 years) the response rate after 2 cycles of GDP was 69.5%.⁹²

Maintenance Therapy Post-Transplant

In a phase III trial, Bv given as consolidation following HDT/ASCR was shown to significantly improve 5-year PFS among adult patients with HL with unfavorable-risk relapsed HL or primary refractory HL.⁹³ In a retrospective study of 67 pediatric patients with R/R HL, Bv consolidation following HDT/ASCR, 3-year PFS was 85%.⁹⁴

In a phase II trial assessing pembrolizumab as a consolidative therapy following HDT/ASCR among adults with R/R HL (n = with evaluable data; age range, 20–69 years), 90% of whom had high-risk disease, 18-month PFS and OS were 82% and 100%, respectively.⁹⁵

Multiple studies support the addition of RT in the transplant setting by showing benefit for local tumor control and improved EFS/OS/disease-free survival (DFS).^{65,96,97}

NCCN Recommendations for Relapsed or Refractory CHL

Histologic confirmation with biopsy is recommended before initiating treatment for relapsed or refractory disease given the risk for transformation and high false-positive rate of FDG-PET/CT. Importantly, false positive findings may be observed on FDG-PET/CT or FDG-PET/MRI following treatment with regimens that include a CPI given their potential for delayed response or pseudo-progression. Clinical judgment is often necessary in such settings. Repeat imaging at short intervals per LYRIC recommendations may be considered prior to obtaining a biopsy.³⁴

Following a negative biopsy, the Panel recommends either observation with short-interval follow-up or additional workup, such as imaging or rebiopsy, if there is a high index of suspicion. Depending on the level of

clinical suspicion, a consultation with radiation oncology may be considered.

Following a positive biopsy and restaging, the Panel recommends enrollment in a clinical trial if available, and referral to or consultation with a center of expertise, as several options exist for the treatment of R/R disease. Cross-toxicity with prior lines of therapy, including but not limited to cumulative anthracycline dose, and prior RT therapy are important considerations in the selection of a re-induction regimen or subsequent therapy regimen. Considerations for RT in the R/R setting include site of relapse, history of prior RT, and eligibility for HDT/ASCT. RT alone may also be considered as consolidation following second-line therapy.⁹⁸

Typically, patients are treated with re-induction therapies, and after an FDG-PET/CT or FDG-PET/MRI assessment, if metabolic CR is observed (Deauville score ≤ 3), treatment can be followed up with HDT/ASCR with or without ISRT and with or without maintenance therapy. Of note, a higher risk of engraftment syndrome may be observed in patients who have been previously treated with CPIs.⁹⁹ In general, RT is performed as consolidation after transplant. If a metabolic CR is not achieved, RT may be used before transplant.

For patients with Deauville score of 4–5 following re-induction therapy, subsequent therapy options should be considered, including re-induction options that were not previously used. Alternatively, HDT/ASCR with or without ISRT may be considered. Of note, therapy following a Deauville 5 response to reinduction for R/R disease is not well established. Individualized treatment is necessary. Referral to or consultation with a center of expertise is strongly encouraged.

In certain cases, patients with Deauville score ≤ 3 following re-induction therapy may avoid ASCR. These include patients with initial disease stages other than IIIB or IVB, patients who have no prior exposure to RT,



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

patients with duration of first CR >1 year, patients with no extranodal disease or B symptoms at relapse, and those with adequate response to <4 cycles of chemotherapy following relapse between 3 and 12 months.^{50,68,100} In these patients, re-induction therapy plus ISRT may be considered for initial treatment of R/R HL.

A CPI-based regimen is preferred for those who are eligible, particularly if a CPI was not utilized in the frontline setting. Bv plus nivolumab, with or without bendamustine is the preferred re-induction therapy option. As previously noted, CPIs should be used with caution in those with active or previously documented autoimmune disease and/or current use of immunosuppressive agents. Other recommended non-CPI containing regimens with data in the pediatric setting include bortezomib + ifosfamide + vinorelbine, Bv + gemcitabine, GDP, GV, and EPIC.

Other regimens that may be useful in certain circumstances, with or without pediatric data, include CPI-based regimens such as pembrolizumab + ICE or GVD, nivolumab + ICE, or single agent pembrolizumab or nivolumab. Non-CPI containing regimens without pediatric data that may be useful in certain circumstances include Bv + bendamustine, DHAP, and ICE. Additional options may be considered for patients >18 years (see NCCN Guidelines for Hodgkin Lymphoma [Adult], available at www.NCCN.org). Referral for allogeneic hematopoietic cell transplantation (HCT) can be considered for appropriate patients.

For patients undergoing HDT/ASCR, in general, two post-transplant treatment options may be considered including: 1) maintenance therapy with Bv (especially useful in patients with disease with high-risk features including progressive disease, refractory disease, or relapse within 1 year of diagnosis)^{93,94}; and 2) RT consolidation after HDT/ASCR. Of note, the role of maintenance BV has not been well-defined in patients who received BV or a CPI prior to maintenance therapy. Similarly, the role of CPI maintenance has not been established.

Management of NLPHL

In this section, data from select clinical trials that are recommended in the Guidelines are reviewed to provide a rationale for their inclusion in the Guidelines.

Stage IA NLPHL with Complete Resection Confirmed by FDG-PET/CT

The presentation of NLPHL is clinically distinct from that of CHL. There is a strong male predominance (~70%) and pediatric patients often present at a younger age than CHL. Most patients are asymptomatic and present with early-stage disease (stage IA), with localized peripheral lymph node involvement. Mediastinal involvement is seen infrequently.^{101,102}

While pediatric patients with NLPHL have historically achieved excellent responses to regimens used for early-stage CHL, the chemotherapy and RT involved in these regimens carry risks of late toxicities, including secondary cancers and transformation to non-Hodgkin lymphoma (NHL).¹⁰¹ Several cooperative groups have published evidence to suggest that complete surgical resection of localized disease, confirmed postoperatively, is an effective option in a select group of patients.¹⁰¹⁻¹⁰⁴

In the Children's Oncology Group AHOD03P1 study, a total of 52 patients with stage IA NLPHL limited to a single lymph node were observed without further therapy following complete resection.¹⁰¹ Five-year EFS was 77.1%. Of the 52 patients in this group, 13 relapsed at a median of 11.5 months from the time of study enrollment. Twelve of the 13 relapses were stage IA, occurring at the initial site of disease or in adjacent nodes/nodal regions.

The European Network on Pediatric Hodgkin Lymphoma collected data from 58 patients with limited-stage disease treated with resection alone.¹⁰² Fifty-four of the patients had stage IA disease, 2 had stage IIA disease, and 2 had stage IIIA disease. At a median follow-up of 43 months, PFS



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

and OS rates were 57% and 100%, respectively. Of the patients who achieved CR after surgery (51 of 58 patients), overall PFS was 67% at 26-month follow-up. All patients who had residual disease after surgery had recurrence at a median of 17 months and 14 of the 51 patients who achieved CR after surgery had recurrence at a median of 26 months. Only 5 patients received RT. With a median follow-up of 52 months, PFS for patients who recurred was 80%.

Shankar and colleagues investigated low-intensity CVbP (cyclophosphamide, vinblastine, prednisone) in patients with early-stage NLP HL and found this to be a safe and effective upfront chemotherapy regimen for patients who were not able to be treated with surgical resection alone.¹⁰⁵ In this study, a total of 55 patients with either previously untreated (n = 45) or relapsed (following surgical resection) (n = 10) stage I or II NLP HL were treated with 3 cycles of CVbP prior to response assessment. Of the 45 patients who received first-line CVbP, 80% (n = 36) achieved either a CR or CRu (complete response unconfirmed) and the remaining (n = 9) patients achieved a PR. Of the 9 patients who experienced PR, all achieved CR with either salvage chemotherapy or IFRT. All 10 of the patients who received CVbP at relapse following surgical resection achieved a CR. For the entire cohort, 40-month freedom from treatment failure (FFTF) and OS were 75.4% and 100%, respectively. With a median follow-up period of 41 months, only 3 patients experienced relapse following CVbP (all 3 with stage IIA disease at diagnosis), and all 3 went on to achieve CR following OEPA +/- COPP. CVbP was well tolerated, with no grade 3 or 4 hematologic toxicities reported, and hair loss and mucositis were rare.

NCCN Recommendations for Stage IA NLP HL with Complete Resection Confirmed by FDG-PET/CT

For all stages of NLP HL, the following risk factors, in addition to stage, may inform decisions about primary treatment: age ≥45 years, hemoglobin <10.5 g/dL, and splenic involvement.¹⁰⁶

For patients with stage IA NLP HL with complete resection confirmed by FDG-PET/CT, the Panel recommends enrollment in an ongoing clinical trial or observation as the preferred strategies. Although AHOD0P31 only included completely resected single node disease for observation, additional retrospective data supports observation after complete resection of stage I, multiple nodes^{101-103,107}; however, extensive node or repeat node dissection is discouraged.

Observation should consist of a clinical exam with or without US of the primary site every 3 to 4 months for the first 2 years. Additional imaging should be pursued as indicated for concern for relapsed disease. Biopsy should also be performed if there is concern for relapse. If biopsy is negative, observation can continue. If biopsy confirms relapse, patients should undergo restaging, with treatment decisions to follow based on stage.

Chemotherapy with CVbP for 3 cycles is another recommended option. For patients >18 years, additional options can be considered as per the NCCN Guidelines for Hodgkin Lymphoma (Adult) (available at www.NCCN.org).

Stage IA or IIA NLP HL with Incomplete Resection and Non-Bulky

Combination chemotherapy regimens used in the treatment of CHL have shown excellent outcomes in children with NLP HL; however, these treatment strategies carry long-term risks associated with chemotherapy and RT; thus, contemporary treatment approaches aim to limit chemotherapy cycles with or without RT.^{101,103}

The COG AHOD03P1 study, as discussed above, not only aimed to evaluate the strategy of surgical resection alone for patients with early-stage NLP HL, but also evaluated the use of AVPC (doxorubicin/vincristine/prednisone/cyclophosphamide) chemotherapy.¹⁰¹ A total of 135 patients with either unresected stage IA or stage IIA disease



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

or with recurrence following complete resection received 3 cycles of AVPC. CR was obtained in 92% of patients. The 8% of patients who experienced less than a CR went on to receive IFRT. Five-year EFS was estimated at 88.8%, with 12 relapses and the development of secondary cancer in 3 patients. AVPC was overall well tolerated, with grade 3 or 4 fever and neutropenia in 5.1% of patients.

CVbP was found to be a safe and effective first-line chemotherapy regimen for children and adolescents with stage IA or IIA NLPHL by Shankar et al, as previously discussed in *Stage IA NLPHL with Complete Resection Confirmed by FDG-PET/CT*.¹⁰⁵ While data for the use of the anti-CD20 antibody rituximab for early-stage NLPHL are limited, it has been safely added to chemotherapy regimens in many subtypes of lymphoma and has yielded favorable PFS and response rates in adult NLPHL when combined with ABVD and CHOP. Shankar et al thus support the addition of rituximab to low-intensity CVbP and postulate that it would improve PFS while adding minimal toxicity.¹⁰³

NCCN Recommendations for Stage IA or IIA NLPHL with Incomplete Resection and Non-Bulky

For patients with stage IA or IIA NLPHL with incomplete resection and non-bulky disease, the Panel recommends enrollment in an ongoing clinical trial, AVPC x 3 cycles (category 1 recommendation), or CVbP with or without rituximab as preferred strategies. An FDA-approved biosimilar is an acceptable substitute for rituximab. With the AVPC approach, if an adequate response is not achieved following 3 cycles, ISRT is recommended. With the CVbP approach, if an adequate response is not achieved following 3 cycles of CVbP with or without rituximab, the Panel recommends consideration of ISRT or a referral to or consultation with a center of expertise for consideration of additional therapy.

Another recommended option is OEPA x 2 cycles as studied in children and adolescents with CHL in the EuroNet-PHL-C1 study.⁶⁰ For those who

experience inadequate response to 2 cycles of OEPA, the Panel recommends consideration of ISRT or a referral to or consultation with a center of expertise for consideration of additional therapy. For patients >18 years, additional options can be considered as per the NCCN Guidelines for Hodgkin Lymphoma (Adult) (available at www.NCCN.org).

Stage IA–IIA with Bulk, IB–IIB, III, IV NLPHL

Intermediate- and high-risk NLPHL is rare in pediatric patients. Although it can occur, particularly in cases with variant morphology, this is uncommon. Therefore, expert hematopathology re-evaluation of original pathology slides is suggested to confirm pathologic diagnosis. Misdiagnosed CHL or THRLBCL are diagnostic considerations in this setting and should be excluded.

In a subsequent report from the AHOD0031 study, the investigators evaluated the outcomes of a subgroup of patients in the study who had lymphocyte-predominant HL (LPHL) (n = 96 of 1711), and found that compared to CHL, patients with LPHL were more likely to achieve RER (93.6% vs 81.0%; $P = .002$) and CR (74.2% vs 49.3%; $P = .000005$) after chemotherapy.¹⁰⁸ In addition, the 5-year EFS was higher in the LPHL subgroup compared to CHL (92.2% vs 83.5%, respectively; $P = .04$).¹⁰⁸ Based on the data, the investigators state that this subgroup may benefit from treatment with chemotherapy alone.

There are limited data on the treatment of intermediate-/high-risk NLPHL. It is commonly treated similar to CHL. Given its rarity, consideration of referral to or consultation with a center of expertise is recommended.

Relapsed or Refractory NLPHL

Histologic confirmation with biopsy is recommended before initiating treatment for relapsed or refractory disease given the risk for transformation and high false-positive rate of FDG-PET/CT. As previously noted, false-positive findings may be observed on FDG-PET/CT after



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

treatment with a regimen that includes a CPI. Repeat imaging at short intervals (per LYRIC recommendations) may be considered prior to obtaining a biopsy.³⁴ In the case of NLPHL, residual FDG-PET-avid disease can represent foci of concurrent or transformed diffuse large B-cell lymphoma (DLBCL) or reactive lymph node with progressive transformation of germinal centers.^{109,110}

Select Principles of Supportive Care

Although variations exist between institutional protocols, the following supportive care issues are important to consider in the treatment of pediatric patients with HL.

Immunotherapy

Prior to initiating CPI containing regimens, patients and caregivers should receive counseling on the wide-ranging warning signs and symptoms of immune-related adverse events (irAEs). A thorough clinical assessment is also recommended, including physical examination alongside patient and relevant family history of autoimmune or organ-specific disease, endocrinopathy, or infectious disease.

A clinical exam with irAE symptom assessment should be performed at each visit, assessing for any new signs or symptoms, including severe fatigue, headache, rash, cough, shortness of breath, chest pain, abdominal bloating, change in bowel pattern, weight loss, vision changes or eye pain, severe muscle weakness, severe muscle or joint pains, and/or mood changes.

Blood work including a CBC, CMP, thyroid-stimulating hormone (TSH), free thyroxine (T4), and serum cortisol (morning preferred), is recommended prior to initiating a CPI-based regimen as well as at regular intervals during therapy. For further details on blood work recommendations, see *Select Principles of Supportive Care* in the algorithm. For other pre-assessment, monitoring, and irAE management

recommendations, see NCCN Guidelines for Management of Immune Checkpoint Inhibitor-Related Toxicities (available at www.NCCN.org).

Peripheral Neuropathy

Several neurotoxic chemotherapy agents are utilized in the treatment of pediatric patients with HL, including Bv, vincristine, and vinblastine, thus providers need to be aware of grading and/or screening methods as well as appropriate management.

Grading and/or screening methods that can be utilized for peripheral neuropathy include the Common Terminology Criteria for Adverse Events (CTCAE) v5.0 or Balis Score, patient reported outcomes, or the Pediatric-modified Total Neuropathy Score (peds-mTNS) and/or the Total Neuropathy Score-pediatric version.^{111,112}

For patients who develop peripheral neuropathy, reducing the dose of or holding the neurotoxic chemotherapy agent based on protocol recommendations may be considered.^{113,114} Early referral to physical therapy should also be considered.

To date, no pharmacologic intervention has shown benefit for peripheral neuropathy in the pediatric and adolescent cancer population. For patients ≥18 years of age, consideration can be made for duloxetine. See the NCCN Guidelines for Adult Cancer Pain (available at www.NCCN.org) for further information.

Risk of Cardiac Toxicity

A COG study examined long-term outcomes among children with newly diagnosed malignancies, including HL, who received dexrazoxane as part of clinical trial therapy.¹¹⁵ Patients who had been randomized to doxorubicin with or without dexrazoxane were assessed with a median follow-up period of 18.6 years. Dexrazoxane was not found to be associated with relapse (HR, 0.84), secondary cancers (HR, 1.19),



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

all-cause mortality (HR, 1.07), or cardiovascular mortality (HR, 1.45), and serious cardiovascular outcomes occurred less commonly when dexrazoxane was used (5.6% vs 17.6%; $P = .02$). In a retrospective study assessing outcomes among children with HL, acute lymphoblastic leukemia, or osteosarcoma who received doxorubicin with or without dexrazoxane, the use of dexrazoxane, particularly among those who received a cumulative doxorubicin dose of ≥ 250 mg/m², was associated with a reduced risk of reduction in lower left ventricular function or ejection fraction at 18.1 ± 2.7 years from cancer diagnosis.¹¹⁶

Based on this study, strong consideration should be made for administering dexrazoxane to children and adolescents expected to receive ≥ 250 mg/m² of cumulative doxorubicin, or an equivalent amount of an alternative anthracycline.¹¹⁷

For additional select supportive care recommendations, including management of Bv-related infusion reactions and growth factor recommendations, see *Select Principles of Supportive Care* in the algorithm. Additional growth factor recommendations can also be found in the NCCN Guidelines for Hematopoietic Growth Factors (available at www.NCCN.org).

Follow-Up After Completion of Treatment

Given the long-term risks of the therapies for HL, including secondary cancers, fatigue, pulmonary toxicity, thyroid dysfunction, and reproductive issues,¹¹⁸⁻¹²⁶ patients should be followed by an oncologist or survivorship specialist who is aware of these risks and complications, in coordination with the primary care provider, especially during the first 2 years after treatment. The follow-up schedule should be individualized, depending on clinical circumstances such as patient's age, sex, stage of disease, and initial treatment modality. It is recommended that the patient be provided with a treatment summary at the completion of therapy, including details of

RT and chemotherapy, including cumulative anthracycline and alkylator dosages given.

The Panel recommends an interim history and physical examination every 3 to 4 months for 1 to 2 years, then every 6 to 12 months until year 3, and then annually until 5 years. Recommended laboratory studies include: CBC with differential, ESR or CRP, and a chemistry profile as clinically indicated. If the patient's neck was treated with RT, TSH should be evaluated annually. If patients were exposed to regimens that contain bleomycin or pulmonary RT, or have significant pulmonary involvement or other clinical concerns, PFTs should be considered. At the end of therapy, an echocardiogram may be considered, with repeat echocardiograms thereafter based on specific risk profile (eg, Children's Oncology Group Long-Term Follow-up Guidelines).

An annual influenza vaccination and other vaccines per the Centers for Disease Control and Prevention (CDC) is recommended for all patients (see the COG Survivorship Guidelines¹²⁷ for more details).

Screening for delayed immune-mediate adverse events (IMAE), such as hypothyroidism, which may present within the first 2 years following completion of treatment, may be needed for those who received a CPI. In a retrospective study among 795 patients treated with CPIs, 10.8% of IMAEs occurred >12 months after initiation of the CPI, with kidney and hematologic IMAEs the most frequent late-onset IMAEs.¹²⁸

Due to the risk of false positives, routine or surveillance FDG-PET scans are discouraged. If relapse is suspected (based on imaging, clinical, and pathologic correlations) imaging studies are recommended. It is acceptable to obtain a posterior-anterior and lateral chest x-ray, CT scan with contrast, or MRI of original sites of disease, followed at 3- to 6-month intervals for up to 2 years following completion of therapy. Although an MRI scan may substitute CT scan for neck, abdomen, and pelvic regions,



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

a diagnostic CT of the chest is required to evaluate lung parenchyma. If the previous FDG-PET was positive (Deauville 3 to 5), an FDG-PET/CT or FDG-PET/MRI scan is recommended to confirm CR at the end of all prescribed therapy, including RT. The Panel notes that once negative, a repeat FDG-PET should not be done unless evaluating suspicious findings on the history and physical, CT, or MRI. In addition, to minimize false-positive results, it is important to wait at least 8 to 12 weeks after the end of RT to perform FDG-PET assessments.

In general, and in terms of monitoring late effects (≥ 2 years after completion of systemic therapy), patients should be encouraged to undergo counseling on issues regarding survivorship, long-term treatment effects (eg, secondary cancers [with special attention to thyroid and breast cancer], cardiac disease, and issues affecting the thyroid, bone, and fertility and reproductive health), health habits, and psychosocial issues. For comprehensive details, see the COG Survivorship Guidelines.¹²⁷

Summary

Pediatric HL is now curable in most patients because of the introduction of more effective and less toxic regimens. However, survivors may experience late treatment-related side effects. For this reason, long-term follow-up is essential after completion of treatment. In addition, improvements in harmonization of staging and response criteria, and risk stratification will improve the therapeutic index.¹⁰ Emerging data will continue to inform the Panel's recommendations and consistent with NCCN philosophy, participation in clinical trials is always encouraged.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

References

1. Allen CE, Kelly KM, Bollard CM. Pediatric lymphomas and histiocytic disorders of childhood. *Pediatr Clin North Am* 2015;62:139–165. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25435117>.
2. Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health Organization classification of haematolymphoid tumours: Lymphoid neoplasms. *Leukemia* 2022;36:1720–1748. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35732829>.
3. Fan Z, Natkunam Y, Bair E, et al. Characterization of variant patterns of nodular lymphocyte predominant Hodgkin lymphoma with immunohistologic and clinical correlation. *Am J Surg Pathol* 2003;27:1346–1356. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14508396>.
4. Jarrett RF. Risk factors for Hodgkin's lymphoma by EBV status and significance of detection of EBV genomes in serum of patients with EBV-associated Hodgkin's lymphoma. *Leuk Lymphoma* 2003;44 Suppl 3:S27–32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15202522>.
5. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin* 2020;70:7–30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31912902>.
6. Swerdlow SH, Campo E, Harris NL, et al. WHO classification of tumours of haematopoietic and lymphoid tissues. Revised 4th ed. IARC Press: Lyon 2017.
7. Siegel RL, Kratzer TB, Giaquinto AN, et al. Cancer statistics, 2025. *CA Cancer J Clin* 2025;75:10–45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39817679>.
8. Ward E, DeSantis C, Robbins A, et al. Childhood and adolescent cancer statistics, 2014. *CA Cancer J Clin* 2014;64:83–103. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24488779>.
9. Miller KD, Fidler-Benaoudia M, Keegan TH, et al. Cancer statistics for adolescents and young adults, 2020. *CA Cancer J Clin* 2020;70:443–459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32940362>.
10. Kelly KM. Hodgkin lymphoma in children and adolescents: improving the therapeutic index. *Hematology Am Soc Hematol Educ Program* 2015;2015:514–521. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26637767>.
11. Mauz-Korholz C, Metzger ML, Kelly KM, et al. Pediatric Hodgkin Lymphoma. *J Clin Oncol* 2015;33:2975–2985. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26304892>.
12. U.S. National Library of Medicine Key MEDLINE® Indicators. Available at: http://www.nlm.nih.gov/bsd/bsd_key.html. Accessed January 24, 2020.
13. Campo E, Jaffe ES, Cook JR, et al. The International Consensus Classification of mature lymphoid neoplasms: a report from the Clinical Advisory Committee. *Blood* 2022;140:1229–1253. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35653592>.
14. Huppmann AR, Nicolae A, Slack GW, et al. EBV may be expressed in the LP cells of nodular lymphocyte-predominant Hodgkin lymphoma (NLPHL) in both children and adults. *Am J Surg Pathol* 2014;38:316–324. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24525501>.
15. Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol* 2014;32:3059–3068. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113753>.
16. El-Galaly TC, d'Amore F, Mylam KJ, et al. Routine bone marrow biopsy has little or no therapeutic consequence for positron emission tomography/computed tomography-staged treatment-naïve patients with Hodgkin lymphoma. *J Clin Oncol* 2012;30:4508–4514. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23150698>.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

17. Salaun PY, Gastinne T, Bodet-Milin C, et al. Analysis of 18F-FDG PET diffuse bone marrow uptake and splenic uptake in staging of Hodgkin's lymphoma: a reflection of disease infiltration or just inflammation? *Eur J Nucl Med Mol Imaging* 2009;36:1813–1821. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19499219>.
18. Purz S, Mauz-Korholz C, Korholz D, et al. [18F]Fluorodeoxyglucose positron emission tomography for detection of bone marrow involvement in children and adolescents with Hodgkin's lymphoma. *J Clin Oncol* 2011;29:3523–3528. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21825262>.
19. Close A, Burns K, Bjornard K, et al. Fertility preservation in pediatric leukemia and lymphoma: A report from the Children's Oncology Group. *Pediatr Blood Cancer* 2023;70:e30407. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37194406>.
20. Carbone PP, Kaplan HS, Musshoff K, et al. Report of the Committee on Hodgkin's Disease Staging Classification. *Cancer Res* 1971;31:1860–1861. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/5121694>.
21. Lister TA, Crowther D, Sutcliffe SB, et al. Report of a committee convened to discuss the evaluation and staging of patients with Hodgkin's disease: Cotswolds meeting. *J Clin Oncol* 1989;7:1630–1636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2809679>.
22. Zijtregtop EAM, Zeal J, Metzger ML, et al. Significance of E-lesions in Hodgkin lymphoma and the creation of a new consensus definition: a report from SEARCH. *Blood Adv* 2023;7:6303–6319. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37522740>.
23. Stoevesandt D, Steglich J, Bartelt JM, et al. CT, MRI, and FDG PET/CT in the assessment of lymph node involvement in pediatric Hodgkin lymphoma: An expert consensus definition by an international collaboration on Staging Evaluation and Response Criteria Harmonization for Children, Adolescent, and Young Adult Hodgkin Lymphoma (SEARCH for CAYAHL). *Radiology* 2025;314:e232650. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39835977>.
24. Flerlage JE, Kelly KM, Beishuizen A, et al. Staging Evaluation and Response Criteria Harmonization (SEARCH) for Childhood, Adolescent and Young Adult Hodgkin Lymphoma (CAYAHL): Methodology statement. *Pediatr Blood Cancer* 2017;64. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28097818>.
25. Kluge R, Kurch L, Georgi T, Metzger M. Current Role of FDG-PET in Pediatric Hodgkin's Lymphoma. *Semin Nucl Med* 2017;47:242–257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28417854>.
26. McCarten KM, Nadel HR, Shulkin BL, Cho SY. Imaging for diagnosis, staging and response assessment of Hodgkin lymphoma and non-Hodgkin lymphoma. *Pediatr Radiol* 2019;49:1545–1564. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31620854>.
27. Voss SD, Cairo MS. Surveillance imaging in pediatric lymphoma. *Pediatr Radiol* 2019;49:1565–1573. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31620855>.
28. Barrington SF, Mikhaeel NG, Kostakoglu L, et al. Role of imaging in the staging and response assessment of lymphoma: consensus of the International Conference on Malignant Lymphomas Imaging Working Group. *J Clin Oncol* 2014;32:3048–3058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113771>.
29. Meignan M, Gallamini A, Haioun C, Polliack A. Report on the Second International Workshop on interim positron emission tomography in lymphoma held in Menton, France, 8-9 April 2010. *Leuk Lymphoma* 2010;51:2171–2180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21077737>.
30. Meignan M, Gallamini A, Itti E, et al. Report on the Third International Workshop on Interim Positron Emission Tomography in Lymphoma held in Menton, France, 26-27 September 2011 and Menton 2011 consensus. *Leuk Lymphoma* 2012;53:1876–1881. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22432519>.
31. Barrington SF, Qian W, Somer EJ, et al. Concordance between four European centres of PET reporting criteria designed for use in



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

multicentre trials in Hodgkin lymphoma. *Eur J Nucl Med Mol Imaging* 2010;37:1824–1833. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20505930>.

32. Barrington SF, Kluge R. FDG PET for therapy monitoring in Hodgkin and non-Hodgkin lymphomas. *Eur J Nucl Med Mol Imaging* 2017;44:97–110. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28411336>.

33. Schaefer NG, Taverna C, Strobel K, et al. Hodgkin disease: diagnostic value of FDG PET/CT after first-line therapy—is biopsy of FDG-avid lesions still needed? *Radiology* 2007;244:257–262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17581905>.

34. Cheson BD, Ansell S, Schwartz L, et al. Refinement of the Lugano Classification lymphoma response criteria in the era of immunomodulatory therapy. *Blood* 2016;128:2489–2496. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27574190>.

35. Petrausch U, Samaras P, Veit-Haibach P, et al. Hodgkin's lymphoma in remission after first-line therapy: which patients need FDG-PET/CT for follow-up? *Ann Oncol* 2010;21:1053–1057. Available at: <https://pubmed.ncbi.nlm.nih.gov/19914963/>.

36. Filippi AR, Ragona R, Piva C, et al. Optimized volumetric modulated arc therapy versus 3D-CRT for early stage mediastinal Hodgkin lymphoma without axillary involvement: a comparison of second cancers and heart disease risk. *Int J Radiat Oncol Biol Phys* 2015;92:161–168. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25863763>.

37. Hoppe BS, Flampouri S, Su Z, et al. Effective dose reduction to cardiac structures using protons compared with 3DCRT and IMRT in mediastinal Hodgkin lymphoma. *Int J Radiat Oncol Biol Phys* 2012;84:449–455. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22386373>.

38. Hoppe BS, Flampouri S, Zaiden R, et al. Involved-node proton therapy in combined modality therapy for Hodgkin lymphoma: results of a phase 2 study. *Int J Radiat Oncol Biol Phys* 2014;89:1053–1059. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24928256>.

39. Lagerlof I, Fohlin H, Enblad G, et al. Limited, but not eliminated, excess long-term morbidity in stage I-IIA Hodgkin lymphoma treated with doxorubicin, bleomycin, vinblastine, and dacarbazine and limited-field radiotherapy. *J Clin Oncol* 2022;40:1487–1496. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35077204>.

40. Lagerlof I, Holte H, Glimelius I, et al. No excess long-term mortality in stage I-IIA Hodgkin lymphoma patients treated with ABVD and limited field radiotherapy. *Br J Haematol* 2020;188:685–691. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31612478>.

41. Xiang M, Chang DT, Pollom EL. Second cancer risk after primary cancer treatment with three-dimensional conformal, intensity-modulated, or proton beam radiation therapy. *Cancer* 2020;126:3560–3568. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32426866>.

42. Hoppe BS, Mailhot-Vega RB, Renfro LA, et al. Comparison of contemporary radiation therapy approaches in combined modality treatment on pediatric high-risk classic Hodgkin lymphoma study: AHOD 1331. *Int J Radiat Oncol Biol Phys* 2025;123:972–979. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40588069>.

43. Lo AC, Milgrom SA, Vega RM, et al. Enhancing radiation therapy quality assurance in lymphoma: A rigorous real-time central review process in AHOD2131. *Int J Radiat Oncol Biol Phys* 2025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41453454>.

44. Hoskin PJ, Diez P, Williams M, et al. Recommendations for the use of radiotherapy in nodal lymphoma. *Clin Oncol (R Coll Radiol)* 2013;25:49–58. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22889569>.

45. Giulino-Roth L, Keller FG, Hodgson DC, Kelly KM. Current approaches in the management of low risk Hodgkin lymphoma in children and adolescents. *Br J Haematol* 2015;169:647–660. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25824371>.

46. Dorffel W, Luders H, Ruhl U, et al. Preliminary results of the multicenter trial GPOH-HD 95 for the treatment of Hodgkin's disease in



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

children and adolescents: analysis and outlook. *Klin Padiatr* 2003;215:139–145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12838937>.

47. Mauz-Korholz C, Hasenclever D, Dorffel W, et al. Procarbazine-free OEPA-COPDAC chemotherapy in boys and standard OPPA-COPP in girls have comparable effectiveness in pediatric Hodgkin's lymphoma: the GPOH-HD-2002 study. *J Clin Oncol* 2010;28:3680–3686. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20625128>.

48. Mauz-Korholz C, Landman-Parker J, Fernandez-Teijeiro A, et al. Response-adapted omission of radiotherapy in children and adolescents with early-stage classical Hodgkin lymphoma and an adequate response to vincristine, etoposide, prednisone, and doxorubicin (EuroNet-PHL-C1): a titration study. *Lancet Oncol* 2023;24:252–261. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36858722>.

49. Keller FG, Castellino SM, Chen L, et al. Results of the AHOD0431 trial of response adapted therapy and a salvage strategy for limited stage, classical Hodgkin lymphoma: A report from the Children's Oncology Group. *Cancer* 2018;124:3210–3219. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29738613>.

50. Hoppe BS, Milgrom SA, Renfro LA, et al. Transplant-free salvage therapy for low-risk relapsed pediatric Hodgkin lymphoma: A nonrandomized clinical trial. *JAMA Oncol* 2025;11:340–342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39745710>.

51. Bhethanabhotla S, Jain S, Kapoor G, et al. Outcome of pediatric advanced Hodgkin lymphoma treated with ABVD and predictors of inferior survival: a multicenter study of 186 patients. *Leuk Lymphoma* 2017;58:1617–1623. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27919174>.

52. Jain S, Kapoor G, Bajpai R. ABVD-Based therapy for Hodgkin lymphoma in children and adolescents: Lessons learnt in a tertiary care oncology center in a developing country. *Pediatr Blood Cancer* 2016;63:1024–1030. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26855007>.

53. Marr KC, Connors JM, Savage KJ, et al. ABVD chemotherapy with reduced radiation therapy rates in children, adolescents and young adults with all stages of Hodgkin lymphoma. *Ann Oncol* 2017;28:849–854. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28327925>.

54. Stieglitz E, Dinh T, Phelps AS, et al. ABVD without radiation for newly diagnosed pediatric and young adult patients with hodgkin lymphoma: A single center retrospective analysis of 28 consecutive patients. *J Pediatr Hematol Oncol* 2018;40:290–294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29432308>.

55. Zubizarreta PA, Alfaro E, Gutter M, et al. Children and adolescent Hodgkin lymphoma in Argentina: Long-term results after combined ABVD and restricted radiotherapy. *J Pediatr Hematol Oncol* 2017;39:602–608. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28902084>.

56. Friedman DL, Chen L, Wolden S, et al. Dose-intensive response-based chemotherapy and radiation therapy for children and adolescents with newly diagnosed intermediate-risk hodgkin lymphoma: a report from the Children's Oncology Group Study AHOD0031. *J Clin Oncol* 2014;32:3651–3658. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25311218>.

57. Castellino SM, Pei Q, Parsons SK, et al. Brentuximab vedotin with chemotherapy in pediatric high-risk Hodgkin's lymphoma. *N Engl J Med* 2022;387:1649–1660. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36322844>.

58. Metzger ML, Link MP, Billett AL, et al. Excellent outcome for pediatric patients with high-risk Hodgkin lymphoma treated with brentuximab vedotin and risk-adapted residual node radiation. *J Clin Oncol* 2021;39:2276–2283. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33826362>.

59. Landman-Parker J, Wallace H, Hasenclever D, et al. First International Inter-Group Study for Classical Hodgkin Lymphoma in Children and Adolescents: EuroNet-PHL-C1 European protocol Euronet PHL-C1; Report of the latest interim Analysis [Abstract# P064]. *Haematologica*; 10th International Symposium on Hodgkin Lymphoma



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Symposium (ISHL10) 2016;101:35. Available at:
<https://haematologica.org/article/view/7868>.

60. Mauz-Korholz C, Landman-Parker J, Balwierz W, et al. Response-adapted omission of radiotherapy and comparison of consolidation chemotherapy in children and adolescents with intermediate-stage and advanced-stage classical Hodgkin lymphoma (EuroNet-PHL-C1): a titration study with an open-label, embedded, multinational, non-inferiority, randomised controlled trial. *Lancet Oncol* 2022;23:125–137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34895479>.

61. Herrera AF, LeBlanc M, Castellino SM, et al. Nivolumab+AVD in advanced-stage classic Hodgkin's lymphoma. *N Engl J Med* 2024;391:1379–1389. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39413375>.

62. Castellino SM, Li H, Herrera AF, et al. Three-Year follow-up of nivolumab-AVD versus brentuximab vedotin-AVD in adolescents with advanced-stage classic Hodgkin lymphoma on S1826. *J Clin Oncol* 2026;44:449–454. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41512237>.

63. Straus DJ, Dlugosz-Danecka M, Connors JM, et al. Brentuximab vedotin with chemotherapy for stage III or IV classical Hodgkin lymphoma (ECHELON-1): 5-year update of an international, open-label, randomised, phase 3 trial. *Lancet Haematol* 2021;8:e410–e421. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34048680>.

64. Daw S, Wynn R, Wallace H. Management of relapsed and refractory classical Hodgkin lymphoma in children and adolescents. *Br J Haematol* 2011;152:249–260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21133886>.

65. Daw S, Hasenclever D, Mascarín M, et al. Risk and Response Adapted Treatment Guidelines for Managing First Relapsed and Refractory Classical Hodgkin Lymphoma in Children and Young People. Recommendations from the EuroNet Pediatric Hodgkin Lymphoma Group. *Hemasphere* 2020;4:e329. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32072145>.

66. Harker-Murray P, Mauz-Korholz C, Leblanc T, et al. Nivolumab and brentuximab vedotin with or without bendamustine for R/R Hodgkin lymphoma in children, adolescents, and young adults. *Blood* 2023;141:2075–2084. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36564047>.

67. Greve P, Beishuizen A, Hagleitner M, et al. Nivolumab plus brentuximab vedotin +/- bendamustine combination therapy: A safe and effective treatment in pediatric recurrent and refractory classical Hodgkin lymphoma. *Front Immunol* 2023;14:1229558. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37583696>.

68. Daw S, Cole PD, Hoppe BS, et al. Transplant-free approach in relapsed Hodgkin lymphoma in children, adolescents, and young adults: A nonrandomized clinical trial. *JAMA Oncol* 2025;11:249–257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39745739>.

69. O'Connor OA, Lue JK, Sawas A, et al. Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin's lymphoma: an international, multicentre, single-arm, phase 1-2 trial. *Lancet Oncol* 2018;19:257–266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29276022>.

70. Forlenza CJ, Gulati N, Mauguen A, et al. Combination brentuximab vedotin and bendamustine for pediatric patients with relapsed/refractory Hodgkin lymphoma. *Blood Adv* 2021;5:5519–5524. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34559223>.

71. Cole PD, McCarten KM, Pei Q, et al. Brentuximab vedotin with gemcitabine for paediatric and young adult patients with relapsed or refractory Hodgkin's lymphoma (AHOD1221): a Children's Oncology Group, multicentre single-arm, phase 1-2 trial. *Lancet Oncol* 2018;19:1229–1238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30122620>.

72. Armand P, Shipp MA, Ribrag V, et al. Programmed death-1 blockade with pembrolizumab in patients with classical Hodgkin lymphoma after brentuximab vedotin failure. *J Clin Oncol* 2016;36:3733–3739. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27354476>.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

73. Younes A, Santoro A, Shipp M, et al. Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial. *Lancet Oncol* 2016;17:1283–1294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27451390>.

74. Mei MG, Lee HJ, Palmer JM, et al. Response-adapted anti-PD-1-based salvage therapy for Hodgkin lymphoma with nivolumab alone or in combination with ICE. *Blood* 2022;139:3605–3616. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35316328>.

75. Chen R, Zinzani PL, Fanale MA, et al. Phase II study of the efficacy and safety of pembrolizumab for relapsed/refractory classic Hodgkin lymphoma. *J Clin Oncol* 2017;35:2125–2132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28441111>.

76. Geoerger B, Kang HJ, Yalon-Oren M, et al. Pembrolizumab in paediatric patients with advanced melanoma or a PD-L1-positive, advanced, relapsed, or refractory solid tumour or lymphoma (KEYNOTE-051): interim analysis of an open-label, single-arm, phase 1-2 trial. *Lancet Oncol* 2020;21:121–133. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31812554>.

77. Bryan LJ, Casulo C, Allen PB, et al. Pembrolizumab added to ifosfamide, carboplatin, and etoposide chemotherapy for relapsed or refractory classic Hodgkin lymphoma: A multi-institutional phase 2 investigator-initiated nonrandomized clinical trial. *JAMA Oncol* 2023;9:683–691. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36928527>.

78. Moskowitz AJ, Shah G, Schoder H, et al. Phase II trial of pembrolizumab plus gemcitabine, vinorelbine, and liposomal doxorubicin as second-line therapy for relapsed or refractory classical Hodgkin lymphoma. *J Clin Oncol* 2021;39:3109–3117. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34170745>.

79. Herrera AF, Moskowitz AJ, Bartlett NL, et al. Interim results of brentuximab vedotin in combination with nivolumab in patients with

relapsed or refractory Hodgkin lymphoma. *Blood* 2018;131:1183–1194. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29229594>.

80. Advani RH, Moskowitz AJ, Bartlett NL, et al. Brentuximab vedotin in combination with nivolumab in relapsed or refractory Hodgkin lymphoma: 3-year study results. *Blood* 2021;138:427–438. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33827139>.

81. Kuruvilla J, Ramchandren R, Santoro A, et al. KEYNOTE-204: Randomized, open-label, phase III study of pembrolizumab (pembro) versus brentuximab vedotin (BV) in relapsed or refractory classic Hodgkin lymphoma (R/R cHL). *J Clin Oncol* 2020;38:8005–8005. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2020.38.15_suppl.8005.

82. Armand P, Engert A, Younes A, et al. Nivolumab for relapsed/refractory classic Hodgkin lymphoma after failure of autologous hematopoietic cell transplantation: Extended follow-up of the multicohort single-arm phase II CheckMate 205 trial. *J Clin Oncol* 2018;36:1428–1439. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29584546>.

83. Cole PD, Schwartz CL, Drachtman RA, et al. Phase II study of weekly gemcitabine and vinorelbine for children with recurrent or refractory Hodgkin's disease: a children's oncology group report. *J Clin Oncol* 2009;27:1456–1461. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19224841>.

84. Mahdy A, Hamoda A, Zaher A, et al. Outcome and toxicity of ifosfamide, carboplatin, and etoposide versus gemcitabine and vinorelbine regimen for pediatric patients with relapsed or refractory Hodgkin's lymphoma. *Front Oncol* 2023;13:1153128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37441423>.

85. Jan MRU, Saeed H, Abubakar M, Wali RM. Response rates, long term outcomes and toxicity profile of gemcitabine and vinorelbine based outpatient chemotherapy regimen in primary progressive and relapsed childhood Hodgkin lymphoma. *Leuk Lymphoma* 2020;61:3369–3377. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32924702>.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

86. Moskowitz CH, Nimer SD, Zelenetz AD, et al. A 2-step comprehensive high-dose chemoradiotherapy second-line program for relapsed and refractory Hodgkin disease: analysis by intent to treat and development of a prognostic model. *Blood* 2001;97:616–623. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11157476>.

87. Trippett TM, Schwartz CL, Guillerman RP, et al. Ifosfamide and vinorelbine is an effective reinduction regimen in children with refractory/relapsed Hodgkin lymphoma, AHOD00P1: a children's oncology group report. *Pediatr Blood Cancer* 2015;62:60–64. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25308760>.

88. Horton TM, Drachtman RA, Chen L, et al. A phase 2 study of bortezomib in combination with ifosfamide/vinorelbine in paediatric patients and young adults with refractory/recurrent Hodgkin lymphoma: a Children's Oncology Group study. *Br J Haematol* 2015;170:118–122. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25833390>.

89. Krishnan A, Bhatia S, Slovak ML, et al. Predictors of therapy-related leukemia and myelodysplasia following autologous transplantation for lymphoma: an assessment of risk factors. *Blood* 2000;95:1588–1593. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10688812>.

90. Josting A, Rudolph C, Reiser M, et al. Time-intensified dexamethasone/cisplatin/cytarabine: an effective salvage therapy with low toxicity in patients with relapsed and refractory Hodgkin's disease. *Ann Oncol* 2002;13:1628–1635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12377653>.

91. Shankar A, Hayward J, Kirkwood A, et al. Treatment outcome in children and adolescents with relapsed Hodgkin lymphoma--results of the UK HD3 relapse treatment strategy. *Br J Haematol* 2014;165:534–544. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24754633>.

92. Baetz T, Belch A, Couban S, et al. Gemcitabine, dexamethasone and cisplatin is an active and non-toxic chemotherapy regimen in relapsed or refractory Hodgkin's disease: a phase II study by the National Cancer Institute of Canada Clinical Trials Group. *Ann Oncol* 2003;14:1762–1767. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14630682>.

93. Moskowitz CH, Walewski J, Nademanee A, et al. Five-year PFS from the AETHERA trial of brentuximab vedotin for Hodgkin lymphoma at high risk of progression or relapse. *Blood* 2018;132:2639–2642. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30266774>.

94. Forlenza CJ, Rosenzweig J, Mauguén A, et al. Brentuximab vedotin after autologous transplantation in pediatric patients with relapsed/refractory Hodgkin lymphoma. *Blood Adv* 2023;7:3225–3231. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36897253>.

95. Armand P, Chen YB, Redd RA, et al. PD-1 blockade with pembrolizumab for classical Hodgkin lymphoma after autologous stem cell transplantation. *Blood* 2019;134:22–29. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30952672>.

96. Kahn S, Flowers C, Xu Z, Esiashvili N. Does the addition of involved field radiotherapy to high-dose chemotherapy and stem cell transplantation improve outcomes for patients with relapsed/refractory Hodgkin lymphoma? *Int J Radiat Oncol Biol Phys* 2011;81:175–180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20732769>.

97. Wilke C, Cao Q, Dusenbery KE, et al. Role of Consolidative Radiation Therapy After Autologous Hematopoietic Cell Transplantation for the Treatment of Relapsed or Refractory Hodgkin Lymphoma. *Int J Radiat Oncol Biol Phys* 2017;99:94–102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28816170>.

98. Mailhot Vega RB, Harker-Murray PD, Forlenza CJ, et al. Radiation therapy use in refractory and relapsed adolescent and young adult Hodgkin lymphoma: A report from the Children's Oncology Group. *Int J Radiat Oncol Biol Phys* 2023;117:400–403. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37116589>.

99. Park JJ, Lin J, Huang E, Schaar DG. The impact of prior salvage treatment with immune checkpoint inhibitors on Hodgkin lymphoma patients undergoing autologous hematopoietic stem cell transplantation: A single-center experience. *Clin Lymphoma Myeloma Leuk* 2021;21:e726–e730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34158267>.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

100. Daw S, Claviez A, Kurch L, et al. Transplant and nontransplant salvage therapy in pediatric relapsed or refractory Hodgkin lymphoma: The EuroNet-PHL-R1 phase 3 nonrandomized clinical trial. *JAMA Oncol* 2025;11:258–267. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39745682>.

101. Appel BE, Chen L, Buxton AB, et al. Minimal treatment of low-risk, pediatric lymphocyte-predominant Hodgkin lymphoma: A report from the Children's Oncology Group. *J Clin Oncol* 2016;34:2372–2379. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27185849>.

102. Mauz-Korholz C, Gorde-Grosjean S, Hasenclever D, et al. Resection alone in 58 children with limited stage, lymphocyte-predominant Hodgkin lymphoma-experience from the European network group on pediatric Hodgkin lymphoma. *Cancer* 2007;110:179–185. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17526010>.

103. Shankar A, Hall GW, McKay P, et al. Management of children and adults with all stages of nodular lymphocyte predominant Hodgkin lymphoma - All StAGes: A consensus-based position paper from the Hodgkin lymphoma subgroup of the UK National Cancer Research Institute. *Br J Haematol* 2022;197:679–690. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35362554>.

104. Pellegrino B, Terrier-Lacombe MJ, Oberlin O, et al. Lymphocyte-predominant Hodgkin's lymphoma in children: therapeutic abstention after initial lymph node resection--a Study of the French Society of Pediatric Oncology. *J Clin Oncol* 2003;21:2948–2952. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12885814>.

105. Shankar A, Hall GW, Gorde-Grosjean S, et al. Treatment outcome after low intensity chemotherapy [CVP] in children and adolescents with early stage nodular lymphocyte predominant Hodgkin's lymphoma - an Anglo-French collaborative report. *Eur J Cancer* 2012;48:1700–1706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22093944>.

106. Binkley MS, Flerlage JE, Savage KJ, et al. International prognostic score for nodular lymphocyte-predominant Hodgkin lymphoma. *J Clin*

Oncol 2024;42:2271–2280. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38531001>.

107. Punnett A, Baxter NN, Hodgson D, et al. Treatment patterns and outcomes in adolescents and young adults with nodular lymphocyte-predominant Hodgkin lymphoma: an IMPACT cohort study. *Br J Haematol* 2023;201:1081–1087. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37015867>.

108. Marks LJ, Pei Q, Bush R, et al. Outcomes in intermediate-risk pediatric lymphocyte-predominant Hodgkin lymphoma: A report from the Children's Oncology Group. *Pediatr Blood Cancer* 2018;65:e27375. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30277639>.

109. Al-Mansour M, Connors JM, Gascoyne RD, et al. Transformation to aggressive lymphoma in nodular lymphocyte-predominant Hodgkin's lymphoma. *J Clin Oncol* 2010;28:793–799. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20048177>.

110. Shaikh F, Ngan BY, Alexander S, Grant R. Progressive transformation of germinal centers in children and adolescents: an intriguing cause of lymphadenopathy. *Pediatr Blood Cancer* 2013;60:26–30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22707034>.

111. Common Terminology Criteria for Adverse Events (CTCAE). 2017. Available at: https://ctep.cancer.gov/protocoldevelopment/electronic_applications/ctc.htm#ctc_50. Accessed February 1, 2024.

112. Smith EML, Kuisell C, Kanzawa-Lee GA, et al. Approaches to measure paediatric chemotherapy-induced peripheral neurotoxicity: a systematic review. *Lancet Haematol* 2020;7:e408–e417. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32359452>.

113. Loprinzi CL, Lacchetti C, Bleeker J, et al. Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: ASCO Guideline Update. *J Clin Oncol* 2020;38:3325–3348. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32663120>.



NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

114. Heneghan MB, Parsons SK, Keller FG, et al. Protocol-stipulated dose modification to manage chemotherapy-induced peripheral neuropathy in children, adolescents, and young adults with high-risk Hodgkin lymphoma. JCO Oncol Pract 2024;OP2400089. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39303168>.

115. Chow EJ, Aplenc R, Vrooman LM, et al. Late health outcomes after dexrazoxane treatment: A report from the Children's Oncology Group. Cancer 2022;128:788–796. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34644414>.

116. Chow EJ, Aggarwal S, Doody DR, et al. Dexrazoxane and long-term heart function in survivors of childhood cancer. J Clin Oncol 2023;41:2248–2257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36669148>.

117. de Baat EC, van Dalen EC, Mulder RL, et al. Primary cardioprotection with dexrazoxane in patients with childhood cancer who are expected to receive anthracyclines: recommendations from the International Late Effects of Childhood Cancer Guideline Harmonization Group. Lancet Child Adolesc Health 2022;6:885–894. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36174614>.

118. Amini A, Murphy B, Cost CR, et al. Cardiac Mortality in Children and Adolescents with Hodgkin's Lymphoma: A Surveillance, Epidemiology and End Results Analysis. J Adolesc Young Adult Oncol 2016;5:181–186. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26959398>.

119. Bhakta N, Liu Q, Yeo F, et al. Cumulative burden of cardiovascular morbidity in paediatric, adolescent, and young adult survivors of Hodgkin's lymphoma: an analysis from the St Jude Lifetime Cohort Study. Lancet Oncol 2016;17:1325–1334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27470081>.

120. Clausen CT, Hasle H, Holmqvist AS, et al. Hyperthyroidism as a late effect in childhood cancer survivors - an Adult Life after Childhood Cancer in Scandinavia (ALiCCS) study. Acta Oncol 2019;58:227–231. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30585514>.

121. Dorffel W, Riepenhausen M, Luders H, et al. Secondary Malignancies Following Treatment for Hodgkin's Lymphoma in Childhood and Adolescence. Dtsch Arztebl Int 2015;112:320–327, i. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26037468>.

122. Fernandez-Pineda I, Davidoff AM, Lu L, et al. Impact of ovarian transposition before pelvic irradiation on ovarian function among long-term survivors of childhood Hodgkin lymphoma: A report from the St. Jude Lifetime Cohort Study. Pediatr Blood Cancer 2018;65:e27232. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29750388>.

123. Inskip PD, Veiga LHS, Brenner AV, et al. Hypothyroidism after Radiation Therapy for Childhood Cancer: A Report from the Childhood Cancer Survivor Study. Radiat Res 2018;190:117–132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29763379>.

124. O'Brien MM, Donaldson SS, Balise RR, et al. Second malignant neoplasms in survivors of pediatric Hodgkin's lymphoma treated with low-dose radiation and chemotherapy. J Clin Oncol 2010;28:1232–1239. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20124178>.

125. Oeffinger KC, Hudson MM, Mertens AC, et al. Increasing rates of breast cancer and cardiac surveillance among high-risk survivors of childhood Hodgkin lymphoma following a mailed, one-page survivorship care plan. Pediatr Blood Cancer 2011;56:818–824. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21370417>.

126. van Dalen EC, Caron HN, Dickinson HO, Kremer LC. Cardioprotective interventions for cancer patients receiving anthracyclines. Cochrane Database Syst Rev 2011;2011:CD003917. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21678342>.

127. The Children's Oncology Group long-term follow-up guidelines for survivors of childhood, adolescent and young adult cancers. Available at: <https://childrensoncologygroup.org/index.php/survivorshipguidelines>. Accessed November 16, 2023.

128. Durbin SM, Zubiri L, Perlman K, et al. Late-onset immune-related adverse events after immune checkpoint inhibitor therapy. JAMA Netw



National
Comprehensive
Cancer
Network®

PLEASE NOTE that use of this NCCN Content is governed by the End-User License Agreement, and you MAY NOT distribute this Content or use it with any artificial intelligence model or tool.
Printed by Marcos Di Natale on 7/11/2026 4:46:18 PM. Copyright © 2026 National Comprehensive Cancer Network, Inc. All Rights Reserved.

NCCN Guidelines Version 1.2026

Pediatric Hodgkin Lymphoma

Open 2025;8:e252668. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/40146104>.