



National Comprehensive
Cancer Network®


NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

Version 2.2026 — March 17, 2026

NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.





NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

***Jeremy Bordeaux, MD, MPH/Chair** ☐
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute

***Paul Nghiem, MD, PhD/Vice Chair** ☐
Fred Hutchinson Cancer Center

Sumaira Z. Aasi, MD ☐
Stanford Cancer Institute

Murad Alam, MD, MBA, MSCI ☐ ☐ ☐
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Arya Amini, MD §
City of Hope National Medical Center

Diana Bolotin, MD, PhD ☐
The UChicago Medicine
Comprehensive Cancer Center

Jennifer Chang, MD ☐
UC San Diego Moores Cancer Center

Pei-Ling Chen, MD, PhD ≠
Moffitt Cancer Center

Carlo M. Contreras, MD ☐
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Jessica M. Donigan, MD ☐ ☐
Huntsman Cancer Institute
at the University of Utah

Daniel Eisen, MD ☐
UC Davis Comprehensive Cancer Center

Jeffrey M. Farma, MD ☐
Fox Chase Cancer Center

Kelly Harms, MD, PhD ☐
University of Michigan Rogel Cancer Center

Nicole LeBoeuf, MD, MPH ☐
Dana-Farber/Brigham and Women's Cancer
Center | Mass General Cancer Center

John Nicholas Lukens, MD §
Abramson Cancer Center
at the University of Pennsylvania

Susan Manber ≠
Publicis Health

Lawrence Mark, MD, PhD ☐
Indiana University Melvin and Bren Simon
Comprehensive Cancer Center

Stacy McMurray, MD ☐
Vanderbilt-Ingram Cancer Center

Theresa Medina, MD †
University of Colorado Cancer Center

Kishwer S. Nehal, MD ☐
Memorial Sloan Kettering Cancer Center

Kelly Olino, MD ☐
Yale Cancer Center/Smilow Cancer Hospital

Gyorgy Paragh, MD, PhD, FAAD ☐
Roswell Park Comprehensive Cancer Center

Soo Park, MD †
UC San Diego Moores Cancer Center

Tejesh Patel, MD ☐
The University of Tennessee Health Science Center

Jason Rich, MD ☐
Siteman Cancer Center at Barnes-Jewish Hospital
and Washington University School of Medicine

Ashok R. Shaha, MD ☐
Memorial Sloan Kettering Cancer Center

Bhavina Sharma, MD, MPH ≠
Fred & Pamela Buffett Cancer Center

Olayemi Sokumbi, MD ☐ ≠
Mayo Clinic Comprehensive Cancer Center

Divya Srivastava, MD ☐
UT Southwestern Simmons Comprehensive Cancer Center

Joel Sunshine, MD, PhD ☐ ≠
Johns Hopkins Kimmel Cancer Center

Valencia Thomas, MD ☐ ≠ ☐
The University of Texas MD Anderson Cancer Center

Puja Venkat, MD §
UCLA Jonsson Comprehensive Cancer Center

Melodi Whitley, MD, PhD ☐
Duke Cancer Institute

Yaohui Gloria Xu, MD, PhD ☐ ☐
University of Wisconsin
Carbone Cancer Center

Siegrid Yu, MD ☐
UCSF Helen Diller Family Comprehensive Cancer Center

***Mehran Yusuf, MD** §
O'Neal Comprehensive Cancer Center at UAB

NCCN
Sara Espinosa, PhD
Beth McCullough, RN, BS

☐ Dermatology	≠ Pathology/ Dermatopathology
☐ Diagnostic/Interventional radiology	≠ Patient advocacy
≠ Hematology/Hematology oncology	§ Radiotherapy/Radiation oncology
Ⓟ Internal medicine	☐ Reconstructive surgery
† Medical oncology	☐ Surgery/Surgical oncology
☐ Otolaryngology	* Discussion Section Writing Committee

[NCCN Guidelines Panel Disclosures](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

[NCCN Squamous Cell Skin Cancer Panel Members](#)
[NCCN Evidence Blocks Definitions \(EB-DEF\)](#)

[Clinical Presentation, Workup, Diagnosis, and Risk Status \(SCC-1\)](#)
[Stratification to Determine Treatment Options for Local CSCC Based on Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#)
[Treatment for Field Cancerization/Confluent Epidermal Dysplasia \(SCC-3\)](#)
[Treatment for Low-Risk Cutaneous Squamous Cell Cancer \(CSCC\) \(SCC-4\)](#)
[Treatment for High-Risk/Very-High-Risk CSCC Where Surgery or RT Has High Likelihood of Cure \(SCC-5\)](#)
[Treatment for Very-High-Risk CSCC with Significant Risk of Extensive Local Recurrence or Nodal Metastasis \(SCC-6\)](#)
[Treatment for Locally Advanced \(laCSCC\) \(SCC-7\)](#)
[Clinical Staging, Preoperative Assessment, and Primary Treatment \(SCC-8\)](#)
[Treatment of Satellitosis/In-Transit Metastasis \(S-ITM\) \(SCC-9\)](#)
[Treatment for Regional Lymph Nodes \(SCC-10\)](#)
[Follow-up \(SCC-11\)](#)

[Principles of Pathology \(SCC-A\)](#)
[Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#)
[Principles of Treatment \(SCC-C\)](#)
[Principles of Radiation Therapy \(SCC-D\)](#)
[Principles of Systemic Therapy \(SCC-E\)](#)
[Principles of PDEMA Technique \(SCC-F\)](#)
[Principles of Cancer Risk Assessment and Counseling \(SCC-G\)](#)

[Evidence Blocks](#)

[Staging \(ST-1\)](#)

[Abbreviations \(ABBR-1\)](#)

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

NCCN Guidelines for Patients®
available at www.nccn.org/patients

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS

5					
4					
3					
2					
1					

E S Q C A

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

Example Evidence Block

5					
4					
3					
2					
1					

E S Q C A

E = 4
S = 4
Q = 3
C = 4
A = 3

Efficacy of Regimen/Agent

5	Highly effective: Cure likely and often provides long-term survival advantage
4	Very effective: Cure unlikely but sometimes provides long-term survival advantage
3	Moderately effective: Modest impact on survival, but often provides control of disease
2	Minimally effective: No, or unknown impact on survival, but sometimes provides control of disease
1	Palliative: Provides symptomatic benefit only

Safety of Regimen/Agent

5	Usually no meaningful toxicity: Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
4	Occasionally toxic: Rare significant toxicities or low-grade toxicities only; little interference with ADLs
3	Mildly toxic: Mild toxicity that interferes with ADLs
2	Moderately toxic: Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
1	Highly toxic: Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

Note: For significant chronic or long-term toxicities, score decreased by 1

Quality of Evidence

5	High quality: Multiple well-designed randomized trials and/or meta-analyses
4	Good quality: One or more well-designed randomized trials
3	Average quality: Low quality randomized trial(s) or well-designed non-randomized trial(s)
2	Low quality: Case reports or extensive clinical experience
1	Poor quality: Little or no evidence

Consistency of Evidence

5	Highly consistent: Multiple trials with similar outcomes
4	Mainly consistent: Multiple trials with some variability in outcome
3	May be consistent: Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
2	Inconsistent: Meaningful differences in direction of outcome between quality trials
1	Anecdotal evidence only: Evidence in humans based upon anecdotal experience

Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)


5	Very inexpensive
4	Inexpensive
3	Moderately expensive
2	Expensive
1	Very expensive



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™



^a [Principles of Pathology \(SCC-A\)](#).

^b [Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#) and [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).

^c Including CSCC in situ (showing full-thickness epidermal atypia).

^d Extensive disease includes deep involvement such as bone, named nerves, and deep soft tissue. If disease of named nerve(s) is suspected, MRI with and without contrast is preferred. If bone disease is suspected, CT with contrast is preferred unless contraindicated.

^e For rare cases that present with distant metastatic disease at diagnosis, treat per distant metastases pathway on [SCC-11](#).

^f Imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is sufficient to diagnose local recurrence, but MRI with and without contrast can be considered to assess extent of local disease. For nodal or distant metastases, histologic analysis and/or other imaging modalities can be used for confirmation and to gauge extent of disease.

^g MRI with and without contrast of the brain may be considered to rule out subclinical cortical involvement in cases with bone invasion.

^h Field cancerization defined as ultraviolet (UV) induced confluent dysplasia clinically manifested as diffuse actinic keratoses and superficial (in situ) SCC. Willenbrink TJ, et al. J Am Acad Dermatol 2020;83:709-717.

ⁱ A cure is unlikely to result from surgery and/or RT or there are concerns of significant functional impairment. Multidisciplinary discussion and multimodality treatment (including neoadjuvant and adjuvant therapy) merits consideration.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

STRATIFICATION TO DETERMINE TREATMENT OPTIONS FOR LOCAL CSCC BASED ON RISK FACTORS FOR LOCAL RECURRENCE, METASTASES, OR DEATH

Risk Group ^j	Low Risk	High Risk	Very High Risk
Treatment options	SCC-4	SCC-5	SCC-5 and SCC-6
H&P			
Location/diameter (cm)	Trunk, extremities <2 cm	Trunk, extremities 2 cm – ≤4 cm Head, neck, hands, feet, pretibia, and anogenital area (any size) ⁿ	>4 cm (any location)
Clinical borders	Well-defined	Poorly-defined	
Primary vs. recurrent	Primary	Recurrent	
Immunosuppression	(-)	(+)	
Site of prior RT or chronic inflammation	(-)	(+)	
Rapid growth tumor	(-)	(+)	
Neurologic symptoms	(-)	(+)	
Pathology (SCC-A)			
Degree of differentiation	Well or moderately differentiated		Poorly differentiated
Histologic subtype ^k	(-)	(+)	(+)
Depth ^{l,m} : Thickness or level of invasion	<2 mm thick and no invasion beyond subcutaneous fat	2–6 mm depth and no invasion beyond subcutaneous fat	>6 mm or invasion beyond subcutaneous fat
Perineural involvement	(-)	(+)	Tumor cells within the nerve sheath of a nerve lying deeper than the dermis or measuring ≥0.1 mm
Lymphatic or vascular involvement	(-)	(-)	(+)

^j Risk category assignment should be based on the highest risk factor present. The high-risk group has elevated risk of local recurrence; the very-high-risk group has elevated risk of local recurrence and elevated risk of metastasis.

^k Adenosquamous or sarcomatoid subtypes in any portion of the tumor.

^l If clinical evaluation of incisional biopsy suggests that microstaging is inadequate, consider narrow margin excisional biopsy.

^m Deep invasion is defined as invasion beyond the subcutaneous fat OR >6 mm (as measured from the granular layer of adjacent normal epidermis to the base of the tumor, consistent with the AJCC Cancer Staging Manual, 8th Edition).

ⁿ Narrow excision margins due to anatomic and functional constraints are associated with increased recurrence rates with standard histologic processing. Complete margin assessment with Mohs/PDEMA is recommended. For tumors <6 mm in size, without other high-risk or very-high-risk features, other treatment modalities may be considered if at least 4-mm clinically tumor-free margins can be obtained without significant anatomic or functional distortions.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

FIELD CANCERIZATION/CONFLUENT EPIDERMAL DYSPLASIA

Field cancerization^h/
Confluent epidermal
dysplasia

PRIMARY TREATMENT^{o,p,q,r}

- **Prevention:**
 - ▶ **Daily sunscreen^s**
 - ▶ **Nicotinamide^t**
- **Accepted treatment modalities**
 - ▶ **Topical:**
 - ◊ **Calcipotriene/Fluorouracil^u (preferred)**
 - ◊ **Fluorouracil (preferred)**
 - ◊ **Imiquimod**
 - ◊ **Tirbanibulin**
 - ◊ **Diclofenac (category 2B)**
 - ▶ **Destructive:**
 - ◊ **Ablative laser vermilionectomy or electrosurgical destruction (may be of value in the treatment of extensive actinic cheilitis)**
 - ◊ **Ablative skin resurfacing (eg, laser, dermabrasion)**
 - ◊ **Chemical peels (Trichloroacetic acid)**
 - ◊ **Cryotherapy^u**
 - ◊ **Curettage and electrodesiccation (C&E)**
 - ▶ **Other modalities that may be considered:**
 - ◊ **Capecitabine^{r,v} (for severe refractory disease that has progressed on oral retinoids)**
 - ◊ **Photodynamic therapy (PDT)^u (eg, topical Aminolevulinic acid [ALA], Porfimer sodium)**
 - ◊ **Systemic retinoids^w (eg, Acitretin, Isotretinoin)**

[See Evidence Blocks on EB-3](#)

^h Field cancerization defined as UV induced confluent dysplasia clinically manifested as diffuse actinic keratoses and superficial (in situ) SCC. Willenbrink TJ, et al. J Am Acad Dermatol 2020;83:709-717.

^o [Principles of Systemic Therapy](#).

^p Actinic keratoses that have an atypical clinical appearance or do not respond to appropriate therapy should be biopsied for histologic evaluation.

^q Actinic keratoses should be treated at first development.

^r Cornejo CM, et al. J Am Acad Dermatol 2020;83:719-730.

^s Green AC, et al. J Clin Oncol 2011;29:257-263.

^t Oral Nicotinamide may be effective in reducing the development of CSCCs. Chen AC, et al. N Engl J Med 2015;373:1618-1626; Allen NC, et al. N Engl J Med 2023;388:804-812; Mainville L, et al. J Cutan Med Surg 2022;26:297-308; Chen AC, et al. Br J Dermatol 2016;175:1073-1075.

^u The longest duration of prophylaxis against SCC has been demonstrated with topical Fluorouracil plus Calcipotriol. Cunningham TJ, et al. J Clin Invest 2017;127:106-116; Jansen MHE, et al. N Engl J Med 2019;380:935-946.

^v Endrizzi B, et al. Dermatol Surg 2013;39:634-645.

^w Use of oral retinoids (eg, Acitretin, Isotretinoin) is a therapeutic option used to reduce the development of actinic keratoses. Side effects of oral retinoids may be significant, especially in patients of childbearing potential, and therapeutic benefits are limited to the duration of the regimen. Topical retinoids were shown not to reduce development of actinic keratosis. Badri O, et al. Dermatol Surg 2021;47:125-126.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026


Squamous Cell Skin Cancer

NCCN Evidence Blocks™

LOW-RISK CSCC

PRIMARY TREATMENT^x

ADDITIONAL TREATMENT^x



^a [Principles of Pathology \(SCC-A\)](#).

^b [Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#) and [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).

^x [Principles of Treatment \(SCC-C\)](#).

^y C&E may have a lower cure rate than excision.

^z Shave removal (shaving of epidermal or dermal lesion) is a sharp removal by transverse bowl-shaped slicing to remove epidermal and dermal lesions (without including fat) and does not require suture closure. Emmett AJ, et al. Plast Reconstr Surg 1987;80:47-54.

^{aa} If tumor appears to extend beyond the dermis, surgical excision should generally be performed rather than C&E or shave removal.

^{bb} When Mohs is being performed and the preoperative biopsy is considered insufficient for providing all the staging information required to properly treat the tumor, submission of the central specimen for vertical paraffin-embedded permanent sections or documentation of staging parameters in Mohs report is recommended.

^{cc} As per other appropriate use Guidelines. Task Force/Committee Members, Vidal CI, Ambreth EA, et al. J Am Acad Dermatol 2019;80:189-207.e11.

^{dd} Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure.

^{ee} PDEMA with permanent section analysis or intraoperative frozen section analysis is an alternative to Mohs. See [Principles of PDEMA Technique \(SCC-F\)](#).

^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).

^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

HIGH-RISK/VERY-HIGH-RISK CSCC



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Footnotes on SCC-5A](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

FOOTNOTES

^a [Principles of Pathology \(SCC-A\)](#).

^b [Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#) and [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).

^x [Principles of Treatment \(SCC-C\)](#).

^{bb} When Mohs is being performed and the preoperative biopsy is considered insufficient for providing all the staging information required to properly treat the tumor, submission of the central specimen for vertical paraffin-embedded permanent sections or documentation of staging parameters in Mohs report is recommended.

^{cc} As per other appropriate use Guidelines. Task Force/Committee Members, Vidal CI, Armbricht EA, et al. J Am Acad Dermatol 2019;80:189-207.e11.

^{dd} Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure.

^{ee} PDEMA with permanent section analysis or intraoperative frozen section analysis is an alternative to Mohs. See [Principles of PDEMA Technique \(SCC-F\)](#).

^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).

^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.

^{hh} Discuss and consider SLNB prior to or at time of PDEMA for patients with very-high-risk CSCCs that are recurrent or have multiple risk factors placing them in the very-high-risk group and have normal exam of draining nodal basin (category 2B).

ⁱⁱ For positive SLNB: Recommend multidisciplinary discussion after obtaining radiologic staging of the neck, chest, abdomen, and pelvis if not yet completed. In the absence of metastatic disease consider completion lymphadenectomy of the affected nodal basin. If surgery is not an option due to patient preference or poor performance status, then consider radiation therapy. Following neck dissection, refer to [SCC-6](#) for additional recommendations.

^{jj} In patients with very-high-risk CSCC and normal exam of nodal basin, discuss and consider radiologic imaging of nodal basin.

^{kk} If invasion to parotid fascia, superficial parotidectomy may be indicated.

^{ll} For tumors being considered for SLNB, delay reconstruction if not able to close primarily with minimal undermining.

^{mm} Appropriate margins should be determined case by case based on tumor and patient-specific factors.

ⁿⁿ Large nerve involvement is defined by the American Joint Committee on Cancer (AJCC) Cancer Staging Manual, 8th Edition for CSCC of the head and neck as ≥ 0.1 mm or nerve involvement deeper than the dermis. Most nerves deep to the dermis are >0.1 mm.

^{oo} Adjuvant RT can be considered for CSCCs with gross clinical radiologic perineural invasion (PNI), multifocal histologic nerve invasion, ≥ 6 cm tumor diameter, recurrent tumors, high risk for regional or distant metastasis, close surgical margins where further surgery cannot be performed, and desmoplastic or infiltrative tumors in patients who are chronically immunosuppressed. Ruiz ES, et al. J Am Acad Dermatol 2022;87:87-94.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

VERY-HIGH-RISK CSCC



[See Evidence Blocks on EB-1](#)

^{pp} Cemiplimab-rwlc was studied in CSCC with very-high-risk for recurrence due to nodal features (extracapsular extension with largest node ≥20 mm in diameter or ≥3 involved nodes) or non-nodal features (in-transit metastases, T4 lesion [with bone invasion], PNI, or locally recurrent tumor with ≥1 additional risk feature). Rischin D, Porceddu S, Day F, et al. Adjuvant Cemiplimab or Placebo in High-Risk Cutaneous Squamous-Cell Carcinoma. N Engl J Med 2025;393:774-785.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Footnotes on SCC-6A](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

FOOTNOTES

- ^a [Principles of Pathology \(SCC-A\)](#).
- ^b [Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#) and [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).
- ^d Extensive disease includes deep involvement such as bone, named nerves, and deep soft tissue. If disease of named nerve(s) is suspected, MRI with and without contrast is preferred. If bone disease is suspected, CT with contrast is preferred unless contraindicated.
- ^f Imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is sufficient to diagnose local recurrence, but MRI with and without contrast can be considered to assess extent of local disease. For nodal or distant metastases, histologic analysis and/or other imaging modalities can be used for confirmation and to gauge extent of disease.
- ^g MRI with and without contrast of the brain may be considered to rule out subclinical cortical involvement in cases with bone invasion.
- ^o [Principles of Systemic Therapy](#).
- ^x [Principles of Treatment \(SCC-C\)](#).
- ^{bb} When Mohs is being performed and the preoperative biopsy is considered insufficient for providing all the staging information required to properly treat the tumor, submission of the central specimen for vertical paraffin-embedded permanent sections or documentation of staging parameters in Mohs report is recommended.
- ^{cc} As per other appropriate use Guidelines. Task Force/Committee Members, Vidal CI, Armbrrect EA, et al. J Am Acad Dermatol 2019;80:189-207.e11.
- ^{dd} Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure.
- ^{ee} PDEMA with permanent section analysis or intraoperative frozen section analysis is an alternative to Mohs. See [Principles of PDEMA Technique \(SCC-F\)](#).
- ^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).
- ^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.
- ^{hh} Discuss and consider SLNB prior to or at time of PDEMA for patients with very-high-risk CSCCs that are recurrent or have multiple risk factors placing them in the very-high-risk group and have normal exam of draining nodal basin (category 2B).
- ⁱⁱ For positive SLNB: Recommend multidisciplinary discussion after obtaining radiologic staging of the neck, chest, abdomen, and pelvis if not yet completed. In the absence of metastatic disease, consider completion lymphadenectomy of the affected nodal basin. If surgery is not an option due to patient preference or poor performance status, then consider radiation therapy. Following neck dissection, refer to [SCC-6](#) for additional recommendations.
- ^{jj} In patients with very-high-risk CSCC and normal exam of nodal basin, discuss and consider radiologic imaging of nodal basin.
- ^{kk} If invasion to parotid fascia, superficial parotidectomy may be indicated.
- ^{ll} For tumors being considered for SLNB, delay reconstruction if not able to close primarily with minimal undermining.
- ^{mm} Appropriate margins should be determined case by case based on tumor and patient-specific factors.
- ⁿⁿ Large nerve involvement is defined by the AJCC Cancer Staging Manual, 8th Edition for CSCC of the head and neck as ≥ 0.1 mm or nerve involvement deeper than the dermis. Most nerves deep to the dermis are >0.1 mm.
- ^{oo} Adjuvant RT can be considered for CSCCs with gross clinical radiologic PNI, multifocal histologic nerve invasion, ≥ 6 cm tumor diameter, recurrent tumors, high risk for regional or distant metastasis, close surgical margins where further surgery cannot be performed, and desmoplastic or infiltrative tumors in patients who are chronically immunosuppressed. Ruiz ES, et al. J Am Acad Dermatol 2022;87:87-94.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026


Squamous Cell Skin Cancer

NCCN Evidence Blocks™

LOCALLY ADVANCED CSCC

PRIMARY TREATMENT^x

ADDITIONAL TREATMENT^x



[See Evidence Blocks on EB-5](#)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Footnotes on SCC-7A](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

FOOTNOTES

- ^a [Principles of Pathology \(SCC-A\)](#).
- ^b [Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#) and [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).
- ^c [Principles of Systemic Therapy](#).
- ^x [Principles of Treatment \(SCC-C\)](#).
- ^{bb} When Mohs is being performed and the preoperative biopsy is considered insufficient for providing all the staging information required to properly treat the tumor, submission of the central specimen for vertical paraffin-embedded permanent sections or documentation of staging parameters in Mohs report is recommended.
- ^{cc} As per other appropriate use Guidelines. Task Force/Committee Members, Vidal CI, Armbrrect EA, et al. J Am Acad Dermatol 2019;80:189-207.e11.
- ^{dd} Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure.
- ^{ee} PDEMA with permanent section analysis or intraoperative frozen section analysis is an alternative to Mohs. See [Principles of PDEMA Technique \(SCC-F\)](#).
- ^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).
- ^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.
- ^{kk} If invasion to parotid fascia, superficial parotidectomy may be indicated.
- ^{ll} For tumors being considered for SLNB, delay reconstruction if not able to close primarily with minimal undermining.
- ^{mm} Appropriate margins should be determined case by case based on tumor and patient-specific factors.
- ⁿⁿ Large nerve involvement is defined by the AJCC Cancer Staging Manual, 8th Edition for CSCC of the head and neck as ≥ 0.1 mm or nerve involvement deeper than the dermis. Most nerves deep to the dermis are >0.1 mm.
- ^{oo} Adjuvant RT can be considered for CSCCs with gross clinical radiologic PNI, multifocal histologic nerve invasion, ≥ 6 cm tumor diameter, recurrent tumors, high risk for regional or distant metastasis, close surgical margins where further surgery cannot be performed, and desmoplastic or infiltrative tumors in patients who are chronically immunosuppressed. Ruiz ES, et al. J Am Acad Dermatol 2022;87:87-94.
- ^{qq} Bertrand N, et al. eClinicalMedicine 2021;5:100844.
- ^{rr} If patient is immunosuppressed, consider modification or reduction of immunosuppression as appropriate.
- ^{ss} Consider neoadjuvant therapy with Cemiplimab-rwlc, after multidisciplinary discussion, if the tumor has very rapid growth, in-transit metastasis, lymphovascular invasion, is borderline resectable, or surgery alone may not be curative or may result in significant functional limitation.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

CLINICAL STAGING AND PREOPERATIVE ASSESSMENT

PRIMARY TREATMENT^x



^g MRI with and without contrast of the brain may be considered to rule out subclinical cortical involvement in cases with bone invasion.

^o [Principles of Systemic Therapy](#).

^x [Principles of Treatment \(SCC-C\)](#).

^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).

^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.

^{rr} If the patient is immunosuppressed, consider modification or reduction of immunosuppression as appropriate.

^{tt} [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).

^{uu} Ultrasound-guided biopsy by a center or physician with expertise is recommended. Core biopsy may be preferred over FNA in cases where primary tumor histology is uncertain or if a larger tissue sample is required for further genetic or other testing.

^{vv} An excisional biopsy may be considered to confirm a negative initial FNA or core lymph node biopsy if clinical suspicion remains high.

^{ww} Regional lymph node dissection is preferred unless the patient is not a surgical candidate.

^{xx} Cemiplimab-rwlc, in one study of 79 patients with CSCC, showed a 51% complete histologic response in the neoadjuvant setting. Therefore it may be considered in patients who are considered borderline resectable, unresectable, or for whom surgery may carry a high morbidity.

^{yy} Consider palliative RT/surgery for symptomatic sites. Stereotactic body RT (SBRT) may also be considered in select patients.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026


Squamous Cell Skin Cancer

NCCN Evidence Blocks™

SATELLITOSIS/IN-TRANSIT METASTASIS (S-ITM)

TREATMENT PLANNING

PRIMARY TREATMENT^x



[See Evidence Blocks on EB-5](#)

^o [Principles of Systemic Therapy](#).

^x [Principles of Treatment \(SCC-C\)](#).

^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).

^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.

^{xx} Cemiplimab-rwlc, in one study of 79 patients with CSCC, showed a 51% complete histologic response in the neoadjuvant setting. Therefore it may be considered in patients who are considered borderline resectable, unresectable, or for whom surgery may carry a high morbidity.

^{zz} Porceddu SV, et al. Int J Clin Oncol 2018;36:1275-1283.

^{aaa} Marti-Marti I, et al. J Am Acad Dermatol 2023;89:119-127.

^{bbb} ILI/ILP is primarily used for patients with limb only disease with progression on/contraindications to standard therapies. This procedure should only be done at centers with experience with ILI/ILP. O'Donoghue C, et al. Ann Surg Oncol 2017;24:3842-3849; Huis In 't Veld EA, et al. Br J Cancer 2018;119:429-434.


**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.**



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™



^o [Principles of Systemic Therapy](#).

^x [Principles of Treatment \(SCC-C\)](#).

^{ff} [Principles of Radiation Therapy \(SCC-D\)](#).

^{gg} Determination of the appropriateness of RT should be performed together with a radiation oncologist.

^{ccc} Observation is not an option when one positive node ≤ 3 cm, no ENE.

^{ddd} Multidisciplinary consultation recommended.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



FOLLOW-UP

Local disease:

- **H&P^{eee,fff,ggg}**
 - For patients who are low risk:
Every 3–12 mo for 2 y, then every 6–12 mo for 3 y, then annually for life^b
 - For patients who are high risk:
Every 3–6 mo for 2 y, then every 6–12 mo for 3 y, then annually for life^b
 - For patients who are very high risk:
Every 3–6 mo for 2 y, then every 6 mo for 3 y, then every 6–12 mo for life^b
- **Consider imaging:**
 - If clinical exam is insufficient for following disease
 - If there is appreciable risk of subclinical local or nodal recurrence^f
- **Patient education**
 - Sun protection
 - Self examination of skin

Regional/S-ITM disease:


- **H&P^{eee,fff,ggg}**
 - Every 2–3 mo for 1 y,
then every 2–4 mo for 1 y,
then every 4–6 mo for 3 y,
then every 6–12 mo for life
- **Consider imaging:**
 - If clinical exam is insufficient for following disease
 - If there is appreciable risk of subclinical local or nodal recurrence^{f,hhh}
- **Patient education**
 - Sun protection
 - Self examination of skin
and lymph nodes

^b [Risk Factors for Local Recurrence, Metastases, or Death \(SCC-2\)](#) and [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \(SCC-B\)](#).

^f Imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is sufficient to diagnose local recurrence, but MRI with and without contrast can be considered to assess extent of local disease. For nodal or distant metastases, histologic analysis and/or other imaging modalities can be used for confirmation and to gauge extent of disease.

^o [Principles of Systemic Therapy](#).

^{yy} Consider palliative RT/surgery for symptomatic sites. SBRT may also be considered in select patients.



^{eee} Including complete skin and regional lymph node exam.

^{fff} Frequency of follow-up should be adjusted based on risk.

^{ggg} Follow-up with a dermatologist is strongly recommended if any of the following criteria are met: past or imminent solid organ, marrow, or stem cell transplant, one or more cutaneous melanomas in the past 5 years, or four or more non-melanoma skin cancers in the past 5 years.

^{hhh} Surveillance imaging of regional nodal basin and to evaluate for distant metastatic disease, ideally based on multidisciplinary board recommendation, or as clinically indicated.

ⁱⁱⁱ Under highly selective circumstances, in the context of multidisciplinary consultation, resection of limited metastases can be considered.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#). All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF PATHOLOGY

Principles of Biopsy Reporting

- Pathologic evaluation of skin biopsies is ideally performed by a dermatologist, pathologist, or dermatopathologist experienced in interpreting cutaneous neoplasms. Reporting of margins and the elements below is not required for biopsy specimens.

Principles of Excision Reporting (including Mohs excisions)

- Specimens from intended complete surgical removal (eg, shave excisions) should be labeled as such so that margin status is reported.
- Since depth of invasion (in mm) may not be ascertained on tangentially cut Mohs specimens, anatomic level of invasion should be reported. Frozen or permanent section analysis of the clinical tumor specimen may be undertaken if needed for complete reporting of the features below to enable AJCC tumor staging.^{1,2}
- Immunohistochemistry may be utilized as needed to help identify lymphovascular or nerve invasion, or to identify single tumor cells or small aggregates.

Recommended Elements for Pathology Reporting of Excisional Specimens (including Mohs excisions)

- NOTE: Tumors less than 2 cm in diameter without perineural invasion (as defined below) that are superficial (<6 mm in depth or confined to skin and fat) are AJCC T1 and do not require specific reporting of the histologic findings below with the exception of grade. However, reporting the presence of any of the prognostic features below is strongly encouraged.
- Elements reported (on requisition form) by the clinician submitting the tissue:
 - ▶ Anatomic location
 - ▶ Clinical pre-excision diameter in cm
 - ▶ Primary or recurrent tumor
 - ▶ Clinical or radiologic nerve invasion, including name of nerve
 - ▶ Other risk factors (optional) (eg, immunosuppression, prior radiation at site)
- Elements reported by the physician reporting the histologic findings:
 - ▶ Margin status (whether or not tumor is present at margins)
 - ▶ Well, moderate, or poor differentiation
 - ▶ Depth of invasion (either Breslow depth [in mm] measured from granular layer of adjacent normal epidermis to the base of the tumor OR tissue plane of deepest invasion (eg, dermis, fat, fascia, muscle, perichondrium/periosteum, cartilage bone, other)
 - ▶ Perineural invasion defined as tumor cells within the nerve sheath of a nerve deep to dermis or with a caliber 0.1 mm or larger
 - ▶ Lymphovascular invasion
 - ▶ High-risk histology (eg, adenosquamous, or sarcomatoid)
 - ▶ Low-risk histology (optional) (eg, verrucous)
 - ▶ Other useful reportable histologic features (eg, acantholytic, infiltrative, and spindle cell)

¹ Kim JYS, Kozlow JH, Mittal B, et al; Invited Reviewers; Work Group. Guidelines of care for the management of cutaneous squamous cell carcinoma. J Am Acad Dermatol 2018;78:560-578.

² Califano JA, Lydiatt WM, Nehal KS, et al. Cutaneous squamous cell carcinoma of the head and neck. In: Amin MB, Edge S, Greene F, et al, eds. AJCC Cancer Staging Manual (Eighth Edition). New York: Springer International Publishing; 2017:171-181.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

IDENTIFICATION AND MANAGEMENT OF PATIENTS AT HIGH RISK FOR MULTIPLE PRIMARY CSCCs

Definition

- Certain patient groups are at high risk for developing multiple CSCCs and tumors that can behave aggressively. These include:
 - ▶ Organ transplant recipients
 - ▶ Other settings of immunosuppression (eg, lymphoma, chronic lymphocytic leukemia [CLL], drug-induced, HIV)
 - ▶ Genetic syndromes predisposing to CSCC formation^a
- Within these high-risk groups, individual patients who are high risk should be identified for closer follow-up.
- Important individual risk factors include:
 - ▶ Total number of tumors
 - ▶ Frequency of development
 - ▶ Occurrence of aggressive tumors (eg, extension beyond cutaneous structures, perineural involvement, large and poorly differentiated, having ≥3 risk factors for recurrence) (See [Stratification to Determine Treatment Options for Local CSCC Based on Risk Factors for Local Recurrence, Metastases, or Death \[SCC-2\]](#)).

Diagnosis

- Skin lesions in these high-risk populations may be difficult to assess clinically. Therefore, a low threshold for performing skin biopsies of suspect lesions is necessary.
- In these patients, urgent diagnosis and treatment of lesions are important, and nodal staging (CT with contrast and/or ultrasound or pathologic evaluation) may be considered in those with significant risk of nodal metastases.

^a Examples include xeroderma pigmentosum, generalized eruptive keratoacanthoma of Grzybowski, Rothmund-Thomson syndrome, dyskeratosis congenita, epidermodysplasia verruciformis, recessive dystrophic epidermolysis bullosa, severe generalized junctional epidermolysis bullosa, KID syndrome (keratitis, ichthyosis, deafness), and Ferguson-Smith disease.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

IDENTIFICATION AND MANAGEMENT OF PATIENTS AT HIGH RISK FOR MULTIPLE PRIMARY CSCCs

Treatment of Skin Cancers

- Because patients in high-risk groups may develop multiple lesions in short periods of time, destructive therapy (eg, C&E, cryotherapy) may be a preferred treatment for clinically low-risk tumors because of the ability to treat multiple lesions at a single patient visit. If C&E has been performed based solely on the clinical appearance of a low-risk tumor, the pathology from the biopsy taken at the time of C&E should be reviewed to make sure there are no high-risk pathologic features that would suggest the need for further therapy beyond C&E.
- In patients who develop multiple adjacent tumors in close proximity, surgical excision of invasive disease sometimes does not include surrounding in situ disease, and tissue rearrangement should be minimized. In situ disease may then be treated with topical approaches similar to actinic keratoses/field cancerization.
- Compared to the low-risk population, RT is used more frequently as an adjuvant therapy in patients who are high risk and for perineural disease.
- Satellite lesions and in-transit cutaneous metastases may occur more frequently in this population. They must be treated aggressively with multidisciplinary consultation.
- In organ transplant recipients and other patients undergoing immunosuppressive therapy, decreasing the level of immunosuppressive therapy and/or incorporating mTOR inhibitors may be considered in cases of life-threatening skin cancer or the rapid development of multiple tumors.

Follow-Up

- Follow-up schedules should be titrated to the frequency of tumor development.

Patient Education

- Individual risk assessment is necessary and should be discussed.
- Both extensive and repetitive patient education regarding sun avoidance and protection is required.
- Sun avoidance and protection methods must be stringent.
- Monthly self examination of all skin surfaces is recommended. If a patient has a history of invasive skin cancer, self examination of the lymph nodes should be taught and performed.
- Rapid entrance into the health care delivery system at the onset of tumor development is critical.
- Patient education should begin, in the case of recipients of organ transplant, at transplantation, and in the case of xeroderma pigmentosum, at birth or diagnosis.

Prevention

- Regular sunscreen use prevents CSCC long term.¹
- Use of oral retinoids (eg, acitretin, isotretinoin) is effective in reducing the development of CSCC in some patients who are high risk. Side effects of oral retinoids may be significant. Therapeutic effects disappear shortly after cessation of the drug. Oral retinoids are teratogenic and must be used with extreme caution in patients of childbearing potential. Topical retinoids have been shown not to reduce development of CSCC. (See [SCC-3](#))
- Use of nicotinamide may be effective in reducing the development of CSCCs. Therapeutic effects disappear shortly after cessation of the drug.
- Aggressive treatment of precancers can prevent the development of subsequent invasive tumors.

[See Evidence Blocks on EB-1](#)

¹ Green AC, Williams GM, Logan V, Strutton GM. Reduced melanoma after regular sunscreen use: randomized trial follow-up. J Clin Oncol 2011;29:257-263.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF TREATMENT

- The primary goals of treatment of CSCCs are the complete removal of the tumor and the maximal preservation of function and cosmesis. All treatment decisions should be customized to account for the particular factors present in the individual case and for the patient's preference.
- Surgical approaches often offer the most effective and efficient means for accomplishing cure, but considerations of function, cosmesis, and patient preference may lead to choosing RT/topical therapy/systemic therapy as primary treatment in order to achieve optimal overall results.
- In certain patients at high risk for multiple primary tumors, increased surveillance and consideration of prophylactic measures may be indicated. (See [Identification and Management of Patients at High Risk for Multiple Primary CSCCs \[SCC-B\]](#)).
- Epidermally limited CSCC may be treated with nonsurgical options. (See [SCC-3](#) and [Principles of Systemic Therapy SCC-E](#)).
- When Mohs with margin assessment is being performed and the preoperative biopsy is considered insufficient for providing all the staging information required to properly treat the tumor, submission of the central specimen for vertical paraffin-embedded permanent sections or documentation of staging parameters in Mohs report is recommended.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY

General Principles

- Refer to the [ASTRO Guideline on Definitive and Postoperative Radiation Therapy for Basal and Squamous Cell Cancers of the Skin](#)¹ for general indications and dose recommendations.
- Protracted fractionation is associated with improved cosmetic results and should be utilized for poorly vascularized or cartilaginous areas.
- For extensive perineural invasion (PNI) or involvement of named nerves (particularly in the head and neck region), consider including the course of the cranial nerve proximally.
- Perineural tumor spread (PNTS), defined as clinically or radiographically apparent macroscopic spread along nerves, is considered higher risk and warrants comprehensive coverage of the involved, and potentially interconnected, cranial nerve pathways.^{2,3}
- For patients with very-high-risk CSCC such as those with PNTS, consider referral to a high-volume center given potential for severe toxicity when irradiating cranial nerves adjacent to critical optic and neurologic structures.
- RT is contraindicated for genetic conditions predisposing to skin cancer (eg, basal cell nevus syndrome [Gorlin syndrome]) and relatively contraindicated for patients with connective tissue diseases (eg, scleroderma).
- Given higher complication rates, reirradiation should not be routinely utilized for recurrent disease within a prior radiation field.
- Consider elective nodal irradiation (ENI) in cases of multiple high-risk pathologic features ([SCC-2](#)), especially in patients who are immunocompromised.
- Recommendations for external beam RT (EBRT) apply to both photons and electrons.
 - ▶ Image-guided radiation therapy (IGRT) is considered best practice when treating with intensity-modulated radiation therapy (IMRT), proton beam radiotherapy, or 3D conformal radiation. The use of IGRT for other types of radiotherapy to treat skin cancer is considered unnecessary.^a
- Application of bolus over the primary tumor should be considered to ensure full dose to the surface. Use of surface dosimeters should be considered to verify delivered dose.
- Radiation treatments should be given by a practicing radiation oncologist with radiation physics support to meet established quality assurance and dosimetric constraints.

[Primary Tumor and Regional Disease RT Dosing Table on SCC-D \(2 of 4\)](#)

[Brachytherapy RT Dosing Table on SCC-D \(3 of 4\)](#)

^a See [Discussion](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References on SCC-D \(4 of 4\)](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY PRIMARY TUMOR, REGIONAL DISEASE, AND S-ITM

Primary Tumor	RT Dosing	Common examples
Adjuvant RT²		
• Conventional fractionation (1.8–2 Gy/fraction)	BED10 of 60–79 Gy	• 60–66 Gy in 2 Gy/fraction⁴ • 60–66 Gy in 1.8 Gy/fraction (reserved for areas with poor wound healing [eg, lower extremity])
• Hypofractionation	BED10 of 48–70 Gy	• 2.5 Gy x 20 fractions⁵ • 3 Gy x 15–17 fractions to 45–51 Gy • 6 Gy x 5 fractions (non-consecutive days)
Definitive EBRT		
• Conventional fractionation (1.8–2 Gy/fraction)	BED10 of 70–84 Gy	• 60 Gy (for small tumors in cosmetically sensitive areas) • 66–70 Gy (for locally advanced tumors involving bone or cartilage)
• Hypofractionation	BED10 of 48–72 Gy ^c	• Tumors <2 cm: 2.5 Gy x 20 fractions • Tumors ≥2 cm: 2.5–3 Gy x 20–22 fractions • 3 Gy x 17–18 fractions to 51–54 Gy⁶ • 4.4 Gy x 10 fractions, 4 fractions per week⁷ • 6–7 Gy x 5 fractions, 2 treatments per week^{d,7,8}
Regional Disease		
• Conventional fractionation (1.8–2 Gy/fraction) ▶ Macroscopic/gross residual lymph nodes ▶ Lymph node + dissected nodal basins ▶ PNTS^{b,3}	66–70 Gy 60 Gy 50–66 Gy	
• Undissected at-risk nodal basins	1.8–2 Gy x 25–30 fractions	
Satellitosis/In-Transit Metastasis (S-ITM)		
• Resected	50–60 Gy over 5 to 6 weeks	
• Unresected	60–70 Gy over 6 to 7 weeks	

- **Conventionally fractionated radiotherapy consists of five daily treatments per week.**
- **Hypofractionated radiotherapy consists of fewer treatments with larger fraction size.**

^b Track involved cranial nerve to base of skull and at-risk nerve interconnections.

^c Given lack of randomized data, the optimal dose/fractionation is not established, but generally, a goal BED10 of 48–72 Gy is recommended.

^d Generally reserved for patients who are older or unable to come for protracted course.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References on SCC-D \(4 of 4\)](#)

SCC-D
2 OF 4



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF RADIATION THERAPY BRACHYTHERAPY

General Principles-Brachytherapy

- Isotope-based brachytherapy can be an effective treatment for certain sites of disease, particularly on the head and neck.^a
- There are insufficient long-term efficacy and safety data to support the routine use of electronic surface brachytherapy.^a
- Consider longer courses of RT (ie, 16 fractions) for sensitive areas such as ears, nose, or near eyes.
- Consider a higher total dose for immunocompromised patients (ie, 3 Gy x 16 fractions).
- Consider a lower dose for low risk, small tumors when using a single channel applicator (ie, 3 Gy x 12 fractions).⁹
- Consider BID fractionation if performing interstitial brachytherapy.
- Consider shielding normal structures when possible (ie, eye shields, internal nose shields).
- Start adjuvant radiation within 4–8 weeks of surgery after incision has healed.
- Surface applicators should only be used for superficial tumors (<5 mm depth).
- If deeper than 5 mm, consider interstitial brachytherapy or EBRT techniques.
- Consider referral to a center with expertise with brachytherapy.

Primary Tumor	RT Dosing	Common Examples
Definitive or Adjuvant Skin Brachytherapy		
• Custom surface mold applicators: Prescription depth ≤5 mm	GOAL BED10 at least 60 Gy	• 3 Gy x 16 fractions (daily) • 5 Gy x 8 fractions (2 times per week [for non-cosmetically sensitive areas])
• Interstitial Brachytherapy: Prescription depth >5 mm		• 4.4 Gy x 10 fractions (BID [6 hours minimum between fractions])
• Shielded single channel surface applicators (Leipzig or Valencia): Prescription depth ≤3 mm		• 3 Gy x 12–16 fractions (daily)

^a See [Discussion](#).

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References on SCC-D \(4 of 4\)](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

REFERENCES

- ¹ Likhacheva A, Awan M, Barker CA, et al. Definitive and postoperative radiation therapy for basal and squamous cell cancers of the skin: Executive summary of an American Society for Radiation Oncology Clinical Practice Guideline. *Pract Radiat Oncol* 2020;10:8-20.
- ² Porceddu SV, Daniels C, Yom SS, et al. Head and Neck Cancer International Group (HNCIG) Consensus Guidelines for the Delivery of Postoperative Radiation Therapy in Complex Cutaneous Squamous Cell Carcinoma of the Head and Neck (cSCCHN). *Int J Radiat Oncol Biol Phys* 2020;107:641-651.
- ³ Bakst RL, Glastonbury CM, Parvathaneni U, et al. Perineural invasion and perineural tumor spread in head and neck cancer. *Int J Radiat Oncol Biol Phys* 2019;103:1109-1124.
- ⁴ McDowell L, Yom SS. Locally advanced non-melanomatous skin cancer: Contemporary radiotherapeutic management. *Oral Oncol* 2019;99:104443.
- ⁵ Hu M, Liu H, Bernard A, et al. Hypofractionated adjuvant radiotherapy in cutaneous squamous cell and basal cell carcinoma of the head and neck: 50(Gy) in 20 study. *Oral Oncol Rep* 2025;13:100732.
- ⁶ van Hezewijk M, Creutzberg CL, Putter H, et al. Efficacy of a hypofractionated schedule in electron beam radiotherapy for epithelial skin cancer: Analysis of 434 cases. *Radiother Oncol* 2010;95:245-249.
- ⁷ Veness M. Hypofractionated radiotherapy in older patients with non-melanoma skin cancer: Less is better. *Australas J Dermatol* 2018;59:124-127.
- ⁸ Dunder Y, Cannon RB, Hunt JP, et al. Radiotherapy regimens in patients with nonmelanoma head and neck skin cancers. *Int J Dermatol* 2018;57:441-448.
- ⁹ Gauden R, Pracy M, Avery AM, et al. HDR brachytherapy for superficial non-melanoma skin cancers. *J Med Imaging Radiat Oncol* 2013;57:212-217.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

Field Cancerization/Confluent Epidermal Dysplasia ([SCC-E 2 of 5](#))

Very-High-Risk Disease ([SCC-E 3 of 5](#))

Primary and Recurrent laCSCC Disease ([SCC-E 3 of 5](#))

S-ITM ([SCC-E 3 of 5](#))

New Regional Disease ([SCC-E 3 of 5](#))

Regional Recurrence or Distant Metastatic Disease ([SCC-E 3 of 5](#))

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

Field Cancerization/Confluent Epidermal Dysplasia (SCC-3)¹

- Actinic keratoses should be treated at first development.
 - ▶ Accepted treatment modalities (in addition to [SCC-3](#)) include topical Imiquimod and topical Tirbanibulin. For hyperkeratotic actinic keratoses, pretreatment with topical Tazarotene, curettage, or topical keratolytics (topical Urea, Lactic acid, and Salicylic acid) prior to above therapies and select therapies listed in Table 1 may be considered. Another modality that may be considered is topical Diclofenac (category 2B).
- Actinic keratoses that have an atypical clinical appearance or do not respond to appropriate therapy should be biopsied for histologic evaluation.
- Use of oral retinoids (eg, Acitretin, Isotretinoin) is a therapeutic option used to reduce the development of actinic keratoses. Side effects of oral retinoids may be significant, especially in patients of childbearing potential, and therapeutic benefits are limited to the duration of the regimen. Topical retinoids were shown not to reduce development of actinic keratosis.
- In patients with CSCC in situ (Bowen disease), therapies such as topical Fluorouracil, topical Imiquimod, and photodynamic therapy (eg, ALA, Porfimer sodium) may be considered.^a
- Vigorous cryotherapy² may be considered for discrete lesions (not field cancerization).
- Focal squamous cell carcinoma in situ arising within actinic keratosis is not appropriate for surgery and should be treated topically.

Table 1: Therapy Options for Field Cancerization/Confluent Epidermal Dysplasia

Preferred	Other Recommended	Useful in Certain Circumstances
• Topical Calcipotriene/Fluorouracil^{b,3,4} • Topical Fluorouracil	• Topical Aminolevulinic acid (in conjunction with PDT)	• Acitretin⁵ • Capecitabine^{1,6} (for severe refractory disease that has progressed on oral retinoids) • Isotretinoin • Porfimer sodium (in conjunction with PDT)

[See Evidence Blocks on EB-2 and EB-3](#)

^a Cure rates are approximately 10% lower than for surgical treatment modalities.^{7,8}

^b The longest duration of prophylaxis against SCC has been demonstrated with topical Fluorouracil plus Calcipotriol.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[References on SCC-E \(5 of 5\)](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF SYSTEMIC THERAPY

Very-High-Risk Disease (SCC-6)

- In patients with extremely high risk nodal and non-nodal features^c and postoperative negative margins, following adjuvant RT^{d,e,f} Cemiplimab-rwlc is category 1 preferred.
- For tumors with nonreactive non-keratoacanthomatous rapid growth, in-transit metastasis, are borderline resectable, or where surgery alone may not be curative or may result in significant functional limitation, neoadjuvant Cemiplimab-rwlc is an option.

Primary and Recurrent IaCSCC Disease^g (SCC-7)

- If surgery is not feasible, recommend RT, and multidisciplinary teams to discuss concurrent systemic therapy in select cases (Table 2).
- If curative surgery and curative RT^f are not feasible, recommend multidisciplinary consultation to discuss systemic therapy alone (Table 3).

S-ITM (SCC-9)

- For S-ITM, multidisciplinary team to discuss systemic therapy alone (Table 3) or in combination with local therapies such as RT or surgery (Table 2).

New Regional Disease (SCC-8 and SCC-10)

- For most cases of fully resected regional disease, adjuvant systemic therapy is not recommended, unless within a clinical trial.
- For resected high-risk regional disease of head and neck, RT ± systemic therapy (Table 2).
- For unresectable, inoperable, or incompletely resected disease, multidisciplinary consultation is needed to discuss:
 - ▶ RT ± systemic therapy (Table 2)
 - ▶ Systemic therapy alone if curative RT^f is not feasible (Table 3).

Regional Recurrence or Distant Metastatic Disease (SCC-11)

- For regional recurrence or distant metastases, multidisciplinary team to discuss systemic therapy alone (Table 3) or in combination with RT (Table 2).

Table 2: Systemic Therapy Options for Use with RT (or Surgery for Regional Recurrence Following Systemic Treatment)

Preferred	Other Recommended	Useful in Certain Circumstances
• Cisplatin^{h,9} • Clinical trial	• Carboplatin • Carboplatin/Paclitaxel^{h,13,14} • EGFR inhibitors (eg, Cetuximab)^{h,15}	• Cisplatin/Fluorouracil^{h,22}

Table 3: Options for Systemic Therapy Alone

Preferred	Other Recommended	Useful in Certain Circumstances
• Cemiplimab-rwlc^{i,j} (if curative RT^f or surgery is not feasible for locally advanced, recurrent, or metastatic disease)^{10,11} • Pembrolizumab^{i,j,k} (if curative RT^f or surgery is not feasible for locally advanced, recurrent, or metastatic disease)¹² • Clinical trial	• Cosibelimab-ipdl (if curative RT^f or surgery is not feasible for locally advanced [category 2B] or metastatic disease)¹⁶ • Nivolumab^l • If ineligible for or progressed on immune checkpoint inhibitors and clinical trials, consider: ▶ Carboplatin/Paclitaxel¹⁷ ▶ Carboplatin/Paclitaxel + Cetuximab¹⁸⁻²¹ ▶ EGFR inhibitors (eg, Cetuximab)^{h,15}	• Neoadjuvant Cemiplimab-rwlc^{j,10} • If ineligible for or progressed on immune checkpoint inhibitors and clinical trials, consider: ▶ Capecitabine^{23,24} ▶ Cisplatin^{h,i} ▶ Cisplatin/Fluorouracil^{h,24}

[See Evidence Blocks on EB-4, EB-5, and EB-6](#)

^c Cemiplimab-rwlc was studied in CSCC with very-high-risk for recurrence due to nodal features (extracapsular extension with largest node ≥20 mm in diameter or ≥3 involved nodes) or non-nodal features (in-transit metastases, T4 lesion [with bone invasion], PNI, or locally recurrent tumor with ≥1 additional risk feature). Rischin D, Porceddu S, Day F, et al. Adjuvant Cemiplimab or Placebo in High-Risk Cutaneous Squamous-Cell Carcinoma. N Engl J Med 2025;393:774-785.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

[Footnotes on SCC-E \(4 of 5\)](#)
[References on SCC-E \(5 of 5\)](#)

SCC-E
3 OF 5



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

FOOTNOTES

^d [Principles of Radiation Therapy \(SCC-D\)](#).

^e Adjuvant RT can be considered for CSCCs with gross clinical radiologic PNI, multifocal histologic nerve invasion, ≥ 6 cm tumor diameter, recurrent tumors, high risk for regional or distant metastasis, close surgical margins where further surgery cannot be performed, and desmoplastic or infiltrative tumors in patients who are chronically immunosuppressed. Ruiz ES, et al. J Am Acad Dermatol 2022;87:87-94.

^f Assessment of feasibility of RT should be made by a radiation oncologist.

^g A cure is unlikely to result from surgery and/or RT or there are concerns of significant functional impairment. Multidisciplinary discussion and multimodality treatment (including neoadjuvant and adjuvant therapy) merits consideration.

^h These options have occasionally produced useful responses, but data supporting efficacy are limited.

ⁱ Recent published phase II trial data support the efficacy and safety of Cemiplimab-rwlc and Pembrolizumab in patients with laCSCC, recurrent, and metastatic CSCC. Migden MR, et al. Lancet Oncol 2020;21:294-305. Rischin D, et al. J Immunother Cancer 2020;8:e000775. Hughes BGM, et al. Ann Oncol 2021;32:1276-1285.

^j In solid organ transplant recipients, potential benefit from immune checkpoint inhibitor therapy has to be weighed against a significant risk of organ rejection. For patients receiving immunosuppressive therapy, in consultation with their treating physician, consider dose reduction of the immunosuppressive agent(s) and/or minimizing the doses of calcineurin inhibitors and/or antimetabolites in favor of mTOR inhibitors where appropriate. Patients with underlying immunodeficiencies, including CLL, were excluded from the phase I-II Cemiplimab-rwlc trial, so the efficacy of Cemiplimab-rwlc in this population is unclear.

^k Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

^l Nivolumab and hyaluronidase-nvhy subcutaneous injection may be substituted for IV Nivolumab. Nivolumab and hyaluronidase-nvhy has different dosing and administration instructions compared to IV Nivolumab.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

REFERENCES

- 1 Cornejo CM, Jambusaria-Pahlajani A, Willenbrink TJ, et al. Field cancerization: Treatment. *J Am Acad Dermatol* 2020;83:719-730.
- 2 Afsar FS, Erkan CD, Karaca S. Clinical practice trends in cryosurgery: a retrospective study of cutaneous lesions. *Postepy Dermatol Alergol* 2015;32:88-93.
- 3 Cunningham TJ, Tabacchi M, Eliane JP, et al. Randomized trial of calcipotriol combined with 5-fluorouracil for skin cancer precursor immunotherapy. *J Clin Invest* 2017;127:106-116.
- 4 Jansen MHE, Kessels JPHM, Nelemans PJ, et al. Randomized trial of four treatment approaches for actinic keratosis. *N Engl J Med* 2019;380:935-946.
- 5 Badri O, Schmults CD, Karia PS, Ruiz ES. Efficacy and cost analysis for acitretin for basal and squamous cell carcinoma prophylaxis in renal transplant recipients. *Dermatol Surg* 2021;47:125-126.
- 6 Endrizzi B, Ahmed RL, Ray T, et al. Capecitabine to reduce nonmelanoma skin carcinoma burden in solid organ transplant recipients. *Dermatol Surg* 2013;39:634-645.
- 7 Jansen MHE, Mosterd K, Arits AHMM, et al. Five-year results of a randomized controlled trial comparing effectiveness of photodynamic therapy, topical Imiquimod, and topical 5-Fluorouracil in patients with superficial basal cell carcinoma. *J Invest Dermatol* 2018;138:527-533.
- 8 Drew BA, Karia PS, Mora AN, et al. Treatment patterns, outcomes, and patient satisfaction of primary epidermally limited nonmelanoma skin cancer. *Dermatol Surg* 2017;43:1423-1430.
- 9 Guthrie TH Jr, Porubsky ES, Luxenberg MN, et al. Cisplatin-based chemotherapy in advanced basal and squamous cell carcinomas of the skin: results in 28 patients including 13 patients receiving multimodality therapy. *J Clin Oncol* 1990;8:342-346.
- 10 Gross ND, Miller DM, Khushalani NI, et al. Neoadjuvant Cemiplimab for Stage II to IV Cutaneous Squamous-Cell Carcinoma. *N Engl J Med* 2022;387:1557-1568.
- 11 Rischin D, Migden MR, Lim AM, et al. Phase 2 study of cemiplimab in patients with metastatic cutaneous squamous cell carcinoma: primary analysis of fixed-dosing, long-term outcome of weight-based dosing. *J Immunother Cancer* 2020;8:e000775.
- 12 Hughes BGM, Munoz-Couselo E, Mortier L, et al. Pembrolizumab for locally advanced and recurrent/metastatic cutaneous squamous cell carcinoma (KEYNOTE-629 study): an open-label, nonrandomized, multicenter, phase II trial [published correction appears in *Ann Oncol* 2022;33:853]. *Ann Oncol* 2021;32:1276-1285.
- 13 Maring S, Elsayad K, Stenner M, et al. Efficacy of carboplatin/paclitaxel-based radiochemotherapy in locally advanced squamous cell carcinoma of head and neck. *Oncol Res Treat* 2018;41:736-743.
- 14 Vlacich G, Diaz R, Thorpe SW, et al. Intensity-modulated radiation therapy with concurrent carboplatin and paclitaxel for locally advanced head and neck cancer: toxicities and efficacy. *Oncologist* 2012;17:673-681.
- 15 Maubec E, Petrow P, Scheer-Senarich I, et al. Phase II study of cetuximab as first-line single-drug therapy in patients with unresectable squamous cell carcinoma of the skin. *J Clin Oncol* 2011;29:3419-3426.
- 16 Clingan P, Ladwa R, Brungs D, et al. Efficacy and safety of cosibelimab, an anti-PD-L1 antibody, in metastatic cutaneous squamous cell carcinoma. *J Immunother Cancer*. 2023;11:e007637.
- 17 Ferrari D, Fiore J, Codecà C, et al. A phase II study of carboplatin and paclitaxel for recurrent or metastatic head and neck cancer. *Anticancer Drugs* 2009;20:185-190.
- 18 Carinato H, Burgy M, Ferry R, et al. Weekly Paclitaxel, Carboplatin, and Cetuximab as First-Line Treatment of Recurrent and/or Metastatic Head and Neck Squamous Cell Carcinoma for Patients Ineligible to Cisplatin-Based Chemotherapy: A Retrospective Monocentric Study in 60 Patients. *Front Oncol* 2021;11:714551.
- 19 Botticelli A, Pomati G, Cirillo A, et al. Weekly chemotherapy as first line treatment in frail head and neck cancer patients in the immunotherapy era. *J Transl Med* 2021;19:303.
- 20 Tahara M, Kiyota N, Yokota T, et al. Phase II trial of combination treatment with paclitaxel, carboplatin and cetuximab (PCE) as first-line treatment in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck (CSPOR-HN02). *Ann Oncol* 2018;29:1004-1009.
- 21 Geraghty L, Schultz TE, Hoffman SE, et al. Weekly vs. 3-weekly paclitaxel, carboplatin, and cetuximab (PCC) in recurrent/metastatic head and neck cancer. *Mol Clin Oncol* 2021;15:240.
- 22 Khansur T, Kennedy A. Cisplatin and 5-fluorouracil for advanced locoregional and metastatic squamous cell carcinoma of the skin. *Cancer* 1991;67:2030-2032.
- 23 Péron J, Poupard M, Ceruse P, et al. Efficacy and safety of capecitabine in heavily pretreated recurrent/metastatic head and neck squamous cell carcinoma. *Anticancer Drugs* 2012;23:1107-1111.
- 24 Martinez-Trufero J, Isla D, Adansa JC, et al. Phase II study of capecitabine as palliative treatment for patients with recurrent and metastatic squamous head and neck cancer after previous platinum-based treatment. *Br J Cancer* 2010;102:1687-1691.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF PDEMA TECHNIQUE

- PDEMA, also known as complete margin assessment, is a descriptive term for surgical techniques that allow high-quality histologic visualization and interpretation of the entire marginal surface of surgically excised tissue. The NCCN Guidelines Panel recognizes that a variety of surgical methods may achieve complete margin assessment. This NCCN appendix is intended to be inclusive of this diversity, while defining the features that are essential to the superior cure rates achieved by these techniques.¹
- The most commonly used form of PDEMA is Mohs. When anatomic structures at the deep margin (eg, major vessels, nerves, bone) preclude complete histologic evaluation of the marginal surface via Mohs or other forms of PDEMA, Mohs or other forms of PDEMA should be used to evaluate as much of the marginal surface as feasible. Treatment considerations for non-visualized areas may be the subject of multidisciplinary discussion.
- A surgical procedure can be described as PDEMA if and only if all of the following criteria are met:
 1. The entire marginal surface of the surgical specimen is microscopically visualized and histopathologically analyzed for the presence of cancer. The marginal surface includes the complete deep and peripheral margin.
 2. The surgical specimen is oriented with respect to the surgical site and marked in a manner such that any positive margin identified in histopathologic analysis can be accurately located and re-excised.
 3. The surgical margin of any re-excised tissue is again entirely visualized and oriented as above. This process is repeated until no further cancer is identified at the surgical margin or until further excision is not anatomically possible or not in the best interest of the patient.
 4. The time interval between the steps of this process is rapid enough to prevent significant size or shape changes in the wound bed (ie, granulation, contraction) that would decrease the accuracy of orientation.

¹ Curtis KK, Fakult NJ, Strunck JL, et al. Establishing Consensus for Mohs Micrographic Surgical Techniques in the Treatment of Melanoma in Situ for Future Clinical Trials: A Modified Delphi Study. J Natl Compr Canc Netw 2024;22:e247036.


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

NCCN Guidelines Version 2.2026
Squamous Cell Skin Cancer
NCCN Evidence Blocks™

PRINCIPLES OF PDEMA TECHNIQUE

- Visualization of the entire marginal surface of an irregular surgical specimen may be challenging, but is critical to the success of PDEMA methods. Typically, this visualization is achieved by flattening topographically complex surfaces onto a single plane and sectioning the specimen *parallel* to this plane (see Figure 1 below or [Figure 2](#)). Sampling methods such as perpendicular sectioning, also known as “breadloafing,” *do not* achieve direct visualization of the entire surgical margin and would prevent a procedure from achieving PDEMA.
- PDEMA can be achieved with either frozen sections or formalin fixation and paraffin embedding. Although it is often helpful for the surgeon to examine the specimen histologically, the surgeon is not required to examine the specimen histopathologically to achieve PDEMA; a trained pathologist or dermatopathologist may communicate results to the surgeon. If a pathologist or dermatopathologist analyzes the specimen, a consistent communication system must be in place to designate the marginal surfaces for examination and to ensure that the three-dimensional orientation of marginal surfaces and of tissue blocks relative to the wound bed are maintained and communicated to the surgeon to enable accurate localization of residual tumor within the wound bed. The use of multiple operating settings and surgeons is also consistent with PDEMA as long as the orientation of the tissue and wound bed are accurately communicated and complete margin assessment is maintained.²

Figure 1
Tubingen muffin technique



² Leigheb M, Zavattaro E, Bellinzona F, et al. Micrographic surgery (Tubingen torte technique) for the treatment of an invasive dermatofibrosarcoma protuberans with muscular involvement. G Ital Dermatol Venereol 2010;145:309-311.


Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.

PRINCIPLES OF PDEMA TECHNIQUE

Figure 2

Tubingen torte technique

Courtesy of Dr. Brooke Walls, DO, FAAD, Aspen Center for Cosmetic Medicine & Dermatology



Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF PDEMA TECHNIQUE

- Published examples of PDEMA include:
 - ▶ Mohs^{3,4}
 - ▶ Tubingen muffin technique^{5,6}
 - ▶ Tubingen torte technique (see <https://esms-mohs.eu/tuebingen/>)
- Examples of techniques that do not achieve PDEMA include:
 - ▶ Wide local excision with “breadloafing” (perpendicular section prevents visualization of the entire margin)
 - ▶ Square procedure,⁷ quadrant technique, moat technique, and perimeter technique⁸ (wherein the deep margin is assessed with vertical sections so complete visualization of the deep margin is absent). As compared to “breadloafing,” these techniques provide more complete peripheral margin evaluation for superficial tumors (eg, melanoma in situ and extramammary Paget disease) that do not involve subcutaneous tissues. However, these techniques do not provide complete deep margin evaluation so are not PDEMA.

PDEMA Checklist	Yes	No
Is the entire peripheral margin of the surgical specimen microscopically visualized?		
Is the entire deep margin of the surgical specimen microscopically visualized?		
Is the surgical specimen oriented to the wound bed and marked such that any positive margin identified in histopathologic analysis can be accurately located and re-excised?		
Is the process of excision and complete histologic examination repeated until no further cancer is identified or until further excision is no longer in the best interest of the patient?		
Is the process rapid enough to prevent distortion of the wound bed that would decrease accuracy of tissue orientation?		

All of the above categories must be marked Yes to achieve PDEMA. If any of the above are marked No, the procedure does not achieve PDEMA.

³ Tromovitch TA, Stegeman SJ. Microscopically controlled excision of skin tumors. Arch Dermatol 1974;110:231-232.

⁴ Behshad R. Mohs Micrographic Surgery. In: Kantor J, ed. Dermatologic Surgery. McGraw-Hill Education; 2018:388-413.

⁵ Möhrle M, Breuninger H. [The Muffin technique--an alternative to Mohs' micrographic surgery]. J Dtsch Dermatol Ges 2006;4:1080-1084.

⁶ Farma JM, Ammori JB, Zager JS, et al. Dermatofibrosarcoma protuberans: How wide should we resect? Ann Surg Oncol 2010;17:2112-2118.

⁷ Johnson TM, Headington JT, Baker SR, Lowe L. Usefulness of the staged excision for lentigo maligna and lentigo maligna melanoma: The “square” procedure. J Am Acad Dermatol 1997;37:758-764.

⁸ Moehrle M, Breuninger H, Röcken M. A confusing world: what to call histology of three-dimensional tumour margins? J Eur Acad Dermatol Venereol 2007;21:591-595.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

- The decision to offer genetic testing involves three related stages:
 - 1) Pre-test counseling prior to ordering testing;
 - 2) Consideration of the most appropriate testing strategy; and
 - 3) Testing result disclosure and post-test counseling.
- There are rare genetic syndromes that can markedly predispose patients to aggressive CSCC formations. These include xeroderma pigmentosum (XP) and recessive dystrophic epidermolysis bullosa (RDEB). Patients with these conditions should be referred to a cancer center with particular expertise in CSCC prevention and prophylaxis.
- It is recommended that a genetic counselor, medical geneticist, endocrinologist, oncologist, surgeon, oncology nurse, or other health professional with expertise and experience in cancer genetics be involved at each stage whenever possible. Clinicians without direct referral access to the appropriate expertise should be aware of the telehealth genetic counseling options available. These resources can be found through the National Society of Genetic Counselors (NSGC) “Find a Genetic Counselor” tool (www.nsgc.org).

See the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate](#) for the following:

- Principles of Cancer Risk Assessment and Counseling (EVAL-A)
- Pedigree: First-, Second-, and Third-Degree Relatives of Proband (EVAL-B)

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).
All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

PREVENTION: THERAPY TO PREVENT DEVELOPMENT OF ACTINIC KERATOSES AND SCC ([See SCC-B 2 of 2](#))

Acitretin	
Isotretinoin	
Nicotinamide	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

TREATMENT FOR ACTINIC KERATOSES

Pretreatment for hyperkeratotic actinic keratoses	
Topical Tazarotene	
Topical keratolytics (topical urea, lactic acid, and salicylic acid)	

ALTERNATIVE THERAPY FOR SCC IN SITU (BOWEN'S DISEASE)

Topical Fluorouracil	
Topical Imiquimod	
Photodynamic therapy with ALA	
Photodynamic therapy with Porfimer sodium	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

4				
3				
2				
1				
	E	S	Q	C

Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

FIELD CANCERIZATION/CONFLUENT EPIDERMAL DYSPLASIA

Preferred Regimens	
Topical Calcipotriene/Fluorouracil	
Topical Fluorouracil	
Other Recommended Regimen	
Topical Aminolevulinic acid (in conjunction with PDT)	
Useful in Certain Circumstances	
Acitretin	
Capecitabine (for severe refractory disease that has progressed on oral retinoids)	
Isotretinoin	
Porfimer sodium (in conjunction with PDT)	

ADDITIONAL TOPICAL TREATMENT MODALITIES

Topical Imiquimod	
Topical Tirbanibulin	
Topical Diclofenac	

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

Efficacy of Regimen/Agent				
4				
3				
2				
1				
E S Q C A				

S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

SYSTEMIC THERAPY FOR SQUAMOUS CELL SKIN CANCER

Systemic Therapy Options for Use with RT

	Locally Advanced Disease in Non-Surgical Candidates	New Regional Disease – resected high-risk	New Regional Disease – inoperable or incompletely resected	Regional Recurrence or Distant Metastatic Disease	Satellitosis/In-transit Metastasis (S-ITM)
Preferred Regimen					
Cisplatin					
Other Recommended Regimens					
Carboplatin					
Carboplatin/Paclitaxel					
Cetuximab					
Useful in Certain Circumstances					
Cisplatin/Fluorouracil					

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

Cable Network, Inc.				
4				
3				
2				
1				
	E	S	Q	C

E = Efficacy of Regimen/Agent
S = Safety of Regimen/Agent
Q = Quality of Evidence
C = Consistency of Evidence
A = Affordability of Regimen/Agent

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

SYSTEMIC THERAPY FOR SQUAMOUS CELL SKIN CANCER

Very-high-risk with significant risk of extensive local recurrence, nodal, or in-transit metastasis

Cemiplimab-rwlc	
Neoadjuvant Cemiplimab-rwlc	

S-ITM -- Unresectable or incompletely resected disease regional therapy

Isolated limb infusion/perfusion (ILI/ILP) with Melphalan-based regimens	
--	--

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#)

SYSTEMIC THERAPY FOR SQUAMOUS CELL SKIN CANCER

Options for Systemic Therapy Alone

	Locally Advanced Disease in Non-Surgical Candidates	New Regional Disease – inoperable or incompletely resected	Regional Recurrence or Distant Metastatic Disease	Satellitosis/In-transit Metastasis (S-ITM)
Preferred Regimens				
Cemiplimab-rwlc				
Pembrolizumab				
Other Recommended Regimens				
Carboplatin/Paclitaxel				
Carboplatin/Paclitaxel + Cetuximab				
Cetuximab				
Cosibelimab-ipdl				
Nivolumab				
Useful in Certain Circumstances				
Capecitabine				
Cisplatin				
Cisplatin/Fluorouracil				
Neoadjuvant Cemiplimab-rwlc				

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-DEF](#).



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

American Joint Committee on Cancer (AJCC)

TNM Staging Classification for Cutaneous Carcinoma of the Head and Neck (8th ed., 2017)^{1,2}

Table 1. Definitions for T, N, M

T	Primary Tumor
TX	Primary tumor cannot be assessed
Tis	Carcinoma <i>in situ</i>
T1	Tumor smaller than or equal to 2 cm in greatest dimension
T2	Tumor larger than 2 cm, but smaller than or equal to 4 cm in greatest dimension
T3	Tumor larger than 4 cm in maximum dimension or minor bone erosion or perineural invasion or deep invasion*
T4	Tumor with gross cortical bone/marrow, skull base invasion and/or skull base foramen invasion
T4a	Tumor with gross cortical bone/marrow invasion
T4b	Tumor with skull base invasion and/or skull base foramen involvement

*Deep invasion is defined as invasion beyond the subcutaneous fat or >6 mm (as measured from the granular layer of adjacent normal epidermis to the base of the tumor); perineural invasion for T3 classification is defined as tumor cells within the nerve sheath of a nerve lying deeper than the dermis or measuring 0.1 mm or larger in caliber, or presenting with clinical or radiographic involvement of named nerves without skull base invasion or transgression.

Clinical N (cN)

cN	Regional Lymph Nodes
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension and ENE(-)
N2	Metastasis in a single ipsilateral node larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(-); or metastases in multiple ipsilateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-); or in bilateral or contralateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-)
N2a	Metastasis in a single ipsilateral node larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(-)
N2b	Metastases in multiple ipsilateral nodes, none larger than 6 cm in greatest dimension and ENE(-)
N2c	Metastases in bilateral or contralateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-)
N3	Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(-); or metastasis in any node(s) and clinically overt ENE [ENE(+)]
N3a	Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(-)
N3b	Metastasis in any node(s) and ENE (+)

Note: A designation of “U” or “L” may be used for any N category to indicate metastasis above the lower border of the cricoid (U) or below the lower border of the cricoid (L). Similarly, clinical and pathological extranodal extension (ENE) should be recorded as ENE(-) or ENE(+).

¹ These staging tables are for cutaneous squamous cell carcinoma, cutaneous carcinoma, basal cell carcinoma of the head and neck, and all other nonmelanoma skin carcinomas of the head and neck (except Merkel cell carcinoma). Anatomic site of external vermilion lip is included because it has a more similar embryologic origin to skin, and its etiology—which is often based on ultraviolet exposure—is more similar to other nonmelanoma skin cancers. The AJCC Staging Manual, Eighth Edition does not include staging for cutaneous carcinoma outside the head and neck.

² ©, 2021, American College of Surgeons, All Rights Reserved. *Used with the permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging System.*

Continued



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

American Joint Committee on Cancer (AJCC) TNM Staging Classification for Cutaneous Carcinoma of the Head and Neck (8th ed., 2017)^{1,2}

Pathological N (pN)

pN Regional Lymph Nodes

NX Regional lymph nodes cannot be assessed

N0 No regional lymph node metastasis

N1 Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension and ENE(–)

N2 Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension and ENE(+);
or larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(–);
or metastases in multiple ipsilateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(–);
or in bilateral or contralateral lymph node(s), none larger than 6 cm in greatest dimension, ENE(–)

N2a Metastasis in single ipsilateral node 3 cm or smaller in greatest dimension and ENE(+);
or a single ipsilateral node larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(–)

N2b Metastases in multiple ipsilateral nodes, none larger than 6 cm in greatest dimension and ENE(–)

N2c Metastases in bilateral or contralateral lymph node(s), none larger than 6 cm in greatest dimension and ENE(–)

N3 Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(–);
or in a single ipsilateral node larger than 3 cm in greatest dimension and ENE(+);
or multiple ipsilateral, contralateral, or bilateral nodes, any with ENE(+);
or a single contralateral node of any size and ENE(+)

N3a Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(–)

N3b Metastasis in a single ipsilateral node larger than 3 cm in greatest dimension and ENE(+);
or multiple ipsilateral, contralateral, or bilateral nodes, any with ENE(+);
or a single contralateral node of any size and ENE(+)

Note: A designation of “U” or “L” may be used for any N category to indicate metastasis above the lower border of the cricoid (U) or below the lower border of the cricoid (L). Similarly, clinical and pathological extranodal extension (ENE) should be recorded as ENE(–) or ENE(+).

M Distant Metastasis

M0 No distant metastasis

M1 Distant metastasis

G Histologic Grade

GX Grade cannot be assessed

G1 Well differentiated

G2 Moderately differentiated

G3 Poorly differentiated

G4 Undifferentiated

Table 2. AJCC Prognostic Stage Groups

	T	N	M
Stage 0	Tis	N0	M0
Stage I	T1	N0	M0
Stage II	T2	N0	M0
Stage III	T3	N0	M0
	T1	N1	M0
	T2	N1	M0
	T3	N1	M0
Stage IV	T1	N2	M0
	T2	N2	M0
	T3	N2	M0
	Any T	N3	M0
	T4	Any N	M0
	Any T	Any N	M1

¹ These staging tables are for cutaneous squamous cell carcinoma, cutaneous carcinoma, basal cell carcinoma of the head and neck, and all other nonmelanoma skin carcinomas of the head and neck (except Merkel cell carcinoma). Anatomic site of external vermilion lip is included because it has a more similar embryologic origin to skin, and its etiology—which is often based on ultraviolet exposure—is more similar to other nonmelanoma skin cancers. The AJCC Staging Manual, Eighth Edition does not include staging for cutaneous carcinoma outside the head and neck.

² ©, 2021, American College of Surgeons, All Rights Reserved. Used with the permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging System.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

ABBREVIATIONS

ALA	aminolevulinic acid	laCSCC	locally advanced cutaneous squamous cell carcinoma
BED	biologically effective dose	NSGC	National Society of Genetic Counselors
C&E	curettage and electrodesiccation	PDEMA	peripheral and deep en face margin assessment
CLL	chronic lymphocytic leukemia	PDT	photodynamic therapy
CSCC	cutaneous squamous cell carcinoma	PNI	perineural invasion
EBRT	external beam RT	PNTS	perineural tumor spread
ENE	extranodal extension	RDEB	recessive dystrophic epidermolysis bullosa
ENI	elective nodal irradiation	SBRT	stereotactic body radiation therapy
FDG	fluorodeoxyglucose	SCC	squamous cell carcinoma
FNA	fine-needle aspiration	S-ITM	satellitosis/in-transit metastasis
H&P	history and physical	SLNB	sentinel lymph node biopsy
HIV	human immunodeficiency virus	UV	ultraviolet
IGRT	image-guided radiation therapy	XP	xeroderma pigmentosum
ILI	isolated limb infusion		
ILP	isolated limb perfusion		
IMRT	intensity-modulated radiation therapy		
KID	keratitis, ichthyosis, deafness		



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

NCCN Evidence Blocks™

NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Discussion

This discussion corresponds to the NCCN Guidelines for Squamous Cell Skin Cancer. Last updated March 10, 2026.

Table of Contents

Overview	MS-2
Guidelines Update Methodology	MS-2
Literature Search Criteria	MS-2
Sensitive/Inclusive Language	MS-2
Risk Factors for SCC	MS-2
Clinical Presentation and Workup	MS-3
Risk Stratification of Local CSCC Based on Risk Factors for Local Recurrence, Metastases, or Death	MS-4
History and Physical	MS-4
Pathology	MS-5
Field Cancerization/Confluent Epidermal Dysplasia	MS-6
Prevention of Actinic Keratoses	MS-6
Treatment of Actinic Keratoses	MS-7
Treatment Modalities for Local SCC	MS-9
Curettage and Electrodesiccation	MS-9
Mohs Micrographic Surgery or Excision with Peripheral and Deep En Face Margin Assessment	MS-9
Shave Removal	MS-10
Standard Excision with Postoperative Margin Assessment	MS-11
Radiation Therapy	MS-12
Sentinel Lymph Node Biopsy	MS-14
Regional Lymph Node Involvement in SCC	MS-15
Workup for Suspicion of Regional Lymph Node Involvement	MS-15
Treatment of SCC with Regional Lymph Node Involvement	MS-15
Satellitosis/In-Transit Metastasis	MS-16
Recurrence and Metastasis	MS-16
Systemic Therapy for Distant Metastatic Disease	MS-17
Systemic Therapy Options	MS-17
Follow-up	MS-19
Care for Patients at High Risk of Developing Multiple SCCs	MS-20

Treatment of SCC in Patients at High Risk	MS-20
Prevention in Patients at High Risk	MS-20
Patient Education for Patients at High Risk	MS-21
References	MS-24



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Overview

Cutaneous squamous cell carcinoma (SCC or CSCC) is the second most common skin cancer.¹⁻³ Numerous population-based studies have demonstrated that the incidence of SCC is rising.^{1,4-8} Some studies show that SCC incidence rates are rising more rapidly than basal cell carcinoma (BCC), reducing the difference in incidence between these two skin cancers.^{2,3,9} Current estimates by the American Cancer Society indicate that 3.3 million individuals in the United States have ≥ 1 basal or squamous cell skin cancers where approximately 20% are SCCs.¹⁰ Although rarely metastatic, SCC can produce substantial local destruction along with disfigurement and may involve extensive areas of soft tissue, cartilage, and bone. SCCs generally have a good prognosis, with 5-year survival of about 98%.^{1,11-13} However, the treatment landscape continues to evolve to improve a patient's quality of life and minimize disease recurrence.

Guidelines Update Methodology

The complete details of the Development and Update of the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) are available at www.NCCN.org.

Literature Search Criteria

Prior to the update of this version of the NCCN Guidelines® for Squamous Cell Skin Cancer, an electronic search of the PubMed database was performed to obtain key literature published since the previous Guidelines update, using the search term: cutaneous squamous cell carcinoma. The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.¹⁴

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase 2; Clinical Trial, Phase 3; Clinical Trial, Phase 4;

Guideline; Meta-Analysis; Practice Guideline; Randomized Controlled Trial; Systematic Reviews; and Validation Studies. The data from key PubMed articles as well as articles from additional sources deemed as relevant to these guidelines as discussed by the Panel during the Guidelines update have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel's review of lower-level evidence and expert opinion.

Sensitive/Inclusive Language

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation.¹⁵ NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-weight biased; and inclusive of individuals of all sexual orientations and gender identities. NCCN Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will continue to use the terms men, women, female, and male when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Risk Factors for SCC

A number of risk factors are associated with the development of SCC. The most recognized environmental carcinogen is sunlight. Evidence reveals that chronic sun exposure, total site-specific exposure, and number of site-



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

specific sunburns are strongly correlated with development of SCC.¹⁶⁻¹⁹ Due to the link with chronic and cumulative sun exposure, SCC rates are higher in occupations involving outdoor work²⁰⁻²² and increase with age, particularly in sun-exposed sites.^{2,9,18} Indoor tanning is also significantly associated with SCC. According to two large meta-analyses, any exposure to indoor tanning increases the risk of SCC by 67%,²³ with the prevalence of indoor tanning much higher than previously thought among U.S. adults and college students.²⁴

Individuals with light skin, hair, and eye color who have received too much sun exposure are at the greatest risk for SCC.²⁵⁻²⁷ The incidence of ultraviolet (UV)-induced SCCs is very low in non-white populations and has been poorly quantified in people of mixed ethnicities. Most SCCs develop on sun-exposed skin sites, especially the head and neck area.^{5,17,28} Actinic keratoses (AKs) and Bowen's disease, if left untreated, can also progress to invasive SCC with the potential for metastasis.^{21,29-31} Furthermore, SCCs are also known to develop in association with scars or chronic wounds (Marjolin's ulcer).^{32,33} These types of SCCs occur at similar rates in people of all racial and ethnic groups. Such SCC lesions tend to be difficult to treat and have higher risk of recurrence.³⁴⁻³⁶

Lastly, certain genetic syndromes greatly predispose affected individuals to SCC formation, such as albinism³⁷⁻³⁹ and xeroderma pigmentosum.⁴⁰⁻⁴² Certain settings of immunosuppression (eg, organ transplantation, lymphoma, chronic lymphocytic leukemia, drug-induced immunosuppression, HIV) also predispose affected individuals to UV-induced SCC.⁴³⁻⁵⁰ Most notably, analyses of transplant registries have reported a 5-fold to 113-fold increase in incidence of SCC in patients with transplants compared to the general population.^{45,51-53} These patient groups are also at high risk of developing multiple CSCCs and tumors that can behave aggressively.^{42,54-61} Within these high-risk groups, individual patients should be identified for closer follow-up (see *Identification and*

Management of Patients at High Risk for Multiple Primary CSCCs in the algorithm).

Clinical Presentation and Workup

On clinical presentation of a suspicious lesion, workup for SCC begins with a history and physical examination. A skin biopsy that extends into the dermis is then performed on any suspicious lesion. This procedure is preferred because an infiltrative histology may sometimes be present only at the deeper, advancing margins of a tumor and superficial biopsies will frequently miss this component.^{62,63} Skin lesions in high-risk populations may be difficult to assess clinically; therefore, a low threshold for performing skin biopsies in these patients is necessary. Basosquamous carcinoma may behave as aggressively as CSCC.⁶⁴⁻⁶⁶

After a CSCC diagnosis is confirmed, additional workup includes a complete skin, regional lymph node (LN) examination, and imaging studies. A full skin examination is recommended because individuals with a skin cancer often have additional, concurrent precancers or cancers located at other, usually sun-exposed skin sites. These individuals are also at increased risk of developing cutaneous melanoma.^{60,67,68} Imaging studies of the area of interest should be done when locally extensive or metastatic disease is suspected, which includes deep structural involvement such as bone, perineural disease, and deep soft tissue.^{69,70} Due to its higher sensitivity, MRI with and without contrast is preferred for perineural disease or deep soft tissue involvement.⁷¹⁻⁷³ If bone disease is suspected, CT with contrast is preferred unless contraindicated. Imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is sufficient to diagnose local recurrence, but MRI with and without contrast can be considered to assess extent of local disease. For nodal or distant metastases, histologic analysis and/or other imaging modalities can be used for confirmation and to gauge extent of



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

disease. For rare instances that present with distant metastatic disease at diagnosis, the distant metastases pathway should be followed (see *Follow-up* in the algorithm).

Risk Stratification of Local CSCC Based on Risk Factors for Local Recurrence, Metastases, or Death

After the additional workup, a risk assessment of the primary tumor should be performed (see *Stratification to Determine Treatment Options for Local CSCC Based on Risk Factors for Local Recurrence, Metastases, or Death* in the algorithm). Risk category assignment should be based on the highest risk factor present. The high-risk group has elevated risk of local recurrence while the very-high-risk group has elevated risks of local recurrence and metastasis. Other staging systems, including the AJCC 8th edition staging system of CSCC, have been formulated and independently tested to define high-risk groups among patients with localized disease, and can act as additional sources of reference.^{11,54,55,60,74-81}

History and Physical

Location and Diameter

Anatomic location has been known to be a risk factor for SCC recurrence and metastasis for many years.^{34,55,82} In general, SCCs that develop in the head and neck area, particularly the ears and vermillion lips, are more likely to recur and metastasize than those developing on the trunk and extremities.^{11,34,54,55,75,82-85} Besides the head and neck, SCCs that develop on the hands, feet, pretibial, and anogenital areas are also at greater risk of local recurrence and nodal metastasis, independent of diameter.^{11,86,87}

Tumor diameter has also been shown to be a risk factor for SCC recurrence and metastasis.^{34,55,85,88,89} Although different divisions have been used, robust data support that tumors >2 cm are at higher risk of recurrence, metastasis, and poor disease-specific survival (DSS).^{11,34-36,54,55,74,90,91} Taken together, the NCCN Panel recommends that low-risk

location (trunk and extremities) and <2 cm constitute low-risk CSCC. Low-risk location (trunk and extremities) and size ≥2 cm (but ≤4 cm), or high-risk locations (head, neck, hands, feet, pretibia, and anogenital area), constitute high-risk CSCC.

Regardless of location, the NCCN Panel recommends that a tumor diameter of >4 cm warrants the very-high-risk designation. This is based on data from a large prospective study, which demonstrate that lesions <4 cm and ≥4 cm are associated with 3-year DSS rates of 93% and 67% ($P = .0003$), respectively.⁸⁸ Studies have also reported that a mean lesion diameter of 4.2 cm (± 3.4)⁹² or >5 cm⁸⁵ is significantly associated with LN metastases.

Clinical Borders

The risk factor of well-defined versus poorly defined clinical tumor borders has been reported in the context of BCC and extrapolated to the SCC population based on clinical experience of the NCCN Panel.⁹³⁻⁹⁵ The NCCN Panel considers well-defined clinical borders as a low-risk SCC feature.

Primary Versus Recurrent Disease

The higher risk of recurrence and metastasis for recurrent versus primary disease has been extensively documented in the literature.^{34,88,90,92,96,97} The NCCN Panel notes that primary disease is a feature of low risk of recurrence.

Immunosuppression

In addition to increasing the risk of SCC development, immunosuppression has been shown to be associated with recurrence, metastasis, and death in multiple reports.^{34,54,55,58-60,74,98,99} Studies from the organ transplant literature have further elucidated features linked with SCCs in this unique population of patients who are immunocompromised.^{58,77,100,101} A retrospective review confirmed that



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

patients with transplants with CSCC had more primary tumors and were more likely to have deep tissue spread and perineural and lymphatic invasion.⁵⁸ Other studies found diffuse/focal spindle cell morphology, evidence of human papillomavirus (HPV) infection, and aggressive subclinical extension to be more likely in SCCs from patients with transplants.^{100,102} Two large retrospective studies reported high rates of SCC recurrence and metastasis among patients with transplants despite the fact that most SCCs were stage I/II at presentation.^{77,101}

Site of Prior Radiation Therapy or Chronic Inflammation

Tumors developing in sites of prior radiation therapy (RT) refer to primary CSCCs arising in areas previously irradiated for unrelated conditions. All recurrent tumors are defined as high risk irrespective of prior therapy. Data support that prior RT for unrelated (frequently benign) conditions is a risk factor for SCC recurrence or metastasis.^{90,103} Retrospective studies and meta-analyses have also documented increased rates of metastasis for SCC arising from sites of chronic scarring or inflammation.^{34,35,92,96,104-106}

Rapid Growth Tumor

The evidence for growth rate and prognosis is lacking in CSCC. Based on clinical experience, the NCCN Panel included rapid growth rate as a high-risk factor. A Japanese study reported tumor size and rapid growth as prognostic factors for SCC.¹⁰⁷ In a small retrospective series, tumor growth rate of >4 mm/month exhibits a higher risk of nodal progression and a shorter progression time to LN metastasis.¹⁰⁸ There is also evidence that CSCCs in immunosuppressed individuals are often characterized by aggressive behavior and rapid growth.^{109,110}

Neurologic Symptoms

In tumors with perineural involvement (PNI), clinical symptoms suggesting possible involvement of sensory or motor nerves are commonly absent but may occur. Symptoms include pain, burning, stinging, anesthesia, paresthesia, facial paralysis, diplopia, and blurred vision.^{111,112} Any

suggestion of neurologic involvement in the region of a CSCC should place that tumor in a high-risk category, as PNI is associated with recurrence, metastasis, and poor outcomes.^{11,34,54,74,82,88,90,106,113-115} Poorer outcomes are associated with the presence of clinical symptoms and extent of neuronal involvement.^{74,116-119}

Pathology

Degree of Differentiation

Although Broders originally divided CSCC histologically into four grades in 1920, the NCCN Panel has adopted the current trend to reduce the divisions to two groups: 1) well or moderately differentiated; and 2) poorly differentiated.^{75,120} Many studies, including some very large retrospective studies (N > 1000) provide evidence that poor differentiation is correlated with CSCC recurrence, metastasis, DSS, and overall survival (OS).^{11,34,35,54,74,82,85,90,92,97,106,107,121}

Histology

Examples of reportable histologic subtypes are classified as 1) high-risk: adenosquamous or sarcomatoid; 2) low-risk: verrucous; and 3) other useful features: acantholytic, infiltrative, and spindle cell.¹²²⁻¹²⁸ Only case reports and case series document the outcomes of patients with these subtypes, and thus their prognostic significance is debated. Since these tumors may have a high risk of recurrence and likely would not be included in the high-risk category based on their degree of differentiation, the Panel decided to list them as separate risk factors.

Depth

Data from many large studies support that risk of recurrence and metastasis increases with increasing lesion depth.^{11,34,54,74,75,85,89-91,97,129} CSCC lesion depth can be quantified as thickness in millimeters (mm)¹³⁰ or by anatomic layer(s) invaded, both of which have been included in the T classification of the AJCC 7th and 8th Edition staging for CSCC.^{120,131}



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Prospective data from Brantsch and colleagues reported metastasis rates of 0% of tumors ≤ 2.0 mm in thickness, 4% of tumors 2.1 to 6.0 mm in thickness, and 16% of tumors > 6.0 mm thickness, with depth measured as the greatest vertical distance from the top to the bottom of the tumor.⁵⁵

Other studies show that the risk of recurrence and metastasis is significantly higher for lesions with thickness > 2 mm.^{54,85,91} Meta-analyses have shown that 4-mm and 6-mm thickness cutoffs are prognostic for recurrence and metastasis,^{34,54} and one retrospective study showed that risk for recurrence and metastasis increases significantly for every 1-mm increase in tumor depth.⁷⁸ Regarding anatomic level of invasion, some studies showed significantly higher risk of recurrence or metastasis for CSCC lesions with Clark levels IV–V, corresponding to invasion of the deep reticular dermis or subcutaneous fat, respectively.^{34,89} Other studies have shown that lesions with invasion into the subcutaneous fat significantly increases rates of recurrence and metastasis.^{11,54,74,75,88,90,91}

The NCCN Panel has chosen thickness < 2 mm and no invasion beyond subcutaneous fat as low risk while > 6 mm or invasion beyond subcutaneous fat is considered very high risk. If clinical evaluation of incisional biopsy suggests that microstaging is inadequate, the Panel recommends considering narrow margin excisional biopsy to obtain accurate measurement of thickness and anatomic level of invasion.

Perineural Involvement

PNI is uncommon in any non-melanoma skin cancer (NMSC), but develops more frequently and is more aggressive in CSCC versus BCC.^{117,118,132,133} PNI poses increased risks of recurrence, metastasis (nodal and distant), and death, is more common in recurrent versus primary tumors, and is associated with other risk factors, including larger lesion size, poor differentiation, and adenosquamous or sarcomatoid subtypes.^{11,34,54,74,75,82,88,90,106,113-115,134-138} Specifically, PNI with tumor cells within the nerve sheath of a nerve lying deeper than the dermis or

measuring ≥ 0.1 mm has been associated with metastasis and DSS.^{75,79,131} If large nerve involvement is suspected, MRI should be considered to evaluate extent and/or rule out skull involvement in those with head and neck tumors.^{71,72,112,119,139}

Lymphatic or Vascular Involvement

Significant association between lymphovascular invasion (LVI) and LN metastasis has been reported in prospective^{92,115} and retrospective studies.¹¹⁴ One retrospective study showed that in high-risk CSCC populations with PNI or neurotropism, LVI was significantly associated with DSS and all-cause death.⁷⁹ Based on these data, the Panel considers lymphatic or vascular involvement an indication of very-high-risk SCC.

Field Cancerization/Confluent Epidermal Dysplasia

Field cancerization is defined as UV-light-induced confluent dysplasia clinically manifested as diffuse AKs and superficial (in situ) SCC.¹⁴⁰ AKs should be treated at first development and biopsied for histologic evaluation if they have an atypical clinical appearance or do not respond to treatment. Treatment is particularly important in patients with diffuse AKs and/or field cancerization, as these patients are at high risk of developing multiple primary CSCCs.¹⁴¹ Given the limited penetration beyond the epidermis and lower cure rates than with surgical techniques, superficial therapies should be reserved for those patients with SCC in situ.¹⁴²⁻¹⁴⁵ The NCCN Panel's experience indicates that they may be effective for anatomically challenging locations, and recurrences are often small and manageable.

Prevention of Actinic Keratoses

The regular use of sunscreen can significantly reduce the rate of development of new AKs and CSCCs, as well as increase remission rates of AKs.¹⁴⁶⁻¹⁴⁹ Application of sunscreen is particularly important for areas such as the head, neck, hands, and forearms. Oral nicotinamide may also



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

be an effective way to reduce the development of CSCC. Nicotinamide is a form of vitamin B3 that enhances UV-induced DNA damage repair and reduces UV-induced immunity suppression.¹⁵⁰ A phase III randomized controlled trial (N = 386) concluded that the incidence of new AKs and CSCCs was 13% and 30% lower, respectively, compared to the control placebo group at 12 months.¹⁵¹ While a meta-analysis by Mainville et al considers that oral nicotinamide is an effective preventive option for healthy patients and those with organ transplants, two other clinical trials concluded that AK rate was not significantly reduced in immunocompromised patients.¹⁵²⁻¹⁵⁴

Treatment of Actinic Keratoses

Cryotherapy has been used to treat AK for many decades, despite lack of prospective randomized trials comparing them with non-treatment. Large prospective randomized trials in patients with AKs (N > 100) have shown that each of the following therapies provides better complete clearance rates compared with placebo: topical fluorouracil with or without calcipotriol,¹⁵⁵⁻¹⁶⁰ topical imiquimod,¹⁶¹⁻¹⁶⁴ topical tirbanibulin,¹⁶⁵ and photodynamic therapy (PDT).¹⁶⁶⁻¹⁷⁴ Prospective randomized trials have reported pair-wise comparisons of the above treatments, but results are not consistent. These comparisons include PDT versus cryotherapy,^{166,168,171,175-177} imiquimod,^{178,179} fluorouracil,¹⁸⁰⁻¹⁸² or ingenol mebutate¹⁸³⁻¹⁸⁵; cryotherapy versus fluorouracil or imiquimod¹⁸⁶⁻¹⁸⁸; and fluorouracil versus imiquimod¹⁸⁹ or ingenol mebutate.^{190,191} Meta-analyses of randomized trials have attempted to determine an order of preference for these treatments.¹⁹²⁻¹⁹⁵ Hyperkeratotic AKs pretreatment may include topical tazarotene, curettage, or topical keratolytics such as topical urea, lactic acid, and salicylic acid prior to treatment with fluorouracil with or without calcipotriol, imiquimod, or tirbanibulin. The NCCN Panel separates accepted treatment modalities for field cancerization or confluent epidermal dysplasia into topical, destructive, and other modalities to be considered.

Topical Treatment Modalities

Fluorouracil is considered as a preferred topical regimen, along with calcipotriene/fluorouracil, based on data from a randomized trial that reported the cumulative probability of remaining free from disease progression was significantly higher for fluorouracil (74.7%) than imiquimod (53.9%), MAL-PDT (37.7%), or ingenol mebutate (28.9%).¹⁹⁶ The longest duration for CSCC prophylaxis has been demonstrated with the combination of 5-FU and calcipotriol.¹⁶⁰ It was demonstrated that more participants who received topical calcipotriol/fluorouracil for AK remained disease-free over the >1500-day period compared to those receiving petroleum jelly-based skin product plus fluorouracil.¹⁹⁷ Moreover, significantly fewer participants in the test cohort developed CSCC on the treated face and scalp within 3 years (7% vs. 28% in control group; hazard ratio [HR], 0.215; $P = .032$).¹⁹⁷

Retrospective studies, meta-analyses, and a small open-label phase II trial have shown that imiquimod was effective for treating patients with SCC in situ, with high rates of initial clearance (70%–100%) and low rates of recurrence.¹⁹⁸⁻²⁰² One small double-blind randomized trial showed that imiquimod led to the resolution of 73% of lesions compared to 0% with vehicle control ($P < .001$).²⁰³ Clearance rates with fluorouracil tend to be lower than those for topical imiquimod and vary widely, ranging from 27% to 92%.^{199,202,204-206} Toxicities are similar between imiquimod and fluorouracil, being primarily inflammatory skin reactions such as severe eczematous reactions, ulceration, and erosions.^{199,205,206}

Topical tirbanibulin is a recommended treatment for AK based on results from two identically designed double-blind phase III trials in which patients received either tirbanibulin or vehicle ointment for the treatment of AKs on the face or scalp. In both trials, complete clearance by day 57 occurred in significantly more patients in the tirbanibulin group compared to the



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

vehicle group (trial 1: 44% vs. 5%; $P < .001$; trial 2: 54% vs. 13%; $P < .001$).¹⁶⁵

The utility of topical diclofenac is less clear, as efficacy results vary across large randomized trials, with some studies reporting no significant difference between diclofenac/hyaluronan and hyaluronan alone.^{159,207-209}

Diclofenac/hyaluronan has also been shown to be inferior to MAL-PDT and to fluorouracil for the treatment of AKs.^{210,211} The NCCN Panel assigns category 2B for diclofenac in this setting.

Destructive Treatment Modalities

Fewer high-quality data are available regarding the efficacy and safety of other treatments that are sometimes used and may be considered for treating AKs such as ablative laser vermilionectomy or electrosurgical destruction, ablative skin resurfacing (eg, dermabrasion, laser), chemical peels (trichloroacetic acid [TCA]), cryotherapy, and curettage and electrodesiccation (C&E).

AK on the lip, known as actinic cheilitis, may require a different approach. Prospective studies on the treatment of actinic cheilitis are limited.

Therapies tested include surgical vermilionectomy, lip shave, electrodesiccation, laser vermilion ablation, laser resurfacing, fluorouracil, laser + fluorouracil, TCA chemical peel, PDT, PDT + imiquimod, and diclofenac.²¹²⁻²²³ The NCCN Panel considers ablative laser vermilionectomy or electrosurgical destruction to be of value as a primary treatment option for extensive actinic cheilitis.

Studies have confirmed that chemical peel or laser resurfacing significantly reduced AKs, although in some studies they were less effective than PDT or fluorouracil.²²⁴⁻²³¹ The use of chemical peels and ablative skin resurfacing varies widely across NCCN Member Institutions.

Cryotherapy has been used for many years as a fast and cost-effective means for removal of SCCs. Prospective and retrospective studies, including large meta-analyses, have shown recurrence rates of 0% to 4% for invasive SCCs treated with cryotherapy.^{34,232-235} For SCC in situ, recurrence rates range from 1% to 13% in retrospective studies^{143,144,235,236} and 0% to 50% in prospective studies.^{142,206,234,237,238} One prospective study reported that patients were much more likely to experience pain with cryotherapy compared with C&E, and time to complete healing was also significantly longer with cryotherapy.¹⁴² Cryotherapy may also be associated with poorer cosmetic outcomes compared with topical fluorouracil.²⁰⁶

Other Treatment Modalities

PDT with photosensitizing agents including methyl aminolevulinate (MAL) and 5-aminolevulinic acid (ALA) or porfimer sodium is another option for superficial SCC. MAL is no longer produced in the United States. For SCC in situ, rates of initial complete clearance following PDT range between 52% and 98%.^{205,239-250} Durable complete response (CR) rates range from 48% to 89%.^{205,206,239-242,244-248,250,251} It has been shown that differences in PDT techniques can cause significant differences in clearance rate for SCC in situ.^{240,248} Furthermore, results from randomized trials showed fewer treatments required for complete clearance and higher durable CR rates with PDT versus cryotherapy.^{206,237} Compared to fluorouracil, PDT was also associated with higher rates of initial complete clearance and higher durable CR rates.^{205,206} Data suggest that fluorouracil may be associated with lower risk of adverse events (AEs) compared with PDT or cryotherapy, but it is unclear whether risk of toxicity differs between cryotherapy and PDT.^{205,206,237}

The development of AKs may also be reduced by oral retinoid (eg acitretin, isotretinoin) treatment; however, there are significant side effects especially in patients with childbearing potential and the benefits are



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

limited to the time of treatment.²⁵² Another treatment that may be considered for severe refractory disease after progression on oral retinoids is low-dose oral capecitabine, which was investigated in a small cohort of patients with transplants with varying side effects.^{253,254}

Treatment Modalities for Local SCC

Multidisciplinary consultation at a center with specialized expertise is recommended by the NCCN Panel during treatment planning for patients with high-risk, very-high-risk, and locally advanced CSCC (laCSCC).

Curettage and Electrodesiccation

C&E is a fast and cost-effective technique for superficial lesions; however, it does not allow histologic margin assessment. Retrospective and observational data with long-term follow-up (>5 year) indicate that cure rates are between 95% to 100% for patients with primary CSCC lesions treated with C&E.^{34,255-257} These estimates are largely based on patients with low-risk SCC, and there is evidence to suggest that the cure rate is lower for tumors with risk factors. One study reported recurrence rates of 0.4% versus 11% for CSCCs with diameter less than versus greater than 2 cm, and another reported a recurrence rate of 19% for SCCs on the skin of the pinna that were treated with C&E.^{258,259}

The NCCN Panel recommends this technique as a primary treatment option for low-risk CSCCs with three caveats. First, this technique should not be used to treat areas with terminal hair growth such as the scalp, pubic or axillary regions, or beard area in males due to the risk that a tumor extending down follicular structures might not be adequately removed. Second, if the subcutaneous layer is reached during the course of C&E, then surgical excision should generally be performed instead. This change in therapy is necessary as the effectiveness of the C&E technique rests on the ability of the clinician to distinguish between firm, normal dermis, and soft tumor tissue when using a sharp curette. Since

subcutaneous adipose is even softer than tumor tissue, the ability of the curette to distinguish, and therefore to selectively and completely remove tumor cells, diminishes. Third, if C&E has been performed based only on the appearance of a low-risk tumor, biopsy results of the tissue taken at the time of C&E should be reviewed to make sure that there are no high-risk pathologic features that would require additional therapy.

Mohs Micrographic Surgery or Excision with Peripheral and Deep En Face Margin Assessment

Peripheral and deep en face margin assessment (PDEMA), or complete margin assessment, is a term used for a subset of surgical techniques for high-quality histologic visualization and interpretation of the margin surface or surgically excised tissue. Mohs micrographic surgery is the most commonly utilized PDEMA technique. Mohs procedures are particularly successful in non-metastatic basal and squamous cell skin cancers where tissue-sparing and precision microscopic control of margins is a priority and has been associated with lower recurrence rates.^{260,261} While PDEMA is a team procedure that requires the participation of physicians from multiple disciplines, Mohs physicians serve as both the surgeon and pathologist requiring highly specialized training. Efforts have been extended to generate consensus recommendations to offer Mohs surgeons guidance and promote standardization to make data aggregate from multicenter clinical trials possible.²⁶²

Mohs is a primary treatment option for low-risk CSCC and high-risk CSCC, as well as the preferred surgical technique for very-high-risk CSCC because it allows intraoperative analysis of 100% of the excision margin. Additionally, Mohs is a treatment option for patients with laCSCC following a multidisciplinary discussion to consider neoadjuvant systemic therapy and/or participation in a clinical trial. An extensive meta-analysis of studies with long-term follow-up (≥5 years) reported local recurrence rates of 3.1% for primary CSCCs and 10% for recurrences treated with Mohs.³⁴



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Moreover, local recurrence rates have been reported to be significantly less likely with Mohs compared to standard excision.^{263,264} Cure rates for Mohs depended on tumor diameter (<2 vs. ≥2 cm: 98.1% vs. 74.8%) and differentiation (well vs. poorly differentiated: 97.0% vs. 67.4%). For each of these subgroups, cure rates for Mohs were higher than for treatment with non-Mohs modalities.³⁴

Retrospective and prospective observational studies of localized primary SCCs treated with Mohs reported local recurrence rates of 1.2% to 4.1% and rates of metastases between 0% and 6.3%.^{255,256,265-278} Compared with primary tumors, rates of local recurrence or metastasis after Mohs are higher for recurrent tumors (previously treated with a non-Mohs modality).^{90,267} For recurrent CSCCs treated with Mohs, subsequent local reported recurrences occurred in 5.9% to 7.7% of patients; metastasis occurred in 0% to 10% of patients.²⁶⁵⁻²⁷¹ Other risk factors associated with recurrence after Mohs include larger subclinical extension and more Mohs stages required for clearance.²⁶⁷ CSCC with PNI is associated with elevated rates of recurrence (6.8%–32.3%) in studies that occasionally include BCC as well as treatment by RT.^{117,134,136,279-281} Risk factors associated with metastasis after Mohs include: size >2 cm, Clark's level (metastatic CSCC are more likely to be deeper – Clark level III–V), poor differentiation, location in areas of prior radiation, small tumor nests and infiltrative tumor strands, single-cell infiltration, PNI, and acantholysis.⁹⁰

It is not uncommon to find discrepancies between pathology results from preoperative biopsy or initial debulking compared with frozen sections taken during Mohs.²⁸²⁻²⁸⁴ When Mohs with marginal assessment is being performed and the preoperative biopsy is considered insufficient for providing all the staging information required to properly treat the tumor, submission of the central specimen for pathologic evaluation with paraffin sections is recommended. If invasion to the parotid fascia is noted, superficial parotidectomy may be clinically indicated.

Excision with PDEMA (via permanent or frozen section) is acceptable as an alternative to Mohs provided that it includes a complete margin assessment of en face peripheral and horizontal deep margins coupled with close communication between pathologists and surgeons regarding where within the tumor bed further resection is needed. These subsequent specimens must also be processed, and results communicated via the PDEMA method. Low recurrence rates (0%–1%), and specifically lower recurrence rates when compared directly to standard excision,²⁸⁵ have been reported where histologically clear margins are achieved.²⁸⁶ It is important to note that truly histologically negative margins are not necessarily achieved by frozen sectioning alone, without PDEMA. Studies have reported that for CSCC tumors with negative margins upon frozen sectioning, permanent paraffin section analysis indicates positive margins in 10% to 20%.^{283,284,287-289} These discrepancies may be due to unrepresentative sampling of the margins, and instances of frozen sections in which permanent section showed positive margins have reported much higher recurrence rates.²⁸⁹ Overall, the descriptive term PDEMA underscores the Panel's belief that complete assessment of all tissue margins is the key to optimal tumor removal of high-risk tumors. Such effort at local control is particularly important in SCC because one third of deaths occur from local disease alone. Mohs or other forms of PDEMA are also recommended in case of positive margins after standard excision for low-risk CSCC.

Shave Removal

Shave removal, the shaving of epidermal or dermal lesion, is a sharp removal by transverse bowl-shaped slicing to remove epidermal and dermal lesions, without including fat, and does not require suture closure.²⁹⁰ Like C&E, there is concern for inaccurate margin status assessment with shave removal.²⁹¹ However, it is a recommended technique for low-risk SCCs located in the trunk or extremities. Shave removal studies have reported a 0.5% to 30% rate of recurrence over a



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

3- to 5-year follow-up, multiple tumors treated in single visits, and a risk for misdiagnosis of only 1% of patients.²⁹⁰⁻²⁹³ Shave removal is a primary treatment option for low-risk CSCC when the tumor does not extend beyond the dermis.

Standard Excision with Postoperative Margin Assessment

A common therapeutic option for CSCC is standard surgical excision followed by postoperative pathologic evaluation of margins. Retrospective analyses and prospective observational studies have reported rates of recurrence or metastasis ranging from 0% and 14%, with most studies reporting rates of ≤6%.^{34,91,121,129,255,256,270,275,294-298} Distant metastasis was rarely observed, and rates of regional metastasis were highly variable across studies, ranging from 0% to 13%.^{121,129,275,294,298,299} One large meta-analysis found that recurrence rates were lower for primary versus recurrent tumors, both with follow-up of <5 years (5.7% vs. 17.3%) and with longer follow-up (8.1% vs. 23.3%).³⁴ Incomplete excisions can depend on lesion location, thickness, PNI, invasion into the deep fascia, differentiation, surgeons' skills, and primary versus recurrent tumors,^{267,300-302} among other factors.^{275,294,303-310}

The clinical margins chosen by the NCCN Panel for the primary treatment of low-risk CSCC are based on the work of Brodland and colleagues.³¹¹ Their analysis indicated that for well-circumscribed CSCC lesions <2 cm in diameter, excision with 4-mm clinical margins should result in complete removal in >95% of CSCCs. For low-risk lesions >2 cm in diameter, results indicated that 6-mm margins would be needed to achieve histologically clear margins in 95% of instances.

The NCCN Panel also recommends standard excision as the primary treatment for high-risk/very-high-risk CSCC when Mohs micrographic surgery (Mohs) and other forms of PDEMA are not available; however, wider surgical margins than those recommended for low-risk lesions must

be taken and increased recurrence rates should be expected. Standard excision with wider margins is also a recommended treatment option for laCSCC following consideration of neoadjuvant cemiplimab and/or clinical trial after a multidisciplinary discussion to assess the most favorable treatment course.

According to Brodland et al, for CSCCs in high-risk locations (scalp, ears, eyelids, nose, lips) or with other high-risk features (histologic grade ≥2, invasion of subcutaneous tissue), lesions with a diameter <1 cm, 1 to 1.9 cm, and ≥2 cm would require margins of at least 4 mm, 6 mm, and 9 mm, respectively.³¹¹ Other retrospective analyses of CSCCs removed with Mohs further support that larger excision margins are needed to consistently achieve clear margins as tumor diameter increases and when other risk factors are present.^{267,271,300,312} Currently, European Guidelines recommend standard excisions with 6 to 10 mm peripheral clinical margins for high-risk to very-high-risk CSCCs.³¹³⁻³¹⁶ Appropriate margins should be determined case by case based on tumor- and patient-specific factors.

Whenever standard excision is utilized, any peripheral rim of erythema around an SCC must be included in what is assumed to be the tumor. For patients with positive margins from surgical excision and postoperative margin assessment, re-excision often yields clean margins, and in many instances the re-excision specimen contains no tumor cells.^{121,129,295,317-319} Re-excision with postoperative margin assessment is therefore among the recommended treatment options for positive margins after standard excision of low-risk, high-risk/very-high-risk, very-high-risk, and laCSCC. In any case, tissue rearrangement should not be undertaken until clear margins are identified.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Radiation Therapy

Radiation as Primary Therapy

Although surgery is the mainstay of local treatment for SCC, patient preference and other factors may lead to the choice of RT as primary therapy. Patients may receive RT if they are considered nonsurgical candidates due to comorbidities, extent of disease, risk of functional or cosmetic defects, or inability to clear disease with surgery through a multidisciplinary discussion for low-risk, high-risk/very-high-risk, very-high-risk, and laCSCC. In the laCSCC setting, RT may be administered with or without additional systemic therapy. A large meta-analysis reported 5-year recurrence rates of 6.7% and 10% after RT of primary and recurrent SCC, respectively.³⁴ Subsequent retrospective analyses of patients with primary CSCCs have reported a large range of recurrence rates, from 2.8% to 42%, the latter for patients with locally advanced disease (size >2 cm or deeply invasive).^{255,320-326} The risk of recurrence appears to increase with increasing lesion size and T stage.^{323,325,327} A few small studies (n < 20) have reported that for CSCCs that have been previously treated and recurred, treatment with RT results in 16.7% recurrence.^{322,325}

Retrospective analyses have reported recurrence rates ranging from 0% to 10.5% in in situ SCC lesions treated with RT as primary therapy, with most studies reporting local control rates near 100%.^{236,324,325,328-330}

Adjuvant Radiation

For low-risk CSCC, the NCCN Panel recommends adjuvant RT for patients with positive margins after oncologic intent surgery who are not candidates for further resection to clear margins. For high-risk/very-high-risk and very-high-risk CSCC, the NCCN Panel recommends adjuvant RT as a treatment option for patients with positive margins, if further clearance of these positive margins is not feasible. Adjuvant RT may also be considered for patients with laCSCC and positive margins with or without systemic therapy. Adjuvant RT can also be considered for high-risk/very-

high-risk, very-high-risk, and laCSCC with negative margins if there is extensive perineural, large or named nerve involvement, or other poor prognostic features.³³¹

It has been shown that adjuvant RT improved locoregional control and survival outcomes for patients with positive margins after surgery or other high-risk features for recurrence.³³¹⁻³³⁶ Every attempt to clear margins as part of oncologic resection should be made while balancing postoperative recovery for patients with strong indications for adjuvant radiation (eg, perineural spread [PNS], multiple involved nodes, extranodal extension [ENE]).²⁸⁰ The outcome benefit of adjuvant RT following resection of CSCC with negative margins has been estimated to be approximately a 50% reduction in local and nodal recurrence risks,³³⁷ despite older inconclusive data.^{336,338,339} Other retrospective studies combined results for patients treated with other modalities (eg, Mohs/standard excision alone, RT alone, chemotherapy), patients with other types of skin cancer (BCC and metatypical BCC), patients with LN metastases, and a mix of patients with primary and recurrent skin lesions, with and without positive margins.^{116,136,335,340-342} These studies suggest that postoperative RT for patients with high-staged CSCC may improve local and regional control and disease-free survival (DFS), but a survival benefit has not been demonstrated.

Radiation therapy Safety and Administration

The NCCN Panel previously cautioned that RT is often reserved for patients >60 years of age because of concerns about long-term sequelae, including secondary malignancies; however, this statement has been retracted. This is because age is no longer a major factor in determining treatment modality.

Large cohort and population-based studies (N > 1000) have shown that rates of NMSCs are significantly higher in those who received prior RT (either for a benign condition or for cancer) compared with those who have



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

no history of therapeutic RT exposure.³⁴³⁻³⁴⁶ In patients who developed NMSC after prior RT, most NMSC lesions occurred within the radiation field, with elevated risk of NMSC confined to the site of RT exposure. The risk of NMSC was particularly high in patients who received therapeutic RT early in life. RT can result in poor cosmetic outcomes, including telangiectasia, changes in skin pigmentation, and fibrosis. More serious long-term complications can include non-healing ulcers (especially for SCC in situ^{236,328,329}); soft tissue, cartilage, bone, or brain necrosis; decreased sensation; and cataracts (for lesions in the periorbital region).^{325,327,328,330,347-349}

Specifics about the application of RT, including cumulative dose, treatment duration, and contraindications, are described under *Principles of Radiation Therapy* in the algorithm; however, appropriateness of RT treatment should be determined together with a radiation oncologist who has radiation physics support to meet established quality assurance and dosimetric constraints. The NCCN Panel's recommendations for RT dosing are based on clinical practice at their institutions and clinical evidence from studies of various skin cancers. These recommendations are separated into primary tumor (adjuvant RT and definitive EBRT), regional disease, and regarding brachytherapy. In some instances, dose and fractionation schema for conventional fractionation and hypofractionation are provided.^{350,351} Hypofractionation consists of a higher dose of radiation per fraction leading to fewer fractions overall and often a shorter time frame for treatment. For general recommendations regarding radiation doses, refer to the [ASTRO Guideline on Definitive and Postoperative Radiation Therapy for Basal and Squamous Cell Cancers of the Skin](#).³⁵²

RT is contraindicated in patients with genetic conditions predisposing to irradiation-related skin cancer (eg, basal cell nevus syndrome [Gorlin syndrome];³⁵³⁻³⁵⁷ DNA-repair disorders such as Fanconi anemia and

xeroderma pigmentosum), and relatively contraindicated in patients with connective tissue diseases (eg, lupus, scleroderma).³⁵⁸⁻³⁶⁰ Given higher rates of poor cosmesis and complications with increasing cumulative radiation dose,^{327,347,361} reirradiation should not be routinely utilized for recurrent disease within a prior radiation field. Protracted fractionation is associated with improved cosmetic results,^{347,349,362} and should be utilized for poorly vascularized or cartilaginous areas. Retrospective studies have found that for patients with CSCC and high-risk PNI (such as frank perineural spread with clinical or radiographic evidence of nerve involvement or large-caliber nerve involvement, named nerve involvement, or extensive small-caliber nerve involvement), a common pattern of recurrence involves the retrograde pathway of affected named nerves.^{116,363,364} The NCCN Panel recommends including the course of the local nerves retrograde or toward the skull base for extensive PNI, clinically evident PNI, or involvement of named nerves (particularly in the head and neck region).

A variety of external beam options have been shown to be effective for treating CSCC and have similar cosmetic/safety results.^{327,347,362,365-367} Additionally, when treating SCC with intensity-modulated RT (IMRT), proton beam RT, or 3D conformal RT (3D-CRT), it is considered best practice to use image-guided RT (IGRT).³⁶⁸⁻³⁷⁰ IGRT for other types of radiation techniques used to treat SCC is considered unnecessary. Tissue equivalent bolus should be considered for photon/electron treatments when therapeutic targets include the skin surface with use of surface dosimeters to confirm delivered dose at skin surface. Proton radiation may be able to treat superficial targets at therapeutic dose without addition of bolus, but discussion with physics/dosimetry is recommended for individual cases. Considerations for brachytherapy include sensitive treatment areas (eg, ear, nose, near eyes), if immunocompromised, tumor size,³⁷¹ timing of surgical procedures, and optimal brachytherapy technique based on individual patient factors. Isotope-based brachytherapy can be



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

an effective treatment for certain sites of disease, particularly on the head and neck.³⁷²⁻³⁷⁸ A retrospective multicentric analysis of 1676 carcinomas of the skin of the nose and nasal vestibule yielded a local control rate of 93% with a minimum follow-up of 2 years. It was determined in this study that local control depended on tumor size (diameter <2 cm: 96%, 2–3.9 cm: 88%, ≥4 cm: 81%) and tumor site (external surface of the nose: 94%, vestibule: 75%).³⁷⁷ The NCCN Panel recommendations based on their clinical expertise and available data include definitive or adjuvant skin brachytherapy dosing examples for custom surface mold applicators, interstitial brachytherapy, and shielded single-channel surface applications.

On the other hand, there are insufficient long-term efficacy and safety data to support the routine use of electronic surface brachytherapy.^{379,380} Electronic brachytherapy (EB) uses electrically generated x-rays, which are not regulated by the Nuclear Regulatory Commission. One of the apparent advantages of EB devices is the decreased shielding requirements and the portability of the units; however, this may lead to EB use in settings unfamiliar with the hazards of therapeutic radiation delivery.³⁸¹ The [American Association of Physicists in Medicine \(AAPM\) task group 152](#) has provided guidelines for proper and safe EB use including having an authorized medical physicist (AMP) physically present from the initiation through the duration of all treatments involving an EB unit. The AMP is responsible for output calibration, quality assurance, training, and treatment planning.³⁸¹ Furthermore, training and educational requirements for those administering EB vary considerably from state to state, and users should consult their local radiation safety committee to confirm appropriate adherence. The American Brachytherapy Society (ABS) has published a consensus statement for EB, noting a paucity of long-term clinical outcome data and lack of comparison to surgery or standard RT techniques.³⁸² Particularly concerning is the absence of standardized dosimetry (in comparison to high-dose-rate [HDR]

brachytherapy and external beam RT [EBRT]) to ensure adequate target coverage, skin surface dose, and plan quality assurance. The ABS, therefore, recommends that “EB treatment should be performed on clinical registry or trial at this time.” The American Academy of Dermatology (AAD) has published a [Position Statement](#): “The AAD believes additional research is needed on electronic surface brachytherapy particularly on long term outcomes.” The AAD acknowledges concerns regarding aggressive marketing to dermatologists with a focus on revenue streams, potentially undermining quality of care and patient safety. The AAD also cautions dermatologists to be mindful of the Stark physician self-referral law. Recently, image guidance Current Procedural Terminology (CPT) codes have been used for the delivery of EB, despite a lack of evidence demonstrating a benefit in clinical outcomes with image guidance. We do not support the use of image guidance with EB currently. In conclusion, EB should only be performed by radiation oncologists with medical physics support, on registry or clinical trial, with appropriate safeguards, and in adherence with all state and federal regulations. Further research is needed regarding dose deposition and long-term clinical outcomes.

Sentinel Lymph Node Biopsy

The NCCN Panel recommends consideration of sentinel LN biopsy (SLNB) during treatment planning prior to primary treatment when the disease is recurrent or has multiple high-risk features for high-risk/very-high-risk CSCC where surgery or RT has a high likelihood of cure as well as for very-high-risk CSCC with significant risk of extensive local recurrence, nodal or in-transit metastasis. Studies have reported sub-clinical nodal metastases in 7% to 21% of patients with high-risk non-anogenital CSCC who underwent SLNB.^{76,86,114,115,383-389} Although small sample sizes and low rates of SLN positivity limit assessment of prognostic factors, a few studies suggest that risk factors for SLNB positivity include tumor diameter and thickness, LVI, PNI, and the presence of multiple high-risk factors.^{76,114,115,387,390}



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Several studies reported that among patients with localized SCC and a negative SLNB, nodal metastases were later detected in 2% to 15% of patients.^{76,86,114,115,384-386,388,391} In addition to false negatives, some studies documented patients with a negative SLNB who developed local recurrences or metastases outside of the previously biopsied LN basin.^{86,115,384} For positive SLNB, multidisciplinary discussion is recommended after obtaining radiologic staging of the neck, chest, abdomen, and pelvis if not yet completed. In the absence of metastatic disease consider completion lymphadenectomy of the affected nodal basin. If surgery is not an option due to patient preference or poor performance status, then consider RT. Following neck dissection, see *Very-high-risk CSCC* in the algorithm for additional recommendations. It has been shown that despite receiving completion LN dissection, patients with a positive SLN had higher rates of postoperative recurrence/metastases, ranging from 33% to 45%,^{114,115,384,387} and were also more likely to die from SCC, with significantly lower 3-year DSS rates compared to patients who have a negative SLN.^{86,115,387} Therefore, although SLNB may have prognostic value, it is unclear whether SLNB followed by completion LN dissection or adjuvant RT improve patient outcomes.

Regional Lymph Node Involvement in SCC

Regional nodal involvement significantly increases the risk of recurrence and mortality.^{36,85,97} Nodal metastasis also commonly coincides with other adverse histopathologic findings such as LVI, poor differentiation, and PNI.^{11,74,75,82,85,90,92,97,392} About 60% to 82% of patients with nodal disease show involvement in the parotid gland, while cervical neck node disease without parotid invasion is observed in 18% to 41% of patients.³⁹³

Workup for Suspicion of Regional Lymph Node Involvement

The presence of palpable regional LNs or suspicious LNs identified by imaging studies should prompt a fine-needle aspiration (FNA) or core

biopsy of suspicious node(s). If initial pathology results are negative, the NCCN Panel recommends considering re-evaluation by clinical exam, CT with contrast imaging of the nodal basin, and/or pathology on additional LN specimens taken by repeat FNA, core biopsy, or excisional biopsy of the suspicious node(s). For patients with pathologic evidence of LN disease, preoperative imaging of the nodal basin by CT with contrast is recommended to determine the size, number, and location of involved nodes. CT with contrast of the nodal basin can be useful for RT planning. In addition, chest/abdomen/pelvis CT with contrast or FDG-PET/CT are recommended as clinically indicated to rule out distant metastatic disease.

Treatment of SCC with Regional Lymph Node Involvement

The NCCN Panel recommends resection of regional disease over RT or chemotherapy. RT with or without concurrent systemic therapy is reserved for patients who are not surgical candidates. Most studies of patients with regional involvement CSCC focus on treatment of parotid and/or cervical nodes either with surgery alone (parotidectomy and/or neck dissection) or surgery plus adjuvant RT. Some studies included patients receiving concomitant chemotherapy^{56,394-400} or patients who received RT alone.^{96,399,401-403} For studies where the majority of patients receive surgery plus adjuvant RT, recurrence rates are 20% to 35% and estimates of 5-year DFS and DSS are 59% to 83% and 63% to 83%, respectively.^{56,57,394,395,397-399,401,403-408} Many studies support that adjuvant RT improved local regional control, DFS, and OS compared with surgery or RT alone.^{394,395,398,399,401,403-405,408} In contrast, other studies found no significant association between adjuvant RT and improved disease outcomes.^{56,394,406,407} There may be subsets of patients who derive more clinical benefit from adjuvant RT than other patients; however, it is difficult to identify such patients. Results vary for all of the prognostic factors frequently considered such as immunosuppression, primary tumor size, LVI, PNI, differentiation, and features of the regional disease such as ENE and number of involved nodes.^{56,57,394,395,397,398,401,403,405,408,409}



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Several staging systems have been proposed for regional CSCC, as shown in [Table 1](#). O'Brien proposed a staging system that separates parotid involvement from neck LN involvement based on multivariate analysis showing improved local control for P1 compared with P2/P3.⁴¹⁰ Multivariate analyses of survival and locoregional control have yielded favorable^{396,401,403,408} as well as discordant results^{57,396,405} regarding the prognostic value of O'Brien P-stage. O'Brien also showed that survival was significantly better for patients with N0/N1 compared with N2.⁴¹⁰ Two studies supported this result,^{57,408} but several others did not.^{401,403,405} According to the AJCC 7th edition, N1 disease with no ENE had a 5-year cure rate of 92%.¹¹⁶ N2 disease with immunosuppression, in particular patients with transplants or those with hematologic malignancies, on the other hand, had a 5-year survival of 52%, in contrast to 72% for patients who are immunocompetent.⁴¹¹ The AJCC 7th edition staging does not separate parotid from cervical LN involvement and includes both 3-cm and 6-cm cutoffs for largest LN dimension.⁴¹² Forest et al found that LN size was related to ENE, and that 6-cm cutoff and 3-cm cutoff groups performed similarly.³⁹⁴ Risk stratification per the NCCN Guidelines takes into account both ENE^{405,413,414} and margin status after resection^{396,403,405,410,413} as prognostic factors for recurrence and survival. The updated AJCC staging system also includes ENE as a criterion for determining N stage.¹³¹ It should be noted that there are studies that showed no significant association between outcomes and ENE or margin status.^{396,398}

The NCCN-recommended and preferred treatment for CSCC with LN involvement is excision of the primary tumor and regional LN dissection for all surgical candidates. Patients treated with dissection of nodes in the trunk and extremities should consider adjuvant RT of the nodal bed, especially if multiple nodes are involved or if ENE is present. For patients with nodal metastasis to the head and neck, the extent of surgery should depend on the number, location, and size of affected nodes. Postoperative

adjuvant treatment should depend on the pathologic findings after surgery—namely the extent of resection, number of positive nodes, and presence or absence of ENE. Patients with ENE or incompletely excised nodes should receive adjuvant RT and also consider concurrent systemic therapy depending on individual toxicity tolerance. Patients with inoperable nodal disease should be treated with RT of the nodal bed and consider concurrent systemic therapy. Multidisciplinary consultation is recommended for these patients and should consider the systemic therapies used to treat head and neck SCCs as indicated in the NCCN Guidelines for Head and Neck Cancers, available at www.NCCN.org. For symptomatic sites, palliative RT or surgery should be considered. Stereotactic body radiation (SBRT) may be appropriate in select patients.

Satellitosis/In-Transit Metastasis

Satellitosis or in-transit metastasis (S-ITM) is a clinically significant risk factor for the recurrence and defined by the presence of dermal lesions between the primary tumor and lymphatic nodal basins.⁴¹⁵ Due to S-ITM's rarity and therefore scarcity of data, neither the AJCC 8th edition⁴¹⁶ nor the Brigham Women's Hospital (BWH) staging system^{74,75} include it as a CSCC risk factor. The NCCN Panel has strived to delineate treatment planning imaging options as well as primary treatment after a multidisciplinary discussion based on whether the tissue sampling confirmed S-ITM is operable, unresectable, or incompletely resected. Despite these steps S-ITM remains a challenge when encountered in the clinical setting.⁴¹⁷

Recurrence and Metastasis

Metastatic CSCC is rare, estimated at 1.9% to 2.7% of all CSCC, with nodal metastases and distant metastatic disease estimated at 3.7% and 0.4%, respectively.^{11,82} For the management of local tumor recurrence or new regional disease, the algorithm directs clinicians to follow the appropriate pathways for primary treatment. Complicated high-risk tumors,



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

regional recurrence, or the development of distant metastases should be managed by a multidisciplinary tumor board. The NCCN Panel encourages participation in a clinical trial for patients with metastatic CSCC. Unfortunately, such trials are scarce. For symptomatic sites, palliative RT or surgery should be considered. SBRT may be appropriate in select patients. Under highly selective circumstances and in the context of multidisciplinary consultation, resection of limited metastases can be considered.

Systemic Therapy for Distant Metastatic Disease

Unfortunately, evidence regarding systemic therapy for distant metastatic CSCC is limited, except for in the case of the immunotherapy paradigm. Whereas a number of small studies have reported responses to cytotoxic therapy in patients with local or regional CSCC (See *Principles of Systemic Therapy* in the algorithm and *Systemic Therapy Options* below), few of these studies included patients with distant metastatic CSCC.

Systemic Therapy Options

Very-high-risk CSCC with significant risk of extensive local recurrence, nodal, or in-transit metastasis treatment option includes cemiplimab-rwlc as 1) a neoadjuvant therapy after multidisciplinary discussion if the tumor has nonreactive non-keratoacanthomatous rapid growth tumors, in-transit metastasis, is borderline resectable, or if surgery alone may not be curative or result in significant functional limitation; and 2) an adjuvant therapy for patients with extremely high-risk nodal and non-nodal features following adjuvant RT (category 1 preferred).⁴¹⁸

Systemic therapy is also an option that can be considered by a multidisciplinary team for patients with laCSCC, which is defined as primary or recurrent extensive disease where surgery and/or RT may not result in a cure or would possibly produce significant functional impairment.⁴¹⁹ Primary treatment for resectable laCSCC includes

consideration of neoadjuvant cemiplimab-rwlc after multidisciplinary discussion. For patients with laCSCC who cannot undergo surgery due to comorbidities, extent of disease, risk of functional or cosmetic defect, or inability to clear disease with surgery, primary treatment consists of RT with or without systemic therapy or systemic therapy alone if curative RT is not feasible. These primary treatment options also apply to unresectable, inoperable, or incompletely resected palpable regional LNs identified by imaging after a multidisciplinary consultation. Additional treatment for patients with laCSCC and positive margins after resection also includes RT with or without systemic therapy or systemic therapy alone if curative RT is not feasible. Treatment of regional LNs of the head and neck also include RT and consideration of concurrent systemic therapy in certain circumstances.

For patients with confirmed operable S-ITM, neoadjuvant systemic therapy followed by resection or upfront resection with consideration of adjuvant therapy are primary treatment options. Unresectable or incompletely resected S-ITM treatment options include systemic therapy with or without RT or isolated limb infusion/perfusion (ILI/ILP) with melphalan-based regimens (useful in certain circumstances). ILI and ILP are regional chemotherapy techniques used to deliver high-dose treatment to extremities minimizing systemic toxicity.⁴²⁰⁻⁴²² While ILP has comparable higher response rates, ILI is less invasive. A single-institution analysis of a prospective database (N = 163) included 205 ILI procedures with four malignancies treated: melanoma (72% of all ILIs), sarcoma (23%), SCC (2%), and Merkel cell carcinoma (MCC; 3%).⁴²³ Eighty eight percent of patients had a ≤grade II regional Wieberdink toxicity score. The median follow-up was 21.8 months and overall response rate (ORR) was 50.0% for SCC. A retrospective analysis of a prospectively maintained database at two centers identified patients (N = 30) with laCSCC treated with Tumor necrosis factor (TNF) α and melphalan-based ILP (TM-ILP).⁴²⁴ Response could not be determined in three patients. However, after a median follow-



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

up of 25 months, the ORR was 81% with a CR of 59%. Local recurrence occurred in 7 patients, progressive disease was observed in 5 patients, the limb salvage rate was 80%, and OS at 2 years was 67%.

For locoregional disease for which surgery or RT are unlikely to be curative, both cytotoxic and epidermal growth factor receptor (EGFR) inhibitor systemic therapy (monotherapy or combination) have been successfully used to reduce tumor load, which in some cases enabled complete resection or CR with or without concurrent/subsequent RT.⁴²⁵⁻⁴²⁹ In the absence of prospective comparative trial data, it is unclear whether systemic therapy provides additional clinical benefit when used postoperatively with RT. Small retrospective studies were unable to establish definitively that the addition of chemotherapy to postoperative RT significantly improved any disease-related outcome in patients with regional disease,^{397,414,430-432} except for one study.⁴³³ The emergence of anti-programmed cell death protein 1 (PD-1) inhibitors and robust clinical trial data have opened up novel treatment venues for patients with both locally advanced and metastatic CSCC not amenable to surgery and RT. It must be noted that the preferred recommendation for all of these settings is enrollment in a clinical trial.

The preferred systemic therapy for use with RT or surgery for regional recurrence following systemic treatment option recommended by the NCCN Panel is cisplatin.⁴³⁴⁻⁴³⁷ EGFR inhibitors (eg, cetuximab,^{428,436-444} erlotinib,⁴⁴⁵ gefitinib,⁴²⁹ panitumumab⁴⁴⁶) and carboplatin ± paclitaxel^{434,436,447-449} are considered other recommended regimens, while cisplatin + fluorouracil^{425,435,443,444,450} is a regimen useful in certain circumstances. Evidence supporting the efficacy of any of these regimens is mostly limited to case reports and small retrospective studies. In a small (N = 21) prospective phase II study in patients with locally advanced primary or nodal disease who received definitive RT with concurrent cisplatin or carboplatin, the overall CR was reported to be 63%.⁴³⁴ Efficacy

and safety data for cisplatin, cetuximab, or carboplatin + paclitaxel in combination with RT can also be extrapolated from large randomized trials in patients with non-cutaneous head and neck cancers.^{449,451,452} On the other hand, data from a rare, large (N = 321) randomized trial in patients with CSCC of the head and neck testing RT ± carboplatin did not find an added benefit with carboplatin.⁴³⁰

The preferred systemic therapy options if curative RT or surgery is not feasible for locally advanced, recurrent, or metastatic disease for use alone without RT recommended by the NCCN Panel are cemiplimab-rwlc⁴⁵³⁻⁴⁵⁶ and pembrolizumab.^{457,458} Other recommended regimens for systemic therapy alone include nivolumab⁴⁵⁹ or if the patients are ineligible for or progressed on checkpoint inhibitors and clinical trials carboplatin + paclitaxel ± cetuximab⁴⁶⁰⁻⁴⁶⁴ and EGFR inhibitors (eg, cetuximab,^{427,428,437,438,440,442,444,465,466} panitumumab,^{446,467} gefitinib,⁴⁶⁸ dacomitinib,⁴⁶⁹ erlotinib⁴⁷⁰). Regimens that are useful in certain circumstances include neoadjuvant cemiplimab⁴⁵⁶ or if the patients are ineligible for or progressed on checkpoint inhibitors and clinical trials, cisplatin ± fluorouracil^{425,427,435,437,444} and capecitabine.^{471,472}

A phase II, multicenter, nonrandomized study for neoadjuvant cemiplimab-rwlc enrolled 79 patients with CSCC and showed a 51% pathologic CR and 13% pathologic major response in the neoadjuvant setting.⁴⁵⁶ Grade 3 or higher AEs were only observed in 18% of patients. Therefore, the NCCN Panel recommends neoadjuvant cemiplimab-rwlc be discussed as a treatment option for patients whose disease is considered borderline resectable, unresectable, or in which surgery may carry a high morbidity.

Published data reported an ORR of 44% to 54%, CR of 0% to 13%, and partial response (PR) of 31% to 50% to cemiplimab-rwlc in patients with locally advanced, recurrent, or metastatic CSCC.⁴⁵³⁻⁴⁵⁵ Data from the phase II KEYNOTE-629 trial, which included patients with locally advanced, recurrent, or metastatic CSCC, reported an ORR of 34% to



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

50%, a CR of 4% to 17%, and a PR of 25% to 33% for patients treated with pembrolizumab.^{457,458} An open label, single-arm, phase II study evaluated the safety and efficacy of nivolumab as a first-line systemic therapy in individuals with advanced CSCC (N = 24).⁴⁵⁹ A best ORR of 58.3% was achieved, yet no CRs were reported. The study also reported a median follow-up of 17.6 months, and estimated median PFS and OS of 12.7 and 20.7 months, respectively. A median duration of response (DOR) was not reached. Preliminary data and the clinical experience of NCCN Guidelines Panel Members suggest that other anti-PD-1 inhibitors may also be effective in this setting. It was demonstrated in a retrospective study that patients receiving immunotherapy showed a statistically significant better survival compared to those treated with other systemic therapies ($P = .034$)⁴³⁷; the validity of this finding remains to be tested in prospective randomized studies. The use of immune checkpoint inhibitors might perhaps be extended to other indications, with early reports advocating its safety and efficacy concurrently with RT.⁴⁷³

A phase III, randomized trial enrolled 415 patients with local or regional CSCC after postoperative RT at high risk for recurrence due to nodal or non-nodal features.⁴¹⁸ Nodal features were defined as extracapsular extension with largest node ≥ 20 mm in diameter or at least three involved nodes and non-nodal features as in-transit metastases, T4 lesion (with bone invasion), PNI, or locally recurrent tumor with ≥ 1 additional risk factor. Patients received 350 mg of adjuvant cemiplimab or placebo, every 3 weeks for 12 weeks, followed by 700 mg every 6 weeks for up to 36 weeks. At the median follow-up (24 months), the estimated DFS was 87% with cemiplimab and 64% with placebo. Cemiplimab led to lower risks of locoregional recurrence and distant recurrence; however, it also led to higher $\geq$ grade 3 AEs than placebo (23.9% vs. 14.2%).

In the case of EGFR inhibitors, all recommended regimens have been tested in small, single-arm phase II clinical trials among patients with

CSCC nonamenable to surgery and RT. However, low rates of response were documented. Among 25 patients who received cetuximab, 8 PRs and 2 CRs were documented, with a disease control rate of 69%.⁴²⁸ As for gefitinib, among 40 patients treated, the ORR was reported to be 16%, with an additional 35% achieving stable disease at 8 weeks.⁴⁶⁸ In a smaller study of 16 patients with CSCC testing panitumumab, the best ORR (PR and CR) was 31%, with a further 6 patients achieving stable disease.⁴⁶⁷ The response rates reported for 42 patients treated with dacomitinib were 2% CR and 26% PR, with a disease control rate of 86%.⁴⁶⁹ The ORR for 39 patients treated with erlotinib was 10%, with a disease control rate of 71%.⁴⁷⁰ Efficacy data for chemotherapeutic agents are not much better. In a systemic review of 60 patients with metastatic CSCC treated with cisplatin, the CR was reported to be 2%, with an ORR of 45% and median DFS of 14.6 months.⁴⁴⁴ Data supporting carboplatin utility are even more limited, with most studies examining carboplatin combinations and not carboplatin monotherapy.^{437,466}

Follow-up

It has been well-established that 13% to 50% of these patients will develop another SCC within 5 years.⁴⁷⁴⁻⁴⁷⁶ This represents at least a 10-fold increase in risk compared to the general population.^{475,476} Patients with a prior SCC are also at increased risk of developing cutaneous melanoma and BCC, and patients with multiple prior SCCs are at even higher risk.^{68,475} Therefore, continued long-term surveillance of these patients is essential, as is patient education about sun protection and regular self-examination of the skin. Additionally, 70% to 80% of CSCC recurrences develop within 2 years of the initial therapy.^{11,34,60,75,78} Therefore, close follow-up of these patients during this time period is critical.

Patient education is a key component of follow-up for patients who have had CSCC. All patients should be made aware of the various resources that discuss skin cancer prevention. Patients should be educated in strict



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

sun protection and taught how to perform a comprehensive self-examination of the skin. For those who had regional SCC, training in self-examination of LNs is also recommended.

Patients should also be monitored with regular physical exams including complete skin and regional LN examination. The frequency of follow-up should be adjusted based on risk (see *Follow-up* in the algorithm). For following disease, the imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is often sufficient to diagnose local recurrence, but MRI with or without contrast can be considered to assess extent of local disease. For nodal or distant metastases, histologic analysis and/or other imaging modalities can be used for confirmation and to gauge extent of disease. In certain patients at high risk for multiple primary tumors, increased surveillance and consideration of prophylactic measures may be indicated. Follow-up with a dermatologist is strongly recommended if any of the following criteria are met: past or imminent solid organ, marrow, or hematopoietic cell transplant, ≥ 1 cutaneous melanomas in the past 5 years, or ≥ 4 nonmelanoma skin cancers in the past 5 years.

Care for Patients at High Risk of Developing Multiple SCCs

Treatment of SCC in Patients at High Risk

For individuals who rapidly develop multiple CSCC lesions, destructive techniques such as C&E and cryotherapy may be used. Some NCCN Guidelines Panel Members use a combination of shave removal to excise the bulk of the tumor and ensure sufficient material for pathology, and then destructive techniques for margin control. The details of the techniques used to remove CSCC lesions in patients at high risk with multiple lesions vary widely between NCCN Member Institutions and between practitioners at these institutions, and there is no standard language for describing

these methods. Compared to the low-risk population, RT is used more frequently as an adjuvant therapy in patients at high risk and for PNI.⁴⁷⁷ Satellite lesions and in-transit cutaneous metastases may occur more frequently and are more likely to progress in this population.^{478,479}

Analyses of large populations of patients with organ transplants have found that the incidence of new skin cancers is linked to the duration and dose of immunosuppression.⁴⁸⁰⁻⁴⁸² Prospective randomized trials have shown that switching from other immunosuppressants to mTOR inhibitors reduces the risk of developing new CSCC lesions, particularly in patients with a history of one or more CSCCs.⁴⁸³⁻⁴⁹¹ When surgery is impractical due to high CSCC burden, oral capecitabine has been suggested in the transplantation setting, although toxicity is a concern.²⁵⁴

Prevention in Patients at High Risk

Treatment of precancers at first development can help prevent the development of subsequent invasive tumors. Prophylactic treatment may be needed for patients with a history of multiple lesions and/or extensive diffuse AK or field cancerization. Oral retinol and synthetic retinoids (eg, acitretin,²⁵² isotretinoin, etretinate) have been tested in prospective studies in patients at high risk for multiple AKs or SCCs, including patients with transplants,⁴⁹²⁻⁴⁹⁷ xeroderma pigmentosa,⁴⁹⁸ or psoriasis and psoralen plus ultraviolet A (PUVA) exposure.⁴⁹⁹ By comparison with placebo or with SCC incidence during treatment-free periods, data from these studies support that oral retinol and oral retinoids significantly reduce the incidence of new CSCCs in patients at very high risk for multiple lesions.^{493-496,498,499} Outside of these very-high-risk groups, the effectiveness of retinol/retinoid therapy for prophylaxis is less clear.⁵⁰⁰⁻⁵⁰² Side effects may be significant and include mucocutaneous, such as cheilitis, excessive peeling of the skin, and hair disorders,⁴⁹⁷ but musculoskeletal, vascular, hepatic triglyceride, and neurologic AEs have also been reported.^{496,498,501,503} In addition, these



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

agents are teratogenic and must be used with extreme caution in individuals of child-bearing potential.⁵⁰⁴

The NCCN Guidelines do *not* recommend topical retinoids as prophylactic treatment for patients at high risk for multiple AKs or CSCCs. Results of a large randomized trial in patients with a history of ≥ 2 BCCs/SCCs showed that prophylactic topical tretinoin (0.1%) did not reduce the development of new cutaneous BCCs or SCCs compared with vehicle control.⁵⁰⁵ A double-blind randomized study showed that topical tazarotene had a chemopreventative effect in only 6% of patients with basal cell nevus syndrome, a condition associated with frequent development of primary BCCs.⁵⁰⁶

Results from a randomized controlled study suggest that prophylactic nicotinamide may be effective at preventing the development of CSCC recurrence or metastases in patients at high risk.¹⁵¹ Nicotinamide was associated with a 30% reduction in the 12-month rate of new SCCs ($P = .05$), and a 20% reduction in development of new BCCs ($P = .12$) compared to placebo. During the subsequent 6 months after treatment, there was a trend toward increased rates of new SCCs for the nicotinamide arm compared with placebo (-59% relative difference; $P = .07$). Although there are currently no clinical trial data directly comparing nicotinamide with oral retinoids for CSCC prophylaxis, nicotinamide has a much better safety profile. Further clinical research is needed to determine whether nicotinamide provides long-term clinical benefit for patients at risk of developing multiple NMSCs and AK lesions.

Patient Education for Patients at High Risk

Patient education is especially important for those at high risk for CSCC progression or recurrence. Treatment delay is associated with larger tumor size, larger defect size from surgical removal, and more Mohs layers taken to obtain clear margins.^{107,507-510} Significant prognostic factors for patient delay in seeking care include serious comorbidity, low education level,

non-recognition of the seriousness of symptoms, and SCC arising on pre-existing chronic lesions.^{511,512} Low education level is also associated with large NMSC tumor area at presentation.⁵¹³ Educational interventions and physician advice have been shown to increase the likelihood of patients undergoing a complete skin exam, and patients with more knowledge of skin cancer are more likely to get regular complete skin exams.^{390,514,515}

Patient education should begin, in the case of patients with transplants at transplantation and at birth or diagnosis for patients with xeroderma pigmentosum. Education should include discussion of individual risk assessment and the need for stringent sun avoidance and protection methods. Regular sunscreen use can significantly reduce the rate of development of new AKs and CSCCs, and increases remission rates of AKs.¹⁴⁶⁻¹⁴⁹ Knowledge of more than one method for UV protection is associated with higher rates of using some form of protection,⁵¹⁶ as is awareness of susceptibility/risk and overall education level.^{517,518}

Having a prior NMSC has proved to be insufficient motivation for altering patient behavior regarding UV protection and avoidance. Although those with prior NMSC are more likely to use sunscreen and to avoid sun exposure, adoption of preventive measures was low: only 54% used sunscreen, and 20% to 45% used other avoidance/protective methods.⁵¹⁹ Another cross-sectional study showed that tanning bed use was similar among those with and without prior NMSC.⁵²⁰

Randomized trials have shown that educational interventions can effect significant changes in use of solar protection in outdoor workers and patients with transplants.^{515,521-526} While both extensive and repetitive education improve patient knowledge, repetitive education is needed to effect long-term change in patient behavior.⁵²⁷ This is especially important for patients with transplants, as preoccupation with other medical concerns may make them unreceptive to skin cancer education.⁵²⁸ For patients with transplants, an intervention including text messaging reminders was



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

shown to be more effective at improving patient knowledge and changing sun protective behaviors compared with more traditional approaches.⁵²⁹

Monthly self-examination is recommended and should include all skin surfaces and LNs. Patients should be taught the proper method for systematic self-examination of the skin and LNs. A randomized controlled trial has shown that educational intervention increased the frequency and sensitivity of self-examination of the skin among patients with transplants.⁵³⁰ In addition to more frequent and thorough self-examination, follow-up schedules for patients at high risk should be titrated to the frequency of tumor development, and in rare cases may occur weekly.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

Table 1. Staging Systems for Regional CSCC for the Head and Neck

O'Brien 2002 Staging System ⁴¹⁰	
Parotid Stage	
Stage	Criteria
P1	1 LN+ ≤3 cm
P2	1 LN+ >3 and ≤6 cm or ≥2 LN+
P3	1 LN+ >6 cm or Involves VII nerve or skull base
Neck Stage	
Stage	Criteria
N0	No clinical neck disease
N1	1 LN+ ≤3 cm ipsilateral
N2	1 LN+ >3 cm or ≥2 LN+ or ≥1 LN+ contralateral

AJCC 7 th Edition (2009) Regional LN Staging ⁴¹²	
Stage	Criteria
N1	1 LN+ ≤3 cm ipsilateral
N2a	1 LN+ >3 and ≤6 cm ipsilateral
N2b	≥2 LN+ all ≤6 cm ipsilateral
N2c	≥1 LN+ ≤6 cm bilateral/contralateral
N3	≥1 LN+ >6 cm

AJCC 8 th Edition (2017) Regional LN Pathological Staging ¹³¹	
Stage	Criteria ^a
N1	1 LN+ ≤3 cm ENE(-)
N2a	1 LN+, >3 and ≤6 cm ipsilateral ENE(-) or 1 LN+ ≤3 cm ipsilateral ENE(+)
N2b	≥2 LN+ all ≤6 cm ipsilateral ENE(-)
N2c	≥1 LN+ all ≤6 cm bilateral/contralateral ENE(-)
N3a	≥1 LN+ >6 cm ENE(-)
N3b	1 LN+ ≤3 cm ENE(+) contralateral, or ≥1 LN+ >3 cm ipsilateral ENE(+) or ≥2 LN+, any ENE(+)

Forest 2010 N1S3 Staging System ³⁹⁴	
Stage	Criteria
I	1 LN+ ≤3 cm
II	1 LN+ >3 cm or ≥2 LN+ ≤3 cm
III	≥2 LN+ >3 cm

NCCN Guidelines	
Risk Level	Criteria
Low	1 LN+ ≤3 cm ENE(-)
Medium	1 LN+ >3 cm ENE(-) or ≥2 LN+ ENE(-)
High	≥1 LN+ ENE(+) or Incompletely excised disease

ENE(+), with extranodal extension; ENE(-), without extranodal extension; LN+, positive lymph node(s); SCC, squamous cell carcinoma.
^aPathologic criteria.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

References

1. Rubio-Casadevall J, Hernandez-Pujol AM, Ferreira-Santos MC, et al. Trends in incidence and survival analysis in non-melanoma skin cancer from 1994 to 2012 in Girona, Spain: A population-based study. *Cancer Epidemiol* 2016;45:6–10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27639035>.
2. Sella T, Goren I, Shalev V, et al. Incidence trends of keratinocytic skin cancers and melanoma in Israel 2006–11. *Br J Dermatol* 2015;172:202–207. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24976239>.
3. Rudolph C, Schnoor M, Eisemann N, Katalinic A. Incidence trends of nonmelanoma skin cancer in Germany from 1998 to 2010. *J Dtsch Dermatol Ges* 2015;13:788–797. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26213814>.
4. Goon PK, Greenberg DC, Igali L, Levell NJ. Squamous cell carcinoma of the skin has more than doubled over the last decade in the uk. *Acta Derm Venereol* 2016;96:820–821. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26631391>.
5. Robsahm TE, Helsing P, Veierod MB. Cutaneous squamous cell carcinoma in Norway 1963–2011: increasing incidence and stable mortality. *Cancer Med* 2015;4:472–480. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25620456>.
6. Rogers HW, Weinstock MA, Feldman SR, Coldiron BM. Incidence estimate of nonmelanoma skin cancer (keratinocyte carcinomas) in the U.S. population, 2012. *JAMA Dermatol* 2015;151:1081–1086. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25928283>.
7. Callens J, Van Eycken L, Henau K, Garmyn M. Epidemiology of basal and squamous cell carcinoma in Belgium: the need for a uniform and compulsory registration. *J Eur Acad Dermatol Venereol* 2016;30:1912–1918. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27256411>.
8. Muzic JG, Schmitt AR, Wright AC, et al. Incidence and trends of basal cell carcinoma and cutaneous squamous cell carcinoma: A population-based study in Olmsted County, Minnesota, 2000 to 2010. *Mayo Clin Proc* 2017;92:890–898. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28522111>.
9. Perera E, Ganeswaran N, Staines C, et al. Incidence and prevalence of non-melanoma skin cancer in Australia: A systematic review. *Australas J Dermatol* 2015;56:258–267. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25716064>.
10. American Cancer Society. Key statistics for basal and squamous cell skin cancers. Available at: <https://www.cancer.org/cancer/types/basal-and-squamous-cell-skin-cancer/about/key-statistics.html>. Accessed March 2, 2026.
11. Schmults CD, Karia PS, Carter JB, et al. Factors predictive of recurrence and death from cutaneous squamous cell carcinoma: a 10-year, single-institution cohort study. *JAMA Dermatol* 2013;149:541–547. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23677079>.
12. Eisemann N, Jansen L, Castro FA, et al. Survival with nonmelanoma skin cancer in Germany. *Br J Dermatol* 2016;174:778–785. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26676514>.
13. Eigentler TK, Leiter U, Hafner HM, et al. Survival of patients with cutaneous squamous cell carcinoma: Results of a prospective cohort study. *J Invest Dermatol* 2017;137:2309–2315. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28736229>.
14. PubMed Overview. Available at: <https://pubmed.ncbi.nlm.nih.gov/about/>.
15. Freedman-Cass DA, Fischer T, Alpert AB, et al. The value and process of inclusion: Using sensitive, respectful, and inclusive language and images in nccn content. *J Natl Compr Canc Netw* 2023;21:434–441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37156485>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

16. Ramsay HM, Fryer AA, Hawley CM, et al. Factors associated with nonmelanoma skin cancer following renal transplantation in Queensland, Australia. *J Am Acad Dermatol* 2003;49:397–406. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12963901>.
17. Wassberg C, Thorn M, Johansson AM, et al. Increasing incidence rates of squamous cell carcinoma of the skin in Sweden. *Acta Derm Venereol* 2001;81:268–272. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11720174>.
18. Xiang F, Lucas R, Hales S, Neale R. Incidence of nonmelanoma skin cancer in relation to ambient UV radiation in white populations, 1978-2012: empirical relationships. *JAMA Dermatol* 2014;150:1063–1071. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25103031>.
19. Juzeniene A, Grigalavicius M, Baturaite Z, Moan J. Minimal and maximal incidence rates of skin cancer in Caucasians estimated by use of sigmoidal UV dose-incidence curves. *Int J Hyg Environ Health* 2014;217:839–844. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25023193>.
20. Trakatelli M, Barkitzi K, Apap C, et al. Skin cancer risk in outdoor workers: a European multicenter case-control study. *J Eur Acad Dermatol Venereol* 2016;30 Suppl 3:5–11. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26995016>.
21. Xiong MY, Rizzo AE, Cohen TS, et al. Predictors of squamous cell carcinoma in high-risk patients in the VATTC trial. *J Invest Dermatol* 2013;133:1521–1532. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23348836>.
22. Suarez B, Lopez-Abente G, Martinez C, et al. Occupation and skin cancer: the results of the HELIOS-I multicenter case-control study. *BMC Public Health* 2007;7:180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17655745>.
23. Wehner MR, Shive ML, Chren MM, et al. Indoor tanning and non-melanoma skin cancer: systematic review and meta-analysis. *BMJ* 2012;345:e5909. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23033409>.
24. Wehner MR, Chren MM, Nameth D, et al. International prevalence of indoor tanning: A systematic review and meta-analysis. *JAMA Dermatol* 2014;150:390–400. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24477278>.
25. Zanetti R, Rosso S, Martinez C, et al. Comparison of risk patterns in carcinoma and melanoma of the skin in men: A multi-centre case-case-control study. *Br J Cancer* 2006;94:743–751. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16495934>.
26. Green A, Battistutta D, Hart V, et al. Skin cancer in a subtropical Australian population: incidence and lack of association with occupation. The Nambour Study Group. *Am J Epidemiol* 1996;144:1034–1040. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8942434>.
27. English DR, Armstrong BK, Kricger A, et al. Demographic characteristics, pigmentary and cutaneous risk factors for squamous cell carcinoma of the skin: a case-control study. *Int J Cancer* 1998;76:628–634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9610717>.
28. Subramaniam P, Olsen CM, Thompson BS, et al. Anatomical distributions of basal cell carcinoma and squamous cell carcinoma in a population-based study in Queensland, Australia. *JAMA Dermatol* 2016;153:175–182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27892984>.
29. Dika E, Fanti PA, Misciali C, et al. Risk of skin cancer development in 672 patients affected by actinic keratosis. *G Ital Dermatol Venereol* 2016;151:628–633. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26381460>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

30. Criscione VD, Weinstock MA, Naylor MF, et al. Actinic keratoses: Natural history and risk of malignant transformation in the veterans affairs topical tretinoin chemoprevention trial. *Cancer* 2009;115:2523–2530. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19382202>.

31. Eimpunth S, Goldenberg A, Hamman MS, et al. Squamous cell carcinoma in situ upstaged to invasive squamous cell carcinoma: A 5-year, single institution retrospective review. *Dermatol Surg* 2017;43:698–703. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28060173>.

32. Senet P, Combemale P, Debure C, et al. Malignancy and chronic leg ulcers: the value of systematic wound biopsies: a prospective, multicenter, cross-sectional study. *Arch Dermatol* 2012;148:704–708. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22772403>.

33. Combemale P, Bousquet M, Kanitakis J, et al. Malignant transformation of leg ulcers: A retrospective study of 85 cases. *J Eur Acad Dermatol Venereol* 2007;21:935–941. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17659003>.

34. Rowe DE, Carroll RJ, Day CL, Jr. Prognostic factors for local recurrence, metastasis, and survival rates in squamous cell carcinoma of the skin, ear, and lip. Implications for treatment modality selection. *J Am Acad Dermatol* 1992;26:976–990. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1607418>.

35. Eroglu A, Berberoglu U, Berreroğlu S. Risk factors related to locoregional recurrence in squamous cell carcinoma of the skin. *J Surg Oncol* 1996;61:124–130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8606543>.

36. Mullen JT, Feng L, Xing Y, et al. Invasive squamous cell carcinoma of the skin: defining a high-risk group. *Ann Surg Oncol* 2006;13:902–909. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16788750>.

37. Kromberg JG, Castle D, Zwane EM, Jenkins T. Albinism and skin cancer in Southern Africa. *Clin Genet* 1989;36:43–52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2766562>.

38. Mabula JB, Chalya PL, McHembe MD, et al. Skin cancers among Albinos at a University teaching hospital in Northwestern Tanzania: a retrospective review of 64 cases. *BMC Dermatol* 2012;12:5. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22681652>.

39. Yakubu A, Mabogunje OA. Skin cancer in African albinos. *Acta Oncol* 1993;32:621–622. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8260178>.

40. Kraemer KH, Lee MM, Scotto J. Xeroderma pigmentosum. Cutaneous, ocular, and neurologic abnormalities in 830 published cases. *Arch Dermatol* 1987;123:241–250. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3545087>.

41. Bradford PT, Goldstein AM, Tamura D, et al. Cancer and neurologic degeneration in xeroderma pigmentosum: long term follow-up characterises the role of DNA repair. *J Med Genet* 2011;48:168–176. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21097776>.

42. Kraemer KH, Lee MM, Scotto J. DNA repair protects against cutaneous and internal neoplasia: evidence from xeroderma pigmentosum. *Carcinogenesis* 1984;5:511–514. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6705149>.

43. Miranda MB, Lauseker M, Kraus MP, et al. Secondary malignancies in chronic myeloid leukemia patients after imatinib-based treatment: long-term observation in CML Study IV. *Leukemia* 2016;30:1255–1262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26859076>.

44. Kolaitis NA, Duffy E, Zhang A, et al. Voriconazole increases the risk for cutaneous squamous cell carcinoma after lung transplantation. *Transpl Int* 2017;30:41–48. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27678492>.

45. Jensen AO, Svaerke C, Farkas D, et al. Skin cancer risk among solid organ recipients: a nationwide cohort study in Denmark. *Acta Derm Venereol* 2010;90:474–479. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20814621>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

46. Brewer JD, Shanafelt TD, Khezri F, et al. Increased incidence and recurrence rates of nonmelanoma skin cancer in patients with non-Hodgkin lymphoma: a Rochester Epidemiology Project population-based study in Minnesota. *J Am Acad Dermatol* 2015;72:302–309. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25479909>.

47. Gerber SR, Seifert B, Inci I, et al. Exposure to moxifloxacin and cytomegalovirus replication is associated with skin squamous cell carcinoma development in lung transplant recipients. *J Eur Acad Dermatol Venereol* 2015;29:2451–2457. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26403508>.

48. van den Heuvel TR, Wintjens DS, Jeuring SF, et al. Inflammatory bowel disease, cancer and medication: Cancer risk in the Dutch population-based IBDL cohort. *Int J Cancer* 2016;139:1270–1280. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27170593>.

49. Diffey BL, Norval M, Albers PN, Wright CY. The influence of HIV infection on the age dependence of squamous cell carcinoma of the skin in South Africa. *S Afr Med J* 2017;107:127–129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28220738>.

50. Hock BD, McIntosh ND, McKenzie JL, et al. Incidence of cutaneous squamous cell carcinoma in a New Zealand population of chronic lymphocytic leukaemia patients. *Intern Med J* 2016;46:1414–1421. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27664975>.

51. Jensen P, Hansen S, Moller B, et al. Skin cancer in kidney and heart transplant recipients and different long-term immunosuppressive therapy regimens. *J Am Acad Dermatol* 1999;40:177–186. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10025742>.

52. Gupta AK, Cardella CJ, Haberman HF. Cutaneous malignant neoplasms in patients with renal transplants. *Arch Dermatol* 1986;122:1288–1293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3535683>.

53. Garritsen FM, van der Schaft J, van den Reek JM, et al. Risk of non-melanoma skin cancer in patients with atopic dermatitis treated with oral immunosuppressive drugs. *Acta Derm Venereol* 2017;97:724–730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28218345>.

54. Thompson AK, Kelley BF, Prokop LJ, et al. Risk factors for cutaneous squamous cell carcinoma recurrence, metastasis, and disease-specific death: A systematic review and meta-analysis. *JAMA Dermatol* 2016;152:419–428. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26762219>.

55. Brantsch KD, Meisner C, Schonfisch B, et al. Analysis of risk factors determining prognosis of cutaneous squamous-cell carcinoma: a prospective study. *Lancet Oncol* 2008;9:713–720. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18617440>.

56. Schmidt C, Martin JM, Khoo E, et al. Outcomes of nodal metastatic cutaneous squamous cell carcinoma of the head and neck treated in a regional center. *Head Neck* 2015;37:1808–1815. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24995842>.

57. Shao A, Wong DK, McIvor NP, et al. Parotid metastatic disease from cutaneous squamous cell carcinoma: prognostic role of facial nerve sacrifice, lateral temporal bone resection, immune status and P-stage. *Head Neck* 2014;36:545–550. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23780509>.

58. Lott DG, Manz R, Koch C, Lorenz RR. Aggressive behavior of nonmelanotic skin cancers in solid organ transplant recipients. *Transplantation* 2010;90:683–687. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20808266>.

59. Manyam BV, Garsa AA, Chin RI, et al. A multi-institutional comparison of outcomes of immunosuppressed and immunocompetent patients treated with surgery and radiation therapy for cutaneous squamous cell carcinoma of the head and neck. *Cancer* 2017;123:2054–2060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28171708>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

60. Levine DE, Karia PS, Schmults CD. Outcomes of patients with multiple cutaneous squamous cell carcinomas: A 10-year single-institution cohort study. *JAMA Dermatol* 2015;151:1220–1225. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26177278>.
61. Halkud R, Shenoy AM, Naik SM, et al. Xeroderma pigmentosum: clinicopathological review of the multiple oculocutaneous malignancies and complications. *Indian J Surg Oncol* 2014;5:120–124. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25114464>.
62. Maloney ME, Miller SJ. Aggressive vs nonaggressive subtypes (basal cell carcinoma). In: Miller SJ, Maloney ME, eds. *Cutaneous Oncology Pathophysiology, diagnosis, and management*. Malden, MA: Blackwell Science; 1998:609–613.
63. Salasche SJ. Features associated with recurrence (squamous cell carcinoma). In: Miller SJ, Maloney ME, eds. *Cutaneous Oncology Pathophysiology, diagnosis, and management*. Malden, MA: Blackwell Science; 1998:494–499.
64. Akay BN, Saral S, Heper AO, et al. Basosquamous carcinoma: Dermoscopic clues to diagnosis. *J Dermatol* 2017;44:127–134. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27570202>.
65. Wermker K, Roknic N, Goessling K, et al. Basosquamous carcinoma of the head and neck: clinical and histologic characteristics and their impact on disease progression. *Neoplasia* 2015;17:301–305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25810014>.
66. Tan CZ, Rieger KE, Sarin KY. Basosquamous carcinoma: Controversy, advances, and future directions. *Dermatol Surg* 2017;43:23–31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27340741>.
67. Chen J, Ruczinski I, Jorgensen TJ, et al. Nonmelanoma skin cancer and risk for subsequent malignancy. *J Natl Cancer Inst* 2008;100:1215–1222. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18728282>.
68. Banan P, Marvi SK, McMeniman E, De'Ambrosis B. An Australian cohort of 210 patients with multiple invasive squamous cell carcinomas: risk factors and associated increased risk of melanoma and internal malignancies. *Australas J Dermatol* 2016;57:29–32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25970074>.
69. Marrazzo G, Thorpe R, Condie D, et al. Clinical and pathologic factors predictive of positive radiologic findings in high-risk cutaneous squamous cell carcinoma. *Dermatol Surg* 2015;41:1405–1410. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26517320>.
70. Ruiz ES, Karia PS, Morgan FC, Schmults CD. The positive impact of radiologic imaging on high-stage cutaneous squamous cell carcinoma management. *J Am Acad Dermatol* 2017;76:217–225. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27707594>.
71. Gandhi MR, Panizza B, Kennedy D. Detecting and defining the anatomic extent of large nerve perineural spread of malignancy: comparing "targeted" MRI with the histologic findings following surgery. *Head Neck* 2011;33:469–475. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20645285>.
72. Williams LS, Mancuso AA, Mendenhall WM. Perineural spread of cutaneous squamous and basal cell carcinoma: CT and MR detection and its impact on patient management and prognosis. *Int J Radiat Oncol Biol Phys* 2001;49:1061–1069. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11240248>.
73. Nemzek WR, Hecht S, Gandour-Edwards R, et al. Perineural spread of head and neck tumors: how accurate is MR imaging? *AJNR Am J Neuroradiol* 1998;19:701–706. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9576658>.
74. Jambusaria-Pahlajani A, Kanetsky PA, Karia PS, et al. Evaluation of AJCC tumor staging for cutaneous squamous cell carcinoma and a proposed alternative tumor staging system. *JAMA Dermatol* 2013;149:402–410. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23325457>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

75. Karia PS, Jambusaria-Pahlajani A, Harrington DP, et al. Evaluation of american joint committee on cancer, international union against cancer, and brigham and women's hospital tumor staging for cutaneous squamous cell carcinoma. *J Clin Oncol* 2014;32:327–334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24366933>.

76. Schmitt AR, Brewer JD, Bordeaux JS, Baum CL. Staging for cutaneous squamous cell carcinoma as a predictor of sentinel lymph node biopsy results: meta-analysis of American Joint Committee on Cancer criteria and a proposed alternative system. *JAMA Dermatol* 2014;150:19–24. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24226651>.

77. Gonzalez JL, Cunningham K, Silverman R, et al. Comparison of the American Joint Committee on Cancer seventh edition and Brigham and Women's Hospital cutaneous squamous cell carcinoma tumor staging in immunosuppressed patients. *Dermatol Surg* 2017;43:784–791. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28079640>.

78. Roozeboom MH, Lohman BG, Westers-Attema A, et al. Clinical and histological prognostic factors for local recurrence and metastasis of cutaneous squamous cell carcinoma: analysis of a defined population. *Acta Derm Venereol* 2013;93:417–421. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23138613>.

79. Carter JB, Johnson MM, Chua TL, et al. Outcomes of primary cutaneous squamous cell carcinoma with perineural invasion: an 11-year cohort study. *JAMA Dermatol* 2013;149:35–41. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23324754>.

80. Ruiz ES, Karia PS, Besaw R, Schmults CD. Performance of the American Joint Committee on Cancer staging manual, 8th edition vs the Brigham and Women's Hospital tumor classification system for cutaneous squamous cell carcinoma. *JAMA Dermatol* 2019;155:819–825. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30969315>.

81. Karia PS, Morgan FC, Califano JA, Schmults CD. Comparison of tumor classifications for cutaneous squamous cell carcinoma of the head and neck in the 7th vs 8th edition of the AJCC cancer staging manual. *JAMA Dermatol* 2018;154:175–181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29261835>.

82. Brougham ND, Dennett ER, Cameron R, Tan ST. The incidence of metastasis from cutaneous squamous cell carcinoma and the impact of its risk factors. *J Surg Oncol* 2012;106:811–815. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22592943>.

83. Harris BN, Bayoumi A, Rao S, et al. Factors associated with recurrence and regional adenopathy for head and neck cutaneous squamous cell carcinoma. *Otolaryngol Head Neck Surg* 2017;156:863–869. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28322123>.

84. Hollestein LM, de Vries E, Nijsten T. Trends of cutaneous squamous cell carcinoma in the Netherlands: increased incidence rates, but stable relative survival and mortality 1989–2008. *Eur J Cancer* 2012;48:2046–2053. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22342554>.

85. Haisma MS, Plaat BE, Bijl HP, et al. Multivariate analysis of potential risk factors for lymph node metastasis in patients with cutaneous squamous cell carcinoma of the head and neck. *J Am Acad Dermatol* 2016;75:722–730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27473455>.

86. Ross AS, Schmults CD. Sentinel lymph node biopsy in cutaneous squamous cell carcinoma: a systematic review of the English literature. *Dermatol Surg* 2006;32:1309–1321. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17083582>.

87. Lewis KG, Weinstock MA. Nonmelanoma skin cancer mortality (1988–2000): The Rhode Island follow-back study. *Arch Dermatol* 2004;140:837–842. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15262694>.

88. Clayman GL, Lee JJ, Holsinger FC, et al. Mortality risk from squamous cell skin cancer. *J Clin Oncol* 2005;23:759–765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15681519>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

89. Quaedvlieg PJ, Creyten DH, Epping GG, et al. Histopathological characteristics of metastasizing squamous cell carcinoma of the skin and lips. *Histopathology* 2006;49:256–264. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16918972>.

90. Cherpelis BS, Marcusen C, Lang PG. Prognostic factors for metastasis in squamous cell carcinoma of the skin. *Dermatol Surg* 2002;28:268–273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11896781>.

91. Maciburko SJ, Townley WA, Hollowood K, Giele HP. Skin cancers of the hand: a series of 541 malignancies. *Plast Reconstr Surg* 2012;129:1329–1336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22327895>.

92. Moore BA, Weber RS, Prieto V, et al. Lymph node metastases from cutaneous squamous cell carcinoma of the head and neck. *Laryngoscope* 2005;115:1561–1567. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16148695>.

93. Dixon AY, Lee SH, McGregor DH. Histologic features predictive of basal cell carcinoma recurrence: results of a multivariate analysis. *J Cutan Pathol* 1993;20:137–142. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8320358>.

94. Jacobs GH, Rippey JJ, Altini M. Prediction of aggressive behavior in basal cell carcinoma. *Cancer* 1982;49:533–537. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7059912>.

95. de Rosa G, Vetrani A, Zeppa P, et al. Comparative morphometric analysis of aggressive and ordinary basal cell carcinoma of the skin. *Cancer* 1990;65:544–549. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2297645>.

96. de Lima Vazquez V, Sachetto T, Perpetuo NM, Carvalho AL. Prognostic factors for lymph node metastasis from advanced squamous cell carcinoma of the skin of the trunk and extremities. *World J Surg Oncol* 2008;6:73. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18601735>.

97. Wermker K, Kluwig J, Schipmann S, et al. Prediction score for lymph node metastasis from cutaneous squamous cell carcinoma of the external ear. *Eur J Surg Oncol* 2015;41:128–135. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25182221>.

98. Manyam BV, Gastman B, Zhang AY, et al. Inferior outcomes in immunosuppressed patients with high-risk cutaneous squamous cell carcinoma of the head and neck treated with surgery and radiation therapy. *J Am Acad Dermatol* 2015;73:221–227. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26028524>.

99. Tam S, Yao C, Amit M, et al. Association of immunosuppression with outcomes of patients with cutaneous squamous cell carcinoma of the head and neck. *JAMA Otolaryngol Head Neck Surg* 2020;146:128–135. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31804658>.

100. Harwood CA, Proby CM, McGregor JM, et al. Clinicopathologic features of skin cancer in organ transplant recipients: a retrospective case-control series. *J Am Acad Dermatol* 2006;54:290–300. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16443060>.

101. Rabinovics N, Mizrahi A, Hadar T, et al. Cancer of the head and neck region in solid organ transplant recipients. *Head Neck* 2014;36:181–186. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23554129>.

102. Goldenberg A, Ortiz A, Kim SS, Jiang SB. Squamous cell carcinoma with aggressive subclinical extension: 5-year retrospective review of diagnostic predictors. *J Am Acad Dermatol* 2015;73:120–126. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25840729>.

103. Edwards MJ, Hirsch RM, Broadwater JR, et al. Squamous cell carcinoma arising in previously burned or irradiated skin. *Arch Surg* 1989;124:115–117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2910238>.

104. Gul U, Kilic A. Squamous cell carcinoma developing on burn scar. *Ann Plast Surg* 2006;56:406–408. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16557073>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

105. Kowal-Vern A, Criswell BK. Burn scar neoplasms: a literature review and statistical analysis. *Burns* 2005;31:403–413. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15896501>.
106. Kyrgidis A, Tzellos TG, Kechagias N, et al. Cutaneous squamous cell carcinoma (SCC) of the head and neck: risk factors of overall and recurrence-free survival. *Eur J Cancer* 2010;46:1563–1572. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20338745>.
107. Endo Y, Tanioka M, Miyachi Y. Prognostic factors in cutaneous squamous cell carcinoma: is patient delay in hospital visit a predictor of survival? *ISRN Dermatol* 2011;2011:285289. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22363846>.
108. Canueto J, Martin-Vallejo J, Cardenoso-Alvarez E, et al. Rapid growth rate is associated with poor prognosis in cutaneous squamous cell carcinoma. *Clin Exp Dermatol* 2018;43:876–882. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29756221>.
109. Nguyen P, Vin-Christian K, Ming ME, Berger T. Aggressive squamous cell carcinomas in persons infected with the human immunodeficiency virus. *Arch Dermatol* 2002;138:758–763. Available at: <https://pubmed.ncbi.nlm.nih.gov/12056956/>.
110. Winkelhorst JT, Brokelman WJ, Tiggeler RG, Wobbles T. Incidence and clinical course of de-novo malignancies in renal allograft recipients. *Eur J Surg Oncol* 2001;27:409–413. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11417989>.
111. Han A, Ratner D. What is the role of adjuvant radiotherapy in the treatment of cutaneous squamous cell carcinoma with perineural invasion? *Cancer* 2007;109:1053–1059. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17279578>.
112. Galloway TJ, Morris CG, Mancuso AA, et al. Impact of radiographic findings on prognosis for skin carcinoma with clinical perineural invasion. *Cancer* 2005;103:1254–1257. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15693020>.
113. Goepfert H, Dichtel WJ, Medina JE, et al. Perineural invasion in squamous cell skin carcinoma of the head and neck. *Am J Surg* 1984;148:542–547. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6486325>.
114. Durham AB, Lowe L, Malloy KM, et al. Sentinel lymph node biopsy for cutaneous squamous cell carcinoma on the head and neck. *JAMA Otolaryngol Head Neck Surg* 2016;142:1171–1176. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27438434>.
115. Gore SM, Shaw D, Martin RC, et al. Prospective study of sentinel node biopsy for high-risk cutaneous squamous cell carcinoma of the head and neck. *Head Neck* 2016;38 Suppl 1:E884–889. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25965008>.
116. Sapir E, Tolpadi A, McHugh J, et al. Skin cancer of the head and neck with gross or microscopic perineural involvement: Patterns of failure. *Radiother Oncol* 2016;120:81–86. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27475277>.
117. Lin C, Tripcony L, Keller J, et al. Perineural infiltration of cutaneous squamous cell carcinoma and basal cell carcinoma without clinical features. *Int J Radiat Oncol Biol Phys* 2012;82:334–340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21093171>.
118. Jackson JE, Dickie GJ, Wiltshire KL, et al. Radiotherapy for perineural invasion in cutaneous head and neck carcinomas: toward a risk-adapted treatment approach. *Head Neck* 2009;31:604–610. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19132719>.
119. Warren TA, Panizza B, Porceddu SV, et al. Outcomes after surgery and postoperative radiotherapy for perineural spread of head and neck cutaneous squamous cell carcinoma. *Head Neck* 2016;38:824–831. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25546817>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

120. Farasat S, Yu SS, Neel VA, et al. A new American Joint Committee on Cancer staging system for cutaneous squamous cell carcinoma: Creation and rationale for inclusion of tumor (T) characteristics. *J Am Acad Dermatol* 2011;64:1051–1059. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21255868>.

121. Brinkman JN, Hajder E, van der Holt B, et al. The effect of differentiation grade of cutaneous squamous cell carcinoma on excision margins, local recurrence, metastasis, and patient survival: A retrospective follow-up study. *Ann Plast Surg* 2015;75:323–326. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24401812>.

122. Banks ER, Cooper PH. Adenosquamous carcinoma of the skin: a report of 10 cases. *J Cutan Pathol* 1991;18:227–234. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1719048>.

123. Patel NK, McKee PH, Smith NP, Fletcher CD. Primary metaplastic carcinoma (carcinosarcoma) of the skin. A clinicopathologic study of four cases and review of the literature. *Am J Dermatopathol* 1997;19:363–372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9261471>.

124. Nappi O, Pettinato G, Wick MR. Adenoid (acantholytic) squamous cell carcinoma of the skin. *J Cutan Pathol* 1989;16:114–121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2768593>.

125. Ogawa T, Kiuru M, Konia TH, Fung MA. Acantholytic squamous cell carcinoma is usually associated with hair follicles, not acantholytic actinic keratosis, and is not "high risk": Diagnosis, management, and clinical outcomes in a series of 115 cases. *J Am Acad Dermatol* 2017;76:327–333. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27889291>.

126. Garcia C, Crowson AN. Acantholytic squamous cell carcinoma: Is it really a more-aggressive tumor? *Dermatol Surg* 2011;37:353–356. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21410819>.

127. Patel V, Squires SM, Liu DY, Fraga GR. Cutaneous adenosquamous carcinoma: a rare neoplasm with biphasic differentiation. *Cutis* 2014;94:231–233. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25474451>.

128. Cassarino DS, Derienzo DP, Barr RJ. Cutaneous squamous cell carcinoma: a comprehensive clinicopathologic classification--part two. *J Cutan Pathol* 2006;33:261–279. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16630176>.

129. Griffiths RW, Feeley K, Suvarna SK. Audit of clinical and histological prognostic factors in primary invasive squamous cell carcinoma of the skin: assessment in a minimum 5 year follow-up study after conventional excisional surgery. *Br J Plast Surg* 2002;55:287–292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12160533>.

130. College of American Pathologists. Protocol for the examination of specimens from patients with cutaneous squamous cell carcinoma of the head and neck. December 2024; 2024. Available at: https://documents.cap.org/protocols/HN.SCC_1.1.0.0.REL.CAPCP.pdf. Accessed March 2, 2026.

131. Amin MB, Edge S, Greene F, et al., eds. *AJCC cancer staging manual* (ed 8th). New York: Springer International Publishing; 2017.

132. Garcia-Serra A, Hinerman RW, Mendenhall WM, et al. Carcinoma of the skin with perineural invasion. *Head Neck* 2003;25:1027–1033. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14648861>.

133. Lin C, Tripcony L, Keller J, et al. Cutaneous carcinoma of the head and neck with clinical features of perineural infiltration treated with radiotherapy. *Clin Oncol (R Coll Radiol)* 2013;25:362–367. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23489870>.

134. Lawrence N, Cottel WI. Squamous cell carcinoma of skin with perineural invasion. *J Am Acad Dermatol* 1994;31:30–33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8021368>.

135. Kadakia S, Ducic Y, Marra D, Saman M. The role of elective superficial parotidectomy in the treatment of temporal region squamous cell carcinoma. *Oral Maxillofac Surg* 2016;20:143–147. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26685892>.



NCCN Guidelines Version 2.2026 Squamous Cell Skin Cancer

136. Leibovitch I, Huilgol SC, Selva D, et al. Cutaneous squamous cell carcinoma treated with Mohs micrographic surgery in Australia II. Perineural invasion. *J Am Acad Dermatol* 2005;53:261–266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16021121>.

137. Salmon PJ, Hussain W, Geisse JK, et al. Sclerosing squamous cell carcinoma of the skin, an underemphasized locally aggressive variant: a 20-year experience. *Dermatol Surg* 2011;37:664–670. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21269350>.

138. Campoli M, Brodland DG, Zitelli J. A prospective evaluation of the clinical, histologic, and therapeutic variables associated with incidental perineural invasion in cutaneous squamous cell carcinoma. *J Am Acad Dermatol* 2014;70:630–636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24433872>.

139. Balamucki CJ, DeJesus R, Galloway TJ, et al. Impact of radiographic findings on for prognosis skin cancer with perineural invasion. *Am J Clin Oncol* 2015;38:248–251. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23648439>.

140. Willenbrink TJ, Ruiz ES, Cornejo CM, et al. Field cancerization: Definition, epidemiology, risk factors, and outcomes. *J Am Acad Dermatol* 2020;83:709–717. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32387665>.

141. Werner RN, Sammain A, Erdmann R, et al. The natural history of actinic keratosis: A systematic review. *Br J Dermatol* 2013;169:502–518. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23647091>.

142. Ahmed I, Berth-Jones J, Charles-Holmes S, et al. Comparison of cryotherapy with curettage in the treatment of Bowen's disease: a prospective study. *Br J Dermatol* 2000;143:759–766. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11069453>.

143. Hansen JP, Drake AL, Walling HW. Bowen's disease: A four-year retrospective review of epidemiology and treatment at a university center. *Dermatol Surg* 2008;34:878–883. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18363722>.

144. Overmark M, Koskenmies S, Pitkanen S. A retrospective study of treatment of squamous cell carcinoma in situ. *Acta Derm Venereol* 2016;96:64–67. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26073523>.

145. Stewart JR, Lang ME, Brewer JD. Efficacy of nonexcisional treatment modalities for superficially invasive and in situ squamous cell carcinoma: A systematic review and meta-analysis. *J Am Acad Dermatol* 2021;87:131–137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34375669>.

146. Thompson SC, Jolley D, Marks R. Reduction of solar keratoses by regular sunscreen use. *N Engl J Med* 1993;329:1147–1151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8377777>.

147. van der Pols JC, Williams GM, Pandeya N, et al. Prolonged prevention of squamous cell carcinoma of the skin by regular sunscreen use. *Cancer Epidemiol Biomarkers Prev* 2006;15:2546–2548. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17132769>.

148. Ulrich C, Jurgensen JS, Degen A, et al. Prevention of non-melanoma skin cancer in organ transplant patients by regular use of a sunscreen: A 24 months, prospective, case-control study. *Br J Dermatol* 2009;161 Suppl 3:78–84. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19775361>.

149. Green AC, Williams GM, Logan V, Strutton GM. Reduced melanoma after regular sunscreen use: randomized trial follow-up. *J Clin Oncol* 2011;29:257–263. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21135266>.

150. Snidr VA, Damian DL, Halliday GM. Nicotinamide for photoprotection and skin cancer chemoprevention: A review of efficacy and safety. *Exp Dermatol* 2019;28 Suppl 1:15–22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30698874>.

151. Chen AC, Martin AJ, Choy B, et al. A phase 3 randomized trial of nicotinamide for skin-cancer chemoprevention. *N Engl J Med* 2015;373:1618–1626. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26488693>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

152. Allen NC, Martin AJ, Snaird VA, et al. Nicotinamide for skin-cancer chemoprevention in transplant recipients. *N Engl J Med* 2023;388:804–812. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36856616>.

153. Mainville L, Smilga AS, Fortin PR. Effect of nicotinamide in skin cancer and actinic keratoses chemoprophylaxis, and adverse effects related to nicotinamide: A systematic review and meta-analysis. *J Cutan Med Surg* 2022;26:297–308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35134311>.

154. Chen AC, Martin AJ, Dalziel RA, et al. A phase II randomized controlled trial of nicotinamide for skin cancer chemoprevention in renal transplant recipients. *Br J Dermatol* 2016;175:1073–1075. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27061568>.

155. Jorizzo J, Weiss J, Furst K, et al. Effect of a 1-week treatment with 0.5% topical fluorouracil on occurrence of actinic keratosis after cryosurgery: a randomized, vehicle-controlled clinical trial. *Arch Dermatol* 2004;140:813–816. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15262691>.

156. Stockfleth E, von Kiedrowski R, Dominicus R, et al. Efficacy and Safety of 5-Fluorouracil 0.5%/Salicylic Acid 10% in the Field-Directed Treatment of Actinic Keratosis: A Phase III, Randomized, Double-Blind, Vehicle-Controlled Trial. *Dermatol Ther (Heidelb)* 2017;7:81–96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27995485>.

157. Pomerantz H, Hogan D, Eilers D, et al. Long-term efficacy of topical fluorouracil cream, 5%, for treating actinic keratosis: A randomized clinical trial. *JAMA Dermatol* 2015;151:952–960. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25950503>.

158. Rahvar M, Lamel SA, Maibach HI. Randomized, vehicle-controlled trials of topical 5-fluorouracil therapy for actinic keratosis treatment: an overview. *Immunotherapy* 2012;4:939–945. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23046237>.

159. Stockfleth E, Zwingers T, Willers C. Recurrence rates and patient assessed outcomes of 0.5% 5-fluorouracil in combination with salicylic acid treating actinic keratoses. *Eur J Dermatol* 2012;22:370–374. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22494856>.

160. Cunningham TJ, Tabacchi M, Eliane JP, et al. Randomized trial of calcipotriol combined with 5-fluorouracil for skin cancer precursor immunotherapy. *J Clin Invest* 2017;127:106–116. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27869649>.

161. Hadley G, Derry S, Moore RA. Imiquimod for actinic keratosis: systematic review and meta-analysis. *J Invest Dermatol* 2006;126:1251–1255. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16557235>.

162. Gebauer K, Shumack S, Cowen PS. Effect of dosing frequency on the safety and efficacy of imiquimod 5% cream for treatment of actinic keratosis on the forearms and hands: a phase II, randomized placebo-controlled trial. *Br J Dermatol* 2009;161:897–903. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19545297>.

163. Swanson N, Smith CC, Kaur M, Goldenberg G. Imiquimod 2.5% and 3.75% for the treatment of actinic keratoses: two phase 3, multicenter, randomized, double-blind, placebo-controlled studies. *J Drugs Dermatol* 2014;13:166–169. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24509967>.

164. Alomar A, Stockfleth E, Dirschka T, et al. Efficacy and safety of imiquimod 3.75% from Imax in actinic keratosis according to fitzpatrick skin type. *J Drugs Dermatol* 2016;15:285–289. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26954313>.

165. Blauvelt A, Kempers S, Lain E, et al. Phase 3 Trials of Tirbanibulin Ointment for Actinic Keratosis. *N Engl J Med* 2021;384:512–520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33567191>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

166. Freeman M, Vinciullo C, Francis D, et al. A comparison of photodynamic therapy using topical methyl aminolevulinate (Metvix) with single cycle cryotherapy in patients with actinic keratosis: a prospective, randomized study. *J Dermatolog Treat* 2003;14:99–106. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12775317>.

167. Piacquadio DJ, Chen DM, Farber HF, et al. Photodynamic therapy with aminolevulinic acid topical solution and visible blue light in the treatment of multiple actinic keratoses of the face and scalp: investigator-blinded, phase 3, multicenter trials. *Arch Dermatol* 2004;140:41–46. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14732659>.

168. Hauschild A, Stockfleth E, Popp G, et al. Optimization of photodynamic therapy with a novel self-adhesive 5-aminolaevulinic acid patch: results of two randomized controlled phase III studies. *Br J Dermatol* 2009;160:1066–1074. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19222455>.

169. Szeimies RM, Matheson RT, Davis SA, et al. Topical methyl aminolevulinate photodynamic therapy using red light-emitting diode light for multiple actinic keratoses: a randomized study. *Dermatol Surg* 2009;35:586–592. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19309347>.

170. Szeimies RM, Radny P, Sebastian M, et al. Photodynamic therapy with BF-200 ALA for the treatment of actinic keratosis: results of a prospective, randomized, double-blind, placebo-controlled phase III study. *Br J Dermatol* 2010;163:386–394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20518784>.

171. Szeimies RM, Stockfleth E, Popp G, et al. Long-term follow-up of photodynamic therapy with a self-adhesive 5-aminolaevulinic acid patch: 12 months data. *Br J Dermatol* 2010;162:410–414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19804593>.

172. Dirschka T, Radny P, Dominicus R, et al. Long-term (6 and 12 months) follow-up of two prospective, randomized, controlled phase III trials of photodynamic therapy with BF-200 ALA and methyl aminolaevulinate for the treatment of actinic keratosis. *Br J Dermatol* 2013;168:825–836. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23252768>.

173. Pariser DM, Houlihan A, Ferdon MB, et al. Randomized vehicle-controlled study of short drug incubation aminolevulinic acid photodynamic therapy for actinic keratoses of the face or scalp. *Dermatol Surg* 2016;42:296–304. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26863596>.

174. Pariser DM, Lowe NJ, Stewart DM, et al. Photodynamic therapy with topical methyl aminolevulinate for actinic keratosis: results of a prospective randomized multicenter trial. *J Am Acad Dermatol* 2003;48:227–232. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12582393>.

175. Szeimies RM, Karrer S, Radakovic-Fijan S, et al. Photodynamic therapy using topical methyl 5-aminolevulinate compared with cryotherapy for actinic keratosis: A prospective, randomized study. *J Am Acad Dermatol* 2002;47:258–262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12140473>.

176. Morton C, Campbell S, Gupta G, et al. Intraindividual, right-left comparison of topical methyl aminolaevulinate-photodynamic therapy and cryotherapy in subjects with actinic keratoses: a multicentre, randomized controlled study. *Br J Dermatol* 2006;155:1029–1036. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17034536>.

177. Kaufmann R, Spelman L, Weightman W, et al. Multicentre intraindividual randomized trial of topical methyl aminolaevulinate-photodynamic therapy vs. cryotherapy for multiple actinic keratoses on the extremities. *Br J Dermatol* 2008;158:994–999. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18341663>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

178. Serra-Guillen C, Nagore E, Hueso L, et al. A randomized pilot comparative study of topical methyl aminolevulinate photodynamic therapy versus imiquimod 5% versus sequential application of both therapies in immunocompetent patients with actinic keratosis: clinical and histologic outcomes. *J Am Acad Dermatol* 2012;66:e131–137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22226430>.

179. Hadley J, Tristani-Firouzi P, Hull C, et al. Results of an investigator-initiated single-blind split-face comparison of photodynamic therapy and 5% imiquimod cream for the treatment of actinic keratoses. *Dermatol Surg* 2012;38:722–727. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22340282>.

180. Smith S, Piacquadio D, Morhenn V, et al. Short incubation PDT versus 5-FU in treating actinic keratoses. *J Drugs Dermatol* 2003;2:629–635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14711141>.

181. Kurwa HA, Yong-Gee SA, Seed PT, et al. A randomized paired comparison of photodynamic therapy and topical 5-fluorouracil in the treatment of actinic keratoses. *J Am Acad Dermatol* 1999;41:414–418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10459115>.

182. Sotiriou E, Apalla Z, Maliamani F, et al. Intraindividual, right-left comparison of topical 5-aminolevulinic acid photodynamic therapy vs. 5% imiquimod cream for actinic keratoses on the upper extremities. *J Eur Acad Dermatol Venereol* 2009;23:1061–1065. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19470041>.

183. Moggio E, Arisi M, Zane C, et al. A randomized split-face clinical trial analyzing daylight photodynamic therapy with methyl aminolaevulinate vs ingenol mebutate gel for the treatment of multiple actinic keratoses of the face and the scalp. *Photodiagnosis Photodyn Ther* 2016;16:161–165. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27530375>.

184. Zane C, Fabiano A, Arisi M, Calzavara-Pinton P. A randomized split-face clinical trial of photodynamic therapy with methyl aminolevulinate versus ingenol mebutate gel for the treatment of multiple actinic keratoses of the face and scalp. *Dermatology* 2016;232:472–477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27578305>.

185. Berman B, Nestor MS, Newburger J, et al. Treatment of facial actinic keratoses with aminolevulinic acid photodynamic therapy (ALA-PDT) or ingenol mebutate 0.015% gel with and without prior treatment with ALA-PDT. *J Drugs Dermatol* 2014;13:1353–1356. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25607702>.

186. Simon JC, Dominicus R, Karl L, et al. A prospective randomized exploratory study comparing the efficacy of once-daily topical 0.5% 5-fluorouracil in combination with 10.0% salicylic acid (5-FU/SA) vs. cryosurgery for the treatment of hyperkeratotic actinic keratosis. *J Eur Acad Dermatol Venereol* 2015;29:881–889. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25257941>.

187. Krawtchenko N, Roewert-Huber J, Ulrich M, et al. A randomised study of topical 5% imiquimod vs. topical 5-fluorouracil vs. cryosurgery in immunocompetent patients with actinic keratoses: a comparison of clinical and histological outcomes including 1-year follow-up. *Br J Dermatol* 2007;157 Suppl 2:34–40. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18067630>.

188. Foley P, Merlin K, Cumming S, et al. A comparison of cryotherapy and imiquimod for treatment of actinic keratoses: lesion clearance, safety, and skin quality outcomes. *J Drugs Dermatol* 2011;10:1432–1438. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22134568>.

189. Tanghetti E, Werschler P. Comparison of 5% 5-fluorouracil cream and 5% imiquimod cream in the management of actinic keratoses on the face and scalp. *J Drugs Dermatol* 2007;6:144–147. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17373172>.

190. Samorano LP, Torezan LA, Sanches JA. Evaluation of the tolerability and safety of a 0.015% ingenol mebutate gel compared to 5% 5-fluorouracil cream for the treatment of facial actinic keratosis: a prospective randomized trial. *J Eur Acad Dermatol Venereol* 2015;29:1822–1827. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25727104>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

191. Stockfleth E, Sibbring GC, Alarcon I. New topical treatment options for actinic keratosis: A systematic review. *Acta Derm Venereol* 2016;96:17–22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26068001>.
192. Gupta AK, Paquet M. Network meta-analysis of the outcome 'participant complete clearance' in nonimmunosuppressed participants of eight interventions for actinic keratosis: a follow-up on a Cochrane review. *Br J Dermatol* 2013;169:250–259. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23550994>.
193. Gupta AK, Davey V, McPhail H. Evaluation of the effectiveness of imiquimod and 5-fluorouracil for the treatment of actinic keratosis: Critical review and meta-analysis of efficacy studies. *J Cutan Med Surg* 2005;9:209–214. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16502198>.
194. Steeb T, Wessely A, Petzold A, et al. Evaluation of long-term clearance rates of interventions for actinic keratosis: A systematic review and network meta-analysis. *JAMA Dermatol* 2021;157:1066–1077. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34347015>.
195. Steeb T, Wessely A, Schmitz L, et al. Interventions for actinic keratosis in nonscalp and nonface localizations: Results from a systematic review with network meta-analysis. *J Invest Dermatol* 2021;141:345–354 e348. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32645365>.
196. Jansen MHE, Kessels J, Nelemans PJ, et al. Randomized trial of four treatment approaches for actinic keratosis. *N Engl J Med* 2019;380:935–946. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30855743>.
197. Rosenberg AR, Tabacchi M, Ngo KH, et al. Skin cancer precursor immunotherapy for squamous cell carcinoma prevention. *JCI Insight* 2019;4. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30895944>.
198. Warshauer E, Warshauer BL. Clearance of basal cell and superficial squamous cell carcinomas after imiquimod therapy. *J Drugs Dermatol* 2008;7:447–451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18505136>.
199. Love WE, Bernhard JD, Bordeaux JS. Topical imiquimod or fluorouracil therapy for basal and squamous cell carcinoma: a systematic review. *Arch Dermatol* 2009;145:1431–1438. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20026854>.
200. Rosen T, Harting M, Gibson M. Treatment of Bowen's disease with topical 5% imiquimod cream: retrospective study. *Dermatol Surg* 2007;33:427–432. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17430376>.
201. Mackenzie-Wood A, Kossard S, de Launey J, et al. Imiquimod 5% cream in the treatment of Bowen's disease. *J Am Acad Dermatol* 2001;44:462–470. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11209116>.
202. Chitwood K, Etzkorn J, Cohen G. Topical and intralesional treatment of nonmelanoma skin cancer: efficacy and cost comparisons. *Dermatol Surg* 2013;39:1306–1316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23915332>.
203. Patel GK, Goodwin R, Chawla M, et al. Imiquimod 5% cream monotherapy for cutaneous squamous cell carcinoma in situ (Bowen's disease): a randomized, double-blind, placebo-controlled trial. *J Am Acad Dermatol* 2006;54:1025–1032. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16713457>.
204. Bargman H, Hochman J. Topical treatment of Bowen's disease with 5-Fluorouracil. *J Cutan Med Surg* 2003;7:101–105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12447619>.
205. Salim A, Leman JA, McColl JH, et al. Randomized comparison of photodynamic therapy with topical 5-fluorouracil in Bowen's disease. *Br J Dermatol* 2003;148:539–543. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12653747>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

206. Morton C, Horn M, Leman J, et al. Comparison of topical methyl aminolevulinate photodynamic therapy with cryotherapy or Fluorouracil for treatment of squamous cell carcinoma in situ: Results of a multicenter randomized trial. *Arch Dermatol* 2006;142:729–735. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16785375>.

207. McEwan LE, Smith JG. Topical diclofenac/hyaluronic acid gel in the treatment of solar keratoses. *Australas J Dermatol* 1997;38:187–189. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9431711>.

208. Rivers JK, Arlette J, Shear N, et al. Topical treatment of actinic keratoses with 3.0% diclofenac in 2.5% hyaluronan gel. *Br J Dermatol* 2002;146:94–100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11841372>.

209. Gebauer K, Brown P, Varigos G. Topical diclofenac in hyaluronan gel for the treatment of solar keratoses. *Australas J Dermatol* 2003;44:40–43. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12581080>.

210. Zane C, Facchinetti E, Rossi MT, et al. A randomized clinical trial of photodynamic therapy with methyl aminolaevulinate vs. diclofenac 3% plus hyaluronic acid gel for the treatment of multiple actinic keratoses of the face and scalp. *Br J Dermatol* 2014;170:1143–1150. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24506666>.

211. Stockfleth E, Kerl H, Zwingers T, Willers C. Low-dose 5-fluorouracil in combination with salicylic acid as a new lesion-directed option to treat topically actinic keratoses: histological and clinical study results. *Br J Dermatol* 2011;165:1101–1108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21517801>.

212. Whitaker DC. Microscopically proven cure of actinic cheilitis by CO2 laser. *Lasers Surg Med* 1987;7:520–523. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3431329>.

213. Robinson JK. Actinic cheilitis. A prospective study comparing four treatment methods. *Arch Otolaryngol Head Neck Surg* 1989;115:848–852. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2736096>.

214. Laws RA, Wilde JL, Grabski WJ. Comparison of electrodesiccation with CO2 laser for the treatment of actinic cheilitis. *Dermatol Surg* 2000;26:349–353. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10759824>.

215. Satorres Nieto M, Gargallo Albiol J, Gay Escoda C. Surgical management of actinic cheilitis. *Med Oral* 2001;6:205–217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11500638>.

216. Alexiades-Armenakas MR, Geronemus RG. Laser-mediated photodynamic therapy of actinic cheilitis. *J Drugs Dermatol* 2004;3:548–551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15552607>.

217. de Godoy Peres FF, Aigotti Haberbeck Brandao A, Rodarte Carvalho Y, et al. A study of actinic cheilitis treatment by two low-morbidity CO2 laser vaporization one-pass protocols. *Lasers Med Sci* 2009;24:375–385. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18553116>.

218. Sotiriou E, Apalla Z, Chovarda E, et al. Photodynamic therapy with 5-aminolevulinic acid in actinic cheilitis: an 18-month clinical and histological follow-up. *J Eur Acad of Dermatol Venereol* 2009;24:916–920. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20028448>.

219. Lima Gda S, Silva GF, Gomes AP, et al. Diclofenac in hyaluronic acid gel: an alternative treatment for actinic cheilitis. *J Appl Oral Sci* 2010;18:533–537. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21085813>.

220. Sotiriou E, Lallas A, Goussi C, et al. Sequential use of photodynamic therapy and imiquimod 5% cream for the treatment of actinic cheilitis: a 12-month follow-up study. *Br J Dermatol* 2011;165:888–892. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21916880>.

221. Choi SH, Kim KH, Song KH. Efficacy of ablative fractional laser-assisted photodynamic therapy for the treatment of actinic cheilitis: 12-month follow-up results of a prospective, randomized, comparative trial. *Br J Dermatol* 2015;173:184–191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25400009>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

222. Suarez-Perez JA, Lopez-Navarro N, Herrera-Acosta E, et al. Treatment of actinic cheilitis with methyl aminolevulinate photodynamic therapy and light fractionation: a prospective study of 10 patients. *Eur J Dermatol* 2015;25:623–624. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26677980>.

223. Chaves YN, Torezan LA, Lourenco SV, Neto CF. Evaluation of the efficacy of photodynamic therapy for the treatment of actinic cheilitis. *Photodermatol Photoimmunol Photomed* 2017;33:14–21. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27864997>.

224. Zane C, Facchinetti E, Rossi MT, et al. Cryotherapy is preferable to ablative CO2 laser for the treatment of isolated actinic keratoses of the face and scalp: a randomized clinical trial. *Br J Dermatol* 2014;170:1114–1121. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24472087>.

225. Ostertag JU, Quaedvlieg PJ, van der Geer S, et al. A clinical comparison and long-term follow-up of topical 5-fluorouracil versus laser resurfacing in the treatment of widespread actinic keratoses. *Lasers Surg Med* 2006;38:731–739. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16912977>.

226. Scola N, Terras S, Georgas D, et al. A randomized, half-side comparative study of aminolaevulinate photodynamic therapy vs. CO(2) laser ablation in immunocompetent patients with multiple actinic keratoses. *Br J Dermatol* 2012;167:1366–1373. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22708798>.

227. Holzer G, Pinkowicz A, Radakovic S, et al. Randomized controlled trial comparing 35% trichloroacetic acid peel and 5-aminolaevulinic acid photodynamic therapy for treating multiple actinic keratosis. *Br J Dermatol* 2017;176:1155–1161. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28012181>.

228. Di Nuzzo S, Cortelazzi C, Boccaletti V, et al. Comparative study of trichloroacetic acid vs. photodynamic therapy with topical 5-aminolevulinic acid for actinic keratosis of the scalp. *Photodermatol Photoimmunol Photomed* 2015;31:233–238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25660106>.

229. Lawrence N, Cox SE, Cockerell CJ, et al. A comparison of the efficacy and safety of Jessner's solution and 35% trichloroacetic acid vs 5% fluorouracil in the treatment of widespread facial actinic keratoses. *Arch Dermatol* 1995;131:176–181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7857114>.

230. Witheiler DD, Lawrence N, Cox SE, et al. Long-term efficacy and safety of Jessner's solution and 35% trichloroacetic acid vs 5% fluorouracil in the treatment of widespread facial actinic keratoses. *Dermatol Surg* 1997;23:191–196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9145962>.

231. Hantash BM, Stewart DB, Cooper ZA, et al. Facial resurfacing for nonmelanoma skin cancer prophylaxis. *Arch Dermatol* 2006;142:976–982. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16924046>.

232. Kuflik EG, Gage AA. The five-year cure rate achieved by cryosurgery for skin cancer. *J Am Acad Dermatol* 1991;24:1002–1004. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1820761>.

233. Fraunfelder FT, Zacarian SA, Limmer BL, Wingfield D. Cryosurgery for malignancies of the eyelid. *Ophthalmology* 1980;87:461–465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7413134>.

234. Lindemalm-Lundstam B, Dalenback J. Prospective follow-up after curettage-cryosurgery for scalp and face skin cancers. *Br J Dermatol* 2009;161:568–576. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19624544>.

235. Holt PJ. Cryotherapy for skin cancer: results over a 5-year period using liquid nitrogen spray cryosurgery. *Br J Dermatol* 1988;119:231–240. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3166941>.

236. Cox NH, Dyson P. Wound healing on the lower leg after radiotherapy or cryotherapy of Bowen's disease and other malignant skin lesions. *Br J Dermatol* 1995;133:60–65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7669642>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

237. Morton CA, Whitehurst C, Moseley H, et al. Comparison of photodynamic therapy with cryotherapy in the treatment of Bowen's disease. *Br J Dermatol* 1996;135:766–771. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8977678>.

238. Peikert JM. Prospective trial of curettage and cryosurgery in the management of non-facial, superficial, and minimally invasive basal and squamous cell carcinoma. *Int J Dermatol* 2011;50:1135–1138. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22126880>.

239. Cairnduff F, Stringer MR, Hudson EJ, et al. Superficial photodynamic therapy with topical 5-aminolaevulinic acid for superficial primary and secondary skin cancer. *Br J Cancer* 1994;69:605–608. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8123497>.

240. Morton CA, Whitehurst C, Moore JV, MacKie RM. Comparison of red and green light in the treatment of Bowen's disease by photodynamic therapy. *Br J Dermatol* 2000;143:767–772. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11069454>.

241. Morton CA, Whitehurst C, McColl JH, et al. Photodynamic therapy for large or multiple patches of Bowen disease and basal cell carcinoma. *Arch Dermatol* 2001;137:319–324. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11255332>.

242. Varma S, Wilson H, Kurwa HA, et al. Bowen's disease, solar keratoses and superficial basal cell carcinomas treated by photodynamic therapy using a large-field incoherent light source. *Br J Dermatol* 2001;144:567–574. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11260016>.

243. Lui H, Hobbs L, Tope WD, et al. Photodynamic therapy of multiple nonmelanoma skin cancers with verteporfin and red light-emitting diodes: two-year results evaluating tumor response and cosmetic outcomes. *Arch Dermatol* 2004;140:26–32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14732656>.

244. Haddad R, Nesher E, Weiss J, et al. Photodynamic therapy for Bowen's disease and squamous cell carcinoma of the skin. *Photodiagnosis Photodyn Ther* 2004;1:225–230. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25048336>.

245. Calzavara-Pinton PG, Venturini M, Sala R, et al. Methylaminolaevulinate-based photodynamic therapy of Bowen's disease and squamous cell carcinoma. *Br J Dermatol* 2008;159:137–144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18489606>.

246. Lopez N, Meyer-Gonzalez T, Herrera-Acosta E, et al. Photodynamic therapy in the treatment of extensive Bowen's disease. *J Dermatolog Treat* 2012;23:428–430. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21787214>.

247. Truchuelo M, Fernandez-Guarino M, Fleita B, et al. Effectiveness of photodynamic therapy in Bowen's disease: an observational and descriptive study in 51 lesions. *J Eur Acad Dermatol Venereol* 2012;26:868–874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21740466>.

248. Ko DY, Kim KH, Song KH. A randomized trial comparing methyl aminolaevulinate photodynamic therapy with and without Er:YAG ablative fractional laser treatment in Asian patients with lower extremity Bowen disease: results from a 12-month follow-up. *Br J Dermatol* 2014;170:165–172. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24102369>.

249. Choi SH, Kim KH, Song KH. Effect of methyl aminolevulinate photodynamic therapy with and without ablative fractional laser treatment in patients with microinvasive squamous cell carcinoma: A randomized clinical trial. *JAMA Dermatol* 2017;153:289–295. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28199463>.

250. Zaar O, Fouglerberg J, Hermansson A, et al. Effectiveness of photodynamic therapy in Bowen's disease: A retrospective observational study in 423 lesions. *J Eur Acad Dermatol Venereol* 2017;31:1289–1294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28190258>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

251. Souza CS, Felicio LB, Ferreira J, et al. Long-term follow-up of topical 5-aminolaevulinic acid photodynamic therapy diode laser single session for non-melanoma skin cancer. *Photodiagnosis Photodyn Ther* 2009;6:207–213. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19932453>.

252. Badri O, Schmults CD, Karia PS, Ruiz ES. Efficacy and cost analysis for acitretin for basal and squamous cell carcinoma prophylaxis in renal transplant recipients. *Dermatol Surg* 2021;47:125–126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32371782>.

253. Cornejo CM, Jambusaria-Pahlajani A, Willenbrink TJ, et al. Field cancerization: Treatment. *J Am Acad Dermatol* 2020;83:719–730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32387663>.

254. Endrizzi B, Ahmed RL, Ray T, et al. Capecitabine to reduce nonmelanoma skin carcinoma burden in solid organ transplant recipients. *Dermatol Surg* 2013;39:634–645. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23379978>.

255. Lansbury L, Bath-Hextall F, Perkins W, et al. Interventions for non-metastatic squamous cell carcinoma of the skin: systematic review and pooled analysis of observational studies. *BMJ* 2013;347:f6153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24191270>.

256. Chren MM, Linos E, Torres JS, et al. Tumor recurrence 5 years after treatment of cutaneous basal cell carcinoma and squamous cell carcinoma. *J Invest Dermatol* 2013;133:1188–1196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23190903>.

257. Freeman RG, Knox JM, Heaton CL. The treatment of skin cancer: A statistical study of 1,341 skin tumors comparing results obtained with irradiation, surgery, and curettage followed by electrodesiccation. *Cancer* 1964;17:535–538. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14136537>.

258. Honeycutt WM, Jansen GT. Treatment of squamous cell carcinoma of the skin. *Arch Dermatol* 1973;108:670–672. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/4750203>.

259. Shiffman NJ. Squamous cell carcinomas of the skin of the pinna. *Can J Surg* 1975;18:279–283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1125865>.

260. Gniadecki R, Glud M, Mortensen K, et al. Favourable results of Mohs micrographic surgery for basal cell carcinoma. *Dan Med J* 2015;62:A5171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26621396>.

261. Muller FM, Dawe RS, Moseley H, Fleming CJ. Randomized comparison of Mohs micrographic surgery and surgical excision for small nodular basal cell carcinoma: tissue-sparing outcome. *Dermatol Surg* 2009;35:1349–1354. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19500127>.

262. Curtis KK, Fakult NJ, Strunck JL, et al. Establishing consensus for mohs micrographic surgical techniques in the treatment of melanoma in situ for future clinical trials: A modified delphi study. *J Natl Compr Canc Netw* 2024;22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39079545>.

263. Xiong DD, Beal BT, Varra V, et al. Outcomes in intermediate-risk squamous cell carcinomas treated with Mohs micrographic surgery compared with wide local excision. *J Am Acad Dermatol* 2020;82:1195–1204. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31887322>.

264. van Lee CB, Roorda BM, Wakkee M, et al. Recurrence rates of cutaneous squamous cell carcinoma of the head and neck after Mohs micrographic surgery vs. standard excision: a retrospective cohort study. *Br J Dermatol* 2018;181:338–343. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30199574>.

265. Turner RJ, Leonard N, Malcolm AJ, et al. A retrospective study of outcome of Mohs' micrographic surgery for cutaneous squamous cell carcinoma using formalin fixed sections. *Br J Dermatol* 2000;142:752–757. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10792227>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

266. Malhotra R, Huilgol SC, Huynh NT, Selva D. The Australian Mohs database: periocular squamous cell carcinoma. *Ophthalmology* 2004;111:617–623. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15051191>.

267. Leibovitch I, Huilgol SC, Selva D, et al. Cutaneous squamous cell carcinoma treated with Mohs micrographic surgery in Australia I. Experience over 10 years. *J Am Acad Dermatol* 2005;53:253–260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16021120>.

268. Leibovitch I, Huilgol SC, Richards S, et al. Scalp tumors treated with Mohs micrographic surgery: clinical features and surgical outcome. *Dermatol Surg* 2006;32:1369–1374. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17083590>.

269. Silapunt S, Peterson SR, Goldberg LH. Squamous cell carcinoma of the auricle and Mohs micrographic surgery. *Dermatol Surg* 2005;31:1423–1427. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16416611>.

270. van der Eerden PA, Prins ME, Lohuis PJ, et al. Eighteen years of experience in Mohs micrographic surgery and conventional excision for nonmelanoma skin cancer treated by a single facial plastic surgeon and pathologist. *Laryngoscope* 2010;120:2378–2384. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21046543>.

271. Pugliano-Mauro M, Goldman G. Mohs surgery is effective for high-risk cutaneous squamous cell carcinoma. *Dermatol Surg* 2010;36:1544–1553. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21053415>.

272. Skaria AM. Recurrence of basosquamous carcinoma after Mohs micrographic surgery. *Dermatology* 2010;221:352–355. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20924160>.

273. Breuninger H, Schaumburg-Lever G. Control of excisional margins by conventional histopathological techniques in the treatment of skin tumours. An alternative to Mohs' technique. *J Pathol* 1988;154:167–171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2832578>.

274. Vuyk HD, Lohuis PJ. Mohs micrographic surgery for facial skin cancer. *Clin Otolaryngol Allied Sci* 2001;26:265–273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11559334>.

275. Mourouzis C, Boynton A, Grant J, et al. Cutaneous head and neck SCCs and risk of nodal metastasis - UK experience. *J Craniomaxillofac Surg* 2009;37:443–447. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19713116>.

276. Matsumoto A, Li JN, Matsumoto M, et al. Factors predicting outcomes of patients with high-risk squamous cell carcinoma treated with Mohs micrographic surgery. *J Am Acad Dermatol* 2021;85:588–595. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33524409>.

277. Tschetter AJ, Campoli MR, Zitelli JA, Brodland DG. Long-term clinical outcomes of patients with invasive cutaneous squamous cell carcinoma treated with Mohs micrographic surgery: A 5-year, multicenter, prospective cohort study. *J Am Acad Dermatol* 2020;82:139–148. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31279037>.

278. Marrazzo G, Zitelli JA, Brodland D. Clinical outcomes in high-risk squamous cell carcinoma patients treated with Mohs micrographic surgery alone. *J Am Acad Dermatol* 2019;80:633–638. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30244064>.

279. Kropp L, Balamucki CJ, Morris CG, et al. Mohs resection and postoperative radiotherapy for head and neck cancers with incidental perineural invasion. *Am J Otolaryngol* 2013;34:373–377. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23415573>.

280. Ruiz ES, Koyfman SA, Que SKT, et al. Evaluation of the utility of localized adjuvant radiation for node-negative primary cutaneous squamous cell carcinoma with clear histologic margins. *J Am Acad Dermatol* 2020;82:420–429. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31349042>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

281. Miller J, Chang T, Schwartz D, et al. Outcomes of adjuvant radiotherapy following negative surgical margins for cutaneous squamous cell carcinoma. *Dermatol Surg* 2019;45:1111–1116. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30893176>.

282. Koslosky CL, El Tal AK, Workman B, et al. Reliability of skin biopsies in determining accurate tumor margins: a retrospective study after Mohs micrographic surgery. *Dermatol Surg* 2014;40:964–970. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25099294>.

283. Chambers KJ, Kraft S, Emerick K. Evaluation of frozen section margins in high-risk cutaneous squamous cell carcinomas of the head and neck. *Laryngoscope* 2015;125:636–639. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25230253>.

284. Castley AJ, Theile DR, Lambie D. The use of frozen section in the excision of cutaneous malignancy: a Queensland experience. *Ann Plast Surg* 2013;71:386–389. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24025654>.

285. Massey PR, Gupta S, Rothstein BE, et al. Total margin-controlled excision is superior to standard excision for keratinocyte carcinoma on the nose: A veterans affairs nested cohort study. *Ann Surg Oncol* 2021;28:3656–3663. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33742233>.

286. Gayre GS, Hybarger CP, Mannor G, et al. Outcomes of excision of 1750 eyelid and periocular skin basal cell and squamous cell carcinomas by modified en face frozen section margin-controlled technique. *Int Ophthalmol Clin* 2009;49:97–110. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20348860>.

287. de Visscher JG, Gooris PJ, Vermey A, Roodenburg JL. Surgical margins for resection of squamous cell carcinoma of the lower lip. *Int J Oral Maxillofac Surg* 2002;31:154–157. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12102412>.

288. Bogdanov-Berezovsky A, Rosenberg L, Cagniano E, Silberstein E. The role of frozen section histological analysis in the treatment of head and neck skin basal and squamous cell carcinomas. *Isr Med Assoc J* 2008;10:344–345. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18605355>.

289. Moncrieff MD, Shah AK, Igali L, Garioch JJ. False-negative rate of intraoperative frozen section margin analysis for complex head and neck nonmelanoma skin cancer excisions. *Clin Exp Dermatol* 2015;40:834–838. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26290360>.

290. Emmett AJ, Broadbent GD. Shave excision of superficial solar skin lesions. *Plast Reconstr Surg* 1987;80:47–54. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3602160>.

291. Dando EE, Abban C, Shehu Wingrove A, et al. Deep shave removal of suspected basal cell carcinoma: A prospective study. *Dermatol Surg* 2023;49:130–134. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36728062>.

292. Wu X, Elkin EB, Jason Chen CS, Marghoob A. Traditional versus streamlined management of basal cell carcinoma (BCC): A cost analysis. *J Am Acad Dermatol* 2015;73:791–798. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26341142>.

293. Abramson AK, Krasny MJ, Goldman GD. Tangential shave removal of basal cell carcinoma. *Dermatol Surg* 2013;39:387–392. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23279298>.

294. Baker NJ, Webb AA, Macpherson D. Surgical management of cutaneous squamous cell carcinoma of the head and neck. *Br J Oral Maxillofac Surg* 2001;39:87–90. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11286440>.

295. Ang P, Tan AW, Goh CL. Comparison of completely versus incompletely excised cutaneous squamous cell carcinomas. *Ann Acad Med Singapore* 2004;33:68–70. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15008566>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

296. Nemet AY, Deckel Y, Martin PA, et al. Management of periocular basal and squamous cell carcinoma: a series of 485 cases. *Am J Ophthalmol* 2006;142:293–297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16876511>.

297. Seretis K, Thomaidis V, Karpouzis A, et al. Epidemiology of surgical treatment of nonmelanoma skin cancer of the head and neck in Greece. *Dermatol Surg* 2010;36:15–22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19912277>.

298. Jenkins G, Smith AB, Kanatas AN, et al. Anatomical restrictions in the surgical excision of scalp squamous cell carcinomas: does this affect local recurrence and regional nodal metastases? *Int J Oral Maxillofac Surg* 2014;43:142–146. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24128939>.

299. Thomas SS, Matthews RN. Squamous cell carcinoma of the pinna: a 6-year study. *Br J Plast Surg* 1994;47:81–85. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8149063>.

300. Batra RS, Kelley LC. Predictors of extensive subclinical spread in nonmelanoma skin cancer treated with Mohs micrographic surgery. *Arch Dermatol* 2002;138:1043–1051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12164742>.

301. Stewart TJ, Saunders A. Risk factors for positive margins after wide local excision of cutaneous squamous cell carcinoma. *J Dermatolog Treat* 2018;29:1–3. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29448853>.

302. Genders RE, Marsidi N, Michi M, et al. Incomplete excision of cutaneous squamous cell carcinoma; systematic review of the literature. *Acta Derm Venereol* 2020;100:adv00084. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32128598>.

303. Bogdanov-Berezovsky A, Cohen AD, Glesinger R, et al. Risk factors for incomplete excision of squamous cell carcinomas. *J Dermatolog Treat* 2005;16:341–344. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16428157>.

304. Tan PY, Ek E, Su S, et al. Incomplete excision of squamous cell carcinoma of the skin: a prospective observational study. *Plast Reconstr Surg* 2007;120:910–916. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17805118>.

305. Hansen C, Wilkinson D, Hansen M, Soyer HP. Factors contributing to incomplete excision of nonmelanoma skin cancer by Australian general practitioners. *Arch Dermatol* 2009;145:1253–1260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19917954>.

306. Mirshams M, Razzaghi M, Noormohammadpour P, et al. Incidence of incomplete excision in surgically treated cutaneous squamous cell carcinoma and identification of the related risk factors. *Acta Med Iran* 2011;49:806–809. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22174169>.

307. Khan AA, Potter M, Cubitt JJ, et al. Guidelines for the excision of cutaneous squamous cell cancers in the United Kingdom: the best cut is the deepest. *J Plast Reconstr Aesthet Surg* 2013;66:467–471. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23352886>.

308. Carducci M, Bozzetti M, de Marco G, et al. Preoperative margin detection by digital dermoscopy in the traditional surgical excision of cutaneous squamous cell carcinomas. *J Dermatolog Treat* 2013;24:221–226. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22390630>.

309. Ribero S, Osella Abate S, Di Capua C, et al. Squamocellular carcinoma of the skin: Clinicopathological features predicting the involvement of the surgical margins and review of the literature. *Dermatology* 2016;232:279–284. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27028227>.

310. Riml S, Larcher L, Kompatscher P. Complete excision of nonmelanotic skin cancer: a matter of surgical experience. *Ann Plast Surg* 2013;70:66–69. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21712697>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

311. Brodland DG, Zitelli JA. Surgical margins for excision of primary cutaneous squamous cell carcinoma. *J Am Acad Dermatol* 1992;27:241–248. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1430364>.

312. Schell AE, Russell MA, Park SS. Suggested excisional margins for cutaneous malignant lesions based on Mohs micrographic surgery. *JAMA Facial Plast Surg* 2013;15:337–343. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23744451>.

313. Stratigos AJ, Garbe C, Dessinioti C, et al. European interdisciplinary guideline on invasive squamous cell carcinoma of the skin: Part 1. epidemiology, diagnostics and prevention. *Eur J Cancer* 2020;128:60–82. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32113941>.

314. Stratigos AJ, Garbe C, Dessinioti C, et al. European interdisciplinary guideline on invasive squamous cell carcinoma of the skin: Part 2. Treatment. *Eur J Cancer* 2020;128:83–102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32113942>.

315. Keohane SG, Botting J, Budny PG, et al. British Association of Dermatologists guidelines for the management of people with cutaneous squamous cell carcinoma 2020. *Br J Dermatol* 2021;184:401–414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33150585>.

316. Newlands C, Currie R, Memon A, et al. Non-melanoma skin cancer: United kingdom national multidisciplinary guidelines. *J Laryngol Otol* 2016;130:S125–S132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27841126>.

317. Bovill ES, Cullen KW, Barrett W, Banwell PE. Clinical and histological findings in re-excision of incompletely excised cutaneous squamous cell carcinoma. *J Plast Reconstr Aesthet Surg* 2009;62:457–461. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18218349>.

318. Yuan Y, Duff ML, Sammons DL, Wu S. Retrospective chart review of skin cancer presence in the wide excisions. *World J Clin Cases* 2014;2:52–56. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24653985>.

319. Stewart CM, Garlick J, McMullin J, et al. Surgical excision of non-melanoma skin cancer in an elderly veteran's affairs population. *Plast Reconstr Surg Glob Open* 2014;2:e277. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25587511>.

320. Abbattuelli JS, Boulter N, Laforge T, Lozier JC. Radiation therapy of skin carcinomas: results of a hypofractionated irradiation schedule in 675 cases followed more than 2 years. *Radiother Oncol* 1989;14:113–119. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2710943>.

321. Kwan W, Wilson D, Moravan V. Radiotherapy for locally advanced basal cell and squamous cell carcinomas of the skin. *Int J Radiat Oncol Biol Phys* 2004;60:406–411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15380573>.

322. Hernandez-Machin B, Borrego L, Gil-Garcia M, Hernandez BH. Office-based radiation therapy for cutaneous carcinoma: evaluation of 710 treatments. *Int J Dermatol* 2007;46:453–459. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17472670>.

323. Barysch MJ, Eggmann N, Beyeler M, et al. Long-term recurrence rate of large and difficult to treat cutaneous squamous cell carcinomas after superficial radiotherapy. *Dermatology* 2012;224:59–65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22433440>.

324. Cognetta AB, Howard BM, Heaton HP, et al. Superficial x-ray in the treatment of basal and squamous cell carcinomas: a viable option in select patients. *J Am Acad Dermatol* 2012;67:1235–1241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22818756>.

325. Schulte KW, Lippold A, Auras C, et al. Soft x-ray therapy for cutaneous basal cell and squamous cell carcinomas. *J Am Acad Dermatol* 2005;53:993–1001. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16310060>.

326. Krausz AE, Ji-Xu A, Smile T, et al. A systematic review of primary, adjuvant, and salvage radiation therapy for cutaneous squamous cell carcinoma. *Dermatol Surg* 2021;47:587–592. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33577212>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

327. Lovett RD, Perez CA, Shapiro SJ, Garcia DM. External irradiation of epithelial skin cancer. *Int J Radiat Oncol Biol Phys* 1990;19:235–242. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2394605>.

328. Dupree MT, Kiteley RA, Weismantle K, et al. Radiation therapy for Bowen's disease: lessons for lesions of the lower extremity. *J Am Acad Dermatol* 2001;45:401–404. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11511838>.

329. Lukas VanderSpek LA, Pond GR, Wells W, Tsang RW. Radiation therapy for Bowen's disease of the skin. *Int J Radiat Oncol Biol Phys* 2005;63:505–510. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16168842>.

330. Herman JM, Pierce LJ, Sandler HM, et al. Radiotherapy using a water bath in the treatment of Bowen's disease of the digit. *Radiother Oncol* 2008;88:398–402. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18571754>.

331. Zhang J, Wang Y, Wijaya WA, et al. Efficacy and prognostic factors of adjuvant radiotherapy for cutaneous squamous cell carcinoma: A systematic review and meta-analysis. *J Eur Acad Dermatol Venereol* 2021;35:1777–1787. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33930213>.

332. Najim M, Cross S, Gebiski V, et al. Early-stage squamous cell carcinoma of the lip: the Australian experience and the benefits of radiotherapy in improving outcome in high-risk patients after resection. *Head Neck* 2013;35:1426–1430. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22965889>.

333. Babington S, Veness MJ, Cakir B, et al. Squamous cell carcinoma of the lip: is there a role for adjuvant radiotherapy in improving local control following incomplete or inadequate excision? *ANZ J Surg* 2003;73:621–625. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12887533>.

334. Raza SM, Ramakrishna R, Weber RS, et al. Nonmelanoma cutaneous cancers involving the skull base: outcomes of aggressive multimodal management. *J Neurosurg* 2015;123:1–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25909577>.

335. Harris BN, Pipkorn P, Nguyen KNB, et al. Association of adjuvant radiation therapy with survival in patients with advanced cutaneous squamous cell carcinoma of the head and neck. *JAMA Otolaryngol Head Neck Surg* 2018;145:153–158. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30570645>.

336. Canueto J, Jaka A, Corchete LA, et al. Postoperative radiotherapy provides better local control and long-term outcome in selective cases of cutaneous squamous cell carcinoma with perineural invasion. *J Eur Acad Dermatol Venereol* 2020;34:1080–1091. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31587379>.

337. Ruiz ES, Kus KJB, Smile TD, et al. Adjuvant radiation following clear margin resection of high T-stage cutaneous squamous cell carcinoma halves the risk of local and locoregional recurrence: A dual center retrospective study. *J Am Acad Dermatol* 2022;87:87–94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35364211>.

338. Jambusaria-Pahlajani A, Miller CJ, Quon H, et al. Surgical monotherapy versus surgery plus adjuvant radiotherapy in high-risk cutaneous squamous cell carcinoma: a systematic review of outcomes. *Dermatol Surg* 2009;35:574–585. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19415791>.

339. Kim Y, Lehrer EJ, Wirth PJ, et al. Adjuvant radiotherapy may not significantly change outcomes in high-risk cutaneous squamous cell carcinomas with clear surgical margins: A systematic review and meta-analysis. *J Am Acad Dermatol* 2021;86:1246–1257. Available at: <https://pubmed.ncbi.nlm.nih.gov/34890701/>.

340. Cotel WI. Perineural invasion by squamous-cell carcinoma. *J Dermatol Surg Oncol* 1982;8:589–600. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6749931>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

341. Ampil FL, Hardin JC, Peskind SP, Stucker FJ. Perineural invasion in skin cancer of the head and neck: a review of nine cases. *J Oral Maxillofac Surg* 1995;53:34–38. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7799119>.

342. Balamucki CJ, Mancuso AA, Amdur RJ, et al. Skin carcinoma of the head and neck with perineural invasion. *Am J Otolaryngol* 2012;33:447–454. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22185685>.

343. Perkins JL, Liu Y, Mitby PA, et al. Nonmelanoma skin cancer in survivors of childhood and adolescent cancer: a report from the childhood cancer survivor study. *J Clin Oncol* 2005;23:3733–3741. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15923570>.

344. Lichter MD, Karagas MR, Mott LA, et al. Therapeutic ionizing radiation and the incidence of basal cell carcinoma and squamous cell carcinoma. The New Hampshire Skin Cancer Study Group. *Arch Dermatol* 2000;136:1007–1011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10926736>.

345. Karagas MR, Nelson HH, Zens MS, et al. Squamous cell and basal cell carcinoma of the skin in relation to radiation therapy and potential modification of risk by sun exposure. *Epidemiology* 2007;18:776–784. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17917604>.

346. Karagas MR, McDonald JA, Greenberg ER, et al. Risk of basal cell and squamous cell skin cancers after ionizing radiation therapy. For The Skin Cancer Prevention Study Group. *J Natl Cancer Inst* 1996;88:1848–1853. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8961975>.

347. Silva JJ, Tsang RW, Panzarella T, et al. Results of radiotherapy for epithelial skin cancer of the pinna: the Princess Margaret Hospital experience, 1982-1993. *Int J Radiat Oncol Biol Phys* 2000;47:451–459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10802373>.

348. Locke J, Karimpour S, Young G, et al. Radiotherapy for epithelial skin cancer. *Int J Radiat Oncol Biol Phys* 2001;51:748–755. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11697321>.

349. Tsao MN, Tsang RW, Liu FF, et al. Radiotherapy management for squamous cell carcinoma of the nasal skin: The Princess Margaret Hospital experience. *Int J Radiat Oncol Biol Phys* 2002;52:973–979. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11958891>.

350. van Hezewijk M, Creutzberg CL, Putter H, et al. Efficacy of a hypofractionated schedule in electron beam radiotherapy for epithelial skin cancer: Analysis of 434 cases. *Radiother Oncol* 2010;95:245–249. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20334941>.

351. Hu M, Liu H, Bernard A, et al. Hypofractionated adjuvant radiotherapy in cutaneous squamous cell and basal cell carcinoma of the head and neck: 50(Gy) in 20 study. *Oral Oncology Reports* 2025;13:100732. Available at: <https://www.sciencedirect.com/science/article/pii/S2772906025000202>.

352. Likhacheva AO, Devlin PM, Shirvani SM, et al. Skin surface brachytherapy: A survey of contemporary practice patterns. *Brachytherapy* 2017;16:223–229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27908679>.

353. Evans DG, Farndon PA, Burnell LD, et al. The incidence of Gorlin syndrome in 173 consecutive cases of medulloblastoma. *Br J Cancer* 1991;64:959–961. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1931625>.

354. Stavrou T, Bromley CM, Nicholson HS, et al. Prognostic factors and secondary malignancies in childhood medulloblastoma. *J Pediatr Hematol Oncol* 2001;23:431–436. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11878577>.

355. Kimonis VE, Goldstein AM, Pastakia B, et al. Clinical manifestations in 105 persons with nevoid basal cell carcinoma syndrome. *Am J Med Genet* 1997;69:299–308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9096761>.



NCCN Guidelines Version 2.2026 Squamous Cell Skin Cancer

356. Dinehart SM, Anthony JL, Pollack SV. Basal cell carcinoma in young patients after irradiation for childhood malignancy. *Med Pediatr Oncol* 1991;19:508–510. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1961139>.

357. Goldstein AM, Bale SJ, Peck GL, DiGiovanna JJ. Sun exposure and basal cell carcinomas in the nevoid basal cell carcinoma syndrome. *J Am Acad Dermatol* 1993;29:34–41. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8315076>.

358. Spalek M, Jonska-Gmyrek J, Galecki J. Radiation-induced morphea - a literature review. *J Eur Acad Dermatol Venereol* 2015;29:197–202. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25174551>.

359. Lin A, Abu-Isa E, Griffith KA, Ben-Josef E. Toxicity of radiotherapy in patients with collagen vascular disease. *Cancer* 2008;113:648–653. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18506734>.

360. Morris MM, Powell SN. Irradiation in the setting of collagen vascular disease: acute and late complications. *J Clin Oncol* 1997;15:2728–2735. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9215847>.

361. Chao CK, Gerber RM, Perez CA. Reirradiation of recurrent skin cancer of the face. A successful salvage modality. *Cancer* 1995;75:2351–2355. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7712447>.

362. Griep C, Davelaar J, Scholten AN, et al. Electron beam therapy is not inferior to superficial x-ray therapy in the treatment of skin carcinoma. *Int J Radiat Oncol Biol Phys* 1995;32:1347–1350. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7635774>.

363. McNab AA, Francis IC, Benger R, Crompton JL. Perineural spread of cutaneous squamous cell carcinoma via the orbit. Clinical features and outcome in 21 cases. *Ophthalmology* 1997;104:1457–1462. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9307641>.

364. Gluck I, Ibrahim M, Popovtzer A, et al. Skin cancer of the head and neck with perineural invasion: defining the clinical target volumes based on the pattern of failure. *Int J Radiat Oncol Biol Phys* 2009;74:38–46. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18938044>.

365. Holmes ME, Bomford CK. The use of a short distance cobalt unit in the treatment of primary skin tumours. *Br J Radiol* 1982;55:225–228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7066625>.

366. Caccialanza M, Piccinno R, Beretta M, Gnecci L. Results and side effects of dermatologic radiotherapy: a retrospective study of irradiated cutaneous epithelial neoplasms. *J Am Acad Dermatol* 1999;41:589–594. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10495382>.

367. Zablow AI, Eanelli TR, Sanfilippo LJ. Electron beam therapy for skin cancer of the head and neck. *Head Neck* 1992;14:188–195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1587735>.

368. Whalley D, Caine H, McCloud P, et al. Promising results with image guided intensity modulated radiotherapy for muscle invasive bladder cancer. *Radiat Oncol* 2015;10:205. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26407726>.

369. Xing L, Thorndyke B, Schreiber E, et al. Overview of image-guided radiation therapy. *Med Dosim* 2006;31:91–112. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16690451>.

370. Dang A, Kupelian PA, Cao M, et al. Image-guided radiotherapy for prostate cancer. *Transl Androl Urol* 2018;7:308–320. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30050792>.

371. Gauden R, Pracy M, Avery AM, et al. HDR brachytherapy for superficial non-melanoma skin cancers. *J Med Imaging Radiat Oncol* 2013;57:212–217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23551783>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

372. Baris G, Visser AG, van Andel JG. The treatment of squamous cell carcinoma of the nasal vestibule with interstitial iridium implantation. *Radiother Oncol* 1985;4:121–125. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/4070678>.

373. Crook JM, Mazon JJ, Marinello G, et al. Interstitial iridium 192 for cutaneous carcinoma of the external nose. *Int J Radiat Oncol Biol Phys* 1990;18:243–248. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2298627>.

374. McCollough WM, Mendenhall NP, Parsons JT, et al. Radiotherapy alone for squamous cell carcinoma of the nasal vestibule: management of the primary site and regional lymphatics. *Int J Radiat Oncol Biol Phys* 1993;26:73–79. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8482633>.

375. Mendenhall NP, Parsons JT, Cassisi NJ, Million RR. Carcinoma of the nasal vestibule. *Int J Radiat Oncol Biol Phys* 1984;10:627–637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6429094>.

376. Guix B, Finestres F, Tello J, et al. Treatment of skin carcinomas of the face by high-dose-rate brachytherapy and custom-made surface molds. *Int J Radiat Oncol Biol Phys* 2000;47:95–102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10758310>.

377. Mazon JJ, Chassagne D, Crook J, et al. Radiation therapy of carcinomas of the skin of nose and nasal vestibule: a report of 1676 cases by the Groupe Europeen de Curietherapie. *Radiother Oncol* 1988;13:165–173. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3146781>.

378. Mazon JJ, Ghalie R, Zeller J, et al. Radiation therapy for carcinoma of the pinna using iridium 192 wires: a series of 70 patients. *Int J Radiat Oncol Biol Phys* 1986;12:1757–1763. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3759527>.

379. Ballester-Sanchez R, Pons-Llanas O, Candela-Juan C, et al. Two years results of electronic brachytherapy for basal cell carcinoma. *J Contemp Brachytherapy* 2017;9:251–255. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28725249>.

380. Bhatnagar A, Loper A. The initial experience of electronic brachytherapy for the treatment of non-melanoma skin cancer. *Radiat Oncol* 2010;5:87. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20875139>.

381. Eaton DJ. Electronic brachytherapy--current status and future directions. *Br J Radiol* 2015;88:20150002. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25748070>.

382. Tom MC, Hepel JT, Patel R, et al. The american brachytherapy society consensus statement for electronic brachytherapy. *Brachytherapy* 2019;18:292–298. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30497939>.

383. Renzi C, Caggiati A, Mannooranparampil TJ, et al. Sentinel lymph node biopsy for high risk cutaneous squamous cell carcinoma: case series and review of the literature. *Eur J Surg Oncol* 2007;33:364–369. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17129703>.

384. Kwon S, Dong ZM, Wu PC. Sentinel lymph node biopsy for high-risk cutaneous squamous cell carcinoma: clinical experience and review of literature. *World J Surg Oncol* 2011;9:80. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21771334>.

385. Ahmed MM, Moore BA, Schmalbach CE. Utility of head and neck cutaneous squamous cell carcinoma sentinel node biopsy: a systematic review. *Otolaryngol Head Neck Surg* 2014;150:180–187. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24201060>.

386. Fukushima S, Masuguchi S, Igata T, et al. Evaluation of sentinel node biopsy for cutaneous squamous cell carcinoma. *J Dermatol* 2014;41:539–541. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24909214>.

387. Takahashi A, Imafuku S, Nakayama J, et al. Sentinel node biopsy for high-risk cutaneous squamous cell carcinoma. *Eur J Surg Oncol* 2014;40:1256–1262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24953417>.



NCCN Guidelines Version 2.2026 Squamous Cell Skin Cancer

388. Maruyama H, Tanaka R, Fujisawa Y, et al. Availability of sentinel lymph node biopsy for cutaneous squamous cell carcinoma. *J Dermatol* 2017;44:431–437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27666747>.

389. Tejera-Vaquerizo A, Garcia-Doval I, Llombart B, et al. Systematic review of the prevalence of nodal metastases and the prognostic utility of sentinel lymph node biopsy in cutaneous squamous cell carcinoma. *J Dermatol* 2018;45:781–790. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29701281>.

390. Renzi C, Mastroeni S, Mannooranparampil TJ, et al. Skin cancer knowledge and preventive behaviors among patients with a recent history of cutaneous squamous cell carcinoma. *Dermatology* 2008;217:74–80. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18424897>.

391. Allen JE, Stolle LB. Utility of sentinel node biopsy in patients with high-risk cutaneous squamous cell carcinoma. *Eur J Surg Oncol* 2015;41:197–200. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25468745>.

392. Saito Y, Fujikawa H, Takatsuka S, et al. Risk factors for lymph node metastasis in cutaneous squamous cell carcinoma: a long-term retrospective study of Japanese patients. *Int J Clin Oncol* 2021;26:606–612. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33175299>.

393. Gurudutt VV, Genden EM. Cutaneous squamous cell carcinoma of the head and neck. *J Skin Cancer* 2011;2011:502723. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21461387>.

394. Forest VI, Clark JJ, Veness MJ, Milross C. N1S3: a revised staging system for head and neck cutaneous squamous cell carcinoma with lymph node metastases: results of 2 Australian Cancer Centers. *Cancer* 2010;116:1298–1304. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20052712>.

395. Wang JT, Palme CE, Wang AY, et al. In patients with metastatic cutaneous head and neck squamous cell carcinoma to cervical lymph nodes, the extent of neck dissection does not influence outcome. *J Laryngol Otol* 2013;127 Suppl 1:S2–7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23046820>.

396. Goh RY, Bova R, Fogarty GB. Cutaneous squamous cell carcinoma metastatic to parotid - analysis of prognostic factors and treatment outcome. *World J Surg Oncol* 2012;10:117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22731750>.

397. McDowell LJ, Tan TJ, Bressel M, et al. Outcomes of cutaneous squamous cell carcinoma of the head and neck with parotid metastases. *J Med Imaging Radiat Oncol* 2016;60:668–676. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27324298>.

398. Hirshoren N, Danne J, Dixon BJ, et al. Prognostic markers in metastatic cutaneous squamous cell carcinoma of the head and neck. *Head Neck* 2017;39:772–778. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28199044>.

399. Czerwonka L, De Santis RJ, Horowitz G, et al. Staging cutaneous squamous cell carcinoma metastases to the parotid gland. *Laryngoscope* 2017;127:2063–2069. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28295401>.

400. Thom JJ, Moore EJ, Price DL, et al. The role of total parotidectomy for metastatic cutaneous squamous cell carcinoma and malignant melanoma. *JAMA Otolaryngol Head Neck Surg* 2014;140:548–554. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24722863>.

401. Palme CE, O'Brien CJ, Veness MJ, et al. Extent of parotid disease influences outcome in patients with metastatic cutaneous squamous cell carcinoma. *Arch Otolaryngol Head Neck Surg* 2003;129:750–753. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12874077>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

402. Audet N, Palme CE, Gullane PJ, et al. Cutaneous metastatic squamous cell carcinoma to the parotid gland: analysis and outcome. *Head Neck* 2004;26:727–732. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15287040>.

403. Hinerman RW, Indelicato DJ, Amdur RJ, et al. Cutaneous squamous cell carcinoma metastatic to parotid-area lymph nodes. *Laryngoscope* 2008;118:1989–1996. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18849863>.

404. Veness MJ, Morgan GJ, Palme CE, Gebiski V. Surgery and adjuvant radiotherapy in patients with cutaneous head and neck squamous cell carcinoma metastatic to lymph nodes: combined treatment should be considered best practice. *Laryngoscope* 2005;115:870–875. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15867656>.

405. Oddone N, Morgan GJ, Palme CE, et al. Metastatic cutaneous squamous cell carcinoma of the head and neck: the Immunosuppression, Treatment, Extranodal spread, and Margin status (ITEM) prognostic score to predict outcome and the need to improve survival. *Cancer* 2009;115:1883–1891. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19224549>.

406. Sweeny L, Zimmerman T, Carroll WR, et al. Head and neck cutaneous squamous cell carcinoma requiring parotidectomy: prognostic indicators and treatment selection. *Otolaryngol Head Neck Surg* 2014;150:610–617. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24474713>.

407. Chen MM, Roman SA, Sosa JA, Judson BL. Prognostic factors for squamous cell cancer of the parotid gland: an analysis of 2104 patients. *Head Neck* 2015;37:1–7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24339135>.

408. Ch'ng S, Maitra A, Allison RS, et al. Parotid and cervical nodal status predict prognosis for patients with head and neck metastatic cutaneous squamous cell carcinoma. *J Surg Oncol* 2008;98:101–105. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18523982>.

409. Sahovaler A, Krishnan RJ, Yeh DH, et al. Outcomes of cutaneous squamous cell carcinoma in the head and neck region with regional lymph node metastasis: A systematic review and meta-analysis. *JAMA Otolaryngol Head Neck Surg* 2019;145:352–360. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30844021>.

410. O'Brien CJ, McNeil EB, McMahon JD, et al. Significance of clinical stage, extent of surgery, and pathologic findings in metastatic cutaneous squamous carcinoma of the parotid gland. *Head Neck* 2002;24:417–422. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12001070>.

411. Lam JKS, Sundaresan P, Gebiski V, Veness MJ. Immunocompromised patients with metastatic cutaneous nodal squamous cell carcinoma of the head and neck: Poor outcome unrelated to the index lesion. *Head Neck* 2018;40:985–992. Available at: <https://pubmed.ncbi.nlm.nih.gov/29360262/>.

412. Cutaneous squamous cell carcinoma and other cutaneous carcinomas. In: Edge SB, Carducci M, Byrd DR, eds. *AJCC Cancer Staging Manual* (ed 7). New York: Springer-Verlag New York, LLC; 2009.

413. Pramana A, Browne L, Graham PH. Metastatic cutaneous squamous cell carcinoma to parotid nodes: the role of bolus with adjuvant radiotherapy. *J Med Imaging Radiat Oncol* 2012;56:100–108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22339753>.

414. Amoils M, Lee CS, Sunwoo J, et al. Node-positive cutaneous squamous cell carcinoma of the head and neck: Survival, high-risk features, and adjuvant chemoradiotherapy outcomes. *Head Neck* 2017;39:881–885. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28252823>.

415. Smile TD, Ruiz ES, Kus KJB, et al. Implications of satellitosis or in-transit metastasis in cutaneous squamous cell carcinoma: A prognostic omission in cancer staging systems. *JAMA Dermatol* 2022;158:390–394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35195668>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

416. Amin MB, Greene FL, Edge SB, et al. The eighth edition AJCC cancer staging manual: Continuing to build a bridge from a population-based to a more "personalized" approach to cancer staging. *CA Cancer J Clin* 2017;67:93–99. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28094848>.

417. Marti-Marti I, Podlipnik S, Canueto J, et al. Prognostic factors for satellitosis or in-transit metastasis in cutaneous squamous cell carcinoma: A multicentric cohort study. *J Am Acad Dermatol* 2023;89:119–127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36907554>.

418. Rischin D, Porceddu S, Day F, et al. Adjuvant cemiplimab or placebo in high-risk cutaneous squamous-cell carcinoma. *N Engl J Med* 2025;393:774–785. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40454639>.

419. Bertrand N, Guerreschi P, Basset-Seguín N, et al. Vismodegib in neoadjuvant treatment of locally advanced basal cell carcinoma: First results of a multicenter, open-label, phase 2 trial (VISMONEO study): Neoadjuvant Vismodegib in Locally Advanced Basal Cell Carcinoma. *EClinicalMedicine* 2021;35:100844. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33997740>.

420. DePalo DK, Dugan MM, Naqvi SMH, et al. A comparison of isolated limb infusion/perfusion, immune checkpoint inhibitors, and intralesional therapy as first-line treatment for patients with melanoma in-transit metastases. *Cancer* 2025;131:e35636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39522025>.

421. Thompson JF, Kam PC, Waugh RC, Harman CR. Isolated limb infusion with cytotoxic agents: a simple alternative to isolated limb perfusion. *Semin Surg Oncol* 1998;14:238–247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9548607>.

422. Creech O, Jr., Krementz ET, Ryan RF, Winblad JN. Chemotherapy of cancer: Regional perfusion utilizing an extracorporeal circuit. *Ann Surg* 1958;148:616–632. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/13583933>.

423. O'Donoghue C, Perez MC, Mullinax JE, et al. Isolated Limb Infusion: A Single-Center Experience with Over 200 Infusions. *Ann Surg Oncol* 2017;24:3842–3849. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29019175>.

424. Huis In 't Veld EA, Grunhagen DJ, Deroose JP, et al. Isolated limb perfusion for unresectable extremity cutaneous squamous cell carcinoma: an effective limb saving strategy. *Br J Cancer* 2018;119:429–434. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29961756>.

425. Khansur T, Kennedy A. Cisplatin and 5-fluorouracil for advanced locoregional and metastatic squamous cell carcinoma of the skin. *Cancer* 1991;67:2030–2032. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2004320>.

426. Reigneau M, Robert C, Routier E, et al. Efficacy of neoadjuvant cetuximab alone or with platinum salt for the treatment of unresectable advanced nonmetastatic cutaneous squamous cell carcinomas. *Br J Dermatol* 2015;173:527–534. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25704233>.

427. Jarkowski A, 3rd, Hare R, Loud P, et al. Systemic therapy in advanced cutaneous squamous cell carcinoma (CSCC): The Roswell Park experience and a review of the literature. *Am J Clin Oncol* 2016;39:545–548. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24879468>.

428. Maubec E, Petrow P, Scheer-Senyearich I, et al. Phase II study of cetuximab as first-line single-drug therapy in patients with unresectable squamous cell carcinoma of the skin. *J Clin Oncol* 2011;29:3419–3426. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21810686>.

429. Lewis CM, Glisson BS, Feng L, et al. A phase II study of gefitinib for aggressive cutaneous squamous cell carcinoma of the head and neck. *Clin Cancer Res* 2012;18:1435–1446. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22261807>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

430. Porceddu SV, Bressel M, Poulsen MG, et al. Postoperative concurrent chemoradiotherapy versus postoperative radiotherapy in high-risk cutaneous squamous cell carcinoma of the head and neck: The randomized phase III TROG 05.01 trial. *J Clin Oncol* 2018;36:JCO2017770941. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29537906>.

431. Goyal U, Prabhakar NK, Davuluri R, et al. Role of concurrent systemic therapy with adjuvant radiation therapy for locally advanced cutaneous head and neck squamous cell carcinoma. *Cureus* 2017;9:e1784. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29279810>.

432. Trosman SJ, Zhu A, Nicolli EA, et al. High-risk cutaneous squamous cell cancer of the head and neck: Risk factors for recurrence and impact of adjuvant treatment. *Laryngoscope* 2021;131:E136–E143. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32065413>.

433. Tanvetyanon T, Padhya T, McCaffrey J, et al. Postoperative concurrent chemotherapy and radiotherapy for high-risk cutaneous squamous cell carcinoma of the head and neck. *Head Neck* 2015;37:840–845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24623654>.

434. Nottage MK, Lin C, Hughes BG, et al. Prospective study of definitive chemoradiation in locally or regionally advanced squamous cell carcinoma of the skin. *Head Neck* 2017;39:679–683. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28032670>.

435. Guthrie TH, Jr., Porubsky ES, Luxenberg MN, et al. Cisplatin-based chemotherapy in advanced basal and squamous cell carcinomas of the skin: results in 28 patients including 13 patients receiving multimodality therapy. *J Clin Oncol* 1990;8:342–346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2405109>.

436. Lu SM, Lien WW. Concurrent radiotherapy with cetuximab or platinum-based chemotherapy for locally advanced cutaneous squamous cell carcinoma of the head and neck. *Am J Clin Oncol* 2018;41:95–99. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26353121>.

437. Amaral T, Osewold M, Presser D, et al. Advanced cutaneous squamous cell carcinoma: real world data of patient profiles and treatment patterns. *J Eur Acad Dermatol Venereol* 2019;33 Suppl 8:44–51. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31658392>.

438. Preneau S, Rio E, Brocard A, et al. Efficacy of cetuximab in the treatment of squamous cell carcinoma. *J Dermatolog Treat* 2014;25:424–427. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23167307>.

439. Samstein RM, Ho AL, Lee NY, Barker CA. Locally advanced and unresectable cutaneous squamous cell carcinoma: outcomes of concurrent cetuximab and radiotherapy. *J Skin Cancer* 2014;2014:284582. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25136458>.

440. O'Bryan K, Sherman W, Niedt GW, et al. An evolving paradigm for the workup and management of high-risk cutaneous squamous cell carcinoma. *J Am Acad Dermatol* 2013;69:595–602 e591. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23871719>.

441. Joseph K, Alkaabi K, Warkentin H, et al. Cetuximab-radiotherapy combination in the management of locally advanced cutaneous squamous cell carcinoma. *J Med Imaging Radiat Oncol* 2019;63:257–263. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30549229>.

442. Berliner JG, Schulman JM, Lazarova Z, et al. Response of cutaneous squamous cell carcinoma to treatment with cetuximab. *Dermatol Surg* 2019;45:313–316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29957662>.

443. Ogata D, Namikawa K, Otsuka M, et al. Systemic treatment of patients with advanced cutaneous squamous cell carcinoma: response rates and outcomes of the regimes used. *Eur J Cancer* 2020;127:108–117. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32004792>.

444. Trodello C, Pepper JP, Wong M, Wysong A. Cisplatin and cetuximab treatment for metastatic cutaneous squamous cell carcinoma: A systematic review. *Dermatol Surg* 2017;43:40–49. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27618393>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

445. Heath CH, Deep NL, Nabell L, et al. Phase 1 study of erlotinib plus radiation therapy in patients with advanced cutaneous squamous cell carcinoma. *Int J Radiat Oncol Biol Phys* 2013;85:1275–1281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23182701>.

446. Hourbeigt K, Ehret M, Visseaux L, et al. Efficacy and safety of panitumumab alone or in association with radiotherapy in unresectable cutaneous squamous cell carcinoma. *J Eur Acad Dermatol Venereol* 2020;34:2789–2794. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32294281>.

447. Maring S, Elsayad K, Stenner M, et al. Efficacy of carboplatin/paclitaxel-based radiochemotherapy in locally advanced squamous cell carcinoma of head and neck. *Oncol Res Treat* 2018;41:736–743. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30419553>.

448. Vlacich G, Diaz R, Thorpe SW, et al. Intensity-modulated radiation therapy with concurrent carboplatin and paclitaxel for locally advanced head and neck cancer: toxicities and efficacy. *Oncologist* 2012;17:673–681. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22550060>.

449. Suntharalingam M, Haas ML, Conley BA, et al. The use of carboplatin and paclitaxel with daily radiotherapy in patients with locally advanced squamous cell carcinomas of the head and neck. *Int J Radiat Oncol Biol Phys* 2000;47:49–56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10758304>.

450. Fujisawa Y, Umebayashi Y, Ichikawa E, et al. Chemoradiation using low-dose cisplatin and 5-fluorouracil in locally advanced squamous cell carcinoma of the skin: A report of two cases. *J Am Acad Dermatol* 2006;55:S81–85. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17052540>.

451. Cooper JS, Pajak TF, Forastiere AA, et al. Postoperative concurrent radiotherapy and chemotherapy for high-risk squamous-cell carcinoma of the head and neck. *N Engl J Med* 2004;350:1937–1944. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15128893>.

452. Bonner JA, Harari PM, Giralt J, et al. Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *N Engl J Med* 2006;354:567–578. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16467544>.

453. Migden MR, Khushalani NI, Chang ALS, et al. Cemiplimab in locally advanced cutaneous squamous cell carcinoma: results from an open-label, phase 2, single-arm trial. *Lancet Oncol* 2020;21:294–305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31952975>.

454. Migden MR, Rischin D, Schmults CD, et al. PD-1 blockade with cemiplimab in advanced cutaneous squamous-cell carcinoma. *N Engl J Med* 2018;379:341–351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29863979>.

455. Rischin D, Migden MR, Lim AM, et al. Phase 2 study of cemiplimab in patients with metastatic cutaneous squamous cell carcinoma: primary analysis of fixed-dosing, long-term outcome of weight-based dosing. *J Immunother Cancer* 2020;8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32554615>.

456. Gross ND, Miller DM, Khushalani NI, et al. Neoadjuvant cemiplimab for stage II to IV cutaneous squamous-cell carcinoma. *N Engl J Med* 2022;387:1557–1568. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36094839>.

457. Hughes BGM, Munoz-Couselo E, Mortier L, et al. Pembrolizumab for locally advanced and recurrent/metastatic cutaneous squamous cell carcinoma (KEYNOTE-629 study): an open-label, nonrandomized, multicenter, phase II trial. *Ann Oncol* 2021;32:1276–1285. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34293460>.

458. Grob JJ, Gonzalez R, Basset-Seguin N, et al. Pembrolizumab monotherapy for recurrent or metastatic cutaneous squamous cell carcinoma: A single-arm phase II trial (KEYNOTE-629). *J Clin Oncol* 2020;38:2916–2925. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32673170>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

459. Munhoz RR, Nader-Marta G, de Camargo VP, et al. A phase 2 study of first-line nivolumab in patients with locally advanced or metastatic cutaneous squamous-cell carcinoma. *Cancer* 2022;128:4223–4231. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36274573>.

460. Tahara M, Kiyota N, Yokota T, et al. Phase II trial of combination treatment with paclitaxel, carboplatin and cetuximab (PCE) as first-line treatment in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck (CSPOR-HN02). *Ann Oncol* 2018;29:1004–1009. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29408977>.

461. Ferrari D, Fiore J, Codeca C, et al. A phase II study of carboplatin and paclitaxel for recurrent or metastatic head and neck cancer. *Anticancer Drugs* 2009;20:185–190. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19396017>.

462. Botticelli A, Pomati G, Cirillo A, et al. Weekly chemotherapy as first line treatment in frail head and neck cancer patients in the immunotherapy era. *J Transl Med* 2021;19:303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34253248>.

463. Geraghty L, Schultz TE, Hoffman SE, et al. Weekly vs. 3-weekly paclitaxel, carboplatin, and cetuximab (PCC) in recurrent/metastatic head and neck cancer. *Mol Clin Oncol* 2021;15:240. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34650807>.

464. Carinato H, Burgy M, Ferry R, et al. Weekly paclitaxel, carboplatin, and cetuximab as first-line treatment of recurrent and/or metastatic head and neck squamous cell carcinoma for patients ineligible to cisplatin-based chemotherapy: A retrospective monocentric study in 60 patients. *Front Oncol* 2021;11:714551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34778031>.

465. Conen KL, Fischer N, Hofbauer GF, et al. Cetuximab in metastatic squamous cell cancer of the skin: A Swiss case series. *Dermatology* 2014;229:97–101. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24923455>.

466. Cowey CL, Robert NJ, Espirito JL, et al. Clinical outcomes among unresectable, locally advanced, and metastatic cutaneous squamous cell carcinoma patients treated with systemic therapy. *Cancer Med* 2020;9:7381–7387. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32578965>.

467. Foote MC, McGrath M, Guminski A, et al. Phase II study of single-agent panitumumab in patients with incurable cutaneous squamous cell carcinoma. *Ann Oncol* 2014;25:2047–2052. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25091317>.

468. William WN, Jr., Feng L, Ferrarotto R, et al. Gefitinib for patients with incurable cutaneous squamous cell carcinoma: A single-arm phase II clinical trial. *J Am Acad Dermatol* 2017;77:1110–1113 e1112. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28964539>.

469. Cavalieri S, Perrone F, Miceli R, et al. Efficacy and safety of single-agent pan-human epidermal growth factor receptor (HER) inhibitor dacomitinib in locally advanced unresectable or metastatic skin squamous cell cancer. *Eur J Cancer* 2018;97:7–15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29734047>.

470. Gold KA, Kies MS, William WN, Jr., et al. Erlotinib in the treatment of recurrent or metastatic cutaneous squamous cell carcinoma: A single-arm phase 2 clinical trial. *Cancer* 2018;124:2169–2173. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29579331>.

471. Peron J, Poupart M, Ceruse P, et al. Efficacy and safety of capecitabine in heavily pretreated recurrent/metastatic head and neck squamous cell carcinoma. *Anticancer Drugs* 2012;23:1107–1111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22954504>.

472. Martinez-Trufero J, Isla D, Adansa JC, et al. Phase II study of capecitabine as palliative treatment for patients with recurrent and metastatic squamous head and neck cancer after previous platinum-based treatment. *Br J Cancer* 2010;102:1687–1691. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20485287>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

473. Lavaud J, Blom A, Longvert C, et al. Pembrolizumab and concurrent hypo-fractionated radiotherapy for advanced non-resectable cutaneous squamous cell carcinoma. *Eur J Dermatol* 2019;29:636–640. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31903954>.

474. Karagas MR, Stukel TA, Greenberg ER, et al. Risk of subsequent basal cell carcinoma and squamous cell carcinoma of the skin among patients with prior skin cancer. Skin Cancer Prevention Study Group. *JAMA* 1992;267:3305–3310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1597912>.

475. Flohil SC, van der Leest RJ, Arends LR, et al. Risk of subsequent cutaneous malignancy in patients with prior keratinocyte carcinoma: a systematic review and meta-analysis. *Eur J Cancer* 2013;49:2365–2375. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23608733>.

476. Marcil I, Stern RS. Risk of developing a subsequent nonmelanoma skin cancer in patients with a history of nonmelanoma skin cancer: a critical review of the literature and meta-analysis. *Arch Dermatol* 2000;136:1524–1530. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11115165>.

477. Jambusaria-Pahlajani A, Hess SD, Katz KA, et al. Uncertainty in the perioperative management of high-risk cutaneous squamous cell carcinoma among Mohs surgeons. *Arch Dermatol* 2010;146:1225–1231. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21079057>.

478. Martinez JC, Otley CC, Stasko T, et al. Defining the clinical course of metastatic skin cancer in organ transplant recipients: a multicenter collaborative study. *Arch Dermatol* 2003;139:301–306. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12622621>.

479. Carucci JA, Martinez JC, Zeitouni NC, et al. In-transit metastasis from primary cutaneous squamous cell carcinoma in organ transplant recipients and nonimmunosuppressed patients: clinical characteristics, management, and outcome in a series of 21 patients. *Dermatol Surg* 2004;30:651–655. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15061850>.

480. Bordea C, Wojnarowska F, Millard PR, et al. Skin cancers in renal-transplant recipients occur more frequently than previously recognized in a temperate climate. *Transplantation* 2004;77:574–579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15084938>.

481. Fortina AB, Piaserico S, Caforio AL, et al. Immunosuppressive level and other risk factors for basal cell carcinoma and squamous cell carcinoma in heart transplant recipients. *Arch Dermatol* 2004;140:1079–1085. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15381547>.

482. McGeown MG, Douglas JF, Middleton D. One thousand renal transplants at Belfast City Hospital: post-graft neoplasia 1968-1999, comparing azathioprine only with cyclosporin-based regimes in a single centre. *Clin Transpl* 2000:193–202. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11512313>.

483. Euvrard S, Morelon E, Rostaing L, et al. Sirolimus and secondary skin-cancer prevention in kidney transplantation. *N Engl J Med* 2012;367:329–339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22830463>.

484. Campbell SB, Walker R, Tai SS, et al. Randomized controlled trial of sirolimus for renal transplant recipients at high risk for nonmelanoma skin cancer. *Am J Transplant* 2012;12:1146–1156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22420843>.

485. Salgo R, Gossman J, Schofer H, et al. Switch to a sirolimus-based immunosuppression in long-term renal transplant recipients: reduced rate of (pre-)malignancies and nonmelanoma skin cancer in a prospective, randomized, assessor-blinded, controlled clinical trial. *Am J Transplant* 2010;10:1385–1393. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20121752>.

486. Alberu J, Pascoe MD, Campistol JM, et al. Lower malignancy rates in renal allograft recipients converted to sirolimus-based, calcineurin inhibitor-free immunotherapy: 24-month results from the CONVERT trial. *Transplantation* 2011;92:303–310. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21792049>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

487. Campistol JM, Eris J, Oberbauer R, et al. Sirolimus therapy after early cyclosporine withdrawal reduces the risk for cancer in adult renal transplantation. *J Am Soc Nephrol* 2006;17:581–589. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16434506>.

488. Hoogendijk-van den Akker JM, Harden PN, Hoitsma AJ, et al. Two-year randomized controlled prospective trial converting treatment of stable renal transplant recipients with cutaneous invasive squamous cell carcinomas to sirolimus. *J Clin Oncol* 2013;31:1317–1323. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23358973>.

489. Tedesco-Silva H, Peddi VR, Sanchez-Fructuoso A, et al. Open-label, randomized study of transition from tacrolimus to sirolimus immunosuppression in renal allograft recipients. *Transplant Direct* 2016;2:e69. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27500260>.

490. Gu YH, Du JX, Ma ML. Sirolimus and non-melanoma skin cancer prevention after kidney transplantation: a meta-analysis. *Asian Pac J Cancer Prev* 2012;13:4335–4339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23167339>.

491. Lim WH, Russ GR, Wong G, et al. The risk of cancer in kidney transplant recipients may be reduced in those maintained on everolimus and reduced cyclosporine. *Kidney Int* 2017;91:954–963. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28109543>.

492. Shuttleworth D, Marks R, Griffin PJ, Salaman JR. Treatment of cutaneous neoplasia with etretinate in renal transplant recipients. *Q J Med* 1988;68:717–725. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3076965>.

493. Bavinck JN, Tieben LM, Van der Woude FJ, et al. Prevention of skin cancer and reduction of keratotic skin lesions during acitretin therapy in renal transplant recipients: a double-blind, placebo-controlled study. *J Clin Oncol* 1995;13:1933–1938. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7636533>.

494. Gibson GE, O'Grady A, Kay EW, Murphy GM. Low-dose retinoid therapy for chemoprophylaxis of skin cancer in renal transplant recipients. *J Eur Acad Dermatol Venereol* 1998;10:42–47. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9552756>.

495. McKenna DB, Murphy GM. Skin cancer chemoprophylaxis in renal transplant recipients: 5 years of experience using low-dose acitretin. *Br J Dermatol* 1999;140:656–660. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10233316>.

496. George R, Weightman W, Russ GR, et al. Acitretin for chemoprevention of non-melanoma skin cancers in renal transplant recipients. *Australas J Dermatol* 2002;43:269–273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12423433>.

497. de Sevaux RG, Smit JV, de Jong EM, et al. Acitretin treatment of premalignant and malignant skin disorders in renal transplant recipients: clinical effects of a randomized trial comparing two doses of acitretin. *J Am Acad Dermatol* 2003;49:407–412. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12963902>.

498. Kraemer KH, DiGiovanna JJ, Moshell AN, et al. Prevention of skin cancer in xeroderma pigmentosum with the use of oral isotretinoin. *N Engl J Med* 1988;318:1633–1637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3287161>.

499. Nijsten TE, Stern RS. Oral retinoid use reduces cutaneous squamous cell carcinoma risk in patients with psoriasis treated with psoralen-UVA: a nested cohort study. *J Am Acad Dermatol* 2003;49:644–650. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14512910>.

500. Moon TE, Levine N, Cartmel B, et al. Effect of retinol in preventing squamous cell skin cancer in moderate-risk subjects: a randomized, double-blind, controlled trial. Southwest Skin Cancer Prevention Study Group. *Cancer Epidemiol Biomarkers Prev* 1997;6:949–956. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9367069>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

501. Levine N, Moon TE, Cartmel B, et al. Trial of retinol and isotretinoin in skin cancer prevention: a randomized, double-blind, controlled trial. Southwest Skin Cancer Prevention Study Group. *Cancer Epidemiol Biomarkers Prev* 1997;6:957–961. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9367070>.

502. Kadakia KC, Barton DL, Loprinzi CL, et al. Randomized controlled trial of acitretin versus placebo in patients at high-risk for basal cell or squamous cell carcinoma of the skin (North Central Cancer Treatment Group Study 969251). *Cancer* 2012;118:2128–2137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21882176>.

503. Meyskens FL, Jr., Gilmartin E, Alberts DS, et al. Activity of isotretinoin against squamous cell cancers and preneoplastic lesions. *Cancer Treat Rep* 1982;66:1315–1319. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6211233>.

504. Drugs@FDA: FDA-Approved Drugs. U.S. Food & Drug Administration; Available at: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>. Accessed March 2, 2026.

505. Weinstock MA, Bingham SF, Digiovanna JJ, et al. Tretinoin and the prevention of keratinocyte carcinoma (basal and squamous cell carcinoma of the skin): a veterans affairs randomized chemoprevention trial. *J Invest Dermatol* 2012;132:1583–1590. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22318383>.

506. Tang JY, Chiou AS, Mackay-Wiggan JM, et al. Tazarotene: randomized, double-blind, vehicle-controlled, and open-label concurrent trials for basal cell carcinoma prevention and therapy in patients with basal cell nevus syndrome. *Cancer Prev Res (Phila)* 2014;7:292–299. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24441673>.

507. Eide MJ, Weinstock MA, Dufresne RG, Jr., et al. Relationship of treatment delay with surgical defect size from keratinocyte carcinoma (basal cell carcinoma and squamous cell carcinoma of the skin). *J Invest Dermatol* 2005;124:308–314. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15675948>.

508. Renzi C, Mastroeni S, Passarelli F, et al. Factors associated with large cutaneous squamous cell carcinomas. *J Am Acad Dermatol* 2010;63:404–411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20598396>.

509. Diehl J, Choi YM, Liang LJ, Chiu M. Association between mohs surgery wait times and surgical defect size in patients with squamous cell or basal cell carcinoma of the skin. *Dermatol Surg* 2015;41:768–774. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26050214>.

510. Wollina U, Bayyoud Y, Kronert C, Nowak A. Giant epithelial malignancies (Basal cell carcinoma, squamous cell carcinoma): A series of 20 tumors from a single center. *J Cutan Aesthet Surg* 2012;5:12–19. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22557850>.

511. Renzi C, Mastroeni S, Mannooranparampil TJ, et al. Delay in diagnosis and treatment of squamous cell carcinoma of the skin. *Acta Derm Venereol* 2010;90:595–601. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21057742>.

512. Kakagia D, Trypsiannis G, Karanikas M, et al. Patient-related delay in presentation for cutaneous squamous cell carcinoma. A cross-sectional clinical study. *Onkologie* 2013;36:738–744. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24356565>.

513. Amber KT, Bloom R, Abyaneh MY, et al. Patient factors and their association with nonmelanoma skin cancer morbidity and the performance of self-skin exams: A cross-sectional study. *J Clin Aesthet Dermatol* 2016;9:16–22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27878058>.

514. Janda M, Youl P, Neale R, et al. Clinical skin examination outcomes after a video-based behavioral intervention: analysis from a randomized clinical trial. *JAMA Dermatol* 2014;150:372–379. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24553807>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

515. Firooz A, Amin-Nejad R, Bouzari N, et al. Sun protection in Iranian kidney transplant recipients: knowledge, attitude and practice. *J Eur Acad Dermatol Venereol* 2007;21:754–757. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17567302>.

516. Goldenberg A, Nguyen BT, Jiang SI. Knowledge, understanding, and use of preventive strategies against nonmelanoma skin cancer in healthy and immunosuppressed individuals undergoing Mohs surgery. *Dermatol Surg* 2014;40:93–100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24354730>.

517. Borghi A, Corazza M, Battaglia Y, et al. What is the key to improving renal transplant recipients' awareness of skin cancer risk? *Dermatology* 2016;232:715–720. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28343227>.

518. Wohlk IM, Philipsen PA, Wulf HC. Factors associated with cessation of sunbed use among Danish women. *Photodermatol Photoimmunol Photomed* 2016;32:191–198. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27084577>.

519. Fischer AH, Wang TS, Yenokyan G, et al. Sunburn and sun-protective behaviors among adults with and without previous nonmelanoma skin cancer (NMSC): A population-based study. *J Am Acad Dermatol* 2016;75:371–379 e375. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27198078>.

520. Wiznia L, Dai F, Chagpar AB. Do non-melanoma skin cancer survivors use tanning beds less often than the general public? *Dermatol Online J* 2016;22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27617935>.

521. Girgis A, Sanson-Fisher RW, Watson A. A workplace intervention for increasing outdoor workers' use of solar protection. *Am J Public Health* 1994;84:77–81. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8279616>.

522. Robinson JK, Friedewald JJ, Desai A, Gordon EJ. Response across the health-literacy spectrum of kidney transplant recipients to a sun-protection education program delivered on tablet computers: Randomized controlled trial. *JMIR Cancer* 2015;1:e8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28410176>.

523. Loescher LJ, Hansen C, Hepworth JT, et al. A preliminary study of a video intervention to inform solid organ transplant recipients about skin cancer. *Transplant Proc* 2013;45:3187–3189. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23747188>.

524. Tavadia S, Dawn G, Payne C, et al. Skin-cancer awareness in Scottish cardiac transplant recipients. *Clin Exp Dermatol* 2006;31:354–357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16681574>.

525. Ismail F, Mitchell L, Casabonne D, et al. Specialist dermatology clinics for organ transplant recipients significantly improve compliance with photoprotection and levels of skin cancer awareness. *Br J Dermatol* 2006;155:916–925. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17034519>.

526. Seite S, Del Marmol V, Moyal D, Friedman AJ. Public primary and secondary skin cancer prevention, perceptions and knowledge: an international cross-sectional survey. *J Eur Acad Dermatol Venereol* 2017;31:815–820. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28045207>.

527. Clowers-Webb HE, Christenson LJ, Phillips PK, et al. Educational outcomes regarding skin cancer in organ transplant recipients: Randomized intervention of intensive vs standard education. *Arch Dermatol* 2006;142:712–718. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16785373>.

528. Kim NN, Boone SL, Ortiz S, et al. Squamous cell carcinoma in solid organ transplant recipients: influences on perception of risk and optimal time to provide education. *Arch Dermatol* 2009;145:1196–1197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19841415>.



NCCN Guidelines Version 2.2026

Squamous Cell Skin Cancer

529. Robinson JK, Friedewald JJ, Desai A, Gordon EJ. A randomized controlled trial of a mobile medical app for kidney transplant recipients: Effect on use of sun protection. *Transplant Direct* 2016;2. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26900599>.

530. Robinson JK, Turrisi R, Mallett KA, et al. Efficacy of an educational intervention with kidney transplant recipients to promote skin self-examination for squamous cell carcinoma detection. *Arch Dermatol* 2011;147:689–695. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21339418>.