

What are we weighting for?

A mechanistic model for probability weighting

Ole Peters Alexander Adamou Yonatan Berman Mark Kirstein

D-TEA 2020, 16 June 2020



UNIVERSITÄT LEIPZIG



Mathematisches Institut





Mark Kirstein

Main results



- 1 inverse-S shape can be explained by difference in uncertainty
- 2 cautious estimation of probabilities generates such uncertainty

► PW K&T 1979



Mark Kirstein

Main Result


Probability
Weighting

Ergodicity
Question

Estimation

Conclusion

Definition of Probability Weighting (PW)



(TverskyKahneman1992)

- low probabilities treated as higher \rightarrow high probabilities treated as lower
- stable empirical pattern: inverse-S shape

Received wisdom:

- PW = maladaptive irrational cognitive bias

In search of a mechanism

- \hookrightarrow How does this pattern emerge?
- \hookrightarrow Can we derive a functional form (rather than fit a function)?

Task: model payout, x , of a gamble as a random variable.

Disinterested Observer (DO)



DO assigns
probabilities $p(x)$
CDF $F_p(x)$

Decision Maker (DM)



DM assigns different
probabilities $w(x)$ (decision weights)
CDF $F_w(x)$



Mark Kirstein

Main Result

Probability
Weighting




Ergodicity
Question

Estimation

Conclusion

Possible model differences

Locations, Scales, Shapes





Thought experiment: DM assumes greater scale


Main Result

Probability
Weighting

Ergodicity
Question

Estimation

Conclusion



Numerically easy for any pair of distributions (models):


- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$



Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$






Mark Kirstein



Thought experiment: DM assumes greater scale

Different Scales



Numerically easy for any pair of distributions (models):

- 1 list values of DO's CDF, $F_p(x)$, at set x_i
- 2 list values of DM's CDF, $F_w(x)$, at same x_i
- 3 plot $F_w(x)$ vs. $F_p(x)$





Mark Kirstein

Applying the Procedure to the Uncertainty Types





Main Result

Probability
Weighting

Ergodicity
Question

Estimation

Conclusion







Mark Kirstein

Interim conclusion

- DM's greater scale gives inverse-S shape (unimodal distributions)
- difference in locations gives asymmetry
- reproduces observations of probability weighting

Job done. Thank you for your attention ;)

► Functional Forms





Mark Kirstein

The Ergodicity Question

Typical DO concern

What happens on average to
the **ensemble** of subjects?

\neq

Typical DM concern

What happens to me
on average over time?

Main Result

Probability
Weighting

Ergodicity
Question

Estimation

Conclusion



Mark Kirstein

Why DM's greater scale?

- DM has no control over experiment
- experiment may be unclear to DM
- DM may not trust DO
- ...

Main Result

Probability
Weighting

Ergodicity
Question

Estimation

Conclusion



Mark Kirstein

Experiencing probabilities

Main Result

Probability
Weighting

Ergodicity
Question

Estimation

Conclusion

- probabilities are not observable
 - DO encounters probabilities as known frequency in ensemble of experiments
 - DM encounters probabilities as frequencies estimated over time
- ↪ **DM usually has to account for uncertainty in probabilities**



Rare Event

- $p(x) = 0.0001$
- 10 000 observations
- $\sim 99.5\%$ of such time series will contain 0 or 1 events
- Naïve estimation: $\hat{p}(x) = 0$ or $\hat{p}(x) = 0.0001$

↪ either impossible or 1000% (over)estimation

Common Event

- $p(x) = 0.1$
- 10 000 observations
- $\sim 99.5\%$ of time series would contain between 50 and 150 events,
- Naïve estimation: $0.05 < \hat{p}(x) < 0.15$

↪ only $\approx 50\%$ error in $\hat{p}(x)$

↪ small $p(x)$, small count

↪ small count, big uncertainty



Mark Kirstein

Relative estimation error is large for rare events

Asymptotic probability	Most likely count	Standard error in count	Standard error in probability	Relative error in probability
0.1	1000	32	0.003	3%
0.01	100	10	0.001	10%
0.001	10	3	0.0003	30%
0.0001	1	1	0.0001	100%

Table: $T = 10\,000$, assuming Poisson statistics, relative estimation errors $\sim 1/\sqrt{\text{count}}$



To avoid surprises, let's say DMs add estimation uncertainty *err* to every estimated probability, then normalize, s.t.

$$w(x) = \frac{p(x) + \varepsilon[p(x)]}{\int (p(s) + \varepsilon[p(s)]) ds}$$

This allows us to derive a functional form, e.g. for the Gaussian case ... (find in manuscript)

...very similar to function chosen by Kahneman and Tversky.

Not sure we need much more. I'd just have one figure that gives a nice inverse S, for a Gaussian, say, based on estimation error.



Mark Kirstein

Ergodicity Economics and probability weighting

- inverse-S shape appears as neutral indicator of a difference in opinion
- reported observations consistent with DM's extra uncertainty
- may arise from DM estimating probabilities over time

↪ Probability weighting is rational cautious behaviour under uncertainty over time

- See full paper at bit.ly/lml-pw-r1
- links to play with the code are inside

Main Result

Probability
Weighting

Ergodicity
Question

Estimation

Conclusion



Mark Kirstein

Back Up

Thank you for your attention!

I'm looking forward to the discussion
Comments & questions are very welcome, here or to

✉ m.kirstein@lml.org.uk

🐦 [@nonergodicMark](https://twitter.com/nonergodicMark)

WE NEED YOU!



Submit an open peer review to this paper on
bit.ly/lml-pw-r1

RESEARCHERS.ONE



Mark Kirstein

Back Up

BACK UP



Probability Weighting as an Estimation Issue

“It is important to distinguish **overweighting**, which refers to a property of decision weights, from the **overestimation** that is commonly found in the assessment of the probability of rare events. [...] In many real-life situations, overestimation and overweighting may both operate to increase the impact of rare events.” (**KahnemanTversky1979**)

↪ distinguish between

- uncertainty estimation and
- “weighting”

we analyse the former and find very good agreement with the empirical inverse-S pattern


↪ How big is the residual “probability weighting” after accounting for uncertainty estimation?




Mark Kirstein

Estimation Error Explains 99% of Probability Weighting

Tversky & Kahneman (1992)



Tversky & Fox (1995)



- similar fits of Gaussian & t -distributed model

→ How big is the residual “probability weighting” after accounting for estimation errors?

**TverskyKahneman1992**

$$\tilde{F}_w^{TK}(F_p; \gamma) = (F_p)^\gamma \frac{1}{\left[(F_p)^\gamma + (1 - F_p)^\gamma\right]^{1/\gamma}} \quad (1)$$

LattimoreBakerWitte1992

$$\tilde{F}_w^L(F_p; \delta, \gamma) = \frac{\delta F_p^\gamma}{\delta F_p^\gamma + (1 - F_p)^\gamma} \quad (2)$$

We derive decision weight as a function of probability with $(\alpha\sigma)^2$ as the DM's scale

$$w(p) = p^{\frac{1}{\alpha^2}} \frac{(2\pi\sigma^2)^{\frac{1-\alpha^2}{2\alpha^2}}}{\alpha}, \quad (3)$$

which is a power law in p with a pre-factor to ensure normalisation



Linking Probability Weighting to Relative Uncertainties

Decision weight w is the normalised sum of the probability $p(x)$ and its uncertainty $\varepsilon [p(x)]$

$$w(x) = \frac{p(x) + \varepsilon [p(x)]}{\int_{-\infty}^{\infty} (p(s) + \varepsilon [p(s)]) ds} . \quad (4)$$

This can be expressed as

$$w(x) = p(x) \left(\frac{1 + \frac{\varepsilon[p(x)]}{p(x)}}{\int_{-\infty}^{\infty} p(s) \left\{ 1 + \frac{\varepsilon[p(s)]}{p(s)} \right\} ds} \right) , \quad (5)$$

where $\frac{\varepsilon[p(x)]}{p(x)}$ is the relative error, which is large (small) for small (large) probabilities
In the long-time limit $w(x) \rightarrow p(x)$



Mark Kirstein

Back Up

References I