

# New analytic rotations for bifactor modeling and metric invariance in Exploratory Factor Analysis

Marcos Jiménez <sup>1</sup>, Francisco J. Abad <sup>1</sup>,  
Eduardo Garcia-Garzon <sup>2</sup>, Luis E. Garrido <sup>3</sup>, Vithor R. Franco <sup>4</sup>

<sup>1</sup> Universidad Autónoma de Madrid, Spain

<sup>2</sup> Shakers, Spain

<sup>3</sup> Pontificia Universidad Católica Madre y Maestra, Dominican Republic

<sup>4</sup> Universidade São Francisco, Brazil



# Only two kinds of factor rotation?

PSYCHOMETRIKA—VOL. 66, NO. 2, 289–306  
JUNE 2001

## A SIMPLE GENERAL PROCEDURE FOR ORTHOGONAL ROTATION

ROBERT I. JENNRICH

UNIVERSITY OF CALIFORNIA AT LOS ANGELES


PSYCHOMETRIKA—VOL. 67, NO. 1, 7–20  
MARCH 2002

## A SIMPLE GENERAL METHOD FOR OBLIQUE ROTATION

ROBERT I. JENNRICH

UNIVERSITY OF CALIFORNIA AT LOS ANGELES

# The gradient projection method



$$\text{Rotated } (\Lambda) = \text{Unrotated } (U) \times \text{Rotation matrix } (X)$$

$$\begin{pmatrix} \mathbf{0.37} & 0.07 & 0.00 \\ \mathbf{0.57} & 0.01 & 0.02 \\ \mathbf{0.53} & -0.03 & -0.02 \\ -0.02 & \mathbf{0.47} & -0.03 \\ 0.01 & \mathbf{0.44} & 0.01 \\ 0.01 & \mathbf{0.57} & 0.01 \\ -0.05 & 0.00 & \mathbf{0.52} \\ -0.02 & 0.00 & \mathbf{0.56} \\ 0.04 & 0.00 & \mathbf{0.63} \end{pmatrix} = \begin{pmatrix} 0.04 & -0.26 & -0.27 \\ 0.09 & -0.32 & -0.46 \\ 0.04 & -0.27 & -0.46 \\ -0.07 & -0.38 & 0.28 \\ -0.01 & -0.37 & 0.23 \\ -0.02 & -0.48 & 0.31 \\ 0.51 & 0.02 & 0.09 \\ 0.56 & 0.01 & 0.07 \\ 0.63 & -0.01 & 0.03 \end{pmatrix} \begin{pmatrix} -0.13 & -0.06 & 1 \\ 0.55 & -0.83 & 0.01 \\ 0.82 & 0.56 & 0.10 \end{pmatrix}$$

Orthogonal Projection

$$P(G) = G - 0.5(XG^T + X^T G)X$$

$$X^T X = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$


Oblique Projection

$$P(G) = G - X \text{diag}(X^T G)$$

$$X^T X = \begin{pmatrix} 1 & .35 & .44 \\ .35 & 1 & .14 \\ .44 & .14 & 1 \end{pmatrix}$$

# The Partially Oblique Manifold

Rotated ( $\Lambda$ ) = Unrotated ( $\mathbf{U}$ )  $\times$  Rotation matrix ( $\mathbf{X}$ )



$$\begin{pmatrix} \mathbf{0.37} & 0.07 & 0.00 \\ \mathbf{0.57} & 0.01 & 0.02 \\ \mathbf{0.53} & -0.03 & -0.02 \\ -0.02 & \mathbf{0.47} & -0.03 \\ 0.01 & \mathbf{0.44} & 0.01 \\ 0.01 & \mathbf{0.57} & 0.01 \\ -0.05 & 0.00 & \mathbf{0.52} \\ -0.02 & 0.00 & \mathbf{0.56} \\ 0.04 & 0.00 & \mathbf{0.63} \end{pmatrix} = \begin{pmatrix} 0.04 & -0.26 & -0.27 \\ 0.09 & -0.32 & -0.46 \\ 0.04 & -0.27 & -0.46 \\ -0.07 & -0.38 & 0.28 \\ -0.01 & -0.37 & 0.23 \\ -0.02 & -0.48 & 0.31 \\ 0.51 & 0.02 & 0.09 \\ 0.56 & 0.01 & 0.07 \\ 0.63 & -0.01 & 0.03 \end{pmatrix} \begin{pmatrix} -0.13 & -0.06 & 1 \\ 0.55 & -0.83 & 0.01 \\ 0.82 & 0.56 & 0.10 \end{pmatrix}$$




Partially Oblique Projection

$$P(\mathbf{G}) = \mathbf{G} - \mathbf{X} \left( I \circ \mathbf{V} \left( \frac{\mathbf{V}^\top \mathbf{Q} \mathbf{V}}{\mathbf{1d}^\top + \mathbf{d1}^\top} \right) \mathbf{V}^\top \right)$$

$$\mathbf{X}^\top \mathbf{X} = \begin{pmatrix} 1 & \mathbf{0} & .44 \\ \mathbf{0} & 1 & .14 \\ .44 & .14 & 1 \end{pmatrix}$$

## Exploratory Bi-factor Model with Multiple General factors


- [✓] Disentangle the variance due to the general and specific factors.
- [✓] Use any rotation criteria that you want (i.e., target, oblimin, etc.).



Factor Correlation Matrix

$$\begin{pmatrix} 1 & .30 & 0 & 0 & 0 & 0 \\ .30 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & .30 & .30 & .30 \\ 0 & 0 & .30 & 1 & .30 & .30 \\ 0 & 0 & .30 & .30 & 1 & .30 \\ 0 & 0 & .30 & .30 & .30 & 1 \end{pmatrix}$$

## Estimation with the bifactor package



```
efast(Sigma, nfactors = 6, estimator = "ULS",
      rotation = "target", Target = Target,
      projection = "poblq", oblq_factors = c(2, 4))


rotate(Lambda, rotation = "target", Target = Target,
      projection = "poblq", oblq_factors = c(2, 4))
```



## Mixed rotation criteria

[✓] Use Oblimin for the general factors and Target for the specific factors.

	Oblimin		Target			
	Neur	Extr	Ans	Depr	Soc	Aser
X1						
X2						
X3						
X4						
X5						
X6						
X7						
X8						



```
efast(Sigma, nfactors = 6, estimator = "ULS", Target = Target,
      projection = "poblq", oblq_factors = c(2, 4),
      rotation = c("oblimin", "target"), blocks = list(1:2, 3:6))
```

# Multitrait-Multimethod analysis

Trait	Trait	Trait	Method	Method
<b>0.37</b>	0.07	0.00	<b>0.56</b>	0.00
<b>0.57</b>	0.01	0.02	<b>0.56</b>	0.00
<b>0.53</b>	0.03	0.02	0.00	<b>0.51</b>
<b>0.49</b>	0.02	0.03	0.00	<b>0.43</b>
0.02	<b>0.47</b>	0.03	<b>0.61</b>	0.00
0.01	<b>0.44</b>	0.01	<b>0.45</b>	0.00
0.01	<b>0.57</b>	0.01	0.00	<b>0.39</b>
0.08	<b>0.34</b>	0.08	0.00	<b>0.55</b>
0.05	0.00	<b>0.52</b>	<b>0.47</b>	0.00
0.02	0.00	<b>0.56</b>	<b>0.67</b>	0.00
0.04	0.00	<b>0.63</b>	0.00	<b>0.44</b>
0.06	0.07	<b>0.58</b>	0.00	<b>0.50</b>

Loading Matrix

	Trait	Trait	Trait	Method	Method
Trait	1	<b>.33</b>	<b>.45</b>	0	0
Trait	<b>.33</b>	1	<b>.25</b>	0	0
Trait	<b>.45</b>	<b>.25</b>	1	0	0
Method	0	0	0	1	<b>.44</b>
Method	0	0	0	<b>.44</b>	1

Factor Correlation Matrix

```
efast(Sigma, nfactors = 5, estimator = "ULS",
      rotation = "target", Target = Target,
      projection = "poblq", oblq_factors = c(3, 2))
```



## Model identification in CFA

[X] warning: covariance matrix of latent variables is not positive definite

[✓] Solution: Parameterize the covariance matrix of latent variables as  $\mathbf{X}^\top \mathbf{X}$  and optimize it on the partially oblique manifold.

$$\begin{pmatrix} 1 & .33 & .45 & 0 & .68 \\ .33 & 1 & .25 & 0 & .77 \\ .45 & .25 & 1 & .56 & 0 \\ 0 & 0 & .56 & 1 & .44 \\ .68 & .77 & 0 & .44 & 1 \end{pmatrix}$$

Factor correlation matrix optimized  
on the Partially Oblique manifold

# Summary

Applications of the Partially Oblique manifold:

- [✓] Uncorrelate the general and specific factors in Exploratory Bi-factor analysis with either one or multiple general factors.
- [✓] Uncorrelate trait and method factors in multitrait-multimethod analysis.
- [✓] Identify CFA models where the correlations between the latent factors are, at least, positive semi-definite.
- [✓] Similar rotations can be developed to obtain a method for metric invariance in EFA.


# Thank you

Contact: [marcosjnezhquez@gmail.com](mailto:marcosjnezhquez@gmail.com)

 <https://github.com/Marcosjnezh/bifactor>

 @skeptrpsych

Webpage: <https://marcosjnezh.github.io/>



# References

- Jennrich, R. I. (2002). A simple general method for oblique rotation. *Psychometrika*, 67(1), 7–19.  
<https://doi.org/10.1007/BF02294706>
- Jennrich, R. I. (2001). A simple general procedure for orthogonal rotation. *Psychometrika*, 66(2), 289–306. <https://doi.org/10.1007/BF02294840>
- Jiménez, M., Abad, F. J., Garcia-Garzon, E., & Garrido, L. E. (2023). Exploratory bi-factor analysis with multiple general factors. *Multivariate Behavioral Research*, 0(0), 1–18.  
<https://doi.org/10.1080/00273171.2023.2189571>
- Jiménez, M., Abad, F. J., Garcia-Garzon, E., Garrido, L. E., & Franco, V. R. (2022). *Bifactor: Exploratory factor and bi-factor modeling with multiple general factors* [Manual].  
<https://github.com/Marcosjnez/bifactor>
- Jiménez, M., Abad, F. J., Garcia-Garzon, E., Golino, H., Christensen, A. P., & Garrido, L. E. (2023). Dimensionality assessment in bifactor structures with multiple general factors: A network psychometrics approach. *Psychological Methods*, No Pagination Specified–No Pagination Specified. <https://doi.org/10.1037/met0000590>