

# DM549 and DS(K)820

## Lecture 16: Sequences and Summations

Kevin Aguyar Brix

Email: `kabrix@imada.sdu.dk`

University of Southern Denmark

6+7 November, 2024

# Repetition: Recursive Definition

**Recursive Definitions:** They consist of

- base step,
- recursive step.

# Repetition: Recursive Definition

**Recursive Definitions:** They consist of

- base step,
- recursive step.

**We defined recursively:**

- the set of all bitstrings,

# Repetition: Recursive Definition

**Recursive Definitions:** They consist of

- base step,
- recursive step.

**We defined recursively:**

- the set of all bitstrings,
- the set of all palindrome bitstrings,

# Repetition: Recursive Definition

**Recursive Definitions:** They consist of

- base step,
- recursive step.

**We defined recursively:**

- the set of all bitstrings,
- the set of all palindrome bitstrings,
- the set  $\{3n \mid n \in \mathbb{Z}^+\}$ ,

# Repetition: Recursive Definition

**Recursive Definitions:** They consist of

- base step,
- recursive step.

**We defined recursively:**

- the set of all bitstrings,
- the set of all palindrome bitstrings,
- the set  $\{3n \mid n \in \mathbb{Z}^+\}$ ,
- the set of full binary trees,

# Repetition: Recursive Definition

**Recursive Definitions:** They consist of

- base step,
- recursive step.

**We defined recursively:**

- the set of all bitstrings,
- the set of all palindrome bitstrings,
- the set  $\{3n \mid n \in \mathbb{Z}^+\}$ ,
- the set of full binary trees,
- the height of a full binary tree.

# Repetition: Structural Induction

We proved statements on recursively defined structures in the following way:

## Recipe for Proofs by Structural Induction

To show that  $P(S_i)$  holds for all  $i \geq 1$ , prove:

- Basis step: Prove that  $P(S_1)$  holds.
- Inductive step: Prove that

$$\underbrace{P(S_i)}_{\text{inductive hypothesis}} \Rightarrow P(S_{i+1})$$

for all  $i \geq 1$ .

**That is:** Induction on the number of times the recursive step is applied.



# Sequences

## Definition (Definition 2.4.1)

A *sequence* (følge) is a function from a subset of  $\mathbb{N}$  to some set.

### Remarks:

- The domain of the function may be finite or infinite; it is usually  $\{0, 1, 2, \dots\}$  or  $\{1, 2, 3, \dots\}$ .

## Definition (Definition 2.4.1)

A *sequence* (følge) is a function from a subset of  $\mathbb{N}$  to some set.

### Remarks:

- The domain of the function may be finite or infinite; it is usually  $\{0, 1, 2, \dots\}$  or  $\{1, 2, 3, \dots\}$ .
- One also denotes by  $a_n$  the number that the function maps  $n$  to (also called a *term* of the sequence).
  - ▶ Cf. our notation for Fibonacci sequence!

## Definition (Definition 2.4.1)

A *sequence* (følge) is a function from a subset of  $\mathbb{N}$  to some set.

### Remarks:

- The domain of the function may be finite or infinite; it is usually  $\{0, 1, 2, \dots\}$  or  $\{1, 2, 3, \dots\}$ .
- One also denotes by  $a_n$  the number that the function maps  $n$  to (also called a *term* of the sequence).
  - ▶ Cf. our notation for Fibonacci sequence!
- The sequence is then denoted by  $\{a_n\}$ .
  - ▶ Do not confuse with set notation!

## Definition (Definition 2.4.1)

A *sequence* (følge) is a function from a subset of  $\mathbb{N}$  to some set.

### Remarks:

- The domain of the function may be finite or infinite; it is usually  $\{0, 1, 2, \dots\}$  or  $\{1, 2, 3, \dots\}$ .
- One also denotes by  $a_n$  the number that the function maps  $n$  to (also called a *term* of the sequence).
  - ▶ Cf. our notation for Fibonacci sequence!
- The sequence is then denoted by  $\{a_n\}$ .
  - ▶ Do not confuse with set notation!
- One also lists out the terms in order, e.g., for the domain being  $\mathbb{N}$ :

$$\{a_n\} = a_0, a_1, a_2, a_3, \dots$$

## Definition (Definition 2.4.1)

A *sequence* (følge) is a function from a subset of  $\mathbb{N}$  to some set.

### Remarks:

- The domain of the function may be finite or infinite; it is usually  $\{0, 1, 2, \dots\}$  or  $\{1, 2, 3, \dots\}$ .
- One also denotes by  $a_n$  the number that the function maps  $n$  to (also called a *term* of the sequence).
  - ▶ Cf. our notation for Fibonacci sequence!
- The sequence is then denoted by  $\{a_n\}$ .
  - ▶ Do not confuse with set notation!
- One also lists out the terms in order, e.g., for the domain being  $\mathbb{N}$ :

$$\{a_n\} = a_0, a_1, a_2, a_3, \dots$$

- You can also think of a sequence with domain  $D$  as a  $|D|$ -tuple.

# Geometric and Arithmetic Sequences

## Definition (Definition 2.4.2)

An infinite *geometric sequence* (geometrisk følge) is a sequence of the form

$$a_n = c \cdot r^n, \quad n \in \mathbb{N},$$

where  $a \in \mathbb{R}$  is the *initial term* (begyndelsesled) and  $r \in \mathbb{R}$  is the *common ratio* (fælles faktor).

# Geometric and Arithmetic Sequences

## Definition (Definition 2.4.2)

An infinite *geometric sequence* (geometrisk følge) is a sequence of the form

$$a_n = c \cdot r^n, \quad n \in \mathbb{N},$$

where  $a \in \mathbb{R}$  is the *initial term* (begyndelsesled) and  $r \in \mathbb{R}$  is the *common ratio* (fælles faktor). We obtain finite geometric sequences by stopping at some point.



# Geometric and Arithmetic Sequences

## Definition (Definition 2.4.2)

An infinite *geometric sequence* (geometrisk følge) is a sequence of the form

$$a_n = c \cdot r^n, \quad n \in \mathbb{N},$$

where  $a \in \mathbb{R}$  is the *initial term* (begyndelsesled) and  $r \in \mathbb{R}$  is the *common ratio* (fælles faktor). We obtain finite geometric sequences by stopping at some point.

## Definition (Definition 2.4.3)

An infinite *arithmetic sequence* (aritmetisk følge) is a sequence of the form

$$a_n = b + n \cdot d, \quad n \in \mathbb{N},$$

where  $b \in \mathbb{R}$  is the *initial term* (begyndelsesled) and  $d \in \mathbb{R}$  is the *common difference* (fælles forskel).

# Geometric and Arithmetic Sequences

## Definition (Definition 2.4.2)

An infinite *geometric sequence* (geometrisk følge) is a sequence of the form

$$a_n = c \cdot r^n, \quad n \in \mathbb{N},$$

where  $a \in \mathbb{R}$  is the *initial term* (begyndelsesled) and  $r \in \mathbb{R}$  is the *common ratio* (fælles faktor). We obtain finite geometric sequences by stopping at some point.

## Definition (Definition 2.4.3)

An infinite *arithmetic sequence* (aritmetisk følge) is a sequence of the form

$$a_n = b + n \cdot d, \quad n \in \mathbb{N},$$

where  $b \in \mathbb{R}$  is the *initial term* (begyndelsesled) and  $d \in \mathbb{R}$  is the *common difference* (fælles forskel). We obtain finite arithmetic sequences by stopping at some point.

# The Mandelbrot Fractal (not relevant for exam)

## Idea:

- For every  $z \in \mathbb{C}$  ( $\mathbb{C}$  is the set of complex numbers)

# The Mandelbrot Fractal (not relevant for exam)

## Idea:

- For every  $z \in \mathbb{C}$  ( $\mathbb{C}$  is the set of complex numbers), define a sequence as follows:
  - ▶  $x_0 = 0$ ,
  - ▶  $x_n = x_{n-1}^2 + z$  for  $n \geq 1$ .

# The Mandelbrot Fractal (not relevant for exam)


## Idea:

- For every  $z \in \mathbb{C}$  ( $\mathbb{C}$  is the set of complex numbers), define a sequence as follows:
  - ▶  $x_0 = 0$ ,
  - ▶  $x_n = x_{n-1}^2 + z$  for  $n \geq 1$ .
- If  $|x_n|$  does not grow arbitrarily large,  $z$  belongs to the *Mandelbrot set*.

# The Mandelbrot Fractal (not relevant for exam)

## Idea:

- For every  $z \in \mathbb{C}$  ( $\mathbb{C}$  is the set of complex numbers), define a sequence as follows:
  - ▶  $x_0 = 0$ ,
  - ▶  $x_n = x_{n-1}^2 + z$  for  $n \geq 1$ .
- If  $|x_n|$  does not grow arbitrarily large,  $z$  belongs to the *Mandelbrot set*.
- The Mandelbrot set in the complex plane is depicted in black:



(Source: Wikipedia)

The colors encode how many iterations it takes for  $|x_n|$  to surpass 1000.

## Definition

Let

$$a_m, a_{m+1}, \dots, a_n$$

be a sequence. Then there is an associated *series* (række), the sum of all terms in the sequence. It is denoted by

$$\sum_{i=m}^n a_i \quad \text{or} \quad \sum_{m \leq i \leq n} a_i.$$

## Definition

Let

$$a_m, a_{m+1}, \dots, a_n$$

be a sequence. Then there is an associated *series* (række), the sum of all terms in the sequence. It is denoted by

$$\sum_{i=m}^n a_i \quad \text{or} \quad \sum_{m \leq i \leq n} a_i.$$

**Remark:**

- If the domain of the sequence is  $D$ , we also write  $\sum_{i \in D} a_i$ .



## Definition

Let

$$a_m, a_{m+1}, \dots, a_n$$

be a sequence. Then there is an associated *series* (række), the sum of all terms in the sequence. It is denoted by

$$\sum_{i=m}^n a_i \quad \text{or} \quad \sum_{m \leq i \leq n} a_i.$$

## Remark:

- If the domain of the sequence is  $D$ , we also write  $\sum_{i \in D} a_i$ .
- One could also talk about series that are the sum of the infinitely many terms of an infinite sequence.

## Definition

Let

$$a_m, a_{m+1}, \dots, a_n$$

be a sequence. Then there is an associated *series* (række), the sum of all terms in the sequence. It is denoted by

$$\sum_{i=m}^n a_i \quad \text{or} \quad \sum_{m \leq i \leq n} a_i.$$

## Remark:

- If the domain of the sequence is  $D$ , we also write  $\sum_{i \in D} a_i$ .
- One could also talk about series that are the sum of the infinitely many terms of an infinite sequence.
- Here, we focus on finite sequences.

## Definition

Let

$$a_m, a_{m+1}, \dots, a_n$$

be a sequence. Then there is an associated *series* (række), the sum of all terms in the sequence. It is denoted by

$$\sum_{i=m}^n a_i \quad \text{or} \quad \sum_{m \leq i \leq n} a_i.$$

## Remark:

- If the domain of the sequence is  $D$ , we also write  $\sum_{i \in D} a_i$ .
- One could also talk about series that are the sum of the infinitely many terms of an infinite sequence.
- Here, we focus on finite sequences.
- Otherwise, to be completely formal, we would need to talk about a concept from calculus called convergence.

# Geometric Series

## Theorem (Theorem 2.4.1)

For finite *geometric series* (with  $c = 1$ ), the series corresponding to finite geometric sequences, it holds that

$$\sum_{i=0}^n r^i =$$

# Geometric Series

## Theorem (Theorem 2.4.1)

For finite *geometric series* (with  $c = 1$ ), the series corresponding to finite geometric sequences, it holds that

$$\sum_{i=0}^n r^i = \begin{cases} \frac{r^{n+1}-1}{r-1} & \text{if } r \in \mathbb{R} \setminus \{1\}, \end{cases}$$

# Geometric Series

## Theorem (Theorem 2.4.1)

For finite *geometric series* (with  $c = 1$ ), the series corresponding to finite geometric sequences, it holds that

$$\sum_{i=0}^n r^i = \begin{cases} \frac{r^{n+1}-1}{r-1} & \text{if } r \in \mathbb{R} \setminus \{1\}, \\ n+1 & \text{if } r = 1. \end{cases}$$

# Geometric Series

## Theorem (Theorem 2.4.1)

For finite *geometric series* (with  $c = 1$ ), the series corresponding to finite geometric sequences, it holds that

$$\sum_{i=0}^n r^i = \begin{cases} \frac{r^{n+1}-1}{r-1} & \text{if } r \in \mathbb{R} \setminus \{1\}, \\ n+1 & \text{if } r = 1. \end{cases}$$

**Note:** If  $|r| < 1$  and we consider the *infinite* geometric series

# Geometric Series

## Theorem (Theorem 2.4.1)

For finite *geometric series* (with  $c = 1$ ), the series corresponding to finite geometric sequences, it holds that

$$\sum_{i=0}^n r^i = \begin{cases} \frac{r^{n+1}-1}{r-1} & \text{if } r \in \mathbb{R} \setminus \{1\}, \\ n+1 & \text{if } r = 1. \end{cases}$$

**Note:** If  $|r| < 1$  and we consider the *infinite* geometric series, the term  $r^{n+1}$  vanishes as  $n$  grows to  $\infty$ , so

$$\sum_{i=0}^{\infty} r^i = \frac{1}{1-r}.$$



# Arithmetic Series

## Theorem

For finite *arithmetic series*, the series corresponding to finite arithmetic sequences, it holds that

$$\sum_{i=0}^n (b + i \cdot d) =$$

# Arithmetic Series

## Theorem

For finite *arithmetic series*, the series corresponding to finite arithmetic sequences, it holds that

$$\sum_{i=0}^n (b + i \cdot d) = b \cdot (n + 1) + d \cdot \frac{n \cdot (n + 1)}{2}$$

# Arithmetic Series

## Theorem

For finite *arithmetic series*, the series corresponding to finite arithmetic sequences, it holds that

$$\sum_{i=0}^n (b + i \cdot d) = b \cdot (n + 1) + d \cdot \frac{n \cdot (n + 1)}{2}$$

**Additional formulae:** Table 2.4.2.

# A Quiz

Go to [pollev.com/kevs](https://pollev.com/kevs)

