QSD Supplement S7:

Computational Thermodynamics and the Cosmological Constant

A Speculative Extension

Marek Hubka *

February 26, 2026

Abstract

We present a **speculative but quantitative** extension of QSD to the cosmological constant problem [1]. The central observation is that syndrome extraction—the irreversible measurement process at the heart of QSD's error-correcting spacetime—carries a thermodynamic cost governed by Landauer's principle. The entropy production rate per Planck 4-volume is set by the graph strain: $S_{\text{comp}} = n/\varsigma$ nats, where $n = 4$ is the spacetime dimension. The vacuum energy density is then the Planck density suppressed by the Boltzmann factor for this computational entropy cost:

$$\rho_\Lambda = \rho_{\text{Pl}} \times e^{-4/\varsigma}$$

Numerically, $4/\varsigma \approx 279.3$, giving $e^{-4/\varsigma} \approx 10^{-121.3}$, in striking agreement with the observed $\rho_\Lambda \approx 10^{-121.3}\,\rho_{\text{Pl}}$. The residual vacuum energy drives metric expansion as the only available mechanism for a closed system to accommodate monotonically increasing computational entropy. This extension uses no additional free parameters beyond those already present in QSD, but the derivation rests on physical arguments that are not yet rigorous, and we present it as a conjecture meriting further investigation.

Status: Speculative. All results in this supplement are conjectural and do not affect the established predictions of the QSD core framework (S1–S6).

Contents

1 Motivation: Spacetime as a Computer **2**

2 Landauer's Principle and Holographic Bit Density **3**
2.1 Landauer's Principle 3
2.2 Connection to $\mathbb{H}_4$ 3

3 Irreversibility of Syndrome Extraction **3**
3.1 Unitary Evolution vs. Syndrome Measurement 3
3.2 Error Rate and Correction Frequency 4

*Independent Researcher, Czech Republic.
Website: tidesofuncertainty.com.
Email: marek@tidesofuncertainty.com.

4 Entropy Production Rate **4**
4.1 Counting Irreversible Operations 4
4.2 Interpretation 4

5 The Cosmological Constant **4**
5.1 Boltzmann Suppression 4
5.2 Numerical Evaluation 5
5.3 Comparison with Observation 5

6 The Expansion Mechanism **5**
6.1 Why Expansion? 5
6.2 Self-Consistency with de Sitter Temperature 6

7 Dimensional Dependence **6**
7.1 The General Formula 6
7.2 Why $n = 4$? 6

8 The Prefactor **7**

9 Connection to Established QSD Results **7**

10 What This Would Resolve **7**

11 Falsification Criteria **8**

12 Limitations and Open Questions **8**

13 Summary **9**

1 Motivation: Spacetime as a Computer

QSD treats spacetime as a quantum error-correcting code—specifically, the $[[5, 1, 3]]$ perfect code, whose graph structure produces the complete graph K_4 and the cycle graph C_5. Within this framework, physical processes are computational processes: gauge interactions arise from syndrome extraction on K_4, and geometric dynamics arise from the spectral structure of C_5.

Every real computer dissipates heat as it computes. This is not an engineering limitation but a fundamental consequence of the second law of thermodynamics, formalised by Landauer's principle [2]: erasing one bit of information requires at least $k_B T \ln 2$ of energy dissipated as heat. If spacetime *is* a computer—as QSD proposes—then the continuous error-correction process must produce entropy. The question is: where does this computational heat go?

In an ordinary computer, heat flows to the environment. But the universe *has* no environment. For a closed system, the only way to accommodate monotonically increasing entropy is to expand the available phase space. For spacetime, this means metric expansion.

Observation 1.1 (Computational heat → dark energy)**.** If the universe continuously performs error correction on its own quantum-geometric code, the thermodynamic cost of this computation may manifest as the cosmological constant, and the metric expansion of space may be the physical mechanism by which the resulting entropy is accommodated.

This observation connects three of the deepest problems in physics—the cosmological constant, the arrow of time, and the nature of quantum gravity—through a single mechanism: the thermodynamic cost of spacetime error correction. We develop this idea quantitatively in the following sections.

2 Landauer's Principle and Holographic Bit Density

2.1 Landauer's Principle

Landauer's principle [2] states that the erasure of one bit of information in a system at temperature T requires a minimum energy dissipation of:

$$E_{\text{bit}} = k_B T \ln 2 \tag{1}$$

This is not a practical limitation but a *thermodynamic identity*: information erasure is logically irreversible, and the second law demands entropy increase $\Delta S \geq k_B \ln 2$ per bit erased.

2.2 Connection to Hb_4

The holographic bit density in n dimensions (see S2 and the Redefinition Note) is:

$$\text{Hb}_n = \frac{1}{n \ln 2} \tag{2}$$

For $n = 4$:

$$\text{Hb}_4 = \frac{1}{4 \ln 2} \approx 0.3607 \tag{3}$$

The $\ln 2$ in the denominator is *the same* $\ln 2$ that appears in Landauer's principle. This is not a coincidence: Hb_4 was derived from the Bekenstein–Hawking entropy bound [3, 4] $S = A/(4\ell_P^2)$, which counts the maximum number of bits encodable on a surface. The Landauer cost per bit is $k_B T \ln 2$; the information capacity per Planck area per dimension is $1/(n \ln 2)$. The two are conjugate quantities:

$$\text{Hb}_n \times \ln 2 = \frac{1}{n} \tag{4}$$

This identity states that the Landauer energy cost of encoding one holographic bit, measured in units of $k_B T$, is exactly $1/n$ per dimension. The total cost across all n dimensions is unity.

Remark 2.1. Equation (4) is an exact identity, not an approximation. It shows that Hb_n already encodes the thermodynamic cost of information processing at the Planck scale.

3 Irreversibility of Syndrome Extraction

3.1 Unitary Evolution vs. Syndrome Measurement

In the QSD framework, the dynamics of spacetime involve two qualitatively different operations:

1. **Unitary evolution** (Hamiltonian propagation on $K_4 \cup C_5$): This is reversible and generates no entropy.
2. **Syndrome extraction** (measurement of stabiliser eigenvalues): This is *irreversible*—it projects quantum superpositions onto definite syndrome outcomes, collapsing quantum information into classical syndrome bits.

It is the second process—syndrome extraction—that carries Landauer cost. In the $[[5, 1, 3]]$ code, there are 4 stabiliser generators, each yielding one classical bit of syndrome information per measurement cycle. This is the same process that, in the QSD particle-physics sector, produces observable syndromes at $E_{\text{syn}} = 28.24$ GeV (see S5).

3.2 Error Rate and Correction Frequency

The graph strain $\varsigma = 3/(80\varphi^2) \approx 0.01432$ sets the fractional error rate per coding cycle. In each Planck time interval, the probability that a given degree of freedom experiences an error requiring correction is proportional to ς.

The *number* of syndrome extractions per degree of freedom needed to accumulate one full bit of error entropy is therefore:

$$N_{\text{cycle}} \sim \frac{1}{\varsigma} \tag{5}$$

This means the syndrome measurement apparatus is continuously active, but produces irreversible entropy at rate ς per Planck time per degree of freedom.

4 Entropy Production Rate

4.1 Counting Irreversible Operations

Consider one Planck 4-volume ($\ell_P^3 \times t_P$). The error-correcting code operates on this volume, and the entropy production has contributions from each spacetime dimension.

Per dimension, the holographic encoding capacity is Ⱶ_n bits per Planck area. The error rate per cycle is ς. But the relevant quantity for the total entropy cost is not the per-cycle rate but the accumulated entropy barrier: how many nats of entropy separate the "fully corrected" vacuum from the "uncorrected" state.

Proposition 4.1 (Computational entropy per Planck 4-volume)**.** *The total computational entropy cost of maintaining the error-correcting code in n spacetime dimensions is:*

$$S_{comp} = \frac{n}{\varsigma} \quad nats \tag{6}$$

Argument. Each of the n spacetime dimensions requires independent error correction (the code protects geometric information along each axis). Per dimension, the number of syndrome extraction cycles needed to accumulate one nat of entropy is $1/\varsigma$ (since each cycle produces ς nats of irreversible entropy). The total across n dimensions is n/ς. □

For $n = 4$:

$$S_{\text{comp}} = \frac{4}{\varsigma} = \frac{4 \times 80\varphi^2}{3} = \frac{320\varphi^2}{3} \approx 279.26 \text{ nats} \tag{7}$$

4.2 Interpretation

The quantity $S_{\text{comp}} \approx 279$ nats is the *entropy barrier* that separates the Planck-scale vacuum energy density from the observed vacuum energy density. It is large (explaining why Λ is small) but finite and calculable from graph topology (explaining why Λ is not zero).

5 The Cosmological Constant

5.1 Boltzmann Suppression

In statistical mechanics, a state separated from the ground state by an entropy cost S is suppressed by the Boltzmann factor e^{-S}. The vacuum energy density is the Planck density suppressed by the computational entropy barrier:

Conjecture 5.1 (Computational vacuum energy)**.**

$$\boxed{\rho_\Lambda = \rho_{Pl} \times e^{-n/\varsigma}} \tag{8}$$

where $\rho_{Pl} = c^7/(\hbar G^2) \approx 4.63 \times 10^{113}$ J/m^3 is the Planck energy density, $n = 4$ is the spacetime dimension, and $\varsigma = 3/(80\varphi^2)$ is the graph strain.

5.2 Numerical Evaluation

The exponent:

$$\frac{4}{\varsigma} = \frac{320\varphi^2}{3} = \frac{320 \times \frac{3+\sqrt{5}}{2}}{3} = \frac{160(3+\sqrt{5})}{3} \approx 279.26 \tag{9}$$

Converting to base 10:

$$\frac{4/\varsigma}{\ln 10} = \frac{279.26}{2.3026} \approx 121.28 \tag{10}$$

Therefore:

$$e^{-4/\varsigma} \approx 10^{-121.28} \tag{11}$$

5.3 Comparison with Observation

The observed vacuum energy density in Planck units:

$$\rho_\Lambda^{\text{obs}} \approx 5.4 \times 10^{-122}\, \rho_{\text{Pl}} \approx 10^{-121.27}\, \rho_{\text{Pl}} \tag{12}$$

Quantity	**Value (Planck units)**	$\log_{10}$
$e^{-4/\varsigma}$ (QSD prediction)	$\approx 5.3 \times 10^{-122}$	-121.28
$\rho_\Lambda^{\text{obs}}$ (Planck 2018 [5])	$\approx 5.4 \times 10^{-122}$	-121.27
Ratio	≈ 1.03	$\Delta \approx 0.01$

Table 1: Comparison of the QSD computational vacuum energy with observation. The agreement spans 121 orders of magnitude with a residual factor of order unity.

The agreement is within a factor of ~ 1.03 across 121 orders of magnitude.

Remark 5.2. This is the ratio of two independently determined numbers: ς from graph topology (derived in S1), and $\rho_\Lambda^{\text{obs}}$ from CMB and supernovae data [5, 6, 7]. Neither was tuned to match the other.

6 The Expansion Mechanism

6.1 Why Expansion?

The irreversible entropy production from syndrome extraction must be accommodated somewhere. For a system with an external environment, entropy can be exported as heat. But the universe is a closed system—there is no external thermal reservoir.

The second law requires the total entropy of a closed system to be non-decreasing. For a system at maximum entropy density, the only way to accommodate additional entropy is to increase the volume of the system, thereby increasing the total entropy capacity.

Proposition 6.1 (Expansion as entropy accommodation)**.** *For a closed system producing entropy at a rate $\dot{S}_{comp}$ per unit volume, the metric must expand at a rate sufficient to maintain quasi-equilibrium:*

$$\frac{\dot{V}}{V} \geq \frac{\dot{S}_{comp}}{s_{\max}} \tag{13}$$

where $s_{\max}$ is the maximum entropy density (set by the holographic bound).

This identifies the Hubble expansion rate $H = \dot{a}/a$ with the computational entropy production rate. Dark energy is not a mysterious substance—it is the thermodynamic cost of spacetime maintaining its own quantum coherence.

6.2 Self-Consistency with de Sitter Temperature

A de Sitter universe with Hubble parameter H has a horizon temperature:

$$T_{\mathrm{dS}} = \frac{\hbar H}{2\pi k_B} \tag{14}$$

The Landauer cost per bit at this temperature is:

$$E_{\mathrm{Landauer}} = k_B T_{\mathrm{dS}} \ln 2 = \frac{\hbar H \ln 2}{2\pi} \tag{15}$$

And the vacuum energy density from the Friedmann equation is:

$$\rho_\Lambda = \frac{3H^2 M_{\mathrm{Pl}}^2}{8\pi} \tag{16}$$

Equating with our computational expression $\rho_\Lambda = \rho_{\mathrm{Pl}}\, e^{-4/\varsigma}$ gives a self-consistent H:

$$H = \sqrt{\frac{8\pi\,\rho_{\mathrm{Pl}}}{3M_{\mathrm{Pl}}^2}}\; e^{-2/\varsigma} \tag{17}$$

The square root arises because $\rho_\Lambda \propto H^2$ (Friedmann), while $\rho_\Lambda \propto e^{-4/\varsigma}$ (Landauer–Boltzmann). The system selects the self-consistent solution automatically. That H is exponentially small in $1/\varsigma$ is the correct qualitative behaviour.

7 Dimensional Dependence

7.1 The General Formula

For arbitrary spacetime dimension n, Conjecture 5.1 generalises to:

$$\rho_\Lambda(n) = \rho_{\mathrm{Pl},n} \times e^{-n/\varsigma} \tag{18}$$

where $\rho_{\mathrm{Pl},n}$ is the n-dimensional Planck density.

7.2 Why $n = 4$?

The vacuum energy must satisfy two conditions for structure formation:

1. $\rho_\Lambda > 0$ (expansion, preventing gravitational recollapse)
2. $\rho_\Lambda < \rho_{\mathrm{matter}}$ at the epoch of galaxy formation (so gravity can win locally)

The exponential dependence on n makes this an extremely sharp selection. For $\varsigma \approx 0.0143$:

n	$e^{-n/\varsigma}$	$\log_{10}$
2	$\sim 10^{-60.6}$	too large (dominates matter)
3	$\sim 10^{-90.9}$	marginal
4	$\sim \mathbf{10^{-121.3}}$	**observed**
5	$\sim 10^{-151.5}$	too small (no effect)

Table 2: Vacuum energy density as a function of spacetime dimension. Only $n = 4$ gives a value compatible with the observed late-time cosmic acceleration.

This is not a prediction of $n = 4$ *per se*, but an anthropic consistency check: the formula Eq. (18) is compatible with observers only for $n = 4$.

8 The Prefactor

The bare conjecture $\rho_\Lambda = \rho_{\text{Pl}}\, e^{-4/\varsigma}$ gives $10^{-121.28}$ against the observed $10^{-121.27}$. The residual factor is:

$$C = \frac{\rho_\Lambda^{\text{obs}}}{\rho_{\text{Pl}}\, e^{-4/\varsigma}} \approx 1.03 \tag{19}$$

This is remarkably close to unity, but a complete derivation should account for it. Several candidate contributions, each of order unity:

1. **Stabiliser generators:** The $[[5, 1, 3]]$ code has $n_{\text{stab}} = 4$ stabiliser generators. The entropy cost might be $(n - c)/\varsigma$ where c is a small correction from the code redundancy.
2. **Holographic prefactor:** The Bekenstein–Hawking formula [3] includes a factor $1/4$ (i.e. $S = A/(4\ell_P^2)$). This is already captured in $\text{Hb}_4 = 1/(4 \ln 2)$, but its interplay with the exponential could produce an $O(1)$ correction.
3. **Edge multiplicity:** K_4 has 6 edges; the ratio $6/4 = 3/2$ or its inverse might appear as a combinatorial correction to the entropy counting.

At the present speculative stage, we note that $C \approx 1$ and defer the prefactor analysis to future work. The key result—121 orders of magnitude from a single topological expression—is robust against $O(1)$ corrections.

9 Connection to Established QSD Results

The Landauer mechanism does not disturb any existing QSD predictions. It introduces one new formula (Eq. 8) whose only inputs are quantities already defined in the core framework:

Input	**Defined in**	**Value**
n (spacetime dimension)	Assumed	4
ς (graph strain)	S1, Main Preprint §3	$3/(80\varphi^2) \approx 0.01432$
ρ_{Pl} (Planck density)	Standard physics	4.63×10^{113} J/m^3

Table 3: All inputs to the vacuum energy formula.

The following existing QSD structures acquire new roles:

- $\text{Hb}_4 = 1/(4 \ln 2)$ encodes the Landauer cost per dimension (Eq. 4).
- ς controls both the particle-physics error rate (syndromes, coupling corrections) and the cosmological entropy barrier.
- The syndrome extraction process (S5) is simultaneously the origin of the 28 GeV excess *and* of the vacuum energy.
- The n-dimensional generalisation $\text{Hb}_n = 1/(n \ln 2)$ (Redefinition Note) provides the framework for exploring the dimensional dependence.

10 What This Would Resolve

If Conjecture 5.1 is correct, it addresses three longstanding problems simultaneously:

1. **The cosmological constant problem [1].** The standard question is: why is $\rho_\Lambda/\rho_{\text{Pl}} \sim 10^{-122}$ rather than $O(1)$? In QFT, this requires cancellation of $\sim 10^{122}$ zero-point contributions to 122 decimal places. In QSD, the smallness is an exponential Boltzmann suppression: $e^{-4/\varsigma}$ is naturally small because the entropy barrier $4/\varsigma \approx 279$ is naturally large (but finite and calculable).

2. **The coincidence problem.** Why is $\rho_\Lambda \sim \rho_{\text{matter}}$ today? In QSD, both the matter sector and the vacuum energy arise from the same graph strain ς: matter masses come from the code-scale hierarchy (Main Preprint §4), while ρ_Λ comes from the computational entropy cost. The "coincidence" reflects the fact that a single parameter governs both scales.

3. **The arrow of time.** Irreversible syndrome extraction produces entropy monotonically. Metric expansion accommodates this entropy. Time's arrow is thus grounded in the computational irreversibility of error correction, not in special initial conditions.

11 Falsification Criteria

Despite being speculative, this extension makes specific claims that are in principle testable:

Prediction	**QSD (S7)**	**Falsified if**
$\rho_\Lambda/\rho_{\text{Pl}}$	$e^{-4/\varsigma} \approx 10^{-121.3}$	Measured ρ_Λ deviates by $>10\times$
w (equation of state)	-1 (exactly)	$w \neq -1$ confirmed
Time variation of Λ	Constant	$\dot{\Lambda}/\Lambda \neq 0$ observed
Dimensional dependence	$\rho_\Lambda \propto e^{-n/\varsigma}$	Inconsistent with string landscape

Table 4: Falsification criteria for the computational vacuum energy conjecture.

The prediction $w = -1$ exactly follows from the Boltzmann-suppression mechanism: the computational entropy production rate is set by topology, which does not evolve with the scale factor. This distinguishes the QSD mechanism from quintessence models (which generically predict $w \neq -1$) and is testable by next-generation surveys (DESI, Euclid, Roman).

12 Limitations and Open Questions

We emphasise that this supplement is speculative. Key limitations:

1. **The entropy-counting argument is heuristic.** Proposition 4.1 gives $S_{\text{comp}} = n/\varsigma$ by dimensional counting, not by explicit computation from the stabiliser algebra. A rigorous derivation should compute the von Neumann entropy production per syndrome extraction cycle from the $[[5, 1, 3]]$ stabiliser group and sum over all stabiliser generators and all Planck cells.

2. **The Boltzmann interpretation is non-standard.** Applying the Boltzmann suppression e^{-S} to vacuum energy density is motivated by analogy with statistical mechanics, but a cosmological "partition function" from which this factor emerges has not been constructed. The argument is plausible but not proven.

3. **The prefactor is undetermined.** The $O(1)$ factor $C \approx 1.03$ might be exactly 1, or might encode additional code-theoretic structure. This requires further analysis.

4. **Backreaction is ignored.** We treat ς as constant, but in principle the computational entropy production could modify the effective graph strain at cosmological scales. Whether ς runs with the scale factor is an open question.

5. **Relation to other approaches.** The Landauer mechanism has conceptual overlap with Padmanabhan's emergent gravity programme [8] and with Verlinde's entropic gravity [9]. Whether QSD provides a microscopic completion of these ideas, or is an independent framework, remains to be established.

13 Summary

We have proposed a speculative but quantitative connection between QSD's error-correcting spacetime and the cosmological constant:

1. Syndrome extraction is irreversible (Landauer's principle applies).
2. The entropy production rate per Planck 4-volume is $4/\varsigma \approx 279$ nats.
3. The vacuum energy is Planck-density suppressed by $e^{-4/\varsigma} \approx 10^{-121.3}$.
4. This matches observation ($10^{-121.3}$) to within a factor of ~ 1.03.
5. The mechanism requires *zero* additional parameters.
6. Metric expansion is the physical mechanism accommodating computational entropy in a closed universe.

If correct, this resolves the cosmological constant problem, the coincidence problem, and the arrow of time within a single framework. The key insight is that $\text{Hb}_4 = 1/(4 \ln 2)$ already contains Landauer's $\ln 2$—the holographic bit density and the thermodynamic cost of computation are two faces of the same constant.

The formula $\rho_\Lambda = \rho_{Pl} e^{-4/\varsigma}$ is the simplest possible statement: the vacuum energy is the Planck density, attenuated by the computational cost of running the universe.

References

[1] Steven Weinberg. The cosmological constant problem. *Rev. Mod. Phys.*, 61:1–23, 1989.

[2] Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM J. Res. Dev.*, 5:183–191, 1961.

[3] Jacob D. Bekenstein. Black holes and entropy. *Phys. Rev. D*, 7:2333–2346, 1973.

[4] Stephen W. Hawking. Particle creation by black holes. *Commun. Math. Phys.*, 43:199–220, 1975.

[5] Planck Collaboration, N. Aghanim, et al. Planck 2018 results. VI. cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020.

[6] Saul Perlmutter et al. Measurements of ω and λ from 42 high-redshift supernovae. *Astrophys. J.*, 517:565–586, 1999.

[7] Adam G. Riess et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astron. J.*, 116:1009–1038, 1998.

[8] Thanu Padmanabhan. Thermodynamical aspects of gravity: new insights. *Rep. Prog. Phys.*, 73:046901, 2010.

[9] Erik Verlinde. On the origin of gravity and the laws of Newton. *JHEP*, 1104:029, 2011.