QSD Supplement S6:

Correspondence with String Theory

A Structural Isomorphism and Invitation to Investigation

Marek Hubka *

February 26, 2026

Abstract

We identify a striking structural correspondence between Quantum Spectrodynamics (QSD) and key elements of String Theory. The 6 edges of the K_4 graph map to the 6 compactified dimensions of Calabi-Yau manifolds; the Anchor qubit corresponds to a D-brane; the geometric strain ς plays the role of the string coupling g_s; and the 3-loop termination of QSD renormalization mirrors the modular invariance that renders String Theory UV-finite. We present this correspondence as a dictionary relating QSD concepts to their string-theoretic analogs, and we frame the result as an *invitation* to the string theory community: the code-stability condition of the $[[5, 1, 3]]$ perfect quantum code may provide the long-sought vacuum selection mechanism that resolves the Landscape problem.

This supplement does not claim to *derive* String Theory results. It identifies patterns that suggest a deeper duality, and it poses precise questions that string-theoretic techniques could answer.

Keywords: string theory, Calabi-Yau, D-branes, moduli stabilization, landscape problem, quantum error correction, holography

Contents

1 Preface: Scope and Framing **3**

2 Motivation: Two Paths to the Same Structure **3**
- 2.1 The Landscape Problem . . . 3
- 2.2 The QSD Approach . . . 3
- 2.3 The Structural Coincidence . . . 3

3 Background: String Theory Essentials **4**
- 3.1 From Points to Strings . . . 4
- 3.2 Extra Dimensions and Compactification . . . 4
- 3.3 D-Branes . . . 4
- 3.4 Moduli and the Landscape . . . 5
- 3.5 Modular Invariance and UV Finiteness . . . 5

*Independent Researcher, Czech Republic.
Website: tidesofuncertainty.com.
Email: marek@tidesofuncertainty.com.

4 The Dimensional Correspondence **5**
4.1 The Counting: $4 + 6 = 10$ . . . 5
4.2 Gauge Groups from Internal Structure . . . 5
4.3 Why the Match Is Non-Trivial . . . 5

5 The D-Brane Correspondence **6**
5.1 Branes as Physical Objects . . . 6
5.2 The QSD Anchor as a Brane . . . 6
5.3 Brane Processes in QSD Language . . . 7

6 Moduli Stabilization and the Strain ς **8**
6.1 The Central Problem . . . 8
6.2 QSD Stabilization via Code Fidelity . . . 8
6.3 The Correspondence . . . 8
6.4 Vacuum Selection: From 10^{500} to 1 . . . 8

7 Renormalization: Topological Finiteness **9**
7.1 The Problem of Infinities . . . 9
7.2 String Theory: Modular Invariance . . . 9
7.3 QSD: Finite Homology . . . 9
7.4 The Correspondence . . . 10

8 The Hierarchy: Warp Factors and Information Counting **10**
8.1 The Problem in Both Frameworks . . . 10
8.2 String Theory: Warped Compactification . . . 11
8.3 QSD: Information Reduction . . . 11
8.4 The Correspondence . . . 11
8.5 T-Duality and UV/IR Mixing . . . 11

9 The 28 GeV Anomaly in String Language **12**
9.1 QSD Prediction . . . 12
9.2 Three String-Theoretic Interpretations . . . 12
9.3 Why It Couples Preferentially to b-Quarks . . . 12
9.4 Why the Signal Saturates . . . 12

10 The Complete Dictionary **13**

11 Strategic Implications **13**
11.1 What QSD Provides to String Theory . . . 13
11.2 What String Theory Provides to QSD . . . 14
11.3 The Duality Conjecture . . . 14

12 Open Questions and Research Program **14**
12.1 For String Theorists . . . 14
12.2 For Quantum Information Theorists . . . 15
12.3 For Phenomenologists . . . 15

13 Conclusion: An Invitation **15**

1 Preface: Scope and Framing

Before entering the technical details, a word about what this supplement is—and what it is not.

The author is not a string theorist. QSD was developed from the ground up, starting with quantum error correction and graph geometry, without reference to String Theory. Only after the framework was complete and its predictions tabulated did the structural parallels become apparent: the same counting (4 external, 6 internal), the same hierarchy mechanism, the same finiteness, and the same energy scale appear in both frameworks.

We therefore present this material as a **Rosetta Stone**—a side-by-side translation of QSD concepts into string-theoretic language. We believe the correspondence is too detailed to be coincidental, but we also recognize that confirming (or refuting) a true duality requires expertise we do not possess.

Throughout, we adopt the convention:

Observation = a structural parallel we can demonstrate.
Conjecture = a claim we believe is true but cannot prove.
Open Question = an invitation for the reader to investigate.

The goal is not to convince, but to *equip*: if even one of the conjectures below proves correct, the implications for both frameworks would be substantial.

2 Motivation: Two Paths to the Same Structure

2.1 The Landscape Problem

String Theory provides a mathematically rigorous framework for quantum gravity, unifying general relativity and quantum mechanics at the Planck scale. Yet it suffers from a critical limitation: it does not predict a unique vacuum. Instead, it admits an estimated $\sim 10^{500}$ possible vacuum configurations—the "Landscape"—each corresponding to a different compactification of the extra dimensions and therefore a different set of low-energy parameters (particle masses, coupling constants, cosmological constant).

Without a mechanism to select one vacuum from this vast set, String Theory cannot make predictions about the Standard Model parameters we observe. The anthropic principle has been invoked as a partial answer (we observe the vacuum compatible with our existence), but this explanation is widely regarded as unsatisfying: it replaces prediction with selection bias.

2.2 The QSD Approach

QSD approaches the same problem from the opposite direction. Rather than starting with a high-energy continuum theory and compactifying downward, QSD starts with the combinatorial structure of quantum error correction and builds upward.

The $[[5, 1, 3]]$ perfect quantum code defines a unique graph geometry: four bulk qubits forming K_4 (6 edges) plus one anchor qubit forming C_5 (5 edges in cyclic symmetry). From three derived constants—the holographic bit density $\text{Hb}_4 = 1/(4 \ln 2)$, the golden ratio $\varphi = (1+\sqrt{5})/2$, and the geometric strain $\varsigma = 3/(80\varphi^2)$—QSD derives Standard Model parameters with zero free parameters.

The key point: **the code structure is unique**. There is only one $[[5, 1, 3]]$ perfect code (up to equivalence), and its graph geometry is completely determined. There is no landscape of choices.

2.3 The Structural Coincidence

Despite their radically different starting points, QSD and String Theory arrive at strikingly similar structures:

Feature	QSD	String Theory
External degrees of freedom	4 (nodes)	4 (spacetime)
Internal degrees of freedom	6 (edges)	6 (Calabi-Yau)
Total	10	10
Gauge group origin	S_4 decomposition	Calabi-Yau topology
Finiteness mechanism	Topological (3 cycles)	Modular invariance
Hierarchy mechanism	Information factor 2^{-60}	Warp factor $e^{-k\pi r}$
Fundamental scale	$\mu_0 = 25.3$ GeV	Radion / KK mode

Table 1: Structural parallels between QSD and String Theory. Both frameworks produce the same counting and the same hierarchy.

This suggests that QSD may be the **non-perturbative definition** of the String Theory vacuum—analogous to how Matrix Theory provides a non-perturbative definition of M-Theory, or how lattice QCD defines the continuum theory.

3 Background: String Theory Essentials

For readers unfamiliar with String Theory, this section provides the minimum context needed to follow the correspondence. Experts may skip ahead.

3.1 From Points to Strings

In standard particle physics, fundamental objects are point particles. In String Theory, they are replaced by tiny one-dimensional strings whose vibrational modes produce the spectrum of observed particles. The string has a characteristic length scale $\ell_s = \sqrt{\alpha'}$, where α' is the "Regge slope" or string tension parameter.

Different vibrational modes of the string correspond to different particles: one mode gives the graviton, another gives gauge bosons, others give matter fermions. The crucial property is that all particles emerge from a *single* type of object, providing automatic unification.

3.2 Extra Dimensions and Compactification

Consistency of the quantum string requires the ambient spacetime to have 10 dimensions (for superstrings [1]) or 26 dimensions (for bosonic strings). Since we observe only 4 large dimensions, the remaining 6 must be "compactified"—curled up so tightly that they are invisible at currently accessible energies.

The geometry of the compactified dimensions [2] (typically a Calabi-Yau manifold) determines the low-energy physics. Different Calabi-Yau shapes give different gauge groups, particle spectra, and coupling constants. This is both the power and the curse of String Theory: it *can* produce the Standard Model, but it can also produce 10^{500} other theories [3, 4].

3.3 D-Branes

D-branes (short for Dirichlet membranes) are extended objects on which open strings can end [5]. A stack of N coincident D-branes carries $U(N)$ gauge symmetry. In modern string phenomenology, our 4-dimensional universe is often modeled as a stack of D3-branes embedded in the 10-dimensional "bulk."

3.4 Moduli and the Landscape

The shape and size of the Calabi-Yau manifold are controlled by scalar fields called **moduli**:

- **Kähler moduli**: control the volumes of internal cycles [1].
- **Complex structure moduli**: control the shape.
- **Dilaton**: controls the string coupling g_s (the probability of string splitting/joining).

In the absence of a stabilizing potential, these moduli are flat directions—they can take any value, and each value corresponds to a different universe. "Moduli stabilization" (finding a mechanism that fixes these fields at definite values) is one of the central unsolved problems of string phenomenology.

3.5 Modular Invariance and UV Finiteness

String Theory is UV-finite: it has no ultraviolet divergences. The mechanism is **modular invariance** [1] – the symmetry of one-loop string amplitudes (computed on a torus worldsheet) under modular transformations $\tau \to \tau + 1$ and $\tau \to -1/\tau$. This symmetry forces the high-energy and low-energy contributions to cancel, eliminating the infinities that plague point-particle quantum field theory.

4 The Dimensional Correspondence

4.1 The Counting: $4 + 6 = 10$

The most immediate correspondence is numerical. String Theory requires 10 dimensions: 4 external (spacetime) plus 6 internal (Calabi-Yau). The K_4 graph that forms the bulk of QSD has 4 nodes and $\binom{4}{2} = 6$ edges. The total is again 10.

Observation 4.1 (Dimensional Map)**.** The following identification maps QSD onto String Theory compactification:

$$4 \text{ nodes of } K_4 \longleftrightarrow 4 \text{ spacetime dimensions} \tag{1}$$

$$6 \text{ edges of } K_4 \longleftrightarrow 6 \text{ Calabi-Yau dimensions} \tag{2}$$

This counting is necessary but not sufficient for a duality. What makes it compelling is that the *physics derived from each side* also matches.

4.2 Gauge Groups from Internal Structure

In String Theory, gauge symmetries emerge from the topology of the Calabi-Yau manifold. Strings wrapping non-trivial cycles produce gauge bosons; the number of independent cycles determines the gauge group.

In QSD, the 6 edges of K_4 carry information channels that decompose under the S_4 permutation symmetry:

$$\mathbf{6}_{\text{edges}} = \mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3} \tag{3}$$

producing the Standard Model gauge group $U(1)_Y \times SU(2)_L \times SU(3)_c$.

4.3 Why the Match Is Non-Trivial

One might object that "$4 + 6 = 10$" is a simple coincidence. But the correspondence goes deeper:

1. The *decomposition* of the 6 internal degrees of freedom matches: both produce $1 + 2 + 3$ gauge channels.
2. The *role* of the 4 external degrees matches: both are the arena for observable physics.




Figure 1: Gauge group emergence: K_4 edge decomposition (left) and Calabi-Yau cycle wrapping (right) both produce $U(1) \times SU(2) \times SU(3)$. The color-coding indicates the correspondence between S_4 irreps and cycle types.

3. The *total count of 10* is required by internal consistency in both frameworks (quantum anomaly cancellation in strings, code perfection in QSD).

Open Question 4.2 (Anomaly Matching)**.** Does the condition for anomaly-free gauge groups in 10D string compactification reduce to the S_4 decomposition condition when restricted to the $[[5, 1, 3]]$ code structure?

5 The D-Brane Correspondence

5.1 Branes as Physical Objects

In String Theory, D-branes are not merely mathematical constructs—they are dynamical objects with mass, charge, and tension. A Dp-brane extends in p spatial dimensions plus time. Open strings have their endpoints attached to branes, and the low-energy dynamics of a stack of N Dp-branes is a $U(N)$ gauge theory in $(p+1)$ dimensions.

A popular string phenomenology scenario places the Standard Model on a stack of D3-branes (3 spatial dimensions = our universe) floating in the 10D bulk. Hidden sectors may live on other, separated branes.

5.2 The QSD Anchor as a Brane

In QSD, the Anchor qubit (the 5th qubit of the $[[5, 1, 3]]$ code) connects to all 4 bulk qubits and stores the classical record of quantum processes. During the "Snap," the anchor decouples from the bulk, severing 4 bonds and releasing binding energy.

Observation 5.1 (Brane Map)**.**

$$\text{Bulk } K_4 \text{ (4 qubits)} \longleftrightarrow \text{Stack of D3-branes (visible sector)} \tag{4}$$

$$\text{Anchor qubit } (W\text{-bit}) \longleftrightarrow \text{Probe D0-brane (hidden sector)} \tag{5}$$

The justification for this identification:

- The K_4 bulk carries gauge symmetry, just as a D3-brane stack carries $U(N)$ gauge theory.
- The Anchor connects to the bulk via 4 bonds, corresponding to open strings stretched between a probe brane and the stack.

- The Snap (anchor decoupling) corresponds to brane separation or tachyon condensation.
- The Ledger (classical record on the anchor) corresponds to the D-brane worldvolume theory, which stores information about strings that have been absorbed.




Figure 2: The Snap mechanism as brane separation. Left: pre-Snap configuration with 4 stretched strings connecting the D0-brane (Anchor) to the D3-stack (K_4). Right: post-Snap, the brane separates and the string energy is released as the syndrome signal.

5.3 Brane Processes in QSD Language

QSD Process	**String Theory Process**
Anchor-bulk coupling (K_5 phase)	Open strings stretched between D0 and D3-stack
Snap (anchor decoupling)	Brane annihilation or tachyon condensation
Syndrome energy E_{syn}	String tension $\times$ brane separation distance
Ledger (classical record on anchor)	D-brane worldvolume theory
Infinity bits $\{W, X, Y, Z\}$	Chan-Paton factors (string endpoint labels)
Error correction (logical qubit preserved)	Gauge invariance of brane system

Table 2: Process-level correspondence between QSD dynamics and brane physics.

Remark 5.2. The identification of infinity bits with Chan-Paton factors is particularly suggestive. In String Theory, Chan-Paton factors are discrete labels attached to string endpoints on branes, determining which gauge representation a string state belongs to. In QSD, infinity bits $\{W, X, Y, Z\}$ are discrete labels that route information to gauge sectors. Both are non-dynamical quantum numbers that organize gauge structure.

Open Question 5.3 (Chan-Paton Structure)**.** Do the 4 infinity bits of QSD correspond to specific Chan-Paton assignments in a D3/D0-brane system? If so, is the S_4 decomposition $\mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3}$ derivable from the open-string spectrum?

6 Moduli Stabilization and the Strain ς

6.1 The Central Problem

In String Theory, the "moduli problem" is one of the deepest obstacles to making contact with observation. The Calabi-Yau manifold has dozens to hundreds of moduli—continuous parameters describing its shape and size—and without a mechanism to fix them, the theory has no predictive power.

Several mechanisms have been proposed (KKLT [6], Large Volume Scenario, etc.), but all require careful tuning. No mechanism has been shown to uniquely select the Standard Model vacuum.

6.2 QSD Stabilization via Code Fidelity

In QSD, the analog of moduli stabilization is automatic. The geometric strain

$$\varsigma = \frac{3}{80\varphi^2} \approx 0.01432 \tag{6}$$

is not a free parameter to be fixed—it is *derived* from the requirement that the K_4 and C_5 graph structures coexist within the $[[5, 1, 3]]$ code. Any other value would break the code's error-correction capability.

This is analogous to how the geometry of a crystal is fixed by the requirement that its atoms fit together: there is only one way to tile them, and that determines all lattice parameters.

6.3 The Correspondence

Observation 6.1 (Moduli Map)**.**

$$\text{Geometric strain } \varsigma \longleftrightarrow \text{String coupling } g_s \text{ (dilaton VEV)} \tag{7}$$

$$\text{Radion mass } \mu_0 \longleftrightarrow \text{Kähler modulus (volume mode)} \tag{8}$$

$$\text{Golden ratio } \varphi \longleftrightarrow \text{Complex structure modulus} \tag{9}$$

The justification for the $\varsigma \leftrightarrow g_s$ identification:

- Both are small dimensionless parameters ($\varsigma \approx 0.014$; $g_s \sim 0.1$ in realistic string vacua).
- Both control the strength of quantum corrections: ς enters QSD loop corrections as ς^n at n-loop order, just as g_s enters string loop corrections as g_s^{2n}.
- Both must be stabilized for the theory to be predictive.
- In QSD, ς is uniquely fixed by code geometry. If the correspondence holds, this would fix g_s as well, resolving the dilaton stabilization problem.

6.4 Vacuum Selection: From 10^{500} to 1

Conjecture 6.2 (Vacuum Selection via Error Correction)**.** *The 10^{500} string vacua collapse to a unique vacuum under the requirement that the compactification geometry supports a functioning quantum error-correcting code. Specifically:*

$$V_{\text{eff}}(\text{moduli}) \propto 1 - \mathcal{F}_{\text{code}}(\text{moduli}) \tag{10}$$

where $\mathcal{F}_{\text{code}}$ is the fidelity of the $[[5, 1, 3]]$ code realized on the compactification geometry. The universe sits at the unique minimum where $\mathcal{F}_{\text{code}} = 1$.

The "Swampland" (the set of effective field theories that *cannot* arise from String Theory) would then have a clean characterization [4]: it consists of all geometries incompatible with perfect quantum error correction.




Figure 3: Vacuum selection. The string landscape contains $\sim 10^{500}$ local minima (gray curve). QSD's code fidelity condition selects a unique vacuum (red arrow): the one where quantum error correction functions perfectly.

Open Question 6.3 (Swampland Correspondence)**.** Do the known Swampland criteria (Weak Gravity Conjecture, Distance Conjecture, de Sitter Conjecture) follow from the requirement that the vacuum support a $[[5, 1, 3]]$ code?

7 Renormalization: Topological Finiteness

7.1 The Problem of Infinities

In standard quantum field theory, loop calculations produce ultraviolet divergences that must be removed by renormalization. The procedure works (spectacularly well in QED), but the loop series is infinite—in principle, one needs to compute infinitely many diagrams.

Both String Theory and QSD resolve this problem, but through different (yet structurally parallel) mechanisms.

7.2 String Theory: Modular Invariance

String amplitudes are computed as integrals over the moduli space of Riemann surfaces (the string worldsheet). At one loop, the worldsheet is a torus, parametrized by a complex modulus τ. The key constraint is **modular invariance**: the amplitude must be unchanged under

$$\tau \to \tau + 1 \qquad \text{and} \qquad \tau \to -\frac{1}{\tau} \tag{11}$$

These transformations map the high-energy region of the integration domain to the low-energy region, forcing the UV divergences to cancel exactly.

At higher loops, the worldsheet has higher genus (more "handles"), and the modular group generalizes. The result is the same: topology constrains the integrals and eliminates divergences order by order.

7.3 QSD: Finite Homology

QSD renormalization terminates at 3 loops because the graph $K_4 \cup C_5$ has exactly 3 independent topological cycles (first homology group H_1). Each cycle generates one loop correction; after all cycles are exhausted, no further independent loops can be constructed.

The three corrections and their physical origins (derived in detail in Supplement S4):

$$\text{1-Loop:} \quad \delta_1 = +0.239 \quad \text{(network friction: 5 anchor-bulk bonds)} \tag{12}$$

$$\text{2-Loop:} \quad \delta_2 = -0.022 \quad \text{(radion feedback via } S_4 \text{ channels)} \tag{13}$$

$$\text{3-Loop:} \quad \delta_3 = -0.007 \quad \text{(vacuum drag: zero-point energy)} \tag{14}$$

Together with the tree-level value ${\alpha_{\text{em}}}^{-1}_{\text{tree}} = 4\pi\varphi^4/\text{Hb}_4 = 136.826$, these yield:

$${\alpha_{\text{em}}}^{-1} = 136.826 + 0.239 - 0.022 - 0.007 = 137.036 \tag{15}$$

7.4 The Correspondence

Observation 7.1 (Finiteness Map)**.** Both frameworks achieve UV finiteness through topology:

$$\text{QSD: finite graph homology} \quad \longleftrightarrow \quad \text{String: worldsheet modular invariance} \tag{16}$$

Loop Order	String Theory	QSD
Tree level	Sphere (genus 0)	K_4 tree structure
1-loop	Torus (genus 1)	Network friction (+0.239)
2-loop	Double torus (genus 2)	Radion feedback (−0.022)
3-loop	Triple torus (genus 3)	Vacuum drag (−0.007)
≥4-loop	Higher genus (suppressed by g_s^{2n})	**Exactly zero**

Table 3: Loop structure correspondence. In String Theory, higher-loop contributions are suppressed but nonzero. In QSD, they are exactly zero because no further independent cycles exist.

Remark 7.2 (A Stronger Claim?)**.** QSD's termination is *exact*: there is no 4th loop, period. String Theory's higher loops are merely suppressed. If the duality holds, this suggests that the higher-genus string contributions, while formally present, must cancel exactly in the physical vacuum—a testable prediction within string perturbation theory.




Figure 4: Finiteness mechanisms compared. String Theory achieves UV finiteness through worldsheet modular invariance at each genus. QSD achieves exact termination because the graph $K_4 \cup C_5$ has only 3 independent homology cycles.

8 The Hierarchy: Warp Factors and Information Counting

8.1 The Problem in Both Frameworks

Both String Theory and QSD must explain why the electroweak scale (∼100 GeV) is 10^{17} times smaller than the Planck scale (10^{19} GeV). In the Standard Model, this hierarchy requires fine-tuning to 17 decimal places.

8.2 String Theory: Warped Compactification

The Randall-Sundrum (RS) model [7], originally motivated by string theory, resolves the hierarchy via a "warped" extra dimension. The metric takes the form:

$$ds^2 = e^{-2k|y|}\eta_{\mu\nu}dx^\mu dx^\nu + dy^2 \tag{17}$$

where k is the curvature scale and y parametrizes the extra dimension. The Planck scale at $y = 0$ (the "UV brane") is exponentially redshifted at $y = \pi r$ (the "IR brane"):

$$\Lambda_{\text{IR}} = e^{-k\pi r}\, M_{\text{Pl}} \tag{18}$$

For $k\pi r \approx 37$, this gives $\Lambda_{\text{IR}} \sim 1$ TeV.

8.3 QSD: Information Reduction

QSD derives the hierarchy from the state-space volume of the K_4 Hilbert space:

$$\mu_0 = \frac{M_{\text{Pl}}}{2^{60}} \times \varphi^2 \times (1 - 6\varsigma) = 25.34 \text{ GeV} \tag{19}$$

The factor $2^{60} = 16^{15}$ counts the number of distinguishable micro-configurations per macro-state of the code (see Supplement S2 for the full derivation).

8.4 The Correspondence

Observation 8.1 (Hierarchy Map)**.**

$$\text{QSD information factor } 2^{-60} \quad \longleftrightarrow \quad \text{Randall-Sundrum warp factor } e^{-k\pi r} \tag{20}$$

The numerical match: $2^{-60} \approx 8.7 \times 10^{-19}$, while $e^{-37} \approx 8.5 \times 10^{-17}$. The two are within two orders of magnitude, with the remaining difference absorbed by the geometric factors $\varphi^2(1-6\varsigma)$.

The deeper point is that both mechanisms achieve the hierarchy through *geometric* or *information-theoretic* counting rather than fine-tuning. In the RS model, the warp factor arises from the geometry of the extra dimension. In QSD, the factor 2^{-60} arises from the combinatorics of the code.

8.5 T-Duality and UV/IR Mixing

String Theory possesses a remarkable symmetry called **T-duality**: a compactified dimension of radius R is physically identical to one of radius α'/R [1]. This implies a minimum observable length $L_{\text{min}} \sim \sqrt{\alpha'} = \ell_s$ and connects ultraviolet (small distance) physics to infrared (large distance) physics.

QSD exhibits an analogous structure. The Ledger mechanism stores high-energy information (entering a horizon) as low-energy bits on the boundary. The hierarchy is the necessary consequence of this encoding—the conversion factor between "UV bits" (Planck-scale data) and "IR bits" (electroweak-scale observables) is precisely 2^{60}.

Open Question 8.2 (T-Duality Analog)**.** Is there a precise QSD analog of T-duality—a symmetry that exchanges the K_4 ("internal") and C_5 ("external") graph structures while preserving the physics?

9 The 28 GeV Anomaly in String Language

9.1 QSD Prediction

QSD predicts a syndrome signature at:

$$E_{\text{syn}} = \mu_0(1 + 8\varsigma) = 25.34 \times 1.1146 = 28.24 \text{ GeV} \tag{21}$$

matching the CMS dimuon excess at ~ 28.3 GeV to 0.2% (see Supplement S5 for the complete phenomenological analysis).

9.2 Three String-Theoretic Interpretations

The 28 GeV signal admits at least three natural interpretations in string language:

(a) The Breathing Mode / Radion. In Calabi-Yau compactification, the overall volume of the internal manifold can oscillate. The quantum of this oscillation—the "breathing mode" or radion—is a scalar field whose mass is set by the stabilization potential. In QSD, the radion mass is $\mu_0 = 25.34$ GeV; the syndrome energy $E_{\text{syn}} = 28.24$ GeV includes the binding energy of anchor-bulk bonds.

(b) The Lightest Kaluza-Klein Mode. In theories with extra dimensions, every 4D field has a tower of "Kaluza-Klein" (KK) excitations—higher harmonics on the compactified dimensions. In warped scenarios, the lightest KK mode is brought from the Planck scale to the electroweak scale by the warp factor. The QSD hierarchy formula naturally produces this mode at ~ 25 GeV.

(c) A Stretched String. In the brane picture, a string momentarily stretched between the visible sector (D3-stack) and the hidden sector (D0/Anchor) would have energy proportional to the brane separation times the string tension. The syndrome is the energy released when this string snaps.

9.3 Why It Couples Preferentially to b-Quarks

In warped geometries (Randall-Sundrum), the 5D wavefunctions of Standard Model fermions are localized at different positions along the extra dimension. Heavy fermions (top, bottom) are localized near the IR brane, while light fermions (electron, up-quark) are localized near the UV brane.

The radion field also lives on the IR brane. Therefore:

$$\text{Coupling}(\text{radion}, f) \propto |\Psi_f(y_{\text{IR}})|^2 \propto m_f \tag{22}$$

Heavy quarks have large wavefunction overlap with the radion; light leptons have small overlap. This is precisely the mass-proportional coupling derived in QSD from the radion-matter Lagrangian:

$$\mathcal{L}_{\text{int}} = -\frac{\phi_r}{v_r} \sum_f m_f \bar{\psi}_f \psi_f \tag{23}$$

9.4 Why the Signal Saturates

The QSD prediction that the syndrome significance *saturates* (stops growing with luminosity) has a natural holographic interpretation.

In AdS/CFT, a bulk particle corresponds to a boundary operator [8, 9]. The production rate of bulk excitations is limited by the **information-processing capacity** of the holographic

boundary—the Ledger. You cannot create radion excitations faster than the Ledger can record them.

This is analogous to the Bekenstein bound: there is a maximum amount of information that can be processed per unit area per unit time. The syndrome production rate is bounded by this information bandwidth.

String verdict: The saturation, if confirmed by LHC Run 3, would be evidence of the holographic principle at collider energies—a genuinely novel signature.

10 The Complete Dictionary

We collect all correspondences identified in the preceding sections into a single reference table.

QSD Concept	**String Theory Analog**
Structure	
K_4 nodes (4)	Spacetime dimensions (4)
K_4 edges (6)	Calabi-Yau dimensions (6)
S_4 edge decomposition $\mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3}$	Gauge groups from CY cycles
Anchor qubit (W-bit)	D0-brane / probe brane
Bulk K_4	D3-brane stack
Infinity bits $\{W, X, Y, Z\}$	Chan-Paton factors
C_5 cyclic symmetry ($\mathbb{Z}_5$)	Orbifold symmetry
Parameters	
Geometric strain ς	String coupling g_s / dilaton VEV
Radion mass μ_0	Kähler modulus (volume mode)
Golden ratio φ	Complex structure modulus
$\text{Hb}_4 = 1/(4 \ln 2)$	String tension α' (related)
Hb_4/ς (holographic bit density / strain)	Dilaton / Kähler ratio
Mechanisms	
Snap mechanism	Brane separation / tachyon condensation
Syndrome energy 28.24 GeV	Lightest KK mode / stabilized radion
3-loop termination	Modular invariance
Hierarchy factor 2^{-60}	Warp factor $e^{-k\pi r}$
Code stability condition	Swampland constraints
Signal saturation	Holographic bandwidth limit
$[[5, 1, 3]]$ error correction	Gauge invariance preservation
Logical qubit (Higgs)	Lightest string mode

Table 4: Complete QSD–String Theory dictionary. Structural elements, parameters, and dynamical mechanisms are matched across frameworks.

11 Strategic Implications

If the correspondence identified above proves to be a genuine duality, both frameworks gain significantly.

11.1 What QSD Provides to String Theory

1. **Vacuum Selection.** The code fidelity condition (Conjecture 6.2) picks one vacuum from 10^{500}. This would resolve the Landscape problem that has stymied string phenomenology for two decades.

2. **Predictive Power.** QSD derives ${\alpha_{\text{em}}}^{-1} = 137.036$, $\sin^2\theta_W = 0.2312$, $m_H = 126.7$ GeV—values that String Theory currently takes as input. A duality would mean String Theory *predicts* these values via QSD.

3. **A Testable Low-Energy Prediction.** The 28 GeV syndrome gives String Theory its first prediction below the Planck scale that is accessible to current experiments.

4. **The Target Manifold.** If QSD is correct, string theorists should look for a Calabi-Yau with Euler characteristic $|\chi| = 6$ (since $|\chi|/2 = 3$ generations). This dramatically narrows the search within the Kreuzer-Skarke database [10].

11.2 What String Theory Provides to QSD

1. **Mathematical Rigor.** String Field Theory provides formal tools for the Snap mechanism (brane dynamics, tachyon condensation, open-string field theory).

2. **Conceptual Validation.** T-duality justifies the existence of a minimum length scale; modular invariance justifies finite loop corrections. These are established results in String Theory that would bolster QSD's claims.

3. **Community and Credibility.** Framing QSD as "the non-perturbative vacuum of String Theory" connects it to an established research program with thousands of practitioners, removing the "radical outsider" label.

4. **Gravitational Extension.** String Theory naturally includes gravity; QSD currently addresses only particle physics. The duality could provide QSD's gravitational sector "for free."

11.3 The Duality Conjecture

Conjecture 11.1 (QSD–String Duality)**.** *Quantum Spectrodynamics is to String Theory as Matrix Theory [11] is to M-Theory: a discrete, non-perturbative definition of the vacuum state that the continuum theory describes perturbatively.*

Specifically: there exists a Calabi-Yau threefold X with $\chi(X) = \pm 6$ such that the Type IIB string compactification on X with appropriate flux configuration reproduces all QSD predictions in the low-energy limit.

Remark 11.2. The Matrix Theory analogy is deliberate. Matrix Theory provides a non-perturbative definition of M-Theory using a discrete quantum-mechanical system (a matrix model). It was discovered by computing quantities in both frameworks and finding agreement. We propose the same investigative strategy for QSD and String Theory.

12 Open Questions and Research Program

We close with a structured list of open questions, organized by the expertise required to address them.

12.1 For String Theorists

Open Question 12.1 (Calabi-Yau Search)**.** Does the Kreuzer-Skarke database [10] contain a Calabi-Yau threefold with $|\chi| = 6$ whose intersection numbers reproduce the S_4 decomposition $\mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3}$?

Open Question 12.2 (Modular Invariance Implies Termination?)**.** In the physical vacuum (if it corresponds to QSD), do the genus ≥ 4 string amplitudes cancel exactly, or are they merely suppressed?

Open Question 12.3 (KKLT and Code Fidelity)**.** Can the KKLT moduli stabilization mechanism [6] be reformulated as a code-fidelity maximization problem? Does the resulting potential have a unique minimum?

12.2 For Quantum Information Theorists

Open Question 12.4 (Code from Geometry)**.** Can the $[[5, 1, 3]]$ code be derived from a Calabi-Yau compactification, or more generally, can quantum error-correcting codes be classified by the topology of their "parent" manifolds?

Open Question 12.5 (Holographic Codes)**.** Is the $[[5, 1, 3]]$ code a special case of holographic tensor network codes (e.g., HaPPY codes)? If so, what is the corresponding bulk geometry?

12.3 For Phenomenologists

Open Question 12.6 (KK Tower)**.** If the 28 GeV signal is the lightest KK mode, where are the heavier modes? QSD predicts the next syndrome at $\sim \mu_0(1 + 16\varsigma^2) \approx 25.4$ GeV (two-qubit error), but this may not be observable. What are the signatures of the KK tower in the QSD/string picture?

Open Question 12.7 (Proton Decay Channel)**.** QSD predicts $\tau_p \sim 10^{38-40}$ years from code distance. String Theory predicts proton decay via dimension-6 operators with model-dependent rates. Do the two predictions agree for the candidate Calabi-Yau?

13 Conclusion: An Invitation

This supplement does not claim to derive String Theory results. The author is an information theorist, not a string theorist.

What we have shown is a **structural isomorphism**: the same numbers (4 external, 6 internal), the same finiteness mechanism (topology), the same hierarchy (10^{17}), and the same energy scale (~ 25 GeV) appear in both frameworks. The correspondence extends to dynamics (Snap $\leftrightarrow$ brane separation), parameters ($\varsigma \leftrightarrow g_s$), and even phenomenology (mass-proportional coupling from IR brane localization).

We invite the string theory community to investigate:

1. Can the $[[5, 1, 3]]$ code be derived from a specific Calabi-Yau compactification?
2. Does the code stability condition correspond to known Swampland criteria?
3. Can String Field Theory calculate the syndrome production cross-section?
4. Is there a Calabi-Yau with $|\chi| = 6$ that reproduces QSD's predictions?

If the correspondence holds, QSD provides what String Theory has sought for 40 years: **the principle that selects the Standard Model from the Landscape.**

And if it does not hold, the specific points of failure will sharpen both frameworks.

I am not a String Theorist. I am an Information Theorist. I looked at the data (LHC), I applied the logic of Error Correction, and I found the source code. It turns out the source code looks a lot like what String Theorists have been looking for—but I arrived there from the bottom up, not the top down.

References

[1] Joseph Polchinski. *String Theory, Volume 1: An Introduction to the Bosonic String.* Cambridge University Press, 1998.

[2] Philip Candelas, Gary T. Horowitz, Andrew Strominger, and Edward Witten. Vacuum configurations for superstrings. *Nucl. Phys. B*, 258:46–74, 1985.

[3] Leonard Susskind. The anthropic landscape of string theory. *arXiv preprint*, 2003.

[4] Cumrun Vafa. The string landscape and the swampland. *arXiv preprint*, 2005.

[5] Joseph Polchinski. *String Theory, Volume 2: Superstring Theory and Beyond.* Cambridge University Press, 1998.

[6] Shamit Kachru, Renata Kallosh, Andrei Linde, and Sandip P. Trivedi. De Sitter vacua in string theory. *Phys. Rev. D*, 68:046005, 2003.

[7] Lisa Randall and Raman Sundrum. Large mass hierarchy from a small extra dimension. *Phys. Rev. Lett.*, 83:3370–3373, 1999.

[8] Juan Maldacena. The large-N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998.

[9] Ahmed Almheiri, Xi Dong, and Daniel Harlow. Bulk locality and quantum error correction in AdS/CFT. *JHEP*, 04:163, 2015.

[10] Maximilian Kreuzer and Harald Skarke. Complete classification of reflexive polyhedra in four dimensions. *Adv. Theor. Math. Phys.*, 4:1209–1230, 2000.

[11] Tom Banks, Willy Fischler, Stephen H. Shenker, and Leonard Susskind. M theory as a matrix model: a conjecture. *Phys. Rev. D*, 55:5112–5128, 1997.