Undergraduate Complexity Theory
Lecture 8: NP

Marcythm

July 12, 2022

1 Lecture Notes

Conjecture 1.1. $\mathsf{P} \neq \mathsf{NP}(\iff 3\text{-SAT} \notin \mathsf{P})$

Conjecture 1.2 (Exponential Time Hypothesis). $\exists \delta > 0 : 3\text{-SAT} \notin \mathsf{TIME}((1+\delta)^n)$

Conjecture 1.3 (Strong ETH). $\forall \delta > 0 : \exists k : k\text{-SAT} \notin \mathsf{TIME}((2-\delta)^n)$

$\text{SETH} \implies \text{ETH} \implies \mathsf{P} \neq \mathsf{NP}$.

Theorem 1.4 (ABV '15). *SETH implies $\forall \epsilon > 0$, cannot solve* LCS *in time* $O(n^{2-\epsilon})$.

idea of NP: for many problems $\exists$ brute force algo, enumerate (exp many) "candidate sol"s and check in poly-time if each is a genuine sol. e.g. ST-PATH, HAMILTONIAN-PATH, 3-COL, CIRCUIT-SAT, COMPOSITE.

two features:

0. "candidate sol"s are encodable by strings with polynomial length.

1. $\exists$ poly-time algo to <u>check</u> if a candidate sol is a genuine sol.

Informally, a problem is in NP if a checking algo exists. "candidate sol" are also called "potential sol", "witness", "certificate", etc. A problem highly believed not in NP: UN-3COL.

Definition 1.5. An algorithm(TM) V is a *verifier for language L* if

1. V takes as input a pair $\langle x, y \rangle$

2. $\forall x : x \in L \iff \exists y : V(\langle x, y \rangle)$ accepts.

Definition 1.6. Verifier V is said to be *"polynomial time"* if $V(\langle x, y \rangle)$ runs in time $O(|x|^k)$ for some $k \in \mathbb{N}$.

Remark 1.7. Subtlety: V's runtime is measured in terms of $|x|$.

Definition 1.8. $\mathsf{NP} = \{L : L \text{ has a poly-time verifier}\}$

e.g. $\mathsf{SQUARES} = \{\langle B \rangle : B \in \mathbb{N}, \exists x \in \mathbb{N} : x^2 = B\} \in \mathsf{NP}$. $3\mathsf{COL} \in \mathsf{NP}$.

Subtlety in verifier of SQUARE: mark the input as $\langle x, y \rangle$, after interpreting $x = \langle B \rangle$, only read first $|x|$ symbols of y. If $|y| > |x|$, then reject.

Theorem 1.9. $\mathsf{P} \in \mathsf{NP}$.

Proof. Let $L \in \mathsf{P}$, thus $\exists$ a poly-time TM M s.t. $x \in L \iff M(x)$ accepts. Consider the TM V that runs on input $\langle x, y \rangle$: Do $M(x)$, then …

1. Claim V is poly-time verifier. …

2. Claim V verifies L. …

$\square$

About $\mathsf{P} = \mathsf{NP}$, upcoming: Cook-Levin Theorem: $\mathsf{P} = \mathsf{NP} \iff 3\mathsf{SAT} \in \mathsf{P}$.

2 Reading

2.1 Sipser 7.3 (The Class NP)

definition of verifier, NP (by verifiers), and NTIME.