

1 probability review

Exercise 1.1

Let X be a random variable taking values between 0 and π with pdf given by $f(x) = c \sin(x)$, $x \in [0, \pi]$. What is the value of c ?

Solution: Since the integral of the pdf is always 1 (by definition),

$$1 = \int_0^\pi f(x) dx = \int_0^\pi c \sin x dx = -c \Big|_0^\pi \cos x = -c(\cos \pi - \cos 0) = -c(-2) = 2c.$$

And so, $2c = 1$ and $c = \frac{1}{2}$. B. ■

Exercise 1.2

What is $\mathbb{E}[X]$?

Solution: By the definition of expectation,

$$\mathbb{E}[X] = \int_0^\pi x f(x) dx = \int_0^\pi c x \sin x dx = c \Big|_0^\pi (\sin x - x \cos x) = c(-\pi \cos \pi + \sin \pi - 0 \cos 0 + \sin 0) = \pi c.$$

We know from the previous problem that $c = \frac{1}{2}$, so $\mathbb{E}[X] = \frac{\pi}{2}$. A. ■

Exercise 1.3

Let X be a Gaussian random variable with mean $\mu > 0$ and variance μ^2 . What is $\mathbb{E}[X]$?

Solution: The mean, that is $\mathbb{E}[X]$, is μ by definition. B. ■

Exercise 1.4

What is $\mathbb{E}[X^2]$?

Solution: By definition of variance,

$$\mu^2 = \text{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \mathbb{E}[X^2] - \mu^2,$$

so $\mathbb{E}[X^2] = 2\mu^2$. C. ■

Exercise 1.5

What is $\mathbb{E}[X^3]$?

Solution: Using the binomial theorem,

$$\begin{aligned} \mathbb{E}[X^3] &= \mathbb{E}[(X - \mu) + \mu]^3 \\ &= \mathbb{E}[(X - \mu)^3] + 3\mathbb{E}[(X - \mu)^2\mu] + 3\mathbb{E}[(X - \mu)\mu^2] + \mathbb{E}[\mu^3] \\ &= 3\mu\mathbb{E}[(X - \mu)^2] + \mu^3. \end{aligned}$$

Since $X - \mu$ is a Gaussian random variable with mean 0 and variance μ^2 , $\mathbb{E}[(X - \mu)^2] = \mu^2$,

$$\mathbb{E}[X^3] = 3\mu^3 + \mu^3 = 4\mu^3.$$

C. ■

Exercise 1.6

What is $\text{Var}[X^2]$?

Solution: For a normal random variable with mean μ and standard deviation σ , $\mathbb{E}[X^4] = 3\sigma^4 + 6\mu^2\sigma^2 + \mu^4$. Since our mean and standard deviation are both μ in this case, $\mathbb{E}[X^4] = 10\mu^4$. Using this,

$$\begin{aligned}\text{Var}[X^2] &= \mathbb{E}[X^4] - \mathbb{E}[X^2]^2 \\ &= 10\mu^4 - (2\mu^2)^2 \\ &= 6\mu^4.\end{aligned}$$

\boxed{B} . ■

Exercise 1.7

What is $\mathbb{P}[X > 0]$ in terms of the CDF Φ of the standard Gaussian distribution?

Solution: By definition of the cdf,

$$\mathbb{P}[X > 0] = \mathbb{P}\left[\frac{X - \mu}{\sigma} > -\frac{\mu}{\sigma}\right] = \mathbb{P}\left[\frac{X - \mu}{\sigma} > -1\right] = \mathbb{P}\left[\frac{X - \mu}{\sigma} < 1\right] = \Phi(1).$$

\boxed{B} . ■

Exercise 1.8

Let X be a random variable such that

$$X = \begin{cases} 1 & \text{with probability } p \\ -1 & \text{with probability } 1 - p \end{cases}$$

for some $p \in [0, 1]$. What is $\mathbb{E}[X]$?

Solution: Routine algebra shows that $\mathbb{E}[X] = 1 \cdot p + (-1) \cdot (1 - p) = -1 + 2p$. \boxed{D} . ■

Exercise 1.9

What is $\text{Var}[X]$?

Solution: The variance of X is $\mathbb{E}[X^2] - \mathbb{E}[X]^2 = 1 - (1 - 2p)^2 = 4p - 4p^2 = 4p(1 - p)$. \boxed{C} . ■

Exercise 1.10

For what p is $\text{Var}[X]$ maximized?

Solution: We know from the previous problem that $\text{Var}[X] = 4p - 4p^2$, which has derivative $4 - 8p$. The variance is maximized when that derivative is 0, so when $4 - 8p = 0 \implies p = \frac{1}{2}$. \boxed{C} . ■

Exercise 1.11

What is $\mathbb{E}[X^k]$?

Solution: The expected value is $\mathbb{E}[X^k] = 1^k \cdot p + (-1)^k \cdot (1 - p) = p + (-1)^k \cdot (1 - p)$. \boxed{D} . ■

Exercise 1.12

Let X and Y be two independent standard Gaussian random variables. What is $\mathbb{E}[X^2Y]$?

Solution: Since X and Y are independent, $\mathbb{E}[X^2Y] = \mathbb{E}[X^2] \cdot \mathbb{E}[Y] = 0$. \boxed{A} . ■

Exercise 1.13

What is $\text{Var}(X + Y)$?

Solution: Since X and Y are independent, $\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y] = 1 + 1 = 2$. \boxed{C} . ■

Exercise 1.14

What is $\text{Var}[XY]$?

Solution: The variance is $\text{Var}[XY] = \mathbb{E}[(XY)^2] - \mathbb{E}[XY]^2 = \mathbb{E}[X^2]\mathbb{E}[Y^2] = 1$. \boxed{B} . ■

Exercise 1.15

What is $\text{Cov}[X, X + Y]$?

Solution: The covariance is $\text{Cov}[X, X + Y] = \text{Cov}[X, X] + \text{Cov}[X, Y] = 1 + 0 = 1$. \boxed{B} . ■

Exercise 1.16

What is $\text{Cov}[X, XY]$?

Solution: The covariance is $\text{Cov}[X, XY] = \mathbb{E}[X^2Y] - \mathbb{E}[X]\mathbb{E}[XY] = 0$. \boxed{A} . ■

Exercise 1.17**Exercise 1.18****Exercise 1.19**

Let X_1, \dots, X_n be i.i.d. with mean μ and variance σ^2 . What is $\mathbb{E}[\sum_{i=1}^n X_i]$?

Solution: By linearity of expectation,

$$\mathbb{E}\left[\sum_{i=1}^n X_i\right] = \sum_{i=1}^n \mathbb{E}[X_i] = n\mu.$$

\boxed{C} . ■

Exercise 1.20

What is $\text{Var}[\sum_{i=1}^n X_i]$?

Solution: Since each of the X_i s are independent,

$$\text{Var} \left[\sum_{i=1}^n X_i \right] = \sum_{i=1}^n \text{Var}[X_i] = n\sigma^2.$$

\square ■

Exercise 1.21

What is $\mathbb{E}[(\sum_{i=1}^n X_i)^2]$?

Solution: Let $Y = \sum_{i=1}^n X_i$. Then,

$$n\sigma^2 = \text{Var}[Y] = \mathbb{E}[Y^2] - \mathbb{E}[Y]^2 = \mathbb{E}[Y^2] - (n\mu)^2,$$

so $\mathbb{E}[Y^2] = n^2\mu^2 + n\sigma^2$. \square ■

Exercise 1.22

What is $\text{Var} \left[\frac{1}{n} \sum_{i=1}^n X_i \right]$?

Solution: Since $\text{Var}[aX] = a^2\text{Var}[X]$,

$$\text{Var} \left[\frac{1}{n} \sum_{i=1}^n X_i \right] = \frac{1}{n^2} \text{Var} \left[\sum_{i=1}^n X_i \right] = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}.$$

\square ■

2 new concepts

Exercise 2.1

Let $X_n \sim \text{Unif}(-1/n, 1/n)$ and let X be a random variable such that $\mathbb{P}[X = 0] = 1$.

1. Compute and draw the CDF $F_n(x)$ and $F(x)$ of X_n and X respectively.
2. Does $X_n \xrightarrow{P} X$? (prove or disprove)
3. Does $X_n \rightsquigarrow X$ (prove or disprove)

Solution:

1. The CDF of X is

$$F(x) = \begin{cases} 0 & x < 0 \\ 1 & x \geq 0 \end{cases}$$

and the CDF of X_n is

$$F_n(x) = \begin{cases} 0 & x < -\frac{1}{n} \\ \frac{nt+1}{2} & -\frac{1}{n} \leq \frac{1}{n} \\ 1 & x > \frac{1}{n} \end{cases}.$$




Figure 1: This is the cdf for X_n , when $n = 4$.




Figure 2: This is the cdf for X .

2. We know that for all ϵ ,

$$\mathbb{P}[|X_n - X| > \epsilon] = \mathbb{P}[|X_n| > \epsilon] = 1 - \mathbb{P}[|X_n| \leq \epsilon] = 1 - \min\left(\frac{2\epsilon}{\frac{2}{n}}, 1\right) = 1 - \min(n\epsilon, 1).$$

Therefore,

$$\lim_{n \rightarrow \infty} \mathbb{P}[|X_n - X| > \epsilon] = \lim_{n \rightarrow \infty} 1 - \min(n\epsilon, 1) = 0,$$

so X_n does converge to X probabilistically.

3. The CDF of X is

$$F(t) = \begin{cases} 0 & x < 0 \\ 1 & x \geq 0 \end{cases}$$

and the CDF of X_n is

$$F_n(t) = \begin{cases} 0 & x < -\frac{1}{n} \\ \frac{nt+1}{2} & -\frac{1}{n} \leq x \leq \frac{1}{n} \\ 1 & x > \frac{1}{n} \end{cases}.$$

As n approaches ∞ , $F_n(t)$ approaches

$$F_n(t) = \begin{cases} 0 & x < 0 \\ \frac{1}{2} & x = \frac{1}{2} \\ 1 & x = 1 \end{cases}.$$

Although $F_n(t)$ and $F(t)$ are not the same functions, they have equal values at all points for which F is continuous. Thus, X_n converges to X in distribution. ■

Exercise 2.2

Let $X \sim \mathcal{N}(1, 2.25)$. Compute the following probabilities (show your work):

1. $\mathbb{P}[X > 1]$
2. $\mathbb{P}[|X - 2| \leq 1]$
3. $\mathbb{P}[|X| < 1]$
4. $\mathbb{P}[X^2 - 2X - 1 > 0]$.

Solution:

1. Since X is normally distributed around 1, by symmetry, $\mathbb{P}[X > 1] = 0.5$.
2. We know that $\mathbb{P}[|X - 2| \leq 1] = \mathbb{P}[1 \leq X \leq 3]$. If we let Y be $\frac{X - \mu}{\sigma} = \frac{X - 1}{1.5}$, then we can rewrite this as $\mathbb{P}[0 \leq Y \leq \frac{4}{3}]$. Since Y is standard normal, we can now reference the standard normal table, which tells us that $\mathbb{P}[Y \leq \frac{4}{3}] \approx 0.9082$ and $\mathbb{P}[Y \leq 0] = 0.5$, so $\mathbb{P}[0 \leq Y \leq \frac{4}{3}] \approx 0.4082$.
3. Again, we can use a similar technique as the previous part and let $Y = \frac{X - \mu}{\sigma} = \frac{X - 1}{1.5}$. Then, it becomes clear that $\mathbb{P}[|X| < 1] = \mathbb{P}[-1 \leq X \leq 1] = \mathbb{P}[-\frac{4}{3} \leq Y \leq 0]$, which by symmetry means that $\mathbb{P}[|X| \leq 1] = \mathbb{P}[0 \leq Y \leq \frac{4}{3}] \approx 0.4082$.
4. Since $(X - 1)^2 - 2 = X^2 - 2X - 1$, $\mathbb{P}[X^2 - 2X - 1 > 0] = \mathbb{P}[(X - 1)^2 - 2 > 0] = \mathbb{P}[(X - 1)^2 > 2] = \mathbb{P}[1 - \sqrt{2} \leq X \leq 1 + \sqrt{2}]$. Letting $Y = \frac{X - \mu}{\sigma}$, we reduce this to $\mathbb{P}[-\frac{\sqrt{2}}{1.5} < Y < \frac{\sqrt{2}}{1.5}]$. Using a standard normal table, we can compute this to be around 0.3472. ■