


Gaussian processes introduction

Victor Kochegarov

February 19, 2016

Problem statement



Training data: $\{(x_i, y_i) : i = \overline{1, N}\}, x_i, y_i \in R$

Task: given new x^* predict y^*

Three ways to go

Linear regression

Bayesian regression

Gaussian Process

Linear regression


Utilize model

$$y = mx,$$

One should solve

$$m^* = \arg \min_{m \in \mathbb{R}} \sum_{i=1}^N (y_i - mx_i)^2$$


and get $m^* = \frac{\sum_i x_i y_i}{\sum_i x_i^2}$



Bayesian regression

Utilize **model**

$$y_i = mx_i + \varepsilon_i, \quad \varepsilon_i \sim_{\text{iid}} N(0, \sigma_\varepsilon^2), \quad m \sim N(0, 1).$$



The goal is to find $p(m|y, x)$

Bayesian formula

$$p(m|y, x) = \frac{p(y|m, x)p(m)}{\int p(y|m, x)p(m)dm}$$
$$\propto p(y|m, x)p(m)$$

Bayesian formula

$$\begin{aligned} p(m|y, x) &= \frac{p(y|m, x)p(m)}{\int p(y|m, x)p(m)dm} \\ &\propto p(y|m, x)p(m) \\ &\propto \left(\prod_i \frac{1}{\sqrt{2\pi}\sigma_\epsilon} e^{-(y_i - mx_i)^2 / (2\sigma_\epsilon^2)} \right) \frac{1}{\sqrt{2\pi}e^{-m^2/2}} \end{aligned}$$

Bayesian formula

$$\begin{aligned}
 p(m|y, x) &= \frac{p(y|m, x)p(m)}{\int p(y|m, x)p(m)dm} \\
 &\propto p(y|m, x)p(m) \\
 &\propto \left(\prod_i \frac{1}{\sqrt{2\pi}\sigma_\epsilon} e^{-(y_i - mx_i)^2 / (2\sigma_\epsilon^2)} \right) \frac{1}{\sqrt{2\pi}e^{-m^2/2}} \\
 &\propto e^{-\sum_i (y_i - mx_i)^2 / (2\sigma_\epsilon^2) - m^2/2}
 \end{aligned}$$

Bayesian formula

$$\begin{aligned}
 p(m|y, x) &= \frac{p(y|m, x)p(m)}{\int p(y|m, x)p(m)dm} \\
 &\propto p(y|m, x)p(m) \\
 &\propto \left(\prod_i \frac{1}{\sqrt{2\pi}\sigma_\epsilon} e^{-(y_i - mx_i)^2 / (2\sigma_\epsilon^2)} \right) \frac{1}{\sqrt{2\pi}e^{-m^2/2}} \\
 &\propto e^{-\sum_i (y_i - mx_i)^2 / (2\sigma_\epsilon^2) - m^2/2} \\
 &\propto e^{-\frac{\sum_i x_i^2 + \sigma_\epsilon^2}{2\sigma_\epsilon^2} \left(m - \frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)^2}
 \end{aligned}$$

Bayesian formula

$$\begin{aligned}
 p(m|y, x) &= \frac{p(y|m, x)p(m)}{\int p(y|m, x)p(m)dm} \\
 &\propto p(y|m, x)p(m) \\
 &\propto \left(\prod_i \frac{1}{\sqrt{2\pi}\sigma_\epsilon} e^{-(y_i - mx_i)^2 / (2\sigma_\epsilon^2)} \right) \frac{1}{\sqrt{2\pi}e^{-m^2/2}} \\
 &\propto e^{-\sum_i (y_i - mx_i)^2 / (2\sigma_\epsilon^2) - m^2/2} \\
 &\propto e^{-\frac{\sum_i x_i^2 + \sigma_\epsilon^2}{2\sigma_\epsilon^2} \left(m - \frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)^2}
 \end{aligned}$$


That is: $m|y, x \sim \mathcal{N}\left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2}\right)$

Bayesian formula

$$\begin{aligned}
 p(m|y, x) &= \frac{p(y|m, x)p(m)}{\int p(y|m, x)p(m)dm} \\
 &\propto p(y|m, x)p(m) \\
 &\propto \left(\prod_i \frac{1}{\sqrt{2\pi}\sigma_\epsilon} e^{-(y_i - mx_i)^2 / (2\sigma_\epsilon^2)} \right) \frac{1}{\sqrt{2\pi}e^{-m^2/2}} \\
 &\propto e^{-\sum_i (y_i - mx_i)^2 / (2\sigma_\epsilon^2) - m^2/2} \\
 &\propto e^{-\frac{\sum_i x_i^2 + \sigma_\epsilon^2}{2\sigma_\epsilon^2} \left(m - \frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)^2}
 \end{aligned}$$


That is: $m|y, x \sim \mathcal{N}\left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2}\right)$

Parameters posterior




$$m|y, x \sim \mathcal{N} \left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)$$

Parameters posterior




$$m|y, x \sim \mathcal{N} \left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)$$

Parameters posterior




$$m|y, x \sim \mathcal{N} \left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)$$

Parameters posterior




$$m|y, x \sim \mathcal{N} \left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)$$

Parameters posterior



$$m|y, x \sim \mathcal{N} \left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)$$

Parameters posterior



$$m|y, x \sim \mathcal{N} \left(\frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2} \right)$$

Output posterior

$$\begin{aligned}
 p(y^*|x^*, x, y) &= \int p(y^*|x^*, m)p(m|x, y)dm \\
 &= \mathcal{N}\left(x^* \frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, (x^*)^2 \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2}\right)
 \end{aligned}$$

In more general case $m \sim \mathcal{N}(\mathbf{0}, \Sigma_p)$ and input space is multidimensional:

$$y^*|x^*, x, y \sim \mathcal{N}\left(\frac{1}{\sigma_\epsilon^2} \mathbf{x}^{*\top} A^{-1} X \mathbf{y}, \mathbf{x}^{*\top} A^{-1} \mathbf{x}^*\right),$$

where $A = \sigma_\epsilon^{-2} X X^\top + \Sigma_p^{-1}$

Output posterior

$$\begin{aligned}
 p(y^*|x^*, x, y) &= \int p(y^*|x^*, m)p(m|x, y)dm \\
 &= \mathcal{N}\left(x^* \frac{\sum_i x_i y_i}{\sum_i x_i^2 + \sigma_\epsilon^2}, (x^*)^2 \frac{\sigma_\epsilon^2}{\sum_i x_i^2 + \sigma_\epsilon^2}\right)
 \end{aligned}$$

In more general case $m \sim \mathcal{N}(\mathbf{0}, \Sigma_p)$ and input space is multidimensional:

$$y^*|x^*, x, y \sim \mathcal{N}\left(\frac{1}{\sigma_\epsilon^2} \mathbf{x}^{*\top} A^{-1} X \mathbf{y}, \mathbf{x}^{*\top} A^{-1} \mathbf{x}^*\right),$$

where $A = \sigma_\epsilon^{-2} X X^\top + \Sigma_p^{-1}$

Input space to feature space

Project input space with transformation $\phi(\mathbf{x}): \mathbb{R}^n \rightarrow \mathbb{R}^s$.

$$\Phi = \Phi(X) = (\phi(\mathbf{x}_1), \phi(\mathbf{x}_2), \dots, \phi(\mathbf{x}_N))$$

Model takes the form:

$$\mathbf{y} = \phi(\mathbf{x})^\top \mathbf{m}$$

Input space to feature space

Project input space with transformation $\phi(\mathbf{x}): \mathbb{R}^n \rightarrow \mathbb{R}^s$.

$$\Phi = \Phi(X) = (\phi(\mathbf{x}_1), \phi(\mathbf{x}_2), \dots, \phi(\mathbf{x}_N))$$

Model takes the form:

$$\mathbf{y} = \phi(\mathbf{x})^\top \mathbf{m}$$

$$y^* | x^*, x, y \sim \mathcal{N} \left(\phi^{*\top} \Sigma_p \Phi^{-1} (K + \sigma_\epsilon^2 I)^{-1} \mathbf{y}, \right. \\ \left. \phi^{*\top} \Sigma_p \phi^* - \phi^{*\top} \Sigma_p \Phi (K + \sigma_\epsilon^2 I)^{-1} \Phi^\top \Sigma_p \phi^* \right),$$

Input space to feature space

Project input space with transformation $\phi(\mathbf{x}): \mathbb{R}^n \rightarrow \mathbb{R}^s$.

$$\Phi = \Phi(X) = (\phi(\mathbf{x}_1), \phi(\mathbf{x}_2), \dots, \phi(\mathbf{x}_N))$$

Model takes the form:

$$\mathbf{y} = \phi(\mathbf{x})^\top \mathbf{m}$$

$$y^* | x^*, \mathbf{x}, \mathbf{y} \sim \mathcal{N} \left(\phi^{*\top} \Sigma_p \Phi^{-1} (K + \sigma_\epsilon^2 I)^{-1} \mathbf{y}, \right. \\ \left. \phi^{*\top} \Sigma_p \phi^* - \phi^{*\top} \Sigma_p \Phi (K + \sigma_\epsilon^2 I)^{-1} \Phi^\top \Sigma_p \phi^* \right),$$

where $\phi^* = \phi(\mathbf{x}^*)$ and $K = \Phi^\top \Sigma_p \Phi$

Input space to feature space

Project input space with transformation $\phi(\mathbf{x}): \mathbb{R}^n \rightarrow \mathbb{R}^s$.

$$\Phi = \Phi(X) = (\phi(\mathbf{x}_1), \phi(\mathbf{x}_2), \dots, \phi(\mathbf{x}_N))$$

Model takes the form:

$$\mathbf{y} = \phi(\mathbf{x})^\top \mathbf{m}$$

$$y^* | x^*, x, y \sim \mathcal{N} \left(\phi^{*\top} \Sigma_p \Phi^{-1} (K + \sigma_\epsilon^2 I)^{-1} \mathbf{y}, \right. \\ \left. \phi^{*\top} \Sigma_p \phi^* - \phi^{*\top} \Sigma_p \Phi (K + \sigma_\epsilon^2 I)^{-1} \Phi^\top \Sigma_p \phi^* \right),$$

where $\phi^* = \phi(\mathbf{x}^*)$ and $K = \Phi^\top \Sigma_p \Phi$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$
 - $y_i | m, \varepsilon_i = mx_i + \varepsilon_i$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$
 - $y_i | m, \varepsilon_i = mx_i + \varepsilon_i$
- This implicitly defined a joint prior on $\{y_i : i = \overline{1, n}\}$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$
 - $y_i | m, \varepsilon_i = mx_i + \varepsilon_i$
- This implicitly defined a joint prior on $\{y_i : i = \overline{1, n}\}$
 - $y_i \sim \mathcal{N}(0, x_i^2 + \sigma_\varepsilon^2)$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$
 - $y_i | m, \varepsilon_i = mx_i + \varepsilon_i$
- This implicitly defined a joint prior on $\{y_i : i = \overline{1, n}\}$
 - $y_i \sim \mathcal{N}(0, x_i^2 + \sigma_\varepsilon^2)$
 - $\text{cov}(y_i, y_j) = x_i x_j, \forall i \neq j$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$
 - $y_i | m, \varepsilon_i = mx_i + \varepsilon_i$
- This implicitly defined a joint prior on $\{y_i : i = \overline{1, n}\}$
 - $y_i \sim \mathcal{N}(0, x_i^2 + \sigma_\varepsilon^2)$
 - $\text{cov}(y_i, y_j) = x_i x_j, \forall i \neq j$

That is (y_1, y_2, \dots, y_n) has multivariate normal distribution:

$$y \sim \mathcal{N}(\mathbf{0}, \mathbf{K}), \quad k_{ij} = x_i x_j + \delta_{ij} \sigma_\varepsilon^2$$

From Bayesian regression to GP

- We defined a Bayesian linear regression model by specifying priors on m and ε_i
 - $m \sim \mathcal{N}$, $\varepsilon_i \sim_{\text{iid}} \mathcal{N}(0, \sigma_\varepsilon^2)$
 - $y_i | m, \varepsilon_i = mx_i + \varepsilon_i$
- This implicitly defined a joint prior on $\{y_i : i = \overline{1, n}\}$
 - $y_i \sim \mathcal{N}(0, x_i^2 + \sigma_\varepsilon^2)$
 - $\text{cov}(y_i, y_j) = x_i x_j, \forall i \neq j$

That is (y_1, y_2, \dots, y_n) has multivariate normal distribution:

$$y \sim \mathcal{N}(\mathbf{0}, \mathbf{K}), \quad k_{ij} = x_i x_j + \delta_{ij} \sigma_\varepsilon^2$$

Gaussian processes

Definition

Gaussian process is a collection of random variables, any finite number of which have a joint Gaussian distribution

We can write this collection of random variables as

$$\{f(x) : x \in \mathcal{X}\}$$

i.e. a function f evaluated at inputs x .

Gaussian processes

Definition

Gaussian process is a collection of random variables, any finite number of which have a joint Gaussian distribution

We can write this collection of random variables as

$$\{f(x): x \in \mathcal{X}\}$$

i.e. a function f evaluated at inputs x .

GP(μ, k) is completely specified by

Gaussian processes

Definition

Gaussian process is a collection of random variables, any finite number of which have a joint Gaussian distribution

We can write this collection of random variables as

$$\{f(x): x \in \mathcal{X}\}$$

i.e. a function f evaluated at inputs x .

GP(μ, k) is completely specified by

- Mean function $\mu(x) = E(f(x))$

Gaussian processes

Definition

Gaussian process is a collection of random variables, any finite number of which have a joint Gaussian distribution

We can write this collection of random variables as

$$\{f(x) : x \in \mathcal{X}\}$$

i.e. a function f evaluated at inputs x .

GP(μ, k) is completely specified by

- Mean function $\mu(x) = E(f(x))$
- Covariance/kernel function $k(x, x') = Cov(f(x), f(x'))$

Gaussian processes

Definition

Gaussian process is a collection of random variables, any finite number of which have a joint Gaussian distribution

We can write this collection of random variables as


$$\{f(x) : x \in \mathcal{X}\}$$

i.e. a function f evaluated at inputs x .

GP(μ, k) is completely specified by


- Mean function $\mu(x) = E(f(x))$
- Covariance/kernel function $k(x, x') = Cov(f(x), f(x'))$

Linear kernel




$$k(x, x') = xx'$$

Exponential kernel




$$k(x, x') = e^{-(x-x')^2}$$


Non-linear kernel




Non-linear kernel




Non-linear kernel




Non-linear kernel




Non-linear kernel




Non-linear kernel




Linear model gone wrong




Linear model gone wrong




Linear model gone wrong




Linear model gone wrong




Linear model gone wrong




Linear model gone wrong




Non-linearity to the rescue




Non-linearity to the rescue




Non-linearity to the rescue




Non-linearity to the rescue



Non-linearity to the rescue



Non-linearity to the rescue



Kernel types

Five base kernels



Squared
exp. (SE)



Periodic
(PER)



Linear
(LIN)



Constant
(C)



White
noise (WN)

Encoding for the following types of functions



Smooth
functions



Periodic
functions



Linear
functions




Constant
functions




Gaussian
noise


Kernels compositions




$\text{LIN} \times \text{LIN}$




quadratic
functions




$\text{SE} \times \text{PER}$




locally
periodic




$\text{LIN} + \text{PER}$



periodic plus
linear trend



$\text{SE} + \text{PER}$



periodic plus
smooth trend

Classification problem

Training data: $\{(\mathbf{x}_i, y_i) : i = \overline{1, N}\}$, $\mathbf{x}_i \in R^s$, $y_i \in C = \{-1, +1\}$.

Task: given new \mathbf{x}^* predict distribution y^* on C .

That is we want to know

$$\pi(\mathbf{x}) = p(y = +1|\mathbf{x}).$$

Classification problem solving

Main idea:

$$\pi(x) = \sigma(f(\mathbf{x})),$$

$f(\cdot)$ — latent variable.

Step 1

$$p(f^*|X, y, \mathbf{x}^*) = \int p(f^*|X, \mathbf{x}^*, \mathbf{f})p(\mathbf{f}|X, y)d\mathbf{f}$$

Step 2

$$p(y^* = +1|X, y, \mathbf{x}^*) = \int \sigma(f^*)p(f^*|X, y, \mathbf{x}^*)df^*$$