

# Infering species distribution from aggregated spatial data

Going from coarse landings data to fine scale species distribution

Baptiste Alglave, Marie-Pierre Etienne, Youen Vermard, Mathieu Woillez, Etienne Rivot

04/2022 - GdR Ecostats

# Spatial data in ecology



## Survey data

Standardized sampling plan  
High quality data



Low amount of samples



## Examples

EVHOE data, Bay of Biscay (marine ecology)

## Citizen science data

Access to more data  
Exact locations usually available



Ebird application (ornithology)

## Declaration data

Mandatory declaration  
Massive data

Aggregated at the scale of administrative units



Harvest data, Wisconsin (hunting)

# How to integrate all these datasources? (especially when they do not have the same spatial resolution)

- ➡ Some specific proposition in the context of fishery science:

Alglaive Baptiste, Rivot Etienne, Etienne Marie-Pierre, Woillez Mathieu, Thorson James T, Vermand Youen (2022). Combining scientific survey and commercial catch data to map fish distribution. ICES Journal of Marine Science IN PRESS.  
<https://doi.org/10.1093/icesjms/fsac032>

# How to integrate all these datasources? (especially when they do not have the same spatial resolution)


➡ Some specific proposition in the context of fishery science:

Alglave Baptiste, Rivot Etienne, Etienne Marie-Pierre, Woillez Mathieu, Thorson James T, Vermaud Youen (2022). **Combining scientific survey and commercial catch data to map fish distribution.** ICES Journal of Marine Science IN PRESS.  
<https://doi.org/10.1093/icesjms/fsac032>

# Commercial catch declarations data in fishery science

## Logbook landings

(Sole, trawlers OTB-DEF, 2018)




Resolution

Catch are daily declared at the resolution of ICES rectangles

## Fishing locations (VMS)

(Trawlers OTB-DEF, 2018)




VMS pings are vessels GPS locations emitted each hour

► Refine landings spatial resolution

# Commercial catch declarations data in fishery science


## Logbook landings (Sole, trawlers OTB-DEF, 2018)



Resolution

Catch are daily declared at the resolution of ICES rectangles


## Fishing locations (VMS) (Trawlers OTB-DEF, 2018)



VMS pings are vessels GPS locations emitted each hour

➡ Refine landings spatial resolution

# Modelling change of support



Actual situation

$$Y_i | S(x_i), x_i \sim \mathcal{L}_Y(S(x_i), \xi, \sigma)$$

$$Y_i = Y_i^*$$


Alternative solution

$$D_j = \sum_{i \in P_j} Y_i$$

$$D_j | S_{P_j}, P_j \sim \mathcal{L}_D(S_{P_j}, \xi, \sigma)$$

Match  $\mathcal{L}_D$  and  $\mathcal{L}_Y$  moments

# Modelling change of support



Actual situation

$$Y_i | S(x_i), x_i \sim \mathcal{L}_Y(S(x_i), \xi, \sigma)$$

$$Y_i = Y_i^*$$


Alternative solution

$$D_j = \sum_{i \in P_j} Y_i$$

$$D_j | S_{P_j}, P_j \sim \mathcal{L}_D(S_{P_j}, \xi, \sigma)$$

Match  $\mathcal{L}_D$  and  $\mathcal{L}_Y$  moments

# Modelling change of support



Actual situation

$$Y_i | S(x_i), x_i \sim \mathcal{L}_Y(S(x_i), \xi, \sigma)$$

$$Y_i = Y_i^*$$

Alternative solution


$$D_j = \sum_{i \in P_j} Y_i$$

$$D_j | S_{P_j}, P_j \sim \mathcal{L}_D(S_{P_j}, \xi, \sigma)$$

Match  $\mathcal{L}_D$  and  $\mathcal{L}_Y$  moments

# Simulation-estimation and case study

## Simulation/estimation



### Simulation

- Latent field (covariate + spatial random effect)
- Commercial data (3000 samples over 2/3 of the area)
- Reallocation process (10 locations per declaration)
- Scientific data (100 samples over the whole the area)

### Estimation

Comparison of 3 model configurations:

1/ Model fitted to scientific data (reference data)

2/ Integrated model with commercial likelihood built on  $Y_i^*$

3/ Integrated model with commercial likelihood built on  $D_j$

Estimation realized through TMB (Template Model Builder)  
100 runs of simulation-estimation

### Model evaluation

1/ Mean square prediction error

$$MSPE = \frac{\sum_x (S(x) - \hat{S}(x))^2}{n}$$

2/ Covariate effect (or species-habitat relationship):

$$\beta_S = 2 \text{ versus } \hat{\beta}_S$$

### Case study: Sole in the Bay of Biscay



Survey data: Orhago


Commercial data: OTB-DEF trawlers (to ease convergence onboard observer data were integrated in fitting)

Fitted models: same as simulation

Covariate: substrate

# Simulation-estimation and case study

## Simulation/estimation



### Simulation

- Latent field (covariate + spatial random effect)
- Commercial data (3000 samples over 2/3 of the area)
- Reallocation process (10 locations per declaration)
- Scientific data (100 samples over the whole the area)

### Estimation

Comparison of 3 model configurations:

1/ Model fitted to scientific data (reference data)

2/ Integrated model with commercial likelihood built on  $Y_i^*$

3/ Integrated model with commercial likelihood built on  $D_j$

Estimation realized through TMB (Template Model Builder)  
100 runs of simulation-estimation

### Model evaluation

1/ Mean square prediction error

$$MSPE = \frac{\sum_x (S(x) - \hat{S}(x))^2}{n}$$

2/ Covariate effect (or species-habitat relationship):

$$\beta_S = 2 \text{ versus } \hat{\beta}_S$$

**Case study:** Sole in the Bay of Biscay



Survey data: Orhago


Commercial data: OTB-DEF trawlers (to ease convergence onboard observer data were integrated in fitting)

Fitted models: same as simulation

Covariate: substrate

# Simulation-estimation and case study

## Simulation/estimation



### Simulation

- Latent field (covariate + spatial random effect)
- Commercial data (3000 samples over 2/3 of the area)
- Reallocation process (10 locations per declaration)
- Scientific data (100 samples over the whole the area)

### Model evaluation

1/ Mean square prediction error

$$MSPE = \frac{\sum_x (S(x) - \hat{S}(x))^2}{n}$$

2/ Covariate effect (or species-habitat relationship):

$$\beta_S = 2 \text{ versus } \hat{\beta}_S$$

### Estimation

Comparison of 3 model configurations:

1/ Model fitted to scientific data (reference data)

2/ Integrated model with commercial likelihood built on  $Y_i^*$

3/ Integrated model with commercial likelihood built on  $D_j$

Estimation realized through TMB (Template Model Builder)  
100 runs of simulation-estimation


**Case study:** Sole in the Bay of Biscay



Survey data: Orhago  
Commercial data: OTB-DEF trawlers (to ease convergence onboard observer data were integrated in fitting)  
Fitted models: same as simulation  
Covariate: substrate

# Simulation-estimation and case study

## Simulation/estimation



### Simulation

- Latent field (covariate + spatial random effect)
- Commercial data (3000 samples over 2/3 of the area)
- Reallocation process (10 locations per declaration)
- Scientific data (100 samples over the whole the area)

### Model evaluation

1/ Mean square prediction error

$$MSPE = \frac{\sum_x (S(x) - \hat{S}(x))^2}{n}$$

2/ Covariate effect (or species-habitat relationship):

$$\beta_S = 2 \text{ versus } \hat{\beta}_S$$

### Estimation

Comparison of 3 model configurations:

1/ Model fitted to scientific data (reference data)

2/ Integrated model with commercial likelihood built on  $Y_i^*$

3/ Integrated model with commercial likelihood built on  $D_j$

Estimation realized through TMB (Template Model Builder)  
100 runs of simulation-estimation

## Case study: Sole in the Bay of Biscay









**Survey data:** Orhago

**Commercial data:** OTB-DEF  
trawlers (to ease convergence  
onboard observer data were  
integrated in fitting)



**Fitted models:** same as simulation

**Covariate:** substrate

# Simulation-estimation



# Case study: sole in the Bay of Biscay



\*n.b.: punctual commercial data (i.e. observer data) needed to be integrated in inference in order to make the model converge and some parameters needed to be step estimated (e.g. range)

# Discussion

- Framework that integrates **catch declarations data** (rough resolution of catch) and **scientific/onboard observer data** (exact locations of catch)
  - ➡ Still question: how to ease convergence ?
- Change of support issues
  - ➡ not so frequent in spatial ecology,
  - ➡ but increasing amount of data available that may face such issues
- Is it a generic framework ?
  - ➡ the overall approach is,  
(i.e. modelling observed aggregated catches as a convolution of latent exact catches)
  - ➡ but need to adapt the observation model to the data  
(here zero-inflated continuous data)
  - ➡ and make the hypothesis that fishing locations ( $\mathcal{P}_j$ ) are known  
(in reality  $\mathcal{P}_j$  is a trajectory and the set of fishing points are a realization of this trajectory)

# Discussion

- Framework that integrates **catch declarations data** (rough resolution of catch) and **scientific/onboard observer data** (exact locations of catch)
  - ➡ **Still question:** how to ease convergence ?
- **Change of support issues**
  - ➡ not so frequent in spatial ecology,
  - ➡ but increasing amount of data available that may face such issues
- **Is it a generic framework ?**
  - ➡ the overall approach is,  
(i.e. modelling observed aggregated catches as a convolution of latent exact catches)
  - ➡ but need to adapt the observation model to the data  
(here zero-inflated continuous data)
  - ➡ and make the hypothesis that fishing locations ( $\mathcal{P}_j$ ) are known  
(in reality  $\mathcal{P}_j$  is a trajectory and the set of fishing points are a realization of this trajectory)

# Discussion

- Framework that integrates **catch declarations data** (rough resolution of catch) and **scientific/onboard observer data** (exact locations of catch)
  - ➡ **Still question:** how to ease convergence ?
- **Change of support issues**
  - ➡ not so frequent in spatial ecology,
  - ➡ but increasing amount of data available that may face such issues
- **Is it a generic framework ?**
  - ➡ the overall approach is,  
(i.e. modelling observed aggregated catches as a convolution of latent exact catches)
  - ➡ but need to adapt the observation model to the data  
(here zeroinflated continuous data)
  - ➡ and make the hypothesis that fishing locations ( $\mathcal{P}_j$ ) are known  
(in reality  $\mathcal{P}_j$  is a trajectory and the set of fishing points are a realization of this trajectory)

# Discussion

- Framework that integrates **catch declarations data** (rough resolution of catch) and **scientific/onboard observer data** (exact locations of catch)
  - ➡ **Still question:** how to ease convergence ?
- **Change of support issues**
  - ➡ not so frequent in spatial ecology,
  - ➡ but increasing amount of data available that may face such issues
- **Is it a generic framework ?**
  - ➡ the overall approach is,  
(i.e. modelling observed aggregated catches as a convolution of latent exact catches)
  - ➡ but need to adapt the observation model to the data  
(here zeroinflated continuous data)
  - ➡ and make the hypothesis that fishing locations ( $\mathcal{P}_j$ ) are known  
(in reality  $\mathcal{P}_j$  is a trajectory and the set of fishing points are a realization of this trajectory)

# Thank you for your attention!

