How the Brain Rewires Itself When Ears Stop Working

Marie Setaire

Research group of the methods of science education and popularization

Summary

This article doesn’t require any knowledge about the brain and hearing. I want this knowledge to be accessible to anyone! Imagine your brain is like a big house with lots of rooms. Each room has a job. One special room is in charge of hearing — it’s called the auditory cortex. But here’s something amazing: if that “hearing room” stops getting sounds, the brain doesn’t just leave it empty. It gives the room a new job, usually helping with seeing instead (Lee et al., 2001; Sadato et al., 2004). Scientists call this “cross-modal reorganization,” which is just a fancy way of saying “the brain swapping one job for another.”

For a long time, scientists thought this swap was a totally good thing — the brain being clever and not wasting space. But it turns out the story is more complicated, and that’s what this article is about: Is the swap helpful, or does it cause problems?

The brain swapping the “hearing room” into a “seeing room” isn’t simply good or bad — it depends on when it starts, how long it lasts, and who it happens to. Catching it early, while the brain is still bendy and willing to change back, gives each person the best chance to hear the world clearly again.

June 25, 2026

This cover page was added for repository indexing.

How the Brain Rewires Itself When Ears Stop Working

Marie Setaire

Research group of the methods of science education and popularization

This article doesn’t require any knowledge about the brain and hearing. I want this knowledge to be accessible to anyone! Imagine your brain is like a big house with lots of rooms. Each room has a job. One special room is in charge of **hearing** — it's called the auditory cortex. But here's something amazing: if that "hearing room" stops getting sounds, the brain doesn't just leave it empty. It gives the room a *new* job, usually helping with **seeing** instead (Lee et al., 2001; Sadato et al., 2004). Scientists call this "cross-modal reorganization," which is just a fancy way of saying "the brain swapping one job for another."

For a long time, scientists thought this swap was a totally *good* thing — the brain being clever and not wasting space. But it turns out the story is more complicated, and that's what this article is about: **Is the swap helpful, or does it cause problems?**

Your eyes already help your ears

Even in people who hear perfectly, the eyes and ears are teammates. When someone talks to you, their lips actually start moving a tiny bit *before* the sound comes out — about 100 to 200 milliseconds earlier (that's faster than a blink!). Your brain uses those lip movements to guess what sound is coming next, kind of like a weather forecast for talking (Arnal et al., 2009; Arnal et al., 2011; van Wassenhove et al., 2005).

This guessing is so strong that just watching someone's lips can switch on the hearing room of your brain, even when there's no sound at all (Calvert et al., 1997; Besle et al., 2004). Scientists even did a clever experiment where a face raised its eyebrows, and people's brains "heard" a stressed, emphasized word that was never actually said out loud — the eyebrows tricked the ears! (Strelnikov et al., 2015). The brain did this using a loop that travels from the seeing part to the hearing part and back again (Kilian-Hütten et al., 2011). So your eyes helping your ears is normal and useful.

What happens when sound gets fuzzy

To understand deafness, scientists use a trick called **vocoding** — they make sound all crackly and fuzzy, like a broken radio, to copy what a hearing device sounds like (Shannon et al., 1995; Lorenzi et al., 2006).

Here's the cool part: even when sound is super fuzzy, there's one special brain area — the **temporal voice areas**, or **TVA** — that can still tell "that's a human voice!" versus "that's just a noise" (Belin et al., 2000; Pernet et al., 2015). It does this by listening to the rhythm and beat of speech, which stays the same even when the sound is fuzzy (Massida et al., 2011; Strelnikov et al., 2011; Chandrasekaran et al., 2009).

And importantly: when sound is just temporarily fuzzy, people's eyes don't take over their ears. If you show them a face that doesn't match the voice, they don't get confused (Barone et al., 2016). So short-term fuzzy sound is no big deal — the hearing brain stays in charge.

But long-term deafness is different

Now here's the twist. When someone can't hear for a very long time, the TVA "voice room" gets bored with no sound to listen to. So it slowly starts doing visual jobs instead — like reading sign language, lip-reading, and even remembering faces (Sadato et al., 2004; Rouger et al., 2012; Stropahl et al., 2015).

That sounds helpful, right? And it is — for a while. But there's a catch. Doctors can give deaf people a special device called a **cochlear implant (CI)** that lets them hear again. But scientists found something surprising: the more the hearing room had switched over to vision, the harder it was for that person to hear well with their new implant (Lee et al., 2007; Doucet et al., 2006; Lazard et al., 2014). The room got so good at its new "seeing" job that it forgot how to do its old "hearing" job!

Even people whose hearing comes back really well with the implant keep a strange habit: when figuring out things like whether a voice sounds like a man or a woman, they trust their **eyes** more than their **ears** — something people with normal hearing never do (Barone et al., 2016). Scientists think this happens because the "face" part and "voice" part of the brain become extra-wired-together during deafness, and that wiring stays even after the implant (Blank et al., 2011; von Kriegstein et al., 2005).

There's a "window of time" that really matters

Here's something hopeful. Scientists studying cats (and then children) discovered that the brain has a special window of time when it's young and bendy. If a child gets help with hearing during this window, the brain can fix itself much better (Kral et al., 2013; Kral & O'Donoghue, 2010).

For example, in kids who got their two implants more than about 1.5 years apart, one ear became the "favorite" and stayed unbalanced forever. But kids who got both implants closer together stayed nicely balanced (Gordon et al., 2012; Lee et al., 2020). So getting help early makes a huge difference.

Interestingly, the eye-takeover mostly affects things like figuring out who is talking or how they feel, not understanding the actual words (Barone et al., 2016). That's because words have rhythm the implant can copy, but voice details (like the exact pitch) are too delicate for the implant to send clearly (Marx et al., 2015).

How scientists keep an eye on the brain

To watch all this happening, scientists use special brain-scanning machines like fMRI and PET (Lee et al., 2007; Lazard et al., 2014). But those big machines are hard to use with little kids because you have to hold very still and they take a long time.

So scientists made a kid-friendly tool called **fNIRS** — it's gentle, safe, and works even on babies as young as four months old (Wiggins et al., 2016; Minagawa-Kawai et al., 2011). There's even a special test designed just for children aged 5 to 16 who are deaf in one ear, to measure how balanced their brain is and catch problems early (Calmels et al., 2022; Chen et al., 2016).

The big idea: three stages

Stage 1 — Everything normal. The eyes and ears are good teammates, and the voice room does its hearing job just fine (Strelnikov et al., 2015; van Wassenhove et al., 2005). Even fuzzy sound doesn't break this (Strelnikov et al., 2011; Barone et al., 2016).

Stage 2 — The helpful swap. With deafness, the hearing room starts helping with vision, which is useful for lip-reading and communicating. And if hearing is restored soon enough, the brain can mostly switch back.

Stage 3 — The stuck swap. After a very long time, the swap becomes permanent. The eyes-over-ears habit gets locked in and won't go away, even with a great implant (Barone et al., 2016). There's a window between Stage 2 and Stage 3 where doctors can still help — which is why acting early matters so much.

The takeaway

The brain swapping the "hearing room" into a "seeing room" isn't simply good or bad — it depends on **when** it starts, **how long** it lasts, and **who** it happens to. Catching it early, while the brain is still bendy and willing to change back, gives each person the best chance to hear the world clearly again.

References

1. Arnal, L. H., Morillon, B., Kell, C. A., & Giraud, A. L. (2009). Dual neural routing of visual facilitation in speech processing. Journal of Neuroscience, 29(43), 13445–13453. https://doi.org/10.1523/JNEUROSCI.3194-09.2009
2. Arnal, L. H., Wyart, V., & Giraud, A. L. (2011). Transitions in neural oscillations reflect prediction errors generated in audiovisual speech. Nature Neuroscience, 14(6), 797–801. https://doi.org/10.1038/nn.2810
3. Barone, P., Chambaudie, L., Strelnikov, K., Fraysse, B., Marx, M., Belin, P., & Deguine, O. (2016). Crossmodal interactions during non-linguistic auditory processing in cochlear-implanted deaf patients. Cortex, 83, 259–270. https://doi.org/10.1016/j.cortex.2016.08.005
4. Belin, P., Zatorre, R. J., Lafaille, P., Ahad, P., & Pike, B. (2000). Voice-selective areas in human auditory cortex. Nature, 403(6767), 309–312. https://doi.org/10.1038/35002078
5. Besle, J., Fort, A., Delpuech, C., & Giard, M. H. (2004). Bimodal speech: Early suppressive visual effects in human auditory cortex. European Journal of Neuroscience, 20(8), 2225–2234. https://doi.org/10.1111/j.1460-9568.2004.03670.x
6. Blank, H., Anwander, A., & von Kriegstein, K. (2011). Direct structural connections between voice- and face-recognition areas. Journal of Neuroscience, 31(36), 12906–12915. https://doi.org/10.1523/JNEUROSCI.2091-11.2011
7. Calvert, G. A., Bullmore, E. T., Brammer, M. J., Campbell, R., Williams, S. C., McGuire, P. K., Woodruff, P. W., Iversen, S. D., & David, A. S. (1997). Activation of auditory cortex during silent lipreading. Science, 276(5312), 593–596. https://doi.org/10.1126/science.276.5312.593
8. Calmels, M.-N., Gallois, Y., Marx, M., Deguine, O., Taoui, S., Arnaud, E., Strelnikov, K., & Barone, P. (2022). Functional reorganization of the central auditory system in children with single-sided deafness: A protocol using fNIRS. Brain Sciences, 12(4), 423. https://doi.org/10.3390/brainsci12040423
9. Chandrasekaran, C., Trubanova, A., Stillittano, S., Caplier, A., & Ghazanfar, A. A. (2009). The natural statistics of audiovisual speech. PLoS Computational Biology, 5(7), e1000436. https://doi.org/10.1371/journal.pcbi.1000436
10. Chen, L. C., Sandmann, P., Thorne, J. D., Bleichner, M. G., & Debener, S. (2016). Cross-modal functional reorganization of visual and auditory cortex in adult cochlear implant users identified with fNIRS. Neural Plasticity, 2016, 4382656. https://doi.org/10.1155/2016/4382656

11. Doucet, M. E., Bergeron, F., Lassonde, M., Ferron, P., & Lepore, F. (2006). Cross-modal reorganization and speech perception in cochlear implant users. Brain, 129(12), 3376–3383. https://doi.org/10.1093/brain/awl264

12. Gordon, K. A., Salloum, C., Toor, G. S., van Hoesel, R., & Papsin, B. C. (2012). Binaural interactions develop in the auditory brainstem of children who are deaf: Effects of place and level of bilateral electrical stimulation. Journal of Neuroscience, 32(12), 4212–4223. https://doi.org/10.1523/JNEUROSCI.5741-11.2012

13. Kilian-Hütten, N., Vroomen, J., & Formisano, E. (2011). Brain activation during audiovisual exposure anticipates future perception of ambiguous speech. NeuroImage, 57(4), 1601–1607. https://doi.org/10.1016/j.neuroimage.2011.05.043

14. Kral, A., Hubka, P., Heid, S., & Tillein, J. (2013). Single-sided deafness leads to unilateral aural preference within an early sensitive period. Brain, 136(1), 180–193. https://doi.org/10.1093/brain/aws305

15. Kral, A., & O'Donoghue, G. M. (2010). Profound deafness in childhood. New England Journal of Medicine, 363(15), 1438–1450. https://doi.org/10.1056/NEJMra0911225

16. Lazard, D. S., Innes-Brown, H., & Barone, P. (2014). Adaptation of the communicative brain to post-lingual deafness: Evidence from functional imaging. Hearing Research, 307, 136–143. https://doi.org/10.1016/j.heares.2013.07.011

17. Lee, D. S., Lee, J. S., Oh, S. H., Kim, S.-K., Kim, J.-W., Chung, J.-K., Lee, M. C., & Kim, C. S. (2001). Cross-modal plasticity and cochlear implants. Nature, 409(6817), 149–150. https://doi.org/10.1038/35051653

18. Lee, H. J., Giraud, A. L., Kang, E., Oh, S. H., Kang, H., Kim, C. S., & Lee, D. S. (2007). Cortical activity at rest predicts cochlear implantation outcome. Cerebral Cortex, 17(4), 909–917. https://doi.org/10.1093/cercor/bhl001

19. Lee, H.-J., Smieja, D., Polonenko, M. J., Cushing, S. L., Papsin, B. C., & Gordon, K. A. (2020). Consistent and chronic cochlear implant use partially reverses cortical effects of single sided deafness in children. Scientific Reports, 10(1), 21526. https://doi.org/10.1038/s41598-020-78371-6

20. Lorenzi, C., Gilbert, G., Carn, H., Garnier, S., & Moore, B. C. (2006). Speech perception problems of the hearing impaired reflect inability to use temporal fine structure. Proceedings of the National Academy of Sciences, 103(49), 18866–18869. https://doi.org/10.1073/pnas.0607364103

21. Marx, M., James, C., Foxton, J., Capber, A., Fraysse, B., Barone, P., & Deguine, O. (2015). Speech prosody perception in cochlear implant users with and without residual hearing. Ear and Hearing, 36(2), 239–248. https://doi.org/10.1097/AUD.0000000000000105

22. Massida, Z., Belin, P., James, C., Rouger, J., Fraysse, B., Barone, P., & Deguine, O. (2011). Voice discrimination in cochlear-implanted deaf subjects. Hearing Research, 275(1–2), 120–129. https://doi.org/10.1016/j.heares.2010.12.013

23. Minagawa-Kawai, Y., van der Lely, H., Ramus, F., Sato, Y., Mazuka, R., & Dupoux, E. (2011). Optical brain imaging reveals general auditory and language-specific processing in early infant development. Cerebral Cortex, 21(2), 254–261. https://doi.org/10.1093/cercor/bhq082

24. Pernet, C. R., McAleer, P., Latinus, M., Gorgolewski, K. J., Charest, I., Bestelmeyer, P. E., & Belin, P. (2015). The human voice areas: Spatial organization and inter-individual variability in temporal and extra-temporal cortices. NeuroImage, 119, 164–174. https://doi.org/10.1016/j.neuroimage.2015.06.050

25. Rouger, J., Lagleyre, S., Démonet, J.-F., Fraysse, B., Deguine, O., & Barone, P. (2012). Evolution of crossmodal reorganization of the voice area in cochlear-implanted deaf patients. Human Brain Mapping, 33(8), 1929–1940. https://doi.org/10.1002/hbm.21331

26. Sadato, N., Yamada, H., Okada, T., Yoshida, M., Hasegawa, T., Matsuki, K., Yonekura, Y., & Itoh, H. (2004). Age-dependent plasticity in the superior temporal sulcus in deaf humans: A functional MRI study. BMC Neuroscience, 5, 56. https://doi.org/10.1186/1471-2202-5-56

27. Shannon, R. V., Zeng, F.-G., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech recognition with primarily temporal cues. Science, 270(5234), 303–304. https://doi.org/10.1126/science.270.5234.303

28. Strelnikov, K., Massida, Z., Rouger, J., Belin, P., & Barone, P. (2011). Effects of vocoding and intelligibility on the cerebral response to speech. BMC Neuroscience, 12, 122. https://doi.org/10.1186/1471-2202-12-122

29. Strelnikov, K., Foxton, J., Marx, M., & Barone, P. (2015). Brain prediction of auditory emphasis by facial expressions during audiovisual continuous speech. Brain Topography, 28(3–4), 494–505. https://doi.org/10.1007/s10548-013-0338-2

30. Stropahl, M., Plotz, K., Schönfeld, R., Lenarz, T., Sandmann, P., & Yovel, G. (2015). Cross-modal reorganization in cochlear implant users: Auditory cortex contributes to visual face processing. NeuroImage, 121, 159–170. https://doi.org/10.1016/j.neuroimage.2015.07.062

31. van Wassenhove, V., Grant, K. W., & Poeppel, D. (2005). Visual speech speeds up the neural processing of auditory speech. Proceedings of the National Academy of Sciences, 102(4), 1181–1186. https://doi.org/10.1073/pnas.0408949102

32. von Kriegstein, K., Kleinschmidt, A., Sterzer, P., & Giraud, A. L. (2005). Interaction of face and voice areas during speaker recognition. Journal of Cognitive Neuroscience, 17(3), 367–376. https://doi.org/10.1162/0898929053279577

33. Wiggins, I. M., Anderson, C. A., Kitterick, P. T., & Hartley, D. E. H. (2016). Speech-evoked activation in adult temporal cortex measured using functional near-infrared spectroscopy (fNIRS): Are the measurements reliable? Hearing Research, 339, 142–154. https://doi.org/10.1016/j.heares.2016.07.007