

wax and of dermatitis due to Bisphenol A sensitivity other than in epoxy resin workers.

### References

1. Camarasa G. Occupational dermatitis from beeswax. *Contact Dermatitis* 1975; 1: 124.
2. Schwartz L. Cutaneous hazards in the citrus fruit industry. *Arch Dermatol* 1938; 37: 631-638.
3. Thorgeirsson A, Fregert S, Ramnas O. Sensitisation capacity of epoxy resin oligomers in the guinea pig. *Acta Dermato-venereologica* 1978; 58: 17-21.
4. Krajewska D, Rudzki E. Sensitivity to epoxy resins and triethylenetetramine. *Contact Dermatitis* 1976; 2: 135-138.
5. Fregert S, Rorsman H. Hypersensitivity to diethylstilboestrol. Cross sensitisation to dinestrol, hexestrol, Bisphenol-A, p-benzyl-phenol, hydroquinone monobenzylether and p-hydroxybenzoic-bezylester. *Acta Dermato-venereologica* 1960; 40: 206-209.
6. Fregert S, Rorsman H. Hypersensitivity to epoxy resins with reference to the rôle played by Bisphenol-A. *J Invest Dermatol* 1962; 39: 471-472.

## Phototoxicity of a weedkiller: a correction

T. C. HINDSON<sup>1</sup> AND B. L. DIFFEY<sup>2</sup>

<sup>1</sup>Dermatology Department, Sunderland Royal Infirmary, New Durham Road, Sunderland SR2 7JE, England

<sup>2</sup>Regional Medical Physics Department, Dryburn Hospital, Durham DH1 5TW, England

**Key words:** Phototoxicity - weedkiller - glyphosate - benzisothiazolone.

In a previous communication (1), we reported a phototoxic reaction in a patient to the weedkiller 'Tumbleweed' (Murphys Ltd.) which we attributed to its active ingredient, glyphosate. Subsequent information received from the manufacturer indicated that, in addition to glyphosate, the weedkiller contained the preservative benzisothiazolone. The patient was re-phototested as described previously (1) to the separate constituents glyphosate and benzisothiazolone, both at concentrations of 1% in aqueous solution. No phototoxic reactions were observed with glyphosate, but the combination of UV-A and benzisothiazolone

produced delayed erythema at doses in excess of 7 Jcm<sup>-2</sup>; this dose is about one third of the minimal erythema dose on patients' unsensitised skin. We conclude that benzisothiazolone is the phototoxic agent in 'Tumbleweed' and not glyphosate as we originally implied.

### Reference

1. Hindson T C, Diffey B L. Phototoxicity of glyphosate in a weedkiller. *Contact Dermatitis* 1984; 10: 51-52.

## Estradiol enhances nickel-induced blast transformation

S. SILVENNÖINEN-KASSINEN<sup>1</sup>, H. ISOTALO<sup>2</sup> AND H. JAKKULA<sup>1</sup>

<sup>1</sup>Department of Medical Microbiology and <sup>2</sup>Department of Clinical Chemistry, University of Oulu, Oulu, Finland

**Key words:** Nickel sensitivity - blast transformation - estradiol - progesterone - cortisol.

Female subjects suffer about 10 times more often from nickel sensitivity than males (1). Therefore we studied the effect of the female sex steroid hormones on nickel-induced lymphocyte blast transformation which is specific to clinical nickel sensitivity (2).

Lymphocytes of 9 nickel sensitive subjects, 6 female and 3 male, were cultured for 7 days in the presence of 6.25 µg/ml NiSO<sub>4</sub> as described earlier (3). Steroids present in the human male AB serum used (10%) in the RPMI-1640 culture medium were removed by

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.