



PREOPERATIVE AND POSTOPERATIVE PHYSICIAN ORDERS

Admitting Diagnosis: _____

Planned Procedure: _____ ☐ RIGHT EYE ☐ LEFT EYE ☐ BILATERAL ☐ _____

Pre-Operative Orders:

- ☐ Proparacaine 0.5% 1 gtt on arrival to both eyes ☐ Acetaminophen 500mg PO
☐ _____

IV: Saline lock

- 1% or 2% lidocaine injection may be used as local anesthetic prior to IV placement.
- 0.9% sodium chloride flush may be used to verify IV placement.

- ☐ Check blood glucose level and report result to CRNA

Post-Operative Orders:

- ☐ erythromycin ophthalmic ointment to incision sites and operative eye(s)
☐ prednisolone acetate 1% 1 drop to operative eye
☐ bacitracin zinc 500unit/ gram 30gram tube to hairline incision site
☐ _____

For Post-Op Pain:

For pain rated 3-5 on pain scale of 1-10 administer: Acetaminophen 500mg PO

For pain rated 6-10 on pain scale of 1-10: consult physician

For nausea/vomiting administer: If IV is present, administer 4mg Ondansetron IV and re-evaluate

If no IV is present, administer 4mg Ondansetron PO and re-evaluate

If nausea persists after administration, consult physician

Diet: remain NPO in postop, then advance as tolerated upon discharge

IV: Discontinue IV when patient is ready for discharge

Review discharge instructions with patient

Verbal Order – Read Back –Verified: _____

RN/LPN signature

Date

Time

MD signature

Date

Time

OPHTHALMIC HISTORY AND PHYSICAL UPDATE

I attest there are no changes to the patient's history or exam as included in the surgical chart and electronic medical record.

_____, MD Date: _____ Time: _____
Thomas Copperman, MD

Patient Sticker