



THE UNIVERSITY OF TEXAS AT AUSTIN  
McCOMBS SCHOOL OF BUSINESS

# Probability Review 2

---

## Lecture 3

STA 371G


## Sample vs Population

Find out the average house price in Austin.

# Sample vs Population

Find out the average house price in Austin.


How would you do that?



## Sample vs Population

Find out the average house price in Austin.

How would you do that?




Look at each house price?

## Sample vs Population

Find out the average house price in Austin.

How would you do that?




Look at each house price?

360,000 houses in Austin!

## Sample vs Population

Find out the average house price in Austin.

How would you do that?



Look at each house price?

360,000 houses in Austin!

Can we do something smarter?

# Sample vs Population

A smarter approach:

# Sample vs Population

A smarter approach:

- Pick  $n$  houses randomly (e.g.  $n = 100$ )



# Sample vs Population

A smarter approach:

- Pick  $n$  houses randomly (e.g.  $n = 100$ )
- Take the average of the prices of these  $n$  houses

# Sample vs Population

A smarter approach:

- Pick  $n$  houses randomly (e.g.  $n = 100$ )
- Take the average of the prices of these  $n$  houses
- Hope that your estimate is close to the true price average.

## Sample vs Population

A smarter approach:

- Pick  $n$  houses randomly (e.g.  $n = 100$ )
- Take the average of the prices of these  $n$  houses
- Hope that your estimate is close to the true price average.

Just like making polls to predict election results!

## Sample vs Population

	Population	Sample
Members	all house prices	prices you picked
Average	population mean	sample mean
Variance	population variance	sample variance

## Sample vs Population


	Population	Sample
Members	all house prices	prices you picked
Average	population mean	sample mean
Variance	population variance	sample variance

Estimating a **population parameter** (population mean) based on a **sample statistic** (sample mean).

# Collecting a sample

Zillow.com, “Austin, TX.”

- Click “More Map”
- Select 15 houses, note their prices in an R script.
- Do not discard any price, use the first 15
- Try to represent different regions



## Collecting a sample

Your R script should look like this

```
# Create a vector of house prices (You should have 15 price data)
sample_house_prices <- c(327000,276000,513000)
# Calculate sample statistics
sample_mean <- mean(sample_house_prices)
sample_variance <- var(sample_house_prices)
sample_standard_deviation <- sd(sample_house_prices)
# Sample mean of first 5 houses
sample_mean_5 <- mean(sample_house_prices[1:5])
# Print them to console
cat("Sample Mean", sample_mean)
cat("Sample Variance", sample_variance)
cat("Sample Standard Deviation", sample_standard_deviation)
cat("Sample Mean of first 5 houses",sample_mean_5)
```

# Sampling Distribution

On Learning Catalytics, enter your results.

And here is what they look like...




# Sampling Distribution


Distribution of your answers → Sampling distribution

Statistic	Population	Sample Mean
Mean	$\mu$	$\mu$
Standard Deviation	$\sigma$	$\sigma/\sqrt{n}$

# Sampling Distribution



# Sampling Distribution



## Sampling Distribution

Assume  $\mu = \$300\text{K}$ ,  $\sigma = \$60\text{K}$ .

	$n$	$\sigma/\sqrt{n}$	3 std. dev. range (99.7%)
Survey 1	25	\$12K	\$264K ..... \$336K
Survey 2	100	\$6K	\$282K ..... \$318K
Survey 3	3600	\$1K	\$297K ... \$303K

# Sampling Distribution

Let's compare sample mean of 5 houses vs 15 houses.

What do you expect to see?

## $t$ Distribution

We often do not know population variance and use sample variance instead.

In that case, the sample mean will have a  $t$  distribution.

# Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

## Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

Your survey on 25 houses: Average house price is \$305K.



# Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

Your survey on 25 houses: Average house price is \$305K.

Questions, questions...

# Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

Your survey on 25 houses: Average house price is \$305K.

Questions, questions...

- Would you reject the hypothesis? Why?

# Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

Your survey on 25 houses: Average house price is \$305K.

Questions, questions...

- Would you reject the hypothesis? Why?
- Is it possible that, out of bad luck, you picked the cheapest houses?

# Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

Your survey on 25 houses: Average house price is \$305K.

Questions, questions...

- Would you reject the hypothesis? Why?
- Is it possible that, out of bad luck, you picked the cheapest houses?
- Would you be more comfortable with your conclusion if you had 1000 houses in your survey?

# Hypothesis Testing

Hypothesis: Average house price in Austin is \$1M.

Your survey on 25 houses: Average house price is \$305K.

Questions, questions...

- Would you reject the hypothesis? Why?
- Is it possible that, out of bad luck, you picked the cheapest houses?
- Would you be more comfortable with your conclusion if you had 1000 houses in your survey?
- When should you reject the hypothesis? When not?

## P-Value

Your sample mean: \$305K.

## P-Value

Your sample mean: \$305K.

$H_0 : \mu = \$1M$  (Null hypothesis)

$H_1 : \mu < \$1M$  (Alternative hypothesis)

## P-Value

Your sample mean: \$305K.

$H_0 : \mu = \$1M$  (Null hypothesis)

$H_1 : \mu < \$1M$  (Alternative hypothesis)

The **P-value** is “the probability of observing such an extreme (\$305K or less) sample statistic given the null hypothesis is true.”



## P-Value

Your sample mean: \$305K.

$H_0 : \mu = \$1M$  (Null hypothesis)

$H_1 : \mu < \$1M$  (Alternative hypothesis)

The **P-value** is “the probability of observing such an extreme (\$305K or less) sample statistic given the null hypothesis is true.”

- $P\text{-value} \leq \alpha$ , reject the null hypothesis

## P-Value

Your sample mean: \$305K.

$H_0 : \mu = \$1M$  (Null hypothesis)

$H_1 : \mu < \$1M$  (Alternative hypothesis)

The **P-value** is “the probability of observing such an extreme (\$305K or less) sample statistic given the null hypothesis is true.”

- $P\text{-value} \leq \alpha$ , reject the null hypothesis
- $P\text{-value} > \alpha$ , do not reject the null hypothesis

## P-Value

Your sample mean: \$305K.

$H_0 : \mu = \$1M$  (Null hypothesis)

$H_1 : \mu < \$1M$  (Alternative hypothesis)

The **P-value** is “the probability of observing such an extreme (\$305K or less) sample statistic given the null hypothesis is true.”

- $P\text{-value} \leq \alpha$ , reject the null hypothesis
- $P\text{-value} > \alpha$ , do not reject the null hypothesis


$\alpha$  is usually chosen as 0.05 prior to sampling.

## P-Value

If the null hypothesis were true...


## P-Value

If the null hypothesis were true...



## P-Value


If the null hypothesis were true...



P-value is smaller than  $10^{-100}$ , while  $\alpha = 0.05$ .

## P-Value

If the null hypothesis were true...



P-value is smaller than  $10^{-100}$ , while  $\alpha = 0.05$ .

Rather than thinking you are cursed, you simply reject the hypothesis!

## P-Value

Learning Catalytics...



## P-Value

Learning Catalytics...

Your sample mean = \$305K.

## P-Value

Learning Catalytics...

Your sample mean = \$305K.

$H_0$ : Average house price is \$390K.

Would you reject the hypothesis?

P-value = 0.01,  $\alpha=0.05$

## P-Value

Learning Catalytics...

Your sample mean = \$305K.

$H_0$ : Average house price is \$390K.

Would you reject the hypothesis?

P-value = 0.01,  $\alpha=0.05$

$H_0$ : Average house price is \$320K.

Would you reject the hypothesis?

P-value = 0.34,  $\alpha=0.01$

## P-Value

Learning Catalytics...

Your sample mean = \$305K.

$H_0$ : Average house price is \$390K.

Would you reject the hypothesis?

P-value = 0.01,  $\alpha=0.05$

$H_0$ : Average house price is \$320K.

Would you reject the hypothesis?

P-value = 0.34,  $\alpha=0.01$

## Confidence Interval

Sample mean is not equal to the population mean, but “close.”

## Confidence Interval


Sample mean is not equal to the population mean, but “close.”

**Confidence interval** is a range that includes the population mean with a certain level of “confidence.”

## Confidence Interval

Sample mean is not equal to the population mean, but “close.”

**Confidence interval** is a range that includes the population mean with a certain level of “confidence.”



# Confidence Interval

Add the following to your R script

```
# Calculate 95% confidence interval (default)
avg_price_ci_95 <- t.test(sample_house_prices)
# Calculate 99% confidence interval
avg_price_ci_99 <- t.test(sample_house_prices, conf.level = 0.99)
# Display results
cat("95% confidence interval is:", avg_price_ci_95$conf.int)
cat("99% confidence interval is:", avg_price_ci_99$conf.int)
```

Enter your results on Learning Catalytics.