



THE UNIVERSITY OF TEXAS AT AUSTIN
McCOMBS SCHOOL OF BUSINESS

Probability Review 1

Lecture 2

STA 371G

Reading assignments

- Sign up at perusal1.com

Reading assignments

- Sign up at perusall.com
- Reading assignments help you build (incomplete) mental structures that we can build on in class

Reading assignments

- Sign up at perusal1.com
- Reading assignments help you build (incomplete) mental structures that we can build on in class
- Your assignment is to read and annotate: ask a question, answer a classmate's question, make an observation, try to explain something better than the text does

Reading assignments

- Sign up at perusal1.com
- Reading assignments help you build (incomplete) mental structures that we can build on in class
- Your assignment is to read and annotate: ask a question, answer a classmate's question, make an observation, try to explain something better than the text does
- Make at least three high-quality annotations per document to get full credit (examples on Canvas)

Reading assignments

- Sign up at perusal1.com
- Reading assignments help you build (incomplete) mental structures that we can build on in class
- Your assignment is to read and annotate: ask a question, answer a classmate's question, make an observation, try to explain something better than the text does
- Make at least three high-quality annotations per document to get full credit (examples on Canvas)
- First assignment is due **January 31 at 11:55 PM**

Probability Theory

The Concept of Probability

What do each of these have in common?

Probability Theory

The Concept of Probability

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

Probability Theory

The Concept of Probability

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

We cannot predict any of these with certainty.

Probability Theory

The Concept of Probability

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

We cannot predict any of these with certainty.

Yet, we can model them using **probability theory** and study the values they might take, associated probabilities etc.

Probability Theory

Definitions

Definition

A **random variable** expresses the outcome of a **random experiment** as a number. It is denoted by an uppercase letter.

Probability Theory

Definitions

Definition

A **random variable** expresses the outcome of a **random experiment** as a number. It is denoted by an uppercase letter.

- Random experiment → Rolling a die

Probability Theory

Definitions

Definition

A **random variable** expresses the outcome of a **random experiment** as a number. It is denoted by an uppercase letter.

- Random experiment \rightarrow Rolling a die
- Random variable $\rightarrow X$: The outcome when a die is rolled

Probability Theory

Definitions

Definition

A **random variable** expresses the outcome of a **random experiment** as a number. It is denoted by an uppercase letter.

- Random experiment \rightarrow Rolling a die
- Random variable $\rightarrow X$: The outcome when a die is rolled
- Random variable \rightarrow

$$Y : \begin{cases} 1, & \text{if outcome is odd number,} \\ 2, & \text{if outcome is even number} \end{cases}$$

Probability Theory

Definitions

Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

Probability Theory

Definitions

Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

Probability Theory

Definitions

Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

Examples

- X : Number of stocks on NYSE whose price change today (discrete)

Probability Theory

Definitions

Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

Examples

- X : Number of stocks on NYSE whose price change today (discrete)
- Y : Average price change of the stocks on NYSE (continuous)

Probability Theory

Exercise

Discrete or continuous?

Probability Theory

Exercise

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year

Probability Theory

Exercise

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year
- Lifetime of your MacBook Air

Probability Theory

Exercise

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year
- Lifetime of your MacBook Air
- Number of unique visitors to Amazon.com over the next week

Probability Theory

Exercise

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year
- Lifetime of your MacBook Air
- Number of unique visitors to Amazon.com over the next week
- S&P500 index at the end of January

Probability Theory

Definitions

Definition

Probability is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed.

Probability Theory

Definitions

Definition

Probability is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed.

Probability is a number always between 0 and 1.

Probability Theory

Definitions

Definition

Probability is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed.

Probability is a number always between 0 and 1.

0 implies impossibility, 1 implies certainty.

Probability Theory

Definitions

Definition

Probability is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed.

Probability is a number always between 0 and 1.

0 implies impossibility, 1 implies certainty.

Examples

- X : S&P500 at the end of 2017, $P(X > 2270) = 0.85$

Probability Theory

Definitions

Definition

Probability is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed.

Probability is a number always between 0 and 1.

0 implies impossibility, 1 implies certainty.

Examples

- X : S&P500 at the end of 2017, $P(X > 2270) = 0.85$
- Y : Lifetime of your MacBook, $P(Y > 15 \text{ years}) = 0.05$

Probability Distributions

So, we have defined a random variable. How do we know what the probabilities are?

Probability Distributions

So, we have defined a random variable. How do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is, $P(5 < X < 7) = ?$

Probability Distributions

So, we have defined a random variable. How do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is, $P(5 < X < 7) = ?$

Definition

The **probability distribution** of a random variable X is a description of the probabilities associated with the possible values of X .

Probability Distributions

So, we have defined a random variable. How do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is, $P(5 < X < 7) = ?$

Definition

The **probability distribution** of a random variable X is a description of the probabilities associated with the possible values of X .

Discrete random variable \rightarrow Probability Mass Function (p.m.f.)

Probability Distributions

So, we have defined a random variable. How do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is, $P(5 < X < 7) = ?$

Definition

The **probability distribution** of a random variable X is a description of the probabilities associated with the possible values of X .

Discrete random variable \rightarrow Probability Mass Function (p.m.f.)

Continuous random variable \rightarrow Probability Density Function (p.d.f.)

Probability Distributions

Discrete Random Variables

Example

X : The outcome when you roll an n -sided fair die.

Probability Distributions

Discrete Random Variables

Example

X : The outcome when you roll an n -sided fair die.

Since this is a fair die, the corresponding probability mass function is

$$f(x) = \begin{cases} \frac{1}{n} & x = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

Probability Distributions

Discrete Random Variables

Example

X : The outcome when you roll an n -sided fair die.

Since this is a fair die, the corresponding probability mass function is

$$f(x) = \begin{cases} \frac{1}{n} & x = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

- $f(2) = P(X = 2)$, which is the probability of observing a “2.”

Probability Distributions

Discrete Random Variables

Example

X : The outcome when you roll an n -sided fair die.

Since this is a fair die, the corresponding probability mass function is

$$f(x) = \begin{cases} \frac{1}{n} & x = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

- $f(2) = P(X = 2)$, which is the probability of observing a “2.”
- The probabilities always sum to 1. ($n \times \frac{1}{n} = 1$).

Probability Distributions

Discrete Random Variables

Example

X : The outcome when you roll an n -sided fair die.

Since this is a fair die, the corresponding probability mass function is

$$f(x) = \begin{cases} \frac{1}{n} & x = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

- $f(2) = P(X = 2)$, which is the probability of observing a “2.”
- The probabilities always sum to 1. ($n \times \frac{1}{n} = 1$).
- This is an example of a **Discrete Uniform Distribution**.

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

Let's assume Y has a **Continuous Uniform Distribution** with a maximum of 20 years. Its probability distribution is then given by the probability density function

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

Let's assume Y has a **Continuous Uniform Distribution** with a maximum of 20 years. Its probability distribution is then given by the probability density function

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

Let's assume Y has a **Continuous Uniform Distribution** with a maximum of 20 years. Its probability distribution is then given by the probability density function

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

What is $P(Y = 5) = ?$ or $P(Y = 5.5) = ?$ or $P(Y = 5.551234123) = ?$

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

Let's assume Y has a **Continuous Uniform Distribution** with a maximum of 20 years. Its probability distribution is then given by the probability density function

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

What is $P(Y = 5) = ?$ or $P(Y = 5.5) = ?$ or $P(Y = 5.551234123) = ?$

They are all 0.

Probability Distributions

Continuous Random Variables

Warning!

For a continuous random variable, $P(Y = a)$ is always zero, regardless of a .

Probability Distributions

Continuous Random Variables

Warning!

For a continuous random variable, $P(Y = a)$ is always zero, regardless of a .

For this reason, for continuous random variables we ask questions like " $P(a \leq Y \leq b) = ?$ "

Probability Distributions

Continuous Random Variables

Warning!

For a continuous random variable, $P(Y = a)$ is always zero, regardless of a .

For this reason, for continuous random variables we ask questions like " $P(a \leq Y \leq b) = ?$ "

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

$$P(5 < Y < 7) =$$

Probability Distributions

Continuous Random Variables

Example

Y : Lifetime of your MacBook (in years)

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

$$P(5 < Y < 7) = \int_5^7 \frac{1}{20} dy = \frac{y}{20} \Big|_5^7 = \frac{7}{20} - \frac{5}{20} = \frac{1}{10}$$

Probability Distributions

Continuous Random Variables

Example


Y : Lifetime of your MacBook (in years)

$$f(y) = \begin{cases} \frac{1}{20} & 0 \leq y \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

$$P(5 < Y < 7) = \int_5^7 \frac{1}{20} dy = \frac{y}{20} \Big|_5^7 = \frac{7}{20} - \frac{5}{20} = \frac{1}{10}$$

In general, $P(a \leq Y \leq b) = \int_a^b f(y) dy$.

Probability Distributions



Mean, Variance and Standard Deviation

Definition

Mean or **Expected Value** of a random variable X is a measure of the center of its probability distribution. It is a weighted average of all possible values X can take, where the weights are the corresponding probabilities.

Mean, Variance and Standard Deviation

Definition

Mean or **Expected Value** of a random variable X is a measure of the center of its probability distribution. It is a weighted average of all possible values X can take, where the weights are the corresponding probabilities.

Discrete random variable X

$$\begin{aligned}\mu_X = E[X] &= \sum_x xf(x) \\ &= \sum_x xP(X = x)\end{aligned}$$

Continuous random variable Y

$$\mu_Y = E[Y] = \int_y yf(y) dy$$

Mean, Variance and Standard Deviation

Definition

Variance of a random variable X is a measure of the spread, or variability in its distribution. **Standard Deviation** of X is the square root of its variance.

Mean, Variance and Standard Deviation

Definition

Variance of a random variable X is a measure of the spread, or variability in its distribution. **Standard Deviation** of X is the square root of its variance.

Discrete random variable X

$$\sigma_X^2 = \text{Var}(X) = E[(X - \mu_X)^2] = \sum_x (x - \mu_X)^2 f(x)$$

Mean, Variance and Standard Deviation

Definition

Variance of a random variable X is a measure of the spread, or variability in its distribution. **Standard Deviation** of X is the square root of its variance.

Discrete random variable X

$$\sigma_X^2 = \text{Var}(X) = E[(X - \mu_X)^2] = \sum_x (x - \mu_X)^2 f(x)$$

Continuous random variable Y

$$\sigma_Y^2 = \text{Var}(Y) = E[(Y - \mu_Y)^2] = \int_y (y - \mu_Y)^2 f(y) dy$$

Law of Large Numbers

Definition

For a random variable X , the average of X_1, X_2, \dots, X_n gets very close to the expected value of X ($E[X]$) for large n .

Law of Large Numbers

Definition

For a random variable X , the average of X_1, X_2, \dots, X_n gets very close to the expected value of X ($E[X]$) for large n .

Example

A die is rolled $n = 4$ times: $x_1 = 4, x_2 = 6, x_3 = 1, x_4 = 1$. The average is

$$\frac{x_1 + x_2 + x_3 + x_4}{4} = \frac{4 + 6 + 1 + 1}{4} = 3$$

Law of Large Numbers

Definition

For a random variable X , the average of X_1, X_2, \dots, X_n gets very close to the expected value of X ($E[X]$) for large n .

Example

A die is rolled $n = 4$ times: $x_1 = 4, x_2 = 6, x_3 = 1, x_4 = 1$. The average is

$$\frac{x_1 + x_2 + x_3 + x_4}{4} = \frac{4 + 6 + 1 + 1}{4} = 3$$

For large n , the average will be around 3.5; because $E[X] = 3.5$.

Law of Large Numbers

R Exercise

Go to R Studio...

```
# Simulate rolling a die 1 time
sample(c(1, 2, 3, 4, 5, 6), 1, replace=T)
# Simulate rolling a die 4 times
sample(c(1, 2, 3, 4, 5, 6), 4, replace=T)
# Take the average
mean(sample(c(1, 2, 3, 4, 5, 6), 4, replace=T))
# Let's increase the number of dice
mean(sample(c(1, 2, 3, 4, 5, 6), 10, replace=T))
```

Normal Distribution a.k.a. the Bell Curve

A very common continuous probability distribution.

The mean and variance together uniquely define the distribution:


$$N(\mu, \sigma^2)$$

Normal Distribution a.k.a. the Bell Curve

A very common continuous probability distribution.

The mean and variance together uniquely define the distribution:

$$N(\mu, \sigma^2)$$



Central Limit Theorem

Definition

The sum (and average) of large number of independent random variables will be approximately normally distributed.

Central Limit Theorem

Definition

The sum (and average) of large number of independent random variables will be approximately normally distributed.

Example

Exam grades tend to be normally distributed because they are the combination of independent random variables:

a student's exam grade = preparedness + focus + learning + . . .

Central Limit Theorem

Definition

The sum (and average) of large number of independent random variables will be approximately normally distributed.

Example

Exam grades tend to be normally distributed because they are the combination of independent random variables:

a student's exam grade = preparedness + focus + learning + . . .

Central Limit Theorem

R Exercise

X : Average house price in a particular zip code. Assume house prices in each zip code are uniformly distributed in $[100, 400]$ (\$K)

Central Limit Theorem

R Exercise

X : Average house price in a particular zip code. Assume house prices in each zip code are uniformly distributed in $[100, 400]$ (\$K)

We will simulate k different zip codes, each containing n houses.

Central Limit Theorem

R Exercise

X : Average house price in a particular zip code. Assume house prices in each zip code are uniformly distributed in $[100, 400]$ (\$K)

We will simulate k different zip codes, each containing n houses.

```
# Simulating a zip code with n=3 houses
runif(3, min=100, max=400)
# Repeat this for k=5 zip codes.
house_prices <- replicate(5, runif(3, min=100, max=400))
# Find the average house price in each zip code
avg_house_prices <- colMeans(house_prices)
# Make a histogram of the average house price in your zip codes
hist(avg_house_prices)
# Increase n and k and try again!
```