



THE UNIVERSITY OF TEXAS AT AUSTIN  
McCOMBS SCHOOL OF BUSINESS

# Probability Review 3

---

## Lecture 4

STA 371G

## Who are these people?



Zack



Kelly




Screech



Slater



Jessie



Lisa

## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character

## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)

## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*


## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*


## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*

## Some events




Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*

$P(M) =$



## Some events




Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*

$$P(M) = 3/6 \quad P(L) =$$

## Some events




Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*

$$P(M) = 3/6 \qquad P(L) = 3/6$$

$$P(M|L) =$$

## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*

$$P(M) = 3/6 \qquad P(L) = 3/6$$

$$P(M|L) = 1/3 \qquad P(L|M) =$$

## Some events



Suppose we pick a *Saved by the Bell* character at random. Each of these are events:

- $M$  = we select a male character
- $L$  = we select someone with lush hair, where “lush” means “long curly hair” (mullet included!)
- $M|L$  = we select a male *from among the lush-haired characters*
- $L|M$  = we select a lush-haired character *from among the male characters*

$$P(M) = 3/6$$

$$P(L) = 3/6$$

$$P(M|L) = 1/3$$

$$P(L|M) = 1/3$$

## Joint probability



$P(M \text{ and } L)$  = probability of selecting a lush-haired male

## Joint probability



$P(M \text{ and } L) = \text{probability of selecting a lush-haired male}$   
 $= 1/6$

## Joint probability



$P(M \text{ and } L)$  = probability of selecting a lush-haired male

$$= 1/6$$

$$= P(M)P(L|M) = \frac{3}{6} \cdot \frac{1}{3}$$

$$= P(L)P(M|L) = \frac{3}{6} \cdot \frac{1}{3}$$

## Joint probability



$P(M \text{ and } L)$  = probability of selecting a lush-haired male

$$= 1/6$$

$$= P(M)P(L|M) = \frac{3}{6} \cdot \frac{1}{3}$$

$$= P(L)P(M|L) = \frac{3}{6} \cdot \frac{1}{3}$$

$$\neq P(M)P(L)$$



## One or the other



$P(M \text{ or } L)$  = probability of selecting a lush-haired character, a male character, or both

## One or the other



$P(M \text{ or } L)$  = probability of selecting a lush-haired  
character, a male character, or both  
 $= 5/6$

We have to subtract off  $P(M \text{ and } L)$  because otherwise  
we are double-counting Slater!

## One or the other



$P(M \text{ or } L)$  = probability of selecting a lush-haired character, a male character, or both

$$= 5/6$$

$$= P(M) + P(L) - P(M \text{ and } L)$$

$$= 3/6 + 3/6 - 1/6$$

We have to subtract off  $P(M \text{ and } L)$  because otherwise we are double-counting Slater!

## Conditional probability

When we say  $P(A|B)$ , what we mean is: “In a world where we know B has already happened, how likely is it that A also happened?”





Now that you've found a strange pair of underwear in your dresser,  
what is the probability that your partner is cheating on you?

$$P(\text{partner is cheating} \mid \text{underwear found})$$

Now that you've found a strange pair of underwear in your dresser,  
what is the probability that your partner is cheating on you?

$$P(\text{partner is cheating} \mid \text{underwear found})$$

It's hard to know how to estimate this directly!



Let's come up with estimates for the following:

- $P(\text{underwear found} \mid \text{partner is cheating})$

Let's come up with estimates for the following:

- $P(\text{underwear found} \mid \text{partner is cheating})$
- $P(\text{underwear found} \mid \text{partner is not cheating})$

Let's come up with estimates for the following:

- $P(\text{underwear found} \mid \text{partner is cheating})$
- $P(\text{underwear found} \mid \text{partner is not cheating})$
- $P(\text{partner is cheating})$  (before we found the underwear!)

## Bayes' Rule

Bayes' Rule lets us reverse the conditional probabilities!

$U$  = underwear found,  $C$  = partner is cheating

$$P(C|U) = \frac{P(U|C)P(C)}{P(U|C)P(C) + P(U|\bar{C})P(\bar{C})}$$

Think of Bayes' Rule as a way to update our thinking based on new information:

$P(C)$       ← Prior probability

$P(C|U)$    ← Posterior probability (includes new information)

# HIV Testing

- HIV testing is important for public health, but HIV tests are not perfect

## HIV Testing

- HIV testing is important for public health, but HIV tests are not perfect
- The OraQuick ADVANCE Rapid HIV-1/2 Antibody Test has the following properties:

# HIV Testing

- HIV testing is important for public health, but HIV tests are not perfect
- The OraQuick ADVANCE Rapid HIV-1/2 Antibody Test has the following properties:
  - If you **have HIV**, there is a 99.3% chance the test will show a **positive result**



# HIV Testing

- HIV testing is important for public health, but HIV tests are not perfect
- The OraQuick ADVANCE Rapid HIV-1/2 Antibody Test has the following properties:
  - If you **have HIV**, there is a 99.3% chance the test will show a **positive result**
  - If you **do not have HIV**, there is a 99.8% chance the test will show a **negative result**

## HIV Testing

- HIV testing is important for public health, but HIV tests are not perfect
- The OraQuick ADVANCE Rapid HIV-1/2 Antibody Test has the following properties:
  - If you **have HIV**, there is a 99.3% chance the test will show a **positive result**
  - If you **do not have HIV**, there is a 99.8% chance the test will show a **negative result**
- 0.4% of people in the US are HIV-positive

## HIV Testing

We know  $P(TP|HP) = 0.993$ , but we really want to know  $P(HP|TP)$ !

# HIV Testing

- If you have HIV, there is a 99.3% chance the test will show a positive result

# HIV Testing

- If you have HIV, there is a 99.3% chance the test will show a positive result
- If you do not have HIV, there is a 99.8% chance the test will show a negative result

# HIV Testing

- If you have HIV, there is a 99.3% chance the test will show a positive result
- If you do not have HIV, there is a 99.8% chance the test will show a negative result
- But if you test positive there is about a 1/3 chance you do NOT have HIV!

# HIV Testing

- If you have HIV, there is a 99.3% chance the test will show a positive result
- If you do not have HIV, there is a 99.8% chance the test will show a negative result
- But if you test positive there is about a 1/3 chance you do NOT have HIV!
- This is counterintuitive — it's because of the way we are wired (it even has a name: “base rate fallacy”)