

Theoretical formalism of `DielectricKit`

Meng Wu

Contents

| | | |
|----------|--|----------|
| 1 | Introduction | 1 |
| 2 | Many-body perturbation theory | 1 |
| 3 | The <i>GW</i> method and random-phase approximation | 4 |
| 4 | FFT and matrix elements | 6 |
| 5 | Symmetry of ε^{-1} and χ^* | 6 |
| 6 | Real-space ε^{-1} and χ^* | 7 |
| 7 | Parallelism and scaling performance | 8 |
| 7.1 | <code>Chi.x</code> | 8 |
| 7.2 | <code>EpsInv.x</code> | 9 |
| 7.3 | <code>RealSpace.x</code> | 9 |

1 Introduction

In this document, I will describe in detail the theoretical formalism of `DielectricKit`. I will cover the topic of many-body perturbation theory, polarizability and dielectric response functions within the random-phase approximation (RPA), fast Fourier transform (FFT), symmetry of Bloch waves and response functions, and finally how to calculate real-space response functions.

2 Many-body perturbation theory

Electronic structure in solids is a typical many-body problem, involving a huge number (\sim the Avogadro's constant) of electrons, ions, as well as under various external fields. The most systematic and elegant approach to this problem is based on the quantum field theory and the many-body perturbation theory (MBPT) (also called the Green's function method) [1, 2, 3, 4], from which the *GW* and *GW*-BSE methods are derived. MBPT is probably one of the most powerful, predictive and versatile formalisms in physics. This section is just a sketch of important concepts and conclusions most relevant to the development of the *GW* and *GW*-BSE methods, and we only focus on the Green's function formalism for electrons in the following. More details and other topics about MBPT can be found in several excellent reviews and textbooks [4, 5, 6, 7, 8].

We start from a general Hamiltonian \hat{H} for a system of interacting electrons under an external potential $V_{\text{ext}}(\mathbf{x})$ [9, 5, 6, 10]:

$$\hat{H} = \int d\mathbf{x} \hat{\psi}^\dagger(\mathbf{x}) \left[-\frac{\hbar^2}{2m_e} \nabla^2 + V_{\text{ext}}(\mathbf{x}) \right] \hat{\psi}(\mathbf{x}) + \frac{1}{2} \int d\mathbf{x} d\mathbf{x}' v(\mathbf{r}, \mathbf{r}') \hat{\psi}^\dagger(\mathbf{x}) \hat{\psi}^\dagger(\mathbf{x}') \hat{\psi}(\mathbf{x}') \hat{\psi}(\mathbf{x}), \quad (1)$$

where $\mathbf{x} = \{\mathbf{r}\sigma\}$ and $\int d\mathbf{x} = \sum_\sigma \int d\mathbf{r}$. The spin-resolved electron density operator is defined as, $\hat{n}_e(\mathbf{x}) \equiv \hat{\psi}^\dagger(\mathbf{x}) \hat{\psi}(\mathbf{x})$. The Coulomb interaction is given by, $v(\mathbf{r}, \mathbf{r}') = \frac{q_e^2}{4\pi\epsilon_0|\mathbf{r}-\mathbf{r}'|}$, where ϵ_0 is the vacuum permittivity.

The *time-ordered single-particle Green's function* G in the zero-temperature formalism is defined as:

$$\begin{aligned} G(12) &\equiv \left(-\frac{i}{\hbar}\right) \frac{\langle \Psi_0 | T [e^{i\hat{H}t_1/\hbar} \hat{\psi}(\mathbf{x}_1) e^{-i\hat{H}t_1/\hbar} e^{i\hat{H}t_2/\hbar} \hat{\psi}^\dagger(\mathbf{x}_2) e^{-i\hat{H}t_2/\hbar}] | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle} \\ &\equiv \left(-\frac{i}{\hbar}\right) \frac{\langle \Psi_0 | T [\hat{\psi}(1) \hat{\psi}^\dagger(2)] | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle}, \end{aligned} \quad (2)$$

where $1 \equiv \{\mathbf{x}_1 t_1\}$, T is the time-ordering operator, and $|\Psi_0\rangle$ the ground state of an interacting system with the Hamiltonian in Eq. (1). Throughout the following derivation, the time-dependence of the field operator $\hat{\psi}(1)$ comes from the Hamiltonian in Eq. (1),

$$\hat{\psi}(1) \equiv \hat{\psi}(\mathbf{x}_1, t_1) = e^{i\hat{H}t_1/\hbar} \hat{\psi}(\mathbf{x}_1) e^{-i\hat{H}t_1/\hbar}. \quad (3)$$

Note that the following discussion also applies to the finite-temperature G , as shown in Ref. [3]. After a time Fourier transform with respect to $t_1 - t_2$, we get $G(\mathbf{x}_1, \mathbf{x}_2; \omega)$, defined in the frequency domain.

The single-particle Green's function is at the center of MBPT because (i) the Feynman rules are simpler for G than for other operators, and (ii) the expectation value of any single-particle operator in the ground state of the system can be calculated using G [5]. In the following, we will adopt the functional derivative approach [3, 4, 10] to reduce the many-body problem to the solution of a coupled set of nonlinear integral equations. This approach avoids the cumbersome conventional diagrammatic expansion of relevant quantities.

We first add an external time-dependent local potential φ to the Hamiltonian in Eq. (1) in order to perturb the system,

$$\hat{H}'(t) \equiv \int d\mathbf{x} \hat{\psi}^\dagger(\mathbf{x}) \varphi(\mathbf{x}, t) \hat{\psi}(\mathbf{x}). \quad (4)$$

This perturbation potential will go to zero at the end of our derivation. Since there is an extra term in the Hamiltonian,

$$\hat{H}''(t) \equiv \hat{H} + \hat{H}'(t), \quad (5)$$

we adopt the interaction picture from now on,

$$\hat{H}'_I(t) \equiv e^{i\hat{H}t/\hbar} \hat{H}'(t) e^{-i\hat{H}t/\hbar}, \quad (6)$$

and introduce the S matrix,

$$\hat{S} \equiv \exp \left\{ -\frac{i}{\hbar} \int_{-\infty}^{\infty} dt \hat{H}'_I(t) \right\}. \quad (7)$$

The generalized single-particle Green's functions with respect to $\hat{H}''(t)$ is then,

$$G(12) = \left(-\frac{i}{\hbar}\right) \frac{\langle \Psi_0 | T [\hat{S} \hat{\psi}(1) \hat{\psi}^\dagger(2)] | \Psi_0 \rangle}{\langle \Psi_0 | T [\hat{S}] | \Psi_0 \rangle}. \quad (8)$$

When we take the limit of $\hat{H}' \rightarrow 0$, the S matrix will be reduced to an identity operator. We now introduce a total potential V_{tot} averaged over the many-body ground state $|\Psi_0\rangle$ as,

$$V_{\text{tot}}(1) \equiv \varphi(1) + \int d2 v(12) \langle \hat{n}_e(2) \rangle = \varphi(1) - i\hbar \int d2 v(12) G(22^+), \quad (9)$$

where $v(12) = v(\mathbf{r}_1, \mathbf{r}_2) \delta(t_1 - t_2)$, $\langle \hat{n}_e(2) \rangle \equiv \langle \Psi_0 | \hat{n}_e(2) | \Psi_0 \rangle = -i\hbar G(22^+)$, $2 = \{\mathbf{x}_2 t_2^+\}$, and $t_2^+ = t_2 + \eta$ with $\eta \rightarrow 0^+$. We relate V_{tot} to the external potential φ by introducing the *reducible polarizability* χ and the *inverse dielectric function* ε^{-1} as functional derivatives:

$$\chi(12) \equiv \frac{\delta \langle \hat{n}_e(1) \rangle}{\delta \varphi(2)}, \quad (10)$$

$$\varepsilon^{-1}(12) \equiv \frac{\delta V_{\text{tot}}(1)}{\delta \varphi(2)} = \delta(12) + \int d3 v(13) \chi(32). \quad (11)$$

Eq. (11) can be inverted to get,

$$\varepsilon(12) = \delta(12) - \int d3 v(13) \chi^*(32), \quad (12)$$

where we have defined the *irreducible polarizability* χ^* as,

$$\chi^*(12) \equiv \frac{\delta \langle \hat{n}_e(1) \rangle}{\delta V_{\text{tot}}(2)}. \quad (13)$$

The *screened Coulomb interaction* W is defined intuitively as,

$$W(12) \equiv \int d3 \varepsilon^{-1}(13) v(32). \quad (14)$$

Combine Eqs. (11), (13) and (14), and we get the following expressions of W ,

$$\begin{aligned} W(12) &= v(12) + \int d(34) v(13) \chi(34) v(42) \\ &= v(12) + \int d(34) v(13) \chi^*(34) W(42) \end{aligned} \quad (15)$$

The *electron self-energy* Σ , the Hartree term Σ_{H} and the *mass operator* $M(12)$ are defined as follows,

$$\Sigma(12) \equiv \Sigma_{\text{H}}(12) + M(12), \quad (16)$$

$$\Sigma_{\text{H}}(12) \equiv \delta(12) \int d2 v(12) \langle \hat{n}_e(2) \rangle, \quad (17)$$

$$M(12) \equiv i\hbar \int d(34) v(13) \frac{\delta G(14)}{\delta \varphi(3)} G^{-1}(42). \quad (18)$$

Σ drives the equation of motion of G ,

$$\left[i\hbar \frac{\partial}{\partial t_1} + \frac{\hbar^2}{2m_e} \nabla_1^2 \right] G(12) - \int d3 \Sigma(13) G(32) = \delta(12). \quad (19)$$

By defining a noninteracting single-particle Green's function $G_0(12)$ at the absence of the Coulomb interaction, Eq. (19) can be reformulated into the *Dyson's equation*,

$$G(12) = G_0(12) + \int d(34) G_0(13) \Sigma(34) G(42). \quad (20)$$

The *irreducible three-point vertex function* Γ^* is defined as,

$$\Gamma^*(123) \equiv -\frac{\delta G^{-1}(12)}{\delta V_{\text{tot}}(3)} = \delta(12)\delta(13) + \int d(4567) \frac{\delta M(12)}{\delta G(45)} G(46)G(75)\Gamma^*(673), \quad (21)$$

M can then be expressed as an integral involving G , Γ^* , and W :

$$M(12) = i\hbar \int d(34) G(13)W(41)\Gamma^*(324). \quad (22)$$

χ^* is also related to Γ^* according to,

$$\chi^*(12) = -i\hbar \int d(34) G(13)G(41)\Gamma^*(342). \quad (23)$$

Until now, we get the well-known *Hedin's equations* [1, 4] for a system of interacting electrons:

$$G(12) = G_0(12) + \int d(34) G_0(13)\Sigma(34)G(42), \quad (24)$$

$$\Gamma^*(123) = \delta(13)\delta(23) + \int d(4567) \frac{\delta M(12)}{\delta G(45)} G(46)G(75)\Gamma^*(673), \quad (25)$$

$$\chi^*(12) = -i\hbar \int d(34) G(13)G(41)\Gamma^*(342), \quad (26)$$

$$W(12) = v(12) + \int d(34) v(13)\chi^*(34)W(42), \quad (27)$$

$$M(12) = i\hbar \int d(34) G(13)W(41)\Gamma^*(324) = \Sigma(12) - \Sigma_H(12). \quad (28)$$

This set of equations build up a large self-consistent loop involving the single-particle Green's function G , irreducible vertex function Γ^* , irreducible polarizability χ^* , screened Coulomb interaction W and mass operator M .

3 The GW method and random-phase approximation

The most important approximation of the GW method is to take the zeroth order vertex (i.e., no vertex corrections) in Eq. (25),

$$\Gamma^*(123) = \delta(13)\delta(23), \quad (29)$$

which leads to the irreducible polarizability within the *random-phase approximation* (RPA),

$$\chi^*(12) = -i\hbar G(12)G(21), \quad (30)$$

and the iconic GW mass operator,

$$M(12) = i\hbar G(12)W(21). \quad (31)$$

The number of self-consistent Hedin's equations are now reduced to four. Underneath the simple look of the GW method, it is actually one of the conserving approximations, as discussed by Kadanoff and Baym [11, 12], which means it is expected to satisfy the general (number, momentum, and energy) conservation laws.

As discussed before, the band gap problem of DFT was successfully resolved by the development of the first-principles GW method for quasiparticle excitations by Hybertsen and Louie [13, 14]. In real materials, W is usually much weaker than the bare Coulomb interaction, which leads to the laudable accuracy and versatility

of the GW method. Even though the GW method is much simpler than the original Hedin's equations, we still need further approximations for a realistic calculation. In practice, DFT results are usually used as a starting point for the GW method by replacing the many-body wavefunctions by a Slater determinant of Kohn-Sham eigenstates. This allows us to evaluate the GW self-energy (mass operator) as a first-order perturbation with respect to the Kohn-Sham eigenvalues,

$$\epsilon_{n\mathbf{k}}^{\text{QP}} = \epsilon_{n\mathbf{k}}^{\text{KS}} + \langle n\mathbf{k} | M(\epsilon_{n\mathbf{k}}^{\text{QP}}) - V_{\text{xc}} | n\mathbf{k} \rangle. \quad (32)$$

It has been shown that the overlap between DFT-LDA and quasiparticle wavefunctions is greater than 99.9% in some conventional semiconductors and insulators [13]. Moreover, the full self-consistency is often out of reach for real materials and therefore further approximations have to be adopted, such as the one-shot GW method (also called the G_0W_0 method). In the G_0W_0 method, both G_0 and W_0 are constructed using Kohn-Sham eigenvalues and eigenstates in the quasiparticle approximation. G_0 is given by,

$$G_0(\mathbf{x}_1, \mathbf{x}_2; \omega) = \sum_{n\mathbf{k}} \frac{\phi_{n\mathbf{k}}^{\text{KS}}(\mathbf{x}_1)(\phi_{n\mathbf{k}}^{\text{KS}}(\mathbf{x}_2))^*}{\hbar\omega - \epsilon_{n\mathbf{k}}^{\text{KS}} + i\eta \text{sgn}(\epsilon_{n\mathbf{k}}^{\text{KS}} - \epsilon_{\text{F}})}, \quad \eta \rightarrow 0^+, \quad (33)$$

where ϵ_{F} denotes the Fermi level and sgn is the sign function. After space and time Fourier transforms, W_0 can be constructed from the RPA (denoted by the superscript of 0) irreducible polarizability $\chi^{*,0}$ with Kohn-Sham eigenvalues and eigenstates,

$$\begin{aligned} \chi_{\mathbf{G}_1\mathbf{G}_2}^{*,0}(\mathbf{q}; \omega) = \frac{1}{N_k\Omega} \sum_{c\mathbf{k}} \left\{ \frac{\langle v(\mathbf{k}-\mathbf{q}) | e^{-i(\mathbf{q}+\mathbf{G}_1)\cdot\mathbf{r}_1} | c\mathbf{k} \rangle \langle c\mathbf{k} | e^{i(\mathbf{q}+\mathbf{G}_2)\cdot\mathbf{r}_2} | v(\mathbf{k}-\mathbf{q}) \rangle}{\hbar\omega - (\epsilon_{c\mathbf{k}}^{\text{KS}} - \epsilon_{v(\mathbf{k}-\mathbf{q})}^{\text{KS}}) + i\eta} \right. \\ \left. - \frac{\langle c\mathbf{k} | e^{-i(\mathbf{q}+\mathbf{G}_1)\cdot\mathbf{r}_1} | v(\mathbf{k}+\mathbf{q}) \rangle \langle v(\mathbf{k}+\mathbf{q}) | e^{i(\mathbf{q}+\mathbf{G}_2)\cdot\mathbf{r}_2} | c\mathbf{k} \rangle}{\hbar\omega + (\epsilon_{c\mathbf{k}}^{\text{KS}} - \epsilon_{v(\mathbf{k}+\mathbf{q})}^{\text{KS}}) - i\eta} \right\}, \quad \eta \rightarrow 0^+, \end{aligned} \quad (34)$$

where N_k is the number of k -points in the Brillouin zone and it is also equal to the number of unit cells used in the Born-von Kármán boundary condition [9], and Ω is the volume of a unit cell. Equation (34) is the famous Adler-Wiser expression of the RPA polarizability [15, 16].

When the system has time-reversal symmetry, we can prove that Eq. (34) can be further reduced to,

$$\begin{aligned} \chi_{\mathbf{G}_1\mathbf{G}_2}^{*,0}(\mathbf{q}; \omega) = \frac{1}{N_k\Omega} \sum_{c\mathbf{k}} \langle v(\mathbf{k}-\mathbf{q}) | e^{-i(\mathbf{q}+\mathbf{G}_1)\cdot\mathbf{r}_1} | c\mathbf{k} \rangle \langle c\mathbf{k} | e^{i(\mathbf{q}+\mathbf{G}_2)\cdot\mathbf{r}_2} | v(\mathbf{k}-\mathbf{q}) \rangle \left\{ \frac{1}{\hbar\omega - (\epsilon_{c\mathbf{k}}^{\text{KS}} - \epsilon_{v(\mathbf{k}-\mathbf{q})}^{\text{KS}}) + i\eta} \right. \\ \left. - \frac{1}{\hbar\omega + (\epsilon_{c\mathbf{k}}^{\text{KS}} - \epsilon_{v(\mathbf{k}+\mathbf{q})}^{\text{KS}}) - i\eta} \right\}, \quad \eta \rightarrow 0^+, \end{aligned} \quad (35)$$

or,

$$\begin{aligned} \chi_{\mathbf{G}_1\mathbf{G}_2}^{*,0}(\mathbf{q}; \omega) = \frac{1}{N_k\Omega} \sum_{c\mathbf{k}} \langle c\mathbf{k} | e^{-i(\mathbf{q}+\mathbf{G}_1)\cdot\mathbf{r}_1} | v(\mathbf{k}+\mathbf{q}) \rangle \langle v(\mathbf{k}+\mathbf{q}) | e^{i(\mathbf{q}+\mathbf{G}_2)\cdot\mathbf{r}_2} | c\mathbf{k} \rangle \left\{ \frac{1}{\hbar\omega - (\epsilon_{c\mathbf{k}}^{\text{KS}} - \epsilon_{v(\mathbf{k}-\mathbf{q})}^{\text{KS}}) + i\eta} \right. \\ \left. - \frac{1}{\hbar\omega + (\epsilon_{c\mathbf{k}}^{\text{KS}} - \epsilon_{v(\mathbf{k}+\mathbf{q})}^{\text{KS}}) - i\eta} \right\}, \quad \eta \rightarrow 0^+. \end{aligned} \quad (36)$$

In DielectricKit, we can either use Eq. (35) with keyword `k_minus_q` or Eq. (36) with keyword `k_plus_q`.

4 FFT and matrix elements

To calculate the polarizability at a certain q -point, we need to first calculate the matrix element for every \mathbf{k} -points in the reduced Brillouin zone with respect to the symmetry of q -point.

$$\begin{aligned}
\langle n^v(\mathbf{k} + \mathbf{q}) | e^{i(\mathbf{q} + \mathbf{G}) \cdot \mathbf{x}} | n^c \mathbf{k} \rangle &= \sum_{\mathbf{G}_1 \mathbf{G}_2} \frac{1}{V} \int_V d\mathbf{x} e^{-i(\mathbf{k} + \mathbf{q} + \mathbf{G}_1) \cdot \mathbf{x}} C_{n^v(\mathbf{k} + \mathbf{q}) \mathbf{G}_1}^* e^{i(\mathbf{q} + \mathbf{G}) \cdot \mathbf{x}} e^{i(\mathbf{k} + \mathbf{G}_2) \cdot \mathbf{x}} C_{n^c \mathbf{k} \mathbf{G}_2} \\
&= \sum_{\mathbf{G}_1 \mathbf{G}_2} \frac{1}{V} \int_V d\mathbf{x} e^{i(-\mathbf{G}_1 + \mathbf{G} + \mathbf{G}_2) \cdot \mathbf{x}} C_{n^v(\mathbf{k} + \mathbf{q}) \mathbf{G}_1}^* C_{n^c \mathbf{k} \mathbf{G}_2} \\
&= \sum_{\mathbf{G}_1 \mathbf{G}_2} \left(\frac{1}{V} \sum_{\mathbf{R}_l} e^{i(-\mathbf{G}_1 + \mathbf{G} + \mathbf{G}_2) \cdot \mathbf{R}_l} \right) \int_{\Omega} d\boldsymbol{\xi} e^{i(-\mathbf{G}_1 + \mathbf{G} + \mathbf{G}_2) \cdot \boldsymbol{\xi}} C_{n^v(\mathbf{k} + \mathbf{q}) \mathbf{G}_1}^* C_{n^c \mathbf{k} \mathbf{G}_2} \\
&= \sum_{\mathbf{G}_1 \mathbf{G}_2} \frac{1}{\Omega} \Delta \boldsymbol{\xi} \sum_{\boldsymbol{\xi}_m} e^{i(-\mathbf{G}_1 + \mathbf{G} + \mathbf{G}_2) \cdot \boldsymbol{\xi}_m} C_{n^v(\mathbf{k} + \mathbf{q}) \mathbf{G}_1}^* C_{n^c \mathbf{k} \mathbf{G}_2} \\
&= \frac{1}{N_{FFT}} \sum_{\boldsymbol{\xi}_m} e^{i\mathbf{G} \cdot \boldsymbol{\xi}_m} \left(\sum_{\mathbf{G}_1} e^{-i\mathbf{G}_1 \cdot \boldsymbol{\xi}_m} C_{n^v(\mathbf{k} + \mathbf{q}) \mathbf{G}_1}^* \right) \left(\sum_{\mathbf{G}_2} e^{i\mathbf{G}_2 \cdot \boldsymbol{\xi}_m} C_{n^c \mathbf{k} \mathbf{G}_2} \right) \\
&= \frac{1}{N_{FFT}} \text{FFT}[\text{CONJ}[\text{FFT}[C_{n^v(\mathbf{k} + \mathbf{q})}(\{\mathbf{G}_1\})]] \cdot * \text{FFT}[C_{n^c \mathbf{k}}(\{\mathbf{G}_2\})]](\mathbf{G}),
\end{aligned} \tag{37}$$

where **CONJ** is the complex conjugate function in FORTRAN and the **FFT** all refers to **backward** FFT,

$$Y_k = \sum_{j=0}^{n-1} X_j e^{2\pi j k \sqrt{-1}/n}. \tag{38}$$

In the derivation, we have used the following identity,

$$\frac{1}{V} \sum_{\mathbf{R}_l} e^{i(-\mathbf{G}_1 + \mathbf{G} + \mathbf{G}_2) \cdot \mathbf{R}_l} = \frac{N}{V} = \frac{1}{\Omega}. \tag{39}$$

5 Symmetry of ε^{-1} and χ^*

Apply crystal space group symmetry on $W(\mathbf{x}_1, \mathbf{x}_2; \omega)$, and we will get

$$\begin{aligned}
W(\mathbf{x}_1, \mathbf{x}_2; \omega) &= \int d\mathbf{x}_3 \varepsilon^{-1}(\mathbf{x}_1, \mathbf{x}_3; \omega) V(\mathbf{x}_3, \mathbf{x}_2; \omega) \\
\Rightarrow W(\{\alpha|\mathbf{t}\}\mathbf{x}_1, \{\alpha|\mathbf{t}\}\mathbf{x}_2; \omega) &= \int d\mathbf{x}_3 \varepsilon^{-1}(\{\alpha|\mathbf{t}\}\mathbf{x}_1, \mathbf{x}_3; \omega) V(\mathbf{x}_3, \{\alpha|\mathbf{t}\}\mathbf{x}_2; \omega), \\
&= \int d(\{\alpha|\mathbf{t}\}\mathbf{x}_3) \varepsilon^{-1}(\{\alpha|\mathbf{t}\}\mathbf{x}_1, \{\alpha|\mathbf{t}\}\mathbf{x}_3; \omega) V(\{\alpha|\mathbf{t}\}\mathbf{x}_3, \{\alpha|\mathbf{t}\}\mathbf{x}_2; \omega) \equiv W(\mathbf{x}_1, \mathbf{x}_2),
\end{aligned} \tag{40}$$

which leads to the identity,

$$\varepsilon^{-1}(\{\alpha|\mathbf{t}\}\mathbf{x}_1, \{\alpha|\mathbf{t}\}\mathbf{x}_2; \omega) = \varepsilon^{-1}(\mathbf{x}_1, \mathbf{x}_2; \omega). \tag{41}$$

With Eqn. (41), we can prove that, if

$$\mathbf{q}' = \alpha \mathbf{q} + \mathbf{G}_0, \tag{42}$$

then,

$$\begin{aligned}
\varepsilon_{\mathbf{G}_1\mathbf{G}_2}^{-1}(\mathbf{q}'; \omega) &= \frac{1}{N\Omega} \int d\mathbf{x}_1 d\mathbf{x}_2 e^{-i(\mathbf{q}'+\mathbf{G}_1)\cdot\mathbf{x}_1} e^{i(\mathbf{q}+\mathbf{G}_2)\cdot\mathbf{x}_2} \varepsilon^{-1}(\mathbf{x}_1, \mathbf{x}_2; \omega) \\
&= \frac{1}{N\Omega} \int d\mathbf{x}_1 d\mathbf{x}_2 e^{-i(\alpha\mathbf{q}+\mathbf{G}_0+\mathbf{G}_1)\cdot\mathbf{x}_1} e^{i(\alpha\mathbf{q}+\mathbf{G}_0+\mathbf{G}_2)\cdot\mathbf{x}_2} \varepsilon^{-1}(\mathbf{x}_1, \mathbf{x}_2; \omega) \\
&= \frac{1}{N\Omega} \int d\mathbf{x}_1 d\mathbf{x}_2 e^{-i[\alpha(\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_1))]\cdot\mathbf{x}_1} e^{i[\alpha(\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_2))]\cdot\mathbf{x}_2} \varepsilon^{-1}(\mathbf{x}_1, \mathbf{x}_2; \omega) \\
&= \frac{1}{N\Omega} \int d\mathbf{x}_1 d\mathbf{x}_2 e^{-i[\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_1)]\cdot(\alpha^{-1}\mathbf{x}_1)} e^{i[\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_2)]\cdot(\alpha^{-1}\mathbf{x}_2)} \varepsilon^{-1}(\mathbf{x}_1, \mathbf{x}_2; \omega), \quad \mathbf{y} = \alpha^{-1}\mathbf{x} - \alpha^{-1}\mathbf{t} \\
&= \frac{1}{N\Omega} \int d(\alpha\mathbf{y}_1 + \mathbf{t}) d(\alpha\mathbf{y}_2 + \mathbf{t}) e^{-i[\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_1)]\cdot(\mathbf{y}_1+\alpha^{-1}\mathbf{t})} e^{i[\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_2)]\cdot(\mathbf{y}_2+\alpha^{-1}\mathbf{t})} \varepsilon^{-1}(\alpha\mathbf{y}_1 + \mathbf{t}, \alpha\mathbf{y}_2 + \mathbf{t}; \omega) \\
&= e^{-i(\mathbf{G}_1-\mathbf{G}_2)\cdot\mathbf{t}} \frac{1}{N\Omega} \int d\mathbf{y}_1 d\mathbf{y}_2 e^{-i[\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_1)]\cdot\mathbf{y}_1} e^{i[\mathbf{q}+\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_2)]\cdot\mathbf{y}_2} \varepsilon^{-1}(\mathbf{y}_1, \mathbf{y}_2; \omega) \\
&= e^{-i(\mathbf{G}_1-\mathbf{G}_2)\cdot\mathbf{t}} \varepsilon_{\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_1)\alpha^{-1}(\mathbf{G}_0+\mathbf{G}_2)}^{-1}(\mathbf{q}; \omega).
\end{aligned} \tag{43}$$

This identity also applies to $\chi_{\mathbf{G}_1\mathbf{G}_2}^*(\mathbf{q}; \omega)$. In this way, we only need to calculate ε^{-1} or χ^* at symmetry reduced k -points and unfold it to recover the value at symmetry-equivalent k -points. This unfolding scheme has been implemented in `RealSpace.x`.

6 Real-space ε^{-1} and χ^*

There are two requirements for the real-space inverse dielectric function $\varepsilon^{-1}(\mathbf{r}_1, \mathbf{r}_2; \omega)$.

$$\begin{aligned}
\varepsilon^{-1}(\mathbf{r}_1 + \mathbf{R}, \mathbf{r}_2 + \mathbf{R}; \omega) &= \varepsilon^{-1}(\mathbf{r}_1, \mathbf{r}_2; \omega), \\
\varepsilon^{-1}(\mathbf{r}_1 + \mathbf{N}_{\text{BvO}}, \mathbf{r}_2; \omega) &= \varepsilon^{-1}(\mathbf{r}_1, \mathbf{r}_2; \omega).
\end{aligned} \tag{44}$$

where $\mathbf{R} = m\mathbf{a}_1 + n\mathbf{a}_2 + l\mathbf{a}_3$ is a lattice vector and $\mathbf{N}_{\text{BvO}} = N_{\text{BvO}}^i \mathbf{a}_i, \forall i = x, y, z$ is the period of the large BvO supercell. So it is sufficient to define a real-space function $\varepsilon_{\xi_1\xi_2}^{-1}(\mathbf{R}; \omega) \equiv \varepsilon^{-1}(\xi_1 + \mathbf{R}, \xi_2; \omega)$, where ξ_1 and ξ_2 are real-space FFT grid within a primitive cell. The number of ξ_1 (ξ_2) is equal to the number of G -vectors used in doing the Fourier transform. In this section, we use the information $\varepsilon_{\mathbf{G}_1\mathbf{G}_2}^{-1}(\mathbf{q}; \omega)$ to calculate $\varepsilon_{\xi_1\xi_2}^{-1}(\mathbf{R}; \omega)$ with fixed ξ_2 within a primitive cell, while $\xi_1 + \mathbf{R}$ is defined in the BvO cell.

Note that the real-space and reciprocal-space lattice vectors are different. In reciprocal space, \mathbf{q} is defined in the full Brillouin zone, and \mathbf{G}_1 and \mathbf{G}_2 are defined in a sphere around Γ point with a kinetic energy cutoff. We order the \mathbf{G} -vectors using the kinetic energy $\propto |\mathbf{G} + \mathbf{q}|^2$. In real space, however, we define ξ_1, ξ_2 and \mathbf{R} using a $[0, 1]$ -like convention. For example, we have FFT grid in the primitive cell $N_{\text{FFT}} = [n_1, n_2, n_3]$. Then the fractional coordinates of ξ_1 (ξ_2) in the basis of \mathbf{a}_i are in the range of $[0, 0, 0] \rightarrow [\frac{n_1-1}{n_1}, \frac{n_2-1}{n_2}, \frac{n_3-1}{n_3}]$. We order $\xi_1 = [\frac{\mathbf{i}}{n_1}, \frac{\mathbf{j}}{n_2}, \frac{\mathbf{k}}{n_3}]$ using the following function, $\text{INDEX_xi}(\xi) = \mathbf{i} + \mathbf{j} * n_1 + \mathbf{k} * n_1 * n_2 + 1$ where $\mathbf{i} = 0, 1, \dots, n_1$. As for $\mathbf{R} = [\mathbf{R}_1, \mathbf{R}_2, \mathbf{R}_3]$, it is defined with the same fractional coordinates in the range of $[0, 0, 0] \rightarrow [N_1^{\text{BvO}}, N_2^{\text{BvO}}, N_3^{\text{BvO}}]$. A corresponding indexing function is defined as, $\text{INDEX_R}(\mathbf{R}) = \mathbf{R}_1 + \mathbf{R}_2 * N_1^{\text{BvO}} + \mathbf{R}_3 * N_1^{\text{BvO}} * N_2^{\text{BvO}} + 1$.

$$\begin{aligned}
\varepsilon_{\xi_1 \xi_2}^{-1}(\mathbf{R}; \omega) &\equiv \varepsilon^{-1}(\xi_1 + \mathbf{R}, \xi_2; \omega) = \frac{1}{N\Omega} \sum_{\mathbf{q} \mathbf{G}_1 \mathbf{G}_2} e^{i(\mathbf{q} + \mathbf{G}_1) \cdot (\xi_1 + \mathbf{R})} e^{-i(\mathbf{q} + \mathbf{G}_2) \cdot \xi_2} \varepsilon_{\mathbf{G}_1 \mathbf{G}_2}^{-1}(\mathbf{q}; \omega) \\
&= \frac{1}{N\Omega} \sum_{\mathbf{q}} e^{i\mathbf{q} \cdot (\xi_1 + \mathbf{R})} \left[\sum_{\mathbf{G}_1} e^{i\mathbf{G}_1 \cdot \xi_1} \left(\sum_{\mathbf{G}_2} \varepsilon_{\mathbf{G}_1 \mathbf{G}_2}^{-1}(\mathbf{q}; \omega) e^{-i(\mathbf{q} + \mathbf{G}_2) \cdot \xi_2} \right) \right] \\
&= \frac{1}{N\Omega} \sum_{\mathbf{q}} e^{i\mathbf{q} \cdot (\xi_1 + \mathbf{R})} \text{ucfft}(\xi_1; \xi_2, \mathbf{q}, \omega)
\end{aligned} \tag{45}$$

Now suppose we take the simplest form of $\varepsilon_{\mathbf{G}_1 \mathbf{G}_2}^{-1}(\mathbf{q}; \omega)$, corresponding to no screening at all,

$$\varepsilon_{\mathbf{G}_1 \mathbf{G}_2}^{-1}(\mathbf{q}; \omega) = \delta_{\mathbf{G}_1 \mathbf{G}_2} \tag{46}$$

And then the real-space $\varepsilon^{-1}(\mathbf{r}_1, \mathbf{r}_2; \omega)$ is given by,

$$\begin{aligned}
\varepsilon^{-1}(\mathbf{r}_1, \mathbf{r}_2; \omega) &= \frac{1}{N\Omega} \sum_{\mathbf{q} \mathbf{G}_1 \mathbf{G}_2} e^{i(\mathbf{q} + \mathbf{G}_1) \cdot \mathbf{r}_1} e^{-i(\mathbf{q} + \mathbf{G}_2) \cdot \mathbf{r}_2} \varepsilon_{\mathbf{G}_1 \mathbf{G}_2}^{-1}(\mathbf{q}; \omega) \\
&= \frac{1}{N\Omega} \sum_{\mathbf{q} \mathbf{G}_1} e^{i(\mathbf{q} + \mathbf{G}_1) \cdot (\mathbf{r}_1 - \mathbf{r}_2)} \\
&= \frac{1}{N\Omega} \frac{1}{\Delta \mathbf{k}} \sum_{\mathbf{k}} \Delta \mathbf{k} e^{i\mathbf{k} \cdot (\mathbf{r}_1 - \mathbf{r}_2)}, \quad \Delta \mathbf{k} = \frac{(2\pi)^3}{N\Omega} \\
&= \frac{1}{(2\pi)^3} \int d\mathbf{k} e^{i\mathbf{k} \cdot (\mathbf{r}_1 - \mathbf{r}_2)} = \delta(\mathbf{r}_1 - \mathbf{r}_2)
\end{aligned} \tag{47}$$

But wait, what units should we use for Ω ? It actually depends on what unit we use for $(\mathbf{r}_1 - \mathbf{r}_2)$. Since we use angstrom as the length unit in `RealSpace.x`, we will also use \AA^3 for Ω here. In practice, we will partition \mathbf{q} 's in FBZ among different processors. We then fix ξ_2 and ω . For each FBZ \mathbf{q} we will determine the corresponding RBZ \mathbf{q} . $\varepsilon_{\mathbf{G}_1 \mathbf{G}_2}^{-1}(\mathbf{q}; \omega)$ for this RBZ \mathbf{q} will be read and unfold.

7 Parallelism and scaling performance

7.1 Chi.x

In `Chi.x`, we parallel over k -points and calculate each q -point sequentially. We need wavefunction files `WFn` ($|\mathbf{k}\rangle$ for both valence and conduction bands), `WFnmq` ($|\mathbf{k} - \mathbf{q}\rangle$ for valence bands) or `WFnq` ($|\mathbf{k} + \mathbf{q}\rangle$ for valence bands) on a reduced BZ. That is, in Quantum ESPRESSO calculation, we can use crystal symmetry to reduced the number of k points in the wavefunction files. We will use symmetry operation to unfold the wavefunctions from reduced BZ to full BZ.

To benchmark the scaling performance of `Chi.x`, we consider the example of bulk Si. We use a $16 \times 16 \times 16$ k -grid with 145 k -points in the reduced BZ. Interband transitions between 4 valence bands and 196 conduction bands are used to construct the polarizability function matrix. Kinetic energy cutoff of the polarizability function is set to 20 Ry. We tested the wall time on NERSC Cori Haswell nodes (32 CPUs per node). We use 1 OpenMP thread in the following.

| NNode | Wall times (s) | Speedup | Wall time w/o IO | Speedup w/o IO |
|-------|----------------|---------|------------------|----------------|
| 1 | 4083.40 | 1.00 | 4029.73 | 1.00 |
| 2 | 2051.02 | 1.99 | 2018.97 | 2.00 |
| 4 | 1145.31 | 3.57 | 1061.70 | 3.80 |
| 8 | 560.50 | 7.29 | 514.62 | 7.83 |
| 16 | 324.02 | 12.60 | 258.05 | 15.62 |
| 32 | 180.05 | 22.68 | 134.94 | 29.86 |
| 64 | 100.13 | 40.78 | 73.25 | 55.01 |

Table 1: Scaling performance of `Chi.x`

7.2 EpsInv.x

In `EpsInv.x`, we parallel over \mathbf{G}_1 and \mathbf{G}_2 vectors in $\chi_{\mathbf{G}_1\mathbf{G}_2}^*(\mathbf{q};\omega)$ and calculate each q -point sequentially. We use SCALAPACK to invert the dielectric function $\varepsilon_{\mathbf{G}_1\mathbf{G}_2}(\mathbf{q};\omega) = \delta_{\mathbf{G}_1\mathbf{G}_2} - v_{\mathbf{G}_1}(\mathbf{q})\chi_{\mathbf{G}_1\mathbf{G}_2}^*(\mathbf{q};\omega)$. The matrix inversion step is fairly quick for the benchmark case of bulk Si, and the wall time is dominated by the output of HDF5 files.

7.3 RealSpace.x

In `RealSpace.x`, we parallel over \mathbf{q} -points in full BZ and calculate each ξ_2 sequentially.

References

- [1] Hedin, L. New method for calculating the one-particle Green's function with application to the electron-gas problem. *Phys. Rev.* **139**, A796–A823 (1965).
- [2] Schwinger, J. On the Green's functions of quantized fields. I. *Proc. Natl. Acad. Sci. U.S.A* **37**, 452–455 (1951).
- [3] Baym, G. Field-theoretic approach to the properties of the solid state. *Ann. Phys.* **14**, 1–42 (1961).
- [4] Hedin, L. & Lundqvist, S. Effects of electron-electron and electron-phonon interactions on the one-electron states of solids. In *Solid State Physics*, vol. 23, 1–181 (Elsevier, 1970).
- [5] Fetter, A. L. & Walecka, J. D. *Quantum Theory of Many-Particle Systems* (Courier Corporation, 2012).
- [6] Mahan, G. D. *Many-Particle Physics* (Springer Science & Business Media, 2013).
- [7] Nozières, P. *Theory of Interacting Fermi Systems* (Westview Press, 1997).
- [8] Stefanucci, G. & van Leeuwen, R. *Nonequilibrium Many-Body Theory of Quantum Systems: A Modern Introduction* (Cambridge University Press, 2013).
- [9] Cohen, M. L. & Louie, S. G. *Fundamentals of Condensed Matter Physics* (Cambridge University Press, 2016).
- [10] Strinati, G. Application of the Green's functions method to the study of the optical properties of semiconductors. *Riv. Nuovo Cimento* **11**, 1–86 (1988).
- [11] Baym, G. & Kadanoff, L. P. Conservation laws and correlation functions. *Phys. Rev.* **124**, 287–299 (1961).

- [12] Baym, G. Self-consistent approximations in many-body systems. *Phys. Rev.* **127**, 1391–1401 (1962).
- [13] Hybertsen, M. S. & Louie, S. G. Electron correlation in semiconductors and insulators: Band gaps and quasiparticle energies. *Phys. Rev. B* **34**, 5390–5413 (1986).
- [14] Hybertsen, M. S. & Louie, S. G. First-principles theory of quasiparticles: Calculation of band gaps in semiconductors and insulators. *Phys. Rev. Lett.* **55**, 1418–1421 (1985).
- [15] Adler, S. L. Quantum theory of the dielectric constant in real solids. *Phys. Rev.* **126**, 413–420 (1962).
- [16] Wiser, N. Dielectric constant with local field effects included. *Phys. Rev.* **129**, 62–69 (1963).