

θ = e_q(−βu²), max 1
β fixed at ½ (the support shrinks as q → −∞)
q = −100 (→ −∞: the box)
q = −1 (semicircle)
q = 0 (parabola)
q = 1 (Gaussian)
θ = e_q(−βu²), max 1
β(q) = 1/(2(2−q)): escort variance held (dotted: closed forms)
q = −100 (→ −∞: the box)
q = −1 (semicircle)
q = 0 (parabola)
q = 1 (Gaussian)
closed forms
q: q → −∞ … −1
q: −1 … 0
ψ = ∂θ (q-g1), slope −1 at 0
β fixed at ½ (the support shrinks as q → −∞)
q = −100 (→ −∞: the box)
q = −1 (semicircle)
q = 0 (parabola)
q = 1 (Gaussian)
ψ = ∂θ (q-g1), slope −1 at 0
β(q) = 1/(2(2−q)): escort variance held (dotted: closed forms)
q = −100 (→ −∞: the box)
q = −1 (semicircle)
q = 0 (parabola)
q = 1 (Gaussian)
closed forms
q: 0 … 1
−Δθ (q-Mexican hat), value 1 at 0; q > ½
β fixed at ½ (the support shrinks as q → −∞)
q = 1 (Gaussian)
u = x/σ (σ of the fixed-β Gaussian)
−Δθ (q-Mexican hat), value 1 at 0; q > ½
β(q) = 1/(2(2−q)): escort variance held (dotted: closed forms)
q = 1 (Gaussian)
closed forms
u = x/σ (σ of the fixed-β Gaussian)
q: 1 … 2