



typically 150 mV, an example of which is the trans-epithelial potential (TEP) found in skin, lens, and cornea.<sup>14,20</sup> The TEP mechanism in vertebrate lens is particularly interesting, as it creates a complex pattern of current loops with magnitudes around 200 mA/cm<sup>2</sup>. These currents flow inward at the anterior and posterior poles of the lens, and outward at the equator of the lens, controlling the migration and differentiation of epithelial cells.<sup>7,22</sup> Another important function of TEP can be observed in injuries of the epithelium. A wound short circuits the potential difference between the two sides of the epithelium and, therefore, gives rise to an EF (0.04–0.2 V/mm)<sup>8,9,14,16,20,24</sup> that, in turn, controls the orientation and frequency of cell division<sup>24</sup> and induces directional migration toward the injury.<sup>8,16,21</sup> This has been shown to override any other influence on the cell functioning, such as chemical gradients and population pressure.<sup>14,21</sup> In regeneration and embryonic development, intra- and extracellular EFs play a pivotal role in regulating cellular behavior<sup>14</sup> and the development of spatial patterns, such as left-right organ asymmetry.<sup>12,14,27</sup> Disruption of these fields in embryos has been reported to result in serious defects, for example, the absence of the cranium, a malformed head and the loss of eyes.<sup>14</sup> In studies conducted on partially amputated *Xenopus* tails, electrical currents were observed to occur as a result of the injury. Disrupting these electrical currents reduced regeneration, while artificially reversing its polarity completely blocked the healing process.<sup>29</sup>

The significance of bioelectricity is emphasized by its role in one of the rare examples of human regeneration. Illingworth and Barker<sup>30</sup> reported that the amputated fingertips of children could be fully regenerated as long as the stump is kept clean and hydrated and observed that electrical currents of 30 A/cm<sup>2</sup> were present in wounds. In their article, they theorized that moisture in the wound ensures a continuous electric conductance path, enabling the wound electrical fields to exert their regenerating effect.<sup>30</sup> Streaming potentials, streaming currents, and the piezoelectricity of collagen molecules all contribute to the generation of bioelectricity in bone.<sup>11,31</sup> These electrical phenomena are suggested to be transduced by osteocytes and to play a similar role in remodeling and healing as TEPs.<sup>13,37</sup> Bioelectricity is also suggested to be the ideal mechanism to deliver signals to chondrocyte cells in cartilage, as they usually exist in relative isolation.<sup>40</sup> Other examples of *in vivo* electricity include the rapid action potentials of nerves and muscles; long-lasting DC voltages around damaged nerves, known as injury potentials<sup>14</sup> and the trans-endothelial extracellular potential gradients in arteries and veins.<sup>25,41</sup>

The use of electrical stimulation is well-established in today's medicine. A wide range of electrical therapeutic systems have been developed to treat specific ailments and/or tissue types. These devices include implantable and external bone growth stimulators,<sup>42,43</sup> chronic cutaneous wound-healing systems,<sup>44</sup> functional electrical stimulation devices, to restore muscles in paralyzed patients,<sup>45</sup> and stimulators for pain relief.<sup>46,47</sup>

Transcutaneous nerve stimulation (TENS) units are mainly used to deliver electroanalgesia and have been proven to be greatly effective in the treatment of a wide range of pains.<sup>46,47</sup> In a clinical study examining the effectiveness of these devices, it was shown that TENS treatment reduced

pain by more than a half in 47% of the patients.<sup>48</sup> Additionally, these devices have also been shown to be useful in the treatment of chronic ulcers.<sup>44</sup> In a meta-analysis, where the effect of multiple systems (including TENS units) was assayed upon chronic wound healing, it was found that electrical stimulation produced an enhanced healing rate of 22% per week compared to the 9% of untreated samples. Upon the comparison of the healing effect of the devices involved in this study, no significant difference was observed.<sup>44</sup> In another investigation, examining the effectiveness of electrical therapy in the treatment on pressure ulcers, the wound surface decreased approximately 70% with electrical treatment, while only 36% without.<sup>49</sup> Electrical stimulation systems also have a proven record promoting bone growth. For example, an extensive study, including 175 patients, showed that electrical stimulation was able to induce solid bone union in 83.7% of the cases.<sup>42</sup>

Due to the wide variety of bioelectrical presence and its significant influence on *in vivo* tissue, researchers are now beginning to explore the potential of using this stimulus *in vitro*.

## The Methods of In Vitro Electrical Stimulation

### Types of electrical stimulation

The most basic method of electric stimulation is an applied DC voltage, simply generated by batteries.<sup>50</sup> More complicated stimuli can be in the form of monophasic (DC) or biphasic (AC)<sup>13,32</sup> sinusoidal,<sup>11,39,51,52</sup> saw tooth,<sup>36,53,54</sup> or square wave<sup>28,55</sup> signals, injected in pulses,<sup>36,53,55</sup> pulse bursts,<sup>11,56</sup> or continuously.<sup>5,10,13</sup> These signals can be generated by stimulator chips,<sup>32</sup> signal generators,<sup>28,54,57</sup> or dedicated therapeutic systems,<sup>15,58</sup> such as the Phyback,<sup>59</sup> TENS,<sup>46,47</sup> or the EBI bone-healing systems.<sup>11,56,57</sup>

### Methods of delivering the stimulus

Direct coupling. With direct coupling (Fig. 1A), the electrode is in direct contact with (e.g., inserted into) the cell culture or implanted into the patient or laboratory

FIG. 1. The four main techniques of delivering electrical stimulation: The direct (A) and the indirect capacitive (B), inductive (C) and combined (D) methods. Color images available online at [www.liebertpub.com/teb](http://www.liebertpub.com/teb)

animal.<sup>31,32,37</sup> Although this is the simplest way of delivering electrical stimulation, there are quite a few disadvantages to the approach. The problem of toxicity arises, due to the insufficient biocompatibility of the electrodes,<sup>60</sup> changes in pH,<sup>31</sup> reduced levels of molecular oxygen,<sup>60,61</sup> and the generation of dangerous Faradic byproducts, for example, reactive oxygen species in the culture medium.<sup>13,31,60,61</sup> The facts that the stimuli will be biased, due to the formation of a capacitive bilayer<sup>31,32,61</sup> and that the effects of the stimulation was observed to depend heavily on whether it was measured near the anode or the cathode electrode,<sup>61</sup> further hinder the use of this method.

**Indirect coupling.** To avoid the difficulties inherent in direct coupling, an indirect, noninvasive approach is used in many therapeutic devices and in vitro experimental setups. The three main techniques used are capacitive, inductive, and combined coupling methods (Fig. 1B), where the latter is a combination of a static magnetic and an alternating current generated by a transient electromagnetic field.<sup>11,13,32</sup> A comparison of the effects of the different techniques can be seen in Table 1.

**Capacitive coupling:** In capacitive coupling (CC), a homogenous electromagnetic field is created between two parallel layers of metal (e.g., stainless steel, gold) or carbon (i.e., a plate capacitor) that are placed above and below, the culture medium with a small (0.5 mm) gap between them.<sup>3,13,19,31,32,36,37,40,62,65</sup> CC has the advantage of avoiding the issues with direct coupling and that of generating a homogenous EF, thus equal amount of stimulation for every cell regardless of the position in the culture vessel. However, its usefulness can be limited by the high voltage (e.g., 100 V) required to be generated between the electrodes.<sup>66</sup>

**Inductive coupling:** Inductive coupling utilizes controlled electromagnetic fields generated by coils placed around the cell culture. This has the advantage of generating small-magnitude currents and potentials in the proximity of the targeted cells, rather than delivering the stimulation through electrodes.<sup>25,31,37,51,67,68</sup> In many instances, the coils are used in pairs, placed in the Helmholtz configuration,<sup>39,53,57</sup> meaning that distance between the coils is equal to their radius.<sup>39</sup> This generates a near homogenous magnetic field with uniform electromagnetic field properties across the cell culture. A subtype of this modality is known as pulsed electromagnetic field stimulation or pulsed electromagnetic field stimulation (PEMF), where the stimulus is delivered in pulses (rather than being static or continuously harmonic) to mimic, for example, the natural strain-generated potentials observed in bone.<sup>39,69</sup>

**Scaffolds.** In recent years, electrically conductive polymer scaffolds have been developed as a new means to deliver the electrical stimulus with the most popular materials being polypyrrole (PPy), used either alone<sup>5,18</sup> or in combination with other materials, such as the biodegradable poly-L-lactide (PLA) in a 5:95 ratio.<sup>4</sup> In the future, these multifunctional scaffolds could act as bioactive substrates for cell attachment, while providing a way to better regulate cellular activities through evenly distributed, well-controlled electrical stimuli both on the surface and within the biomaterial scaffold.<sup>4</sup>

The range of parameter variables in electrical stimulation

The level and nature of the voltage applied by the stimulation equipment is, arguably, the most important parameter influencing the cell cultures. The outcome, especially with regard to cellular migration, orientation,<sup>16,27,41,70</sup> and gene expression,<sup>54,71</sup> has been shown to depend upon the amplitude of the voltage applied, with strong indications that a relatively higher setting is favorable in nearly all circumstances.<sup>9,16,27,41,54,70,72</sup> See Table 2. However, effects, such as enhanced proliferation or increased extracellular matrices (ECM) deposition, were witnessed with stimulation currents (or current density) at both relatively high and low settings (Table 3). As such, the value of current does not seem to be important when considering the effectiveness of the stimuli. Indeed, it acts a limiting constrain, since any current above a certain threshold (e.g., 20 mA in case of treating nonunions<sup>42</sup>) will result in cell death.<sup>10,13,31,32,42,73</sup>

The maximum pulse width is limited by current density: the higher the current of the stimulation is, the shorter the duration needs to be to avoid necrosis (i.e., there is a limit on the electrical energy that can be absorbed by the cell safely).<sup>13,73</sup> Nonetheless, none of the temporal durations, from 0.025 to 380 ms, that have already been applied in cell culture experiments, have proved to be ineffective or damaging.<sup>10,13,16,34,39,55,57,62,67,74</sup> A wide range of frequencies (7.5-60000 Hz) have been applied in previous experiments, though most of these were below 100 Hz. This is a logical choice as frequencies in this low range are the ones that are assumed to occur naturally in vivo during normal use and are, therefore, the most likely to have a beneficial effect on the cells.<sup>10,11,13,31,32,34,66,51,63,75</sup> Frequencies between 10 and 30 Hz,<sup>51</sup> and those less than 100 Hz,<sup>39</sup> have been reported as the most effective for stimulating new bone formation, while 7.5 Hz has been suggested as the best frequency to promote the proliferation of primary rat osteoblasts<sup>75</sup> and osteoclast differentiation of bone marrow cells of mice.<sup>54</sup>

#### The Effects of Electrical Stimulation at the Cellular Level: *Calvano-Transduction*

Several theories exist on how EFs affect cells directly. What is known is that the electrical fields seem to be sensed through the same pathways observed to be involved in mechanotransduction and chemotaxis.<sup>9,21,26,27,76</sup> A key component in this process, and one that has been observed to be greatly affected by electrical stimuli, is the intracellular calcium level.

#### Intracellular calcium

Electric signals are believed to be transduced, at least partially, through the calcium/calmodulin pathway.<sup>32</sup> This happens somewhat differently with the various methods of stimulation (Fig. 2). Direct and capacitive coupled stimuli exert their effect mainly on the cellular membrane, as these cannot overcome its high electrical resistance,<sup>20,25</sup> raising the intracellular  $\text{Ca}^{2+}$  concentration and prostaglandin E2 levels by activating the voltage-gated calcium channels in the cell membrane. In addition, inductively and combined coupled electromagnetic fields are theorized to generate potentials and currents in the cytoplasm, releasing intracellular calcium

Table 1. A Summary of Important Research Articles in the Discipline of Electrical Stimulation for Bone Tissue Engineering

Article	Stimulation	Results
Direct coupling Schmidt et al. (1997) <sup>18</sup>	0.1 V with 10 mA for 2 h using a Polypyrrole (PPy) as anode 1 V/mm at 1 Hz	Stimulation on the PPy an produced significantly larger neurite length than in unstimulated and tissue culture plastic control groups.
Serena et al. (2009) <sup>28</sup>	Human ESC line H13	Spontaneous contractions, expression, and sarcometric organization of troponin T suggest cardiac differentiation
Shi et al. (2008) <sup>4</sup>	Human cutaneous fibroblasts	Cells adhered, spread and proliferated on the conductors
Sun et al. (2006) <sup>6</sup>	Rat bone marrow MSCs	Cytokine production enhanced 10-fold by stimulation Changes in morphology observed with both MSCs and fibroblasts, but only the latter aligned in response to the stimulus.
Radisic et al. (2004) <sup>17</sup>	Rat fibroblasts HT1080 Neonatal rat ventricular myocytes	Stimulation resulted in alignment and coupling, increased amplitude contractions, and enhanced ultrastructural organization.
Capacitive coupling Au et al. (2007) <sup>3</sup>	Neonatal rat cardiomyocytes and NIH3T3 fibroblasts	Stimulation enhanced elongation of both cell types and fibroblast alignment on abraded surfaces. Topographical cues were stronger.
Hartig et al. (2000) <sup>36</sup>	Bovine primary osteoblasts	Significant increase in proliferation in nonconductive related protein secretion in conductive cultures.
Kim et al. (2006) <sup>32</sup>	Rat calvarial osteoblasts	Higher proliferation in continuously stimulated samples. No change in ALP, osteopontin, coll. Type I, BMP-2, -4, IGF-2, and TGF- $\beta$ 1 levels.
Kim et al. (2008) <sup>10</sup>	Human bone marrow MSCs	Proliferation increased by 57%. No increase in osteoblast differentiation. Stimulation induced VEGF expression.
Kim et al. (2009) <sup>13</sup>	Human MSCs	Increased proliferation. ALP activity and calcium deposition enhanced after stimulation. Expression of VEGF and BMP-2.
Zhuang et al. (1997) <sup>63</sup>	MC3T3-E1 clonal osteoblastic cells	Stimulation enhanced proliferation and increased the levels of TGF- $\beta$ 1.
Inductive coupling Bodamyali et al. (1998) <sup>57</sup>	Neonatal rat calvarial osteoblasts	Exposure significantly increased the number and size of deposited bone-like nodules. Enhanced BMP-2 and BMP-4 expression.
Lohmann et al. (2003) <sup>53</sup>	MLO-Y4 osteocyte-like cells ROS 17/2.8 cell line	Increased ALP activity, TGF- $\beta$ 1, and prostaglandin E2 expression, while osteocalcin or cell numbers went unchanged.
McLeod et al. (1993) <sup>51</sup>	Osteosarcoma cell line ROS 17/2.8	Exposure limited the normal increase in cell numbers, while enhanced ALP activity. Effects were cell density dependent.
Molen et al. (2000) <sup>35</sup>	Osteosarcoma cell line ROS 17/2.8	Stimulation inhibited cell growth. ALP activity was dependent on gap junctional coupling. Results suggest differential effects.
Comparison of the modalities Brighton et al. (2001) <sup>11</sup>	MC3T3-E1 clonal osteoblastic cells	All three stimulation types increased DNA content in samples, but only the capacitive one did it significantly and in an ever increasing manner. Signal transduction was thorough $\text{Ca}^{2+}$ influx with capacitive coupling, and by the intracellular release of the same ion with inductive and combined coupling.

Direct stimulation was mainly used to induce the morphology and orientation of cells. Capacitive stimulation seems to be more effective in increasing proliferation, while cells treated with inductive stimuli display greater ECM deposition.  
ESC, embryonic stem cell; MSC, mesenchymal stem cell; ROS, reactive oxygen species; ECM, extracellular matrix; TGF, transforming growth factor; ALP, alkaline phosphatase; VEGF, vascular endothelial growth factor.

Table 2. A Comparison of the Effect of Stimulation Delivered with Different Electric Field Strength

The effect of various field strength settings	
0.00000048 V/mm	Reduced osteoclastogenesis in bone marrow cultures compared to controls and higher amplitude settings <sup>54</sup>
0.0000006 V/mm	Reduced proliferation, but increased ALP activity in osteoblasts <sup>51</sup>
0.000002 V/mm	Inhibited cellular growth of osteoblasts <sup>35</sup>
0.002 V/mm	Enhanced proliferation and TGF-B1 expression in osteoblasts <sup>63</sup>
0.01 V/mm	Migration of osteoblasts above this threshold <sup>27</sup>
0.15 V/mm	Orientation of osteoblasts above this threshold <sup>27</sup>
0.4 V/mm	Elongation of osteoblasts above this threshold <sup>27</sup>
6 V/mm	Enhanced mineral formation in osteoblasts compared to controls <sup>34</sup> Enhanced proliferation, ALP activity, and mineral formation in osteoblasts compared to controls <sup>36</sup>

It is also important to note that field strength was not necessarily the only parameter different between the stimuli.

from reservoirs, such as the endoplasmic reticulum.<sup>25,75</sup> The elevated calcium level in both cases activates the cytoskeletal calmodulin, resulting in, for example, enhanced proliferation, increased vascular endothelial growth factor (VEGF), and transforming growth factor (TGF)-b1 expres-

Table 3. A Comparison of the Effect of Stimulation Delivered with Different Electric Current Densities

The effect of various current densities upon osteoblasts and mesenchymal stem cells	
0.015 A/m <sup>2</sup>	Enhanced proliferation, increased VEGF expression, and no upregulation of bone markers in osteoblasts compared to controls <sup>32</sup> Enhanced proliferation, increased VEGF expression, and upregulation of bone markers a week after end of stimulation in MSCs <sup>10</sup> Increased VEGF expression, upregulation of calcium deposition, and AP activity a week after end of stimulation in MSCs compared to controls. Greater proliferation than with 0.15 A/m <sup>2</sup> stimulation. <sup>13</sup>
0.15 A/m <sup>2</sup>	Worse proliferation of MSCs compared to 0.015 A/m <sup>2</sup> samples. <sup>13</sup> Proliferation and TGF-B1 expression of osteoblasts <sup>63</sup>
3 A/m <sup>2</sup>	Enhanced DNA content increase in osteoblasts <sup>11</sup>
4.2 A/m <sup>2</sup>	Enhanced proliferation and gene expression of osteoblasts <sup>31</sup>
5 A/m <sup>2</sup>	Necrosis in osteoblasts above this threshold (DC) <sup>61</sup>

The effect that the stimulation has seems not to depend on current density, but if the electrical current is raised above a certain threshold, for example, 5 A/m<sup>2</sup> in the case of osteoblasts,<sup>61</sup> necrosis will start to occur.

FIG. 2. An illustration of the effect of the different stimulation techniques: Though the signals delivered through the different methods will be transduced along the same intracellular pathway, they exert their effect on different parts of the cells and will result in an alternative cellular response. Color images available online at [www.liebertpub.com/teb](http://www.liebertpub.com/teb)

sion.<sup>5,11,13,25,32,40,63,75</sup> This hypothesis is supported by the fact that blocking of the calcium channels by verapamil and nifedipine,<sup>13,63</sup> the intracellular stores by TMB-8<sup>75</sup> and the calmodulin by W-7,<sup>63,75</sup> actively impaired or completely blocked the stimulation effect.

Growth factors and receptors under electrical stimulation

The significance of growth factor receptors in electrical stimulation transduction has also been highlighted in many instances, particularly those of the epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF- b1, and VEGF.<sup>20,23,41,71</sup>

This is supported by the observation that removing these growth factors from the culture medium disrupted the EF-induced migration of cells.<sup>71</sup> In another study, it was found that, as a result of DC electrical stimulation, EGF receptors on the cell membrane moved to the cathode electrode facing side of the cell.<sup>21</sup>

The mechanisms behind galvanotaxis

How electrical stimulation can induce migration (i.e., galvanotaxis) and orientation is a difficult question. Asymmetric assembly and disassembly of the actin filaments and polarized redistribution of membrane receptors and integrins have been suggested to play a crucial role in this process.<sup>6,9,16,21,25,76,77</sup> On the other hand, intracellular calcium importance in this instance is still debated. Although intermittent discreet Ca<sup>2+</sup> events<sup>9</sup> and waves, originating from the anodal and heading toward the cathodal side of the cells,<sup>76</sup> have been witnessed during galvanotaxis, researchers were unable to link these to changes in cytoskeletal tension



(known to occur in a pattern corresponding to the direction of the applied  $\mathbf{E}$ s) before movement,<sup>78</sup> or any other part of the migration process. Furthermore, inhibiting the calcium channels by nifedipine, gadolinium, nickel, or strontium did not alter the migratory rates, although the latter seemed to influence the directionality of the movement. Similar orientation disrupting effects were witnessed with Type calcium inhibitors. These findings, together with reports that blocking PI3K, a member of an important motility pathway, inhibits the  $\mathbf{E}$ -driven migration to a greater extent than orientation, suggests that these two are regulated by a different mechanism.<sup>9,23</sup>

Cathode-wise polarization of the Golgi apparatus has been proposed as the key factor in defining the direction of the motion, as the presence of this organelle at the anterior side of the nucleus is a known prerequisite of the forward movement of cells. This polarization seems to be an overriding cue during the electrical  $\mathbf{E}$ -driven migration and disrupting it through chemical means was shown to significantly reduce directional motility of cells.<sup>21</sup>

#### Intracellular signaling pathways

There are several intracellular pathways, including the calcium/cadmium pathway mentioned previously, that are believed responsible for coupling the direct effects of electrical  $\mathbf{E}$ s into cellular responses. The polarized activation of PI3K/Pten and its target Akt and rho, (already known to be important in chemotactic reactions),<sup>9,21,26,79</sup> protein kinase C,<sup>20,23,80</sup> and the mitogen-activated protein kinases p38 and Erk<sup>10,13,26,76,77,80</sup> are believed to be involved in this process and responsible for the observed changes in the expression of proliferative and differentiative genes. This is supported by the strong evidence presented by Kim et al., that blocking p38, ERK, PI3K, or calcium signaling pathways inhibited biphasic electric current-induced proliferation and VEGF and BMP-2 expression in mesenchymal stem cells.<sup>10,13</sup>

#### The Effects of Electrical Stimulation at the Tissue Level

##### Galvanotaxis

Galvanotaxis or electrotaxis is the phenomenon of electric gradient-guided migration, where cells move, depending on the particular type, either toward the cathode or the anode of the stimulating electrodes.<sup>21,23,81</sup> Control over the direction and rate of cell movement, orientation, and division through this mechanism has the potential of stimulating or inhibiting tissue construction/reconstruction, immune responses, blood vessel, and neuron growth and could be a powerful new tool for tissue engineering.<sup>3,6,16,26,71</sup>

In vitro studies have already shown the ability of  $\mathbf{E}$ s to influence the behavior of a variety of cell types. Corneal and lens epithelial cells, umbilical vein and aortic endothelial cells, neural crest cells, fibroblasts, osteoblasts, keratinocytes, lymphocytes, mASCs, and mesenchymal stem cells from various species have all been reported to exhibit cathodal migration and perpendicular orientation in a dose-dependant manner.<sup>6,16,22,26,27,41,70,78,82,83</sup> Exceptions to this were the responses of rat primary osteoclasts<sup>83</sup> and human retinal pigment cells<sup>72</sup> who seemed to favor the anode migration.

A recent investigation, conducted by Au et al.,<sup>3</sup> compared the efficiency of topography and electricity in influencing the alignment and elongation of cardiac myocytes and fibroblasts. Their results show that, although cells respond with a greater alignment to topography than to electrical stimulation, the best results can be achieved with the synergistic application of the two techniques.<sup>3</sup>

##### Enhanced wound healing

The usefulness of electrical stimulation in skin tissue engineering and wound treatment is demonstrated by its ability to induce the re-epithelialization of cutaneous and corneal wounds through promoting migration and proliferation of fibroblasts, keratinocytes and epithelial cells, enhancing angiogenesis, improving blood circulation, and blocking edema formation.<sup>15,22,24,49,59,84,85</sup> Furthermore, stimulation of cutaneous fibroblasts through conductive PPy and PLA scaffolds yielded a 10-fold increase in the expression of the genes IL-6 and IL-8, two cytokines known to play an important role in wound repair and promoting the growth of new blood vessels.<sup>4</sup>

##### Improved nerve regeneration

There are also multiple indications that electrical stimulation can promote peripheral and central nervous system regeneration.<sup>50,74</sup> Already used in numerous clinical applications, such as cochlear implants or the treatment of spinal cord injury and disuse atrophy, novel studies report the ability of electrical stimulation to increase nerve fiber and blood vessel density in sciatic nerve models,<sup>50</sup> double the amount of new cells after spinal cord injury,<sup>74</sup> or to significantly increase neurite length when applied through conductive PPy films.<sup>18</sup>

##### Bone for bone

The positive effects of electricity in healing bone fractures has been noted as early as 1812,<sup>11,86</sup> but interest in such treatments increased considerably when Fukada and Yasuda demonstrated the piezoelectric properties of dry bone during the 1950s and 60s, providing a basis for the use of electrical stimulation as an osteoinductive tool.<sup>11,31,36</sup> Such methods have been successful in treating osteoporosis, osteoarthritis, normal, and nonunion fractures and promoting the integration of implanted biomaterials.<sup>11,13,61,68,87</sup>

In vitro studies on osteoblasts and mesenchymal stem cells show a certain disparity between the effects of the various coupling methods. Capacitive stimulation seems to enhance proliferation, alkaline phosphatase activity, and ECM deposition, while delaying differentiation. Expression of osteopontin, BMP-4, IGF-2, and other genes appear unchanged, while mRNA levels of BMP-2 and TGF- $\beta$ 1 have risen in a few instances. VEGF seems to be an exception, where upregulation has been observed, but only in connection to biphasic stimulation.<sup>10,13,31,32,34,36,63</sup> The inductive PEMF method shows a tendency for inhibiting cell growth, while enhancing mineralization and expression of genes, such as BMP-2, BMP-4, TGF- $\beta$ 1, ALP, and prostaglandin E.<sup>35,53,54,57</sup> A comparison of capacitive, inductive, and combined coupling, by Brighton et al.,<sup>11</sup> showed that only the capacitive delivery of the stimulus results in a continuous

increase of DNA levels.<sup>11</sup> However, both methods improved healing during treatment of nonunions and osteoarthritis in animal models as well as clinical trials.<sup>39,65,68,88,90</sup> Both methods also seem to promote chondrogenesis, through enhancing proteoglycan and collagen secretion and the mRNA expression of TGF- $\beta$ 1 and aggrecan in chondrocytes.<sup>56,91</sup>

#### Effects on the cardiovascular system

Electrical stimulation has also been suggested as a promising tool for solving various cardiovascular-related problems.<sup>59,92</sup> The reported ability of electricity to induce VEGF expression both in endothelial and muscle cells (observed *in vitro* as well as *in vivo*), together with its galvanotactic property could be crucial in better controlling angiogenesis, therefore improving tissue perfusion, promoting endothelialization of artificial vascular implants and preventing limb loss.<sup>41,52,59,92</sup> Furthermore, bioelectricity can greatly benefit cardiac tissue engineering by helping to overcome the challenge of insufficient alignment and differentiation in engineered myocardium.<sup>17</sup> This is because greater proliferation and differentiation, together with better coupling between cells and improved contractile and conductive properties, have all been attributed to the positive influence of electrical stimulation.<sup>5,17,19,28</sup>

#### Future Possibilities

Electrical stimulation has the potential of alleviating some of the problems that currently prevail in tissue engineering, cancer treatment, regenerative medicine, and other biomedical sciences. This physical stimulatory method could take advantage of the galvanotactic property of cells, directing, concentrating, and isolating them.<sup>6,9,93</sup> Furthermore, electrical stimulation could help in growing aligned and orientated tissue, opening up new possibilities in 3D tissue and organ engineering, and allowing greater complexity in the design of the tissue-engineered constructs.<sup>9</sup> Problems, such as the inappropriate colonization of lens epithelial cells after lens or cornea transplants or when engineering these organs, could be avoided by replacing the natural EFs with artificial ones.<sup>7,8</sup>

Electric stimulation has the potential to provide better control over the *in vitro* and *in vivo* proliferation and differentiation of cells and the properties of the resultant tissue, both spatially and temporally.<sup>4</sup> Stem cell fate could be defined more precisely, directing them to specific lineages, yielding a more homogenous cell culture, and making their therapeutic application more plausible.<sup>12,25,77</sup> Cell proliferation, differentiation,<sup>6</sup> and ECM deposition could be enhanced or withheld in greater geometrical complexity than with any other stimuli (i.e., chemical, mechanical) giving tissue-engineered products that are more similar to their natural counterparts in architecture and function. The electrical stimulation technique also has the potential to give investigators the ability to regulate their construct not just in culture, but also after implantation, therefore allowing faster and better integration.

Electrical stimulation offers a cheap, simple, and flexible way to deliver growth factors by inducing the cells themselves to produce these materials through natural pathways. Another advantage of electrical stimulation is that, unlike

treatments that involve incorporated growth factors and gene therapy, its effects disappear with the discontinuation of the treatment. This has obvious benefits for clinical usage allowing the clinician better control of the treatment and acting as a safety net to mitigate post-treatment complications. Electrical stimulation can be used synergistically with other techniques, reducing the required levels of expensive growth factors<sup>4,77</sup> and/or other stimuli and, therefore, the cost of the whole process. Furthermore, tissue-engineered products could be created with a greater speed, bringing this scientific discipline one step closer to the ultimate goal of therapeutic application.<sup>21,77,93</sup> It is also possible to create programmable, multiple electrode bioreactor systems that generate complex EFs customized to the needs of the particular treatment case. With increased research and further understanding, electrical stimuli have significant potential for the field of practical tissue engineering.

#### Acknowledgments

We wish to acknowledge the following sources for funding: EPSRC DTA Studentship (The University of Manchester), EPSRC Bridging The Gap 3ME Funding (University of Keele).

#### Disclosure Statement

There is no known actual or potential conflict of interest for any of the authors.

#### References

- Boone, K.J.M., Langelaan, M.L.P., Polak, R.B., van der Schaft, D.W.J., Baaijens, F.P.T., and Post, M.J. Effects of a combined mechanical stimulation protocol: value for skeletal muscle tissue engineering. *J Biomech* 43, 1514, 2010.
- Gauvi, R., Parenteau-Bareil, R., Larouche, D., Marcoux, H., Bisson, F., Bonnet, A., Auger, F.A., Bolduc, S., and Germain, L. Dynamic mechanical stimulations induce anisotropy and improve the tensile properties of engineered tissues produced without exogenous scaffolding. *Acta Biomater* 7, 3294, 2011.
- Au, H.T.H., Cheng, I., Chowdhury, M.F., and Radisic, M. Interactive effects of surface topography and pulsatile electrical field stimulation on orientation and elongation of fibroblasts and cardiomyocytes. *Biomaterials* 28, 4277, 2007.
- Shi, G., Zhang, Z., and Rouabhia, M. The regulation of cell functions electrically using biodegradable polypyrrole/poly lactide conductors. *Biomaterials* 29, 3792, 2008.
- Rowlands, A.S., and Cooper-White, J.J. Directing phenotype of vascular smooth muscle cells using electrically stimulated conducting polymer. *Biomaterials* 29, 4510, 2008.
- Sun, S., Titushkin, I., and Cho, M. Regulation of mesenchymal stem cell adhesion and orientation in 3D collagen scaffold by electrical stimulus. *Bioelectrochemistry* 69, 133, 2006.
- Wang, E., Zhao, M., Forrester, J.V., and McCaig, C.D. Reorientation and faster, directed migration of lens epithelial cells in a physiological electric field. *Exp Eye Res* 71, 91, 2000.
- Wang, E., Zhao, M., Forrester, J.V., and McCaig, C.D. Bidirectional migration of lens epithelial cells in a physiological electrical field. *Exp Eye Res* 76, 29, 2003.
- Hammerick, K.E., Longaker, M.T., and Prinz, F.B. *In vitro* effects of direct current electric fields on adipose-derived

- stromal cells. *Biochem. Biophys Res Commun* 397, 12, 2010.
10. Kim, I.S., Song, J.K., Song, Y.M., Lee, T.H., Cho, T.H., Lim, S.S., Pan, H., Kim, S.J., and Hwang, S.J. Novel action of biphasic electric current in vitro osteogenesis of human bone marrow mesenchymal stromal cells coupled with VEGF production. *Bone* 43, S43, 2008.
11. Brighton, C.T., Wang, W., Seldes, R., Zhang, G., and Pollack, S.R. Signal transduction in electrically stimulated bone cells. *J Bone Joint Surg Am* 83-A, 1514, 2001.
12. Sundelacruz, S., Levin, M., and Kaplan, D.L. Membrane potential controls adipogenic and osteogenic differentiation of mesenchymal stem cells. *PLoS One* 3, 1, 2008.
13. Kim, I.S., Song, J.K., Song, Y.M., Cho, T.H., Lee, T.H., Lim, S.S., Kim, S.J., and Hwang, S.J. Novel effect of biphasic electric current on in vitro osteogenesis and cytokine production in human mesenchymal stromal cells. *Tissue Eng Part A* 15, 1, 2009.
14. McCaig, C.D., Rajnicek, A.M., Song, B., and Zhao, M. Controlling cell behaviour electrically: current views and future potential. *Physiol Rev* 85, 943, 2005.
15. Huo, R., Ma, Q., Wu, J.J., Chin-Nuke, K., Jing, Y., Chen, J., Miyar, M.E., Davis, S.C., and Li, J. Noninvasive electromagnetic fields on keratinocyte growth and migration. *J Surg Res* 162, 299, 2009.
16. Li, X., and Kolega, J. Effects of direct current electric fields on cell migration and actin filament distribution in bovine vascular endothelial cells. *J Vasc Res* 39, 391, 2002.
17. Radisic, M., Park, H., Shing, H., Consi, T., Schoen, F.J., Langer, R., Freed, L.E., and Vunjak-Novakovic, G. Functional assembly of engineered myocardium by electrical stimulation of cardiac myocytes cultured on scaffolds. *Proc Natl Acad Sci U S A* 101, 18129, 2004.
18. Schmidt, C.E., Shastri, V.R., Vacanti, J.P., and Langer, R. Stimulation of neurite outgrowth using an electrically conducting polymer. *Proc Natl Acad Sci USA* 94, 8948, 1997.
19. Sauer, H., Bekhtere, M.M., Hescheler, J., and Wartenberg, M. Redox control of angiogenic factors and CD31-positive vessel-like structures in mouse embryonic stem cells after direct current electrical field stimulation. *Exp Cell Res* 304, 380, 2005.
20. Nuccitelli, R. A role for endogenous electric fields in wound healing. *Curr Top Dev Biol* 58, 1, 2003.
21. Zhao, M. Electrical fields in wound healing: an overriding signal that directs cell migration. *Semin Cell Dev Biol* 20, 674, 2009.
22. Farboud, B., Nuccitelli, R., Schwab, I.R., and Isseroff, R.R. DC Electric fields induce rapid directional migration in cultured human corneal epithelial cells. *Exp Eye Res* 70, 667, 2000.
23. Mycielska, M.E., and Djamgoz, M.B.A. Cellular mechanisms of direct-current electric field effects: galvanotaxis and metastatic disease. *J Cell Sci* 117, 1631, 2004.
24. Song, B., Zhao, M., Forrester, J.V., and McCaig C.D. Electrical cues regulate the orientation and frequency of cell division and the rate of wound healing in vivo. *Proc Natl Acad Sci USA* 99, 13577, 2002.
25. Titushkin, I., and Cho, M. Regulation of cell cytoskeleton and membrane mechanics by electric field: role of linker proteins. *Biophys J* 96, 1717, 2009.
26. Lin, F., Baldessari, F., Gyenge, C.C., Sato, T., Chambers, R.D., Santiago, J.G., and Butcher, E.C. Lymphocyte electro-taxis in vitro and in vivo. *J Immunol* 181, 2465, 2008.
27. Erickson C.A., and Nuccitelli R. Embryonic fibroblast motility and orientation can be influenced by physiological electric fields. *J Cell Biol* 98, 296, 1984.
28. Serena, E., Figallo, E., Tandon, N., Cannizzaro, C., Gerecht, S., Elvassore, N., and Vunjak-Novakovic, G. Electrical stimulation of human embryonic stem cells: cardiac differentiation and the generation of reactive oxygen species. *Exp Cell Res* 315, 3611, 2009.
29. Reid, B., Song, B., and Zhao, M. Electric currents in *Xenopus* tadpole tail regeneration. *Dev Biol* 335, 198, 2009.
30. Illingworth, C.M., and Barker, A.T. Measurement of electrical currents emerging during the regeneration of amputated finger tips in children. *Clin Phys Physiol Meas* 1, 87, 1980.
31. Ercan, B., and Webster, T.J. The effect of biphasic electrical stimulation on osteoblast function at anodized nanotubular titanium surfaces. *Biomaterials* 31, 3684, 2010.
32. Kim, I.S., Song, J.K., Zhang, Y.L., Lee, T.H., Cho, T.H., Song, Y.M., Kim, D.K., Kim, S.J., and Hwang, S.J. Biphasic electric current stimulates proliferation and induces VEGF production in osteoblasts. *Biochim Biophys Acta* 1763, 907, 2006.
33. Petersson, C.J., Holmer, N.G., and Johnell, O. Electrical Stimulation of osteogenesis studies of the cathode effect on rabbit femur. *Acta orthop scand* 53, 727, 1982.
34. Wiesmann, H.P., Hartig, M., Stratmann, U., Meyer, U., and Joos, U. Electrical stimulation influences mineral formation of osteoblast-like cells in vitro. *Biochim Biophys Acta* 1538, 28, 2001.
35. Molen, M.A.V., Donahue, H.J., Rubin, C.T., and McLeod, K.J. Osteoblastic networks with decoupled coupling: differential effects of magnetic and electric field exposure. *Bone* 27, 227, 2000.
36. Hartig, M., Joos, U., and Wiesmann, H.P. Capacitively coupled electric fields accelerate proliferation of osteoblast-like primary cells and increase bone extracellular matrix formation in vitro. *Eur Biophys J* 29, 499, 2000.
37. Huang, C.P., Chen, X.M., and Chen, Z.Q. Osteocyte: the impresario in the electrical stimulation for bone fracture healing. *Med Hypotheses* 70, 287, 2008.
38. Macginitie, L.A., Wu, D.D., and Cochran, G.V.B. Streaming potentials in healing, remodeling, and intact cortical bone. *J Bone Miner Res* 8, 1323, 1993.
39. McLeod, K.J., and Rubin, C.T. The effect of low-frequency electrical fields on osteogenesis. *J Bone Joint Surg Am* 74, 920, 1992.
40. Xu, J., Wang, W., Clark, C.C., and Brighton, C.T. Signal transduction in electrically stimulated articular chondrocytes involves translocation of extracellular calcium through voltage-gated channels. *Osteoarthritis Cartilage* 17, 397, 2009.
41. Zhao, M., Bai, H., Wang, E., Forrester, J.V., and McCaig, C.D. Electrical stimulation directly induces pre-angiogenic responses in vascular endothelial cells by signaling through VEGF receptors. *J Cell Sci* 117, 397, 2004.
42. Brighton, C.T., Black, J., Friedenberg, Z.B., Esterhai, J.L., Day, L.J., and Connolly, J.F. A multicenter study of the treatment of non-union with constant direct current. *J Bone Joint Surg Am* 63, 2, 1981.
43. Evans, R.O.N., Goldberg, J.A., Bruce, W.J.M., and Walsh, W. Reoperated clavicular nonunion treated with osteogenic protein 1 and electrical stimulation. *J Shoulder Elbow Surg* 13, 573, 2004.



44. Gardner, S.E., Franz, R.A., and Schmidt, F.L. Effect of electrical stimulation on chronic wound healing: a meta-analysis. *Wound Repair Regen* 7, 495, 1999.
45. Shimada, Y., Sato, K., Kagaya, H., Konishi, N., Miyamoto, S., and Matsunaga, Y. Clinical use of percutaneous intramuscular electrodes for functional electrical stimulation. *Arch Phys Med Rehabil* 77, 1014, 1996.
46. Janardan, B. Electrical stimulation in medical sciences from basic to applied. *J Appl Basic Med Sci* 3, 49, 2001.
47. Deyo, R.E., Walsh, N.E., Martin, D.C., Shoenfeld, L.S., and Ramamurthy, S. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic back pain. *N Engl J Med* 322, 1627, 1990.
48. Johnson, M.I., Ashton, C.H., and Thompson, J.W. An in-depth study of long-term users of transcutaneous electrical nerve stimulation (TENS). Implications for clinical use of TENS. *Pain* 44, 221, 1991.
49. Houghton, P.E., Campbell, K.E., Fraser, C.H., Harris, C., Keast, D.H., Potter, P.J., Hayes, K.C., and Woodbury, M.G. Electrical stimulation therapy increases rate of healing of pressure ulcers in community-dwelling people with spinal cord injury. *Arch Phys Med Rehabil* 91, 669, 2010.
50. Mendonça A.C., Barbieri, C.H., and Mazzer, N. Directly applied low intensity direct electric current enhances peripheral nerve regeneration in rats. *J Neurosci Methods* 129, 183, 2003.
51. McLeod, K.J., Donahue, H.J., Levin, P.E., Fontaine, M.A., and Rubin, C.T. Electric fields modulate bone cell function in a density-dependent manner. *J Bone Miner Res* 1, 977, 1993.
52. Bouaziz, A., Richert, A., and Caprani, A. Morphological aspects of endothelial cells cultured in the absence of fetal calf serum under controlled electrical charges applied to the support. *Biomaterials* 17, 2281, 1996.
53. Lohmann, C.H., Schwartz, Z., Liu, Y., Li, Z., Simon, B.J., Sylvia, V.L., Dean, D.D., Bonewald, L.F., Donahue, H.J., and Boyan, B.D. Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells. *J Orthop Res* 21, 326, 2003.
54. Chang, K., Chang, W.H.S., Huang, S., Huang, S., and Shih, C. Pulsed electromagnetic fields stimulation affects osteoclast formation by modulation of osteoprotegerin, RANK ligand and macrophage colony-stimulating factor. *J Orthop Res* 23, 1308, 2005.
55. Lam, H., and Qin, Y.X. The effects of frequency-dependent dynamic muscle stimulation on inhibition of trabecular bone loss in a disuse model. *Bone* 43, 1093, 2008.
56. Aaron, R.K., Wang, S., and Ciombor, D. McK. Upregulation of basal TGF $\beta$ 1 levels by EMF coincident with chondrogenesis: implications for skeletal repair and tissue engineering. *J Orthop Res* 20, 233, 2002.
57. Bodamyali, T., Bhatt, B., Hughes, F.J., Winrow, V.R., Kanczler, J.M., Simon, B., Abbott, J., Blake, D.R., and Stevens, C.R. Pulsed electromagnetic fields simultaneously induce osteogenesis and upregulate transcription of bone morphogenetic proteins 2 and 4 in rat osteoblasts in vitro. *Biochem Biophys Res Commun* 250, 458, 1998.
58. Feedar, J.A., Kloth, L.C., and Gentzkow, G.D. Chronic dermal ulcer healing enhanced with monophasic pulsed electrical stimulation. *Phys Ther* 71, 639, 1991.
59. Ferroni P, Roselli M, Guadagni F, Martini F, Mariotti S, Marchitelli E, and Cipriani C. Biological effects of a software-controlled voltage pulse generator (PhyBack PBK-2C) on the release of vascular endothelial growth factor (VEGF). *In Vivo* 19, 949, 2005.
60. Merrill, D.R., Bikson, M., and Jefferys, J.G.R. Electrical stimulation of excitable tissue: design of efficacious and safe protocols. *J Neurosci Methods* 141, 171, 2005.
61. Spadaro, J.A., and Becker, R.O. Function of implanted cathodes in electrode-induced bone growth. *Med Biol Eng Comput* 17, 769, 1979.
62. Antov, Y., Barbul, A., Mantsur, H., and Korenstein, R. Electroendocytosis: exposure of cells to pulsed low electric fields enhances adsorption and uptake of macromolecules. *Biophys J* 88, 2206, 2005.
63. Zhuang, H., Wang, W., Seldes, R.M., Tahernia, A.D., Fan, H., and Brighton, C.T. Electrical stimulation induces the level of TGF- $\beta$ 1 mRNA in osteoblastic cells by a mechanism involving calcium/calmodulin pathway. *Biochem Biophys Res Commun* 237, 225, 1997.
64. Abeed, R.I., Naseer, M., and Abel, E.W. Capacitively coupled electrical stimulation treatment: results from patients with failed long bone fracture unions. *J Orthop Trauma* 12, 510, 1998.
65. Zamora-Navas, P., Borrás Verdera, A., Antelo Lorenzo, R., Saras Ayuso, J.R., and Peña Reina, M.C. Electrical stimulation of bone nonunion with the presence of a gap. *Acta Orthop Belg* 61, 169, 1995.
66. Meng, S., Rouabhia, M., and Zhang, Z. Electrical stimulation in tissue regeneration. In: Gargiulo, G.D., and McEwan, A., eds. *Applied Biomedical Engineering*. Rijeka: InTech, 2011, pp. 37-52.
67. Yonemori, K., Matsunaga, S., Ishidou, Y., Maeda, S., and Yoshida, H. Early effects of electrical stimulation on osteogenesis. *Bone* 19, 173, 1996.
68. Thamsborg G, Florescu A, Oturai P, Fallentin E, Tritsarlis K, and Dissing S. Treatment of knee osteoarthritis with pulsed electromagnetic fields: a randomized, double-blind, placebo-controlled study. *Osteoarthritis Cartilage* 13, 575, 2005.
69. Fu, Y., Lin, C., Chang, J., Wang, G., and Ho, M. A novel single pulsed electromagnetic field stimulates the osteogenesis of human bone marrow derived mesenchymal stem cells. *Bone* 55, S322, 2009.
70. Cooper M.S., and Keller, R.E. Perpendicular orientation and directional migration of amphibian neural crest cells in dc electrical fields. *Proc Natl Acad Sci USA* 81, 160, 1984.
71. Zhao, M., Dick, A., Forrester, J.V., and McCaig, C.D. Electric field-directed cell motility involves up-regulated expression and asymmetric redistribution of the epidermal growth factor receptors and is enhanced by fibronectin and laminin. *Mol Biol Cell* 10, 1259, 1999.
72. Gamboa, O.L., Pu, J., Townend, J., Forrester, J.V., Zhao, M., McCaig, C., and Lois, N. Electrical stimulation of retinal pigment epithelial cells. *Exp Eye Res* 91, 195, 2010.
73. Franek, A., Polak, A., and Kucharzewski, M. Modern application of high voltage stimulation for enhanced healing of venous crural ulceration. *Med Eng Phys* 22, 647, 2000.
74. Becker, D., Gary, D.S., Rosenzweig, E.S., Grill, W.M., and McDonald, J.W. Functional electrical stimulation helps replenish progenitor cells in the injured spinal cord of adult rats. *Exp Neurol* 222, 211, 2010.
75. Li, J.K.J., Lin, J.C.A., Liu, H.C., Sun, J.S., Ruaan, R.C., Shih, C., and Chang, W.H.S. Comparison of ultrasound and electromagnetic field effects on osteoblast growth. *Ultrasound Med Biol* 32, 769, 2006.

76. Funk, R.H.W., Monsees, T., and Özkücur, N. Electromagnetic effects from cell biology to medicine. *Prog Histochem Cytochem* 43, 177, 2009.
77. Titushkin, I., Sun, S., Shin, J., and Cho, M. Physicochemical control of adult stem cell differentiation: shedding light on potential molecular mechanisms. *J Biomed Biotechnol* 2010, 743476, 2010.
78. Curtze, S., Dembo, M., Miron, M., and Jones, D.B. Dynamic changes in traction forces with DC electric field in osteoblast-like cells. *J Cell Sci* 117, 2721, 2004.
79. Levin, M. Large-scale biophysics: ion flows and regeneration. *Trends Cell Biol* 17, 261, 2007.
80. Sun, S., Liu, Y., Lipsky, S., and Cho, M. Physical manipulation of calcium oscillations facilitates osteodifferentiation of human mesenchymal stem cells. *FASEB J* 21, 1472, 2007.
81. Levin, M. Bioelectric mechanisms in regeneration: unique aspects and future perspectives. *Semin Cell Dev Biol* 20, 543, 2009.
82. Zhao, M., Forrester, J.V., and McCaig, C.D. A small, physiological electric field orients cell division. *Proc Natl Acad Sci USA* 96, 4942, 1999.
83. Ferrier, J., Ross, S.M., Kanehisa, J., and Aubin, J.E. Osteoclasts and osteoblasts migrate in opposite directions in response to a constant electrical field. *J Cell Physiol* 129, 283, 1986.
84. Bayat, M., Asgari-Moghadam, Z., Maroufi, M., Rezaie, F.S., Bayat, M., and Rakhshan, M. Experimental wound healing using microamperage electrical stimulation in rabbits. *J Rehabil Res Dev* 43, 219, 2006.
85. Jercinovic, A., Hinsenkamp, M., Scarceriaux, B., Willaert, F., de Graef, Ch., Heenen, M., and Goldshmidt, D. Effects of direct constant current (DC) on keratinocytes in vitro. *Bioelectrochem Bioenerg* 39, 209, 1996.
86. Peltier, L.F. A brief historical note on the use of electricity in the treatment of fractures. *Clin Orthop Relat Res* 161, 4, 1981.
87. Toth, J.M., Seim, H.B., Schwardt, J.D., Humphrey, W.B., Wallskog, J.A., and Turner, A.S. Direct current electrical stimulation increases the fusion rate of spinal fusion cages. *Spine* 25, 2580, 2000.
88. Fini, M., Cadossi, R., Canè, V., Cavani, F., Giavaresi, G., Krajewski, A., Martini, L., Aldini, N.N., Ravaglioli, A., Rimondini, L., Torricelli, P., and Giardino, R. The effect of pulsed electromagnetic fields on the osteointegration of hydroxyapatite implants in cancellous bone: a morphologic and microstructural in vivo study. *J Orthop Res* 20, 756, 2002.
89. Garland, D., Holt, P., Harrington, J.T., Caldwell, J., Zizic, T., and Cholewczynski, J. A 3-month, randomized, double-blind, placebo-controlled study to evaluate the safety and efficacy of a highly optimized, capacitively coupled, pulsed electrical stimulator in patients with osteoarthritis of the knee. *Osteoarthritis Cartilage* 15, 630, 2007.
90. Satter Syed, A., Islam, M.S., Rabbani, K.S., and Talukder, M.S. Pulsed electromagnetic fields for the treatment of bone fractures. *Bangladesh Med Res Counc Bull* 25, 6, 1999.
91. Brighton, C.T., Wang, W., and Clark, C.C. Up-regulation of matrix in bovine articular cartilage explants by electric fields. *Biochem Biophys Res Commun* 342, 556, 2006.
92. Peters, E.J.G., Armstrong, D.G., Wunderlich, R.P., Bosma, J., Stacpoole-Shea, S., and Lavery, L.A. The benefit of electrical stimulation to enhance perfusion in persons with diabetes mellitus. *J Foot Ankle Surg* 37, 396, 1998.
93. Wiesmann, H.P., Joos, U., and Meyer, U. Biological and biophysical principles in extracorporeal bone tissue engineering Part II. *Int J Oral Maxillofac Surg* 33, 523, 2004.

Address correspondence to:

Sarah H. Cartmell, BEng (Hons), PhD  
Materials Science Centre  
University of Manchester  
Grosvenor Street  
Manchester M1 7HS  
United Kingdom

E-mail: sarah.cartmell@manchester.ac.uk

Received: March 21, 2012

Accepted: August 7, 2012

Online Publication Date: October 8, 2012