Lab 8

Quadrature formulas (1)

Problems

1. a) Approximate the integral

$$I = \int_0^1 f(x)dx$$
, for $f(x) = \frac{2}{1+x^2}$,

using trapezium formula.

- b) Plot the graph of the function f and the graph of the trapezium with vertices (0,0),(0,f(0)),(1,f(1)) and (1,0).
 - c) Approximate the integral I using Simpson's formula.
 - 2. Approximate the following double integral

$$\int_{1.4}^{2} \int_{1}^{1.5} \ln(x+2y) dy dx$$

using trapezium formula for double integrals, given in (1). (Result: 0.4295545)

3. Evaluate the integral that arises in electrical field theory:

$$H(p,r) = \frac{60r}{r^2 - p^2} \int_0^{2\pi} \left[1 - \left(\frac{p}{r}\right)^2 \sin x \right]^{1/2} dx,$$

for $r=110,\,p=75,$ using the repeated trapezium formula for two given values of n. (Result: 6.3131)

- **4.** Find the smallest value of n that gives an approximation of the integral $\int_1^2 x \ln(x) dx$ which is correct to three decimals, using the repeated trapezium formula. Apply the repeated trapezium formula for the obtained value of n to approximate the integral. (*Result:* 0.636294368858383)
 - 5. Evaluate the integral

$$\int_0^\pi \frac{dx}{4 + \sin 20x}$$

using the repeated Simpson's formula for n = 10 and 30. (Result: 0.8111579)

6. The error function erf(x) is defined by

$$erf(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt.$$

Use the repeated Simpson's formula to evaluate erf(0.5) with n=4 and n=10. Estimate the accuracy of your result and compare with the correct value erf(0.5)=0.520499876.

Trapezium formula for double integral

Applying succesively trapezium formula with respect to y, and with respect to x, we get

$$\int_{a}^{b} \int_{c}^{d} f(x,y) dy dx \approx \frac{(b-a)(d-c)}{16} \left[f(a,c) + f(a,d) + f(b,c) + f(b,d) \right]$$

$$+ 2f\left(\frac{a+b}{2},c\right) + 2f\left(\frac{a+b}{2},d\right) + 2f\left(a,\frac{c+d}{2}\right)$$

$$+ 2f\left(b,\frac{c+d}{2}\right) + 4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)$$