

COURSE 1

Groups, rings and fields

Definition 1. By a **binary operation** on a set A we understand a map

$$\varphi : A \times A \rightarrow A.$$

Since all the operations considered in this section are binary operations, we briefly call them **operations**. Usually, we denote operations by symbols like $*$, \cdot , $+$, and the image of an arbitrary pair $(x, y) \in A \times A$ is denoted by $x * y$, $x \cdot y$ (multiplicative notation), $x + y$ (additive notation), respectively.

Examples 2. a) The usual addition and multiplication are operations on \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} , \mathbb{C} , but not on the set of irrational numbers.

b) The usual subtraction is an operation on \mathbb{Z} , \mathbb{Q} , \mathbb{R} and \mathbb{C} , but not on \mathbb{N} .

c) The usual division is an operation on \mathbb{Q}^* , \mathbb{R}^* , \mathbb{C}^* , but not on \mathbb{Q} , \mathbb{R} , \mathbb{C} , \mathbb{N} , \mathbb{Z} , \mathbb{N}^* or \mathbb{Z}^* .

Definitions 3. Let $*$ be an operation on A . We say that:

i) $*$ is **associative** if

$$(a_1 * a_2) * a_3 = a_1 * (a_2 * a_3), \quad \forall a_1, a_2, a_3 \in A;$$

ii) $*$ is **commutative** if

$$a_1 * a_2 = a_2 * a_1, \quad \forall a_1, a_2 \in A.$$

iii) $e \in A$ is an **identity element** for $*$ if

$$a * e = e * a = a, \quad \forall a \in A.$$

When using the multiplicative or additive notation, an identity element e is usually denoted by 1 or 0, respectively.

Definition 4. Let A be set and let \cdot be an operation with an identity element 1. An element $a \in A$ **has an inverse** if there exists an element $a' \in A$ such that

$$a \cdot a' = a' \cdot a = e.$$

We say that a' is an **inverse** for a .

When using the multiplicative notation, the inverse of a is denoted by a^{-1} . When using the or additive notation the inverse of a is denoted by $-a$, and it is called **the opposite of a** .

Definitions 5. A pair $(A, *)$ is called **monoid** if $*$ is associative and it has an **identity element**. A monoid with a commutative operation is called **commutative monoid**.

Definition 6. A pair (A, \cdot) is called **group** if it is a monoid in which every element has an inverse. If the operation is commutative as well, the structure is called **commutative** or **Abelian group**.

Examples 7. a) $(\mathbb{N}, +)$ and (\mathbb{Z}, \cdot) are commutative monoids, but they are not groups.

b) (\mathbb{Q}, \cdot) , (\mathbb{R}, \cdot) , (\mathbb{C}, \cdot) are commutative monoids, but they are not groups since 0 has no inverse.

c) $(\mathbb{Z}, +)$, $(\mathbb{Q}, +)$, $(\mathbb{R}, +)$, $(\mathbb{C}, +)$, (\mathbb{Q}^*, \cdot) , (\mathbb{R}^*, \cdot) , (\mathbb{C}^*, \cdot) are Abelian groups.

Remark 8. The group definition can be rewritten: (A, \cdot) is a **group** if and only if it follows the following conditions:

- (i) $(a_1 \cdot a_2) \cdot a_3 = a_1 \cdot (a_2 \cdot a_3)$, $\forall a_1, a_2, a_3 \in A$ (\cdot is associative);
- (ii) $\exists 1 \in A$, $\forall a \in A$: $a \cdot 1 = 1 \cdot a = a$ (there exists an identity element for \cdot);
- (iii) $\forall a \in A$, $\exists a^{-1} \in A$: $a \cdot a^{-1} = a^{-1} \cdot a = 1$ (all the elements of A have inverses).

Definitions 9. Let φ be an operation on the set A and $B \subseteq A$. We say that B is **closed under** φ if

$$b_1, b_2 \in B \Rightarrow \varphi(b_1, b_2) \in B.$$

If B is closed under φ , one can define an operation on B as follows:

$$\varphi' : B \times B \rightarrow B, \quad \varphi'(b_1, b_2) = \varphi(b_1, b_2).$$

We call φ' the **operation induced** by φ on B or, briefly, the **induced operation**. Most of the time, we denote it also by φ .

Remarks 10. a) Let φ be an operation on the set A , $B \subseteq A$ closed under φ and let φ' be the induced operation on B . If φ is associative or commutative, then φ' is associative or commutative, respectively.

b) Let φ_1 and φ_2 be operations on A , let $B \subseteq A$ be closed under φ_1 and φ_2 , and let φ'_1 and φ'_2 be the operations induced by φ_1 and φ_2 on B , respectively. If φ_1 is distributive with respect to φ_2 , i.e.

$$\varphi_1(a_1, \varphi_2(a_2, a_3)) = \varphi_2(\varphi_1(a_1, a_2), \varphi_1(a_1, a_3)), \forall a_1, a_2, a_3 \in A,$$

then φ'_1 is distributive with respect to φ'_2 .

c) The existence of an identity element is not always preserved by induced operations. For instance, \mathbb{N}^* is closed in $(\mathbb{N}, +)$, but $(\mathbb{N}^*, +)$ has no identity element.

Definition 11. Let (G, \cdot) be a group. A subset $H \subseteq G$ is called a **subgroup of G** if:

i) H is closed under the operation of (G, \cdot) , that is,

$$\forall x, y \in H, \quad x \cdot y \in H;$$

ii) H is a group with respect to the induced operation.

Examples 12. a) \mathbb{Z} , \mathbb{Q} , \mathbb{R} are subgroups of $(\mathbb{C}, +)$, \mathbb{Z} , \mathbb{Q} are subgroups of $(\mathbb{R}, +)$ and \mathbb{Z} is a subgroup of $(\mathbb{Q}, +)$.

b) \mathbb{Q}^* , \mathbb{R}^* are subgroups of (\mathbb{C}^*, \cdot) and \mathbb{Q}^* is a subgroup of (\mathbb{R}^*, \cdot) .

c) \mathbb{N} is closed in $(\mathbb{Z}, +)$, but it is not a subgroup.

d) Every non-trivial group (G, \cdot) has two subgroups, namely $\{1\}$ and G . Any other subgroup of (G, \cdot) is called **proper subgroup**.

Definition 13. Let $(G, *)$, (G', \perp) be two groups. A map $f : G \rightarrow G'$ is called **homomorphism** (or **morphism**) if

$$f(x_1 * x_2) = f(x_1) \perp f(x_2), \quad \forall x_1, x_2 \in G.$$

A bijective homomorphism is called **isomorphism**. A homomorphism of $(G, *)$ into itself is called **endomorphism** of $(G, *)$. An isomorphism al lui $(G, *)$ into itself is called **automorphism** of $(G, *)$. If there exists an isomorphism $f : G \rightarrow G'$, we say that the groups $(G, *)$ and (G', \perp) are isomorphic and we denote this by $G \simeq G'$ or $(G, *) \simeq (G', \perp)$.

Let us come back to the multiplicative notation.

Theorem 14. Let (G, \cdot) and (G', \cdot) be groups, and let 1 and $1'$, respectively, be the identity element of (G, \cdot) and (G', \cdot) , respectively. If $f : G \rightarrow G'$ is a group homomorphism, then:

- (i) $f(1) = 1'$;
- (ii) $[f(x)]^{-1} = f(x^{-1}), \forall x \in G$.

Proof.

□