



Specialization report

Heinrich Heine Universität Düsseldorf
Institut für Theoretische Physik I

Presented by:
Marius Theißen
Matrn.: 2163903

22.11.2018

Contents

1	Introduction	2
2	Pictures in Quantum Mechanics	2
2.1	Schrödinger picture and Heisenberg picture	2
2.2	Interaction picture	4
2.3	The In and Out picture: External currents	5
3	Scattering operator	9
3.1	Solutions for the Interaction picture	9
3.2	Solutions for the In and Out picture	12
3.3	Connections	12
3.4	Unitarity of S	15
4	Gell-Mann Low formula	16
4.1	Vacuum expectation value	19
5	Conclusion	20
5.A	Vacuum transition probability under external current	21
5.B	Chronological time ordering	23
5.C	Anti-chronological time ordering	26

1 Introduction

This report has three main goals. First we shall establish the transition between pictures in quantum mechanics and quantum field theory. We go over the more common ones, Schrödinger, Heisenberg and Interaction picture, to derive and justify the two In and Out Pictures. By working out a few handy techniques and methods on the way, we will proof the Gell-Mann Low formula. This important formula allows us transition of polynomial terms of field operators in the Heisenberg picture to multiple different but equivalent picture expressions. These are based on boundary conditions imposing free motion on particles along them. We derive them from experimental and physical point of views. The Gell-Mann Low formula does this to this point without loss of information. In case of calculating multi particle processes the Gell-Mann Low formula allows us to perform perturbation theory on the correlation functions by introducing the scattering operator S which then can be expanded. This S is the key to calculate cross-sections σ and decay rates Γ .

2 Pictures in Quantum Mechanics

2.1 Schrödinger picture and Heisenberg picture

In the Schrödinger picture the operators are time-independent but the wave-functions are time dependent. The time evolution of a state vector is controlled by the Schrödinger equation. Let $|\Psi(t)\rangle$ denote a state vector at time t . It satisfies

$$i\hbar \frac{\partial}{\partial t} |\Psi_S(t)\rangle = \hat{H} |\Psi_S(t)\rangle, \quad (1)$$

where \hat{H} is the Hamiltonian of the system. When assuming it time independent, the solution of eq. (1) can be formally written as

$$|\Psi_S(t)\rangle = \hat{U} |\Psi_S(t_0)\rangle \quad (2)$$

with $\hat{U} = e^{-\frac{i}{\hbar} \hat{H}(t-t_0)}$ is the time evolution operator, which satisfies the differential equation

$$i\hbar \partial_t \hat{U} = \hat{H} \hat{U}. \quad (3)$$

Here $|\Psi_S(t_0)\rangle$ is a ket of $t = t_0$. We shall generally take $t_0 = 0$ and write

$$|\Psi_S(t)\rangle = e^{-\frac{i}{\hbar} \hat{H}t} |\Psi_H\rangle. \quad (4)$$

The state on the right-hand side has no longer time dependence. This defines the state in the Heisenberg picture.

The above two pictures differ between each other in the way of storing the time dependence of the total system. In the Schrödinger picture only the states carry such a dependence, whereas in the Heisenberg picture only operators has this possibility. To verify this statement we study matrix element of an operator in the Schrödinger picture

$$\langle \Psi'_S(t) | \hat{A}_S | \Psi_S(t) \rangle = \langle \Psi'_H | e^{\frac{i}{\hbar} t \hat{H}} \hat{A}_S e^{-\frac{i}{\hbar} t \hat{H}} | \Psi_H \rangle, \quad (5)$$

where Eq.(4) has been used. As a consequence,

$$\hat{A}_H(t) = e^{\frac{i}{\hbar} t \hat{H}} \hat{A}_S e^{-\frac{i}{\hbar} t \hat{H}} = \hat{U}^{-1} \hat{A}_S \hat{U}. \quad (6)$$

This new operator $\hat{A}_H(t)$ in combination with the state $|\Psi_H\rangle$ defines the Heisenberg picture. The choice of a picture always requires to establish the states but also the corresponding operators. observe that the time evolution of $\hat{A}_H(t)$ is dictated by an equation that follows from differentiating the equation above with respect to t :

$$\frac{d}{dt} \hat{A}_H(t) = \frac{i}{\hbar} \hat{H} e^{\frac{i}{\hbar} t \hat{H}} \hat{A}_S e^{-\frac{i}{\hbar} t \hat{H}} + e^{\frac{i}{\hbar} t \hat{H}} \hat{A}_S \left(-\frac{i}{\hbar} \hat{H} \right) e^{-\frac{i}{\hbar} t \hat{H}}. \quad (7)$$

Here we have used the time evolution equation given below (2) derivative of the unitary operator itself:

$$\begin{aligned} \frac{d}{dt} \hat{A}_H(t) &= \frac{i}{\hbar} \hat{U}^{-1} \hat{H} \hat{A}_S \hat{U} - \frac{i}{\hbar} \hat{U}^{-1} \hat{A}_S \hat{H} \hat{U} \\ &= \frac{i}{\hbar} \hat{U}^{-1} \hat{H} \underbrace{\hat{U} \hat{U}^{-1}}_{=1} \hat{A}_S \hat{U} - \frac{i}{\hbar} \hat{U}^{-1} \hat{A}_S \underbrace{\hat{U} \hat{U}^{-1}}_{=1} \hat{H} \hat{U}. \end{aligned} \quad (8)$$

The inserted 1 allows us to express Eq.(8) in term of operators in the Heisenberg picture.

$$\frac{d}{dt} \hat{A}_H(t) = \frac{i}{\hbar} \hat{H}_H(t) \hat{A}_H(t) - \frac{i}{\hbar} \hat{A}_H(t) \hat{H}_H(t), \quad (9)$$

where $\hat{H}_H(t)$ is the respective Hamiltonian in the Heisenberg picture. Therefore:

$$i \frac{d}{dt} \hat{A}_H(t) = \frac{1}{\hbar} \left[\hat{A}_H(t), \hat{H}_H(t) \right]. \quad (10)$$

2.2 Interaction picture

A third picture can be introduced: the Interaction picture (sometimes called the Dirac picture). We will see very shortly that, in the interacting picture both the states and the respective operators are time dependent. Let us suppose that the Hamiltonian in the Schrödinger picture can be splitted as follows $\hat{H} = \hat{H}_0 + \hat{V}$. Normally \hat{H}_0 describe the free motion of a system, whereas \hat{V} represents its interaction, which could be with an external source. Although it often used in a perturbative approach, the Interaction picture does not require \hat{V} to be small as compared with \hat{H}_0 . Inserting this decomposition of \hat{H} in the unitary operator introduced below Eq.(2):

$$\hat{U} = e^{-\frac{i}{\hbar}t\hat{H}} = e^{-\frac{i}{\hbar}t\hat{H}_0 + \hat{V}} = e^{-\frac{i}{\hbar}t\hat{H}_0}\hat{\Omega}(t) \quad (11)$$

This expression helps us to establish a formula from which operators and states in the interaction picture can be defined. For this, consider a matrix element $\langle \Psi'_S(t) | \hat{A}_S | \Psi_S(t) \rangle$. Taking into account Eq.(4) and (11) we find

$$\langle \Psi'_S(t) | \hat{A}_S | \Psi_S(t) \rangle = \langle \Psi'_H | (e^{-\frac{i}{\hbar}t\hat{H}_0}\Omega(t))^\dagger \hat{A}_S e^{-\frac{i}{\hbar}t\hat{H}_0}\Omega(t) | \Psi_H \rangle \quad (12a)$$

$$= \langle \Psi'_H | \Omega(t)^{-1} \hat{A}_I(t) \Omega(t) | \Psi_H \rangle. \quad (12b)$$

Here the operator in the interaction picture reads

$$\hat{A}_I(t) = e^{+\frac{i}{\hbar}t\hat{H}_0} \hat{A}_S e^{-\frac{i}{\hbar}t\hat{H}_0}, \quad (13)$$

whereas a corresponding state in this picture is

$$|\Psi_I(t)\rangle = \Omega(t) |\Psi_H\rangle. \quad (14)$$

At the level of operators, the connection between the Interaction and the Heisenberg picture is established by inverting Eq.(6) and inserting the resulting \hat{A}_S into Eq.(13). This leads to

$$\hat{A}_I(t) = e^{+\frac{i}{\hbar}t\hat{H}_0} \hat{U} \hat{A}_H(t) \hat{U}^{-1} e^{-\frac{i}{\hbar}t\hat{H}_0}. \quad (15a)$$

$$= e^{+\frac{i}{\hbar}t\hat{H}_0} e^{-\frac{i}{\hbar}t\hat{H}} \hat{A}_H(t) e^{\frac{i}{\hbar}t\hat{H}} e^{-\frac{i}{\hbar}t\hat{H}_0}, \quad (15b)$$

resulting in:

$$\hat{A}_I(t) = \hat{\Omega}(t) \hat{A}_H(t) \hat{\Omega}(t)^{-1}. \quad (16)$$

The time evolution equation for $\hat{A}_I(t)$ can be found as done for $\hat{A}_H(t)$ [see below Eq.(6)]:

$$i\hbar \frac{\partial}{\partial t} \hat{A}_I = [\hat{A}_I, \hat{H}_0]. \quad (17)$$

An equation for $\hat{\Omega}(t)$ can be determined. To this end we invert Eq.(11) and express $\hat{\Omega}(t) = e^{\frac{i}{\hbar} t \hat{H}_0} \hat{U}$. Afterwards we differentiate with respect to times:

$$i\hbar \partial_t \hat{\Omega}(t) = e^{\frac{i}{\hbar} t \hat{H}_0} \left(i\hbar \partial_t \hat{U} \right) - \hat{H}_0 e^{\frac{i}{\hbar} t \hat{H}_0} \hat{U} \quad (18a)$$

$$= \hat{H} e^{\frac{i}{\hbar} t \hat{H}_0} \hat{U} - \hat{H}_0 e^{\frac{i}{\hbar} t \hat{H}_0} \hat{U}, \quad (18b)$$

where Eq.(3) has been used. Using the definition of $\hat{\Omega}(t)$ we end up with

$$i\hbar \frac{\partial}{\partial t} \hat{\Omega}(t) = \hat{V}_I(t) \hat{\Omega}(t). \quad (19)$$

2.3 The In and Out picture: External currents

As a motivation for this section, consider the set-up of most experiments especially in elementary particle and nuclear physics. Several particles approach each other from a macroscopic scale, interact in a microscopical section relative to the de Broglie wavelength of the particles. On this scale vacuum fluctuations can accrue, which influence the incoming particle and even make it impossible to distinguish between fluctuation and particle in itself. Afterwards, the products of the interaction spread up to a macroscopic level, again been clearly distinguishable from each other and surrounding space. Therefore, a good description can be derived by taking the states of the system at the beginning and end as direct products of single particle effectively non-interacting states.

To bring this concept on into our formulation let's consider $V = j(t)\hat{q}(t)$ with $\hat{q}(t)$ as the operator of position and j an external source. This will resulting in an unstable vacuum.

In a system with an external current a pure vacuum state can evolve over time into a multi particle state. Starting with the action of a scalar field Φ with mass m coupled to an external source $j(\underline{x}, t)$

$$I = \int d^4x \mathcal{L}(\Phi, \dot{\Phi}, j) = \int d^4x \left(\frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \frac{1}{2} m^2 \Phi^2 - \Phi j \right). \quad (20)$$

Taking the functional derivative with respect to Φ and setting it to zero, we obtain the equation of motion

$$(\partial^2 + m^2) \Phi = j. \quad (21)$$

To proceed, we quantize our field in a box of volume V and length L . The field and its canonical momentum $\Pi = \frac{\partial \mathcal{L}}{\partial \dot{\Phi}(\underline{x}, t)}$ are then promoted to operators $\hat{\Phi}(\underline{x}, t)$ and $\hat{\Pi}(\underline{x}, t)$ in the Heisenberg picture. Satisfying the equal-time commutator:

$$[\hat{\Phi}(\underline{x}, t), \hat{\Pi}(\underline{x}', t)] = i\delta^3(\underline{x} - \underline{x}'). \quad (22)$$

We then expand the field operator as follows:

$$\hat{\Phi}(\underline{x}, t) = \sum_{\underline{k}} \hat{q}_{\underline{k}}(t) u_{\underline{k}}(\underline{x}). \quad (23)$$

The 3 dim. wave vector \underline{k} for the modes is defines as $\underline{k} = \frac{2\pi}{L}(n_x, n_y, n_z)$ with $n_i \in \mathbb{Z}$. In this separated time and space dependency, we choose the Fourier basis for $u_{\underline{k}}(\underline{x})$

$$u_{\underline{k}}(\underline{x}) = \frac{1}{\sqrt{V}} e^{i\underline{k} \cdot \underline{x}}, \quad (24)$$

where the volume V provides the required normalization. We remark that $u_{\underline{k}}(\underline{x})$ constitutes an orthonormalized basis in the Hilbert space

$$\int d^3x u_{\underline{k}'}^*(\underline{x}) u_{\underline{k}}(\underline{x}) = \delta_{\underline{k}, \underline{k}'} \quad (25)$$

$$\sum_{\underline{k}} u_{\underline{k}}^*(\underline{x}) u_{\underline{k}}(\underline{x}') = \delta^3(\underline{x} - \underline{x}'). \quad (26)$$

We now substitute Eq.(24) into the equation of motion (21) :

$$\sum_{\underline{k}} \left[\left(\frac{\partial}{\partial t^2} - \nabla^2 + m^2 \right) \hat{q}_{\underline{k}}(t) u_{\underline{k}}(\underline{x}) \right] = j(\underline{x}, t). \quad (27)$$

The basis will enable us to take the spacial derivative. As a consequence

$$\sum_{\underline{k}} \left[\ddot{\hat{q}}_{\underline{k}}(t) u_{\underline{k}}(\underline{x}) + \underline{k}^2 \hat{q}_{\underline{k}}(t) u_{\underline{k}}(\underline{x}) + m^2 \hat{q}_{\underline{k}}(t) u_{\underline{k}}(\underline{x}) \right] = j(\underline{x}, t). \quad (28)$$

To get an equation for $\hat{q}_{\underline{k}}(t)$ alone we need to get rid of $u_{\underline{k}}(\underline{x})$ and remove the space dependence in the current. Multiplying with $u_{\underline{k}'}^*(\underline{x})$ and integrating over the whole space we find.

$$\sum_{\underline{k}} \left[\int d^3x u_{\underline{k}'}^* u_{\underline{k}} \left(\ddot{\hat{q}}_{\underline{k}}(t) + (\underline{k}^2 + m^2) \hat{q}_{\underline{k}}(t) \right) \right] = \underbrace{\int d^3x j(\underline{x}, t) \frac{1}{\sqrt{V}} e^{i\underline{k} \cdot \underline{x}}}_{=\tilde{j}(\underline{k}, t)}. \quad (29)$$

After using the orthonormality relation (25) this expression reduces to

$$\ddot{\hat{q}}_{\underline{k}}(t) + \omega_{\underline{k}}^2 \hat{q}_{\underline{k}}(t) = \tilde{j}(\underline{k}, t), \quad (30)$$

where $(\underline{k}^2 + m^2) = \omega_{\underline{k}}^2$.

We now make the assumption that the current vanishes outside a finite time interval. As a consequence Eq.(30) allows us to distinguish between early and late times. For early time Eq.(30) reduces to a homogeneous differential equation. We will call its solution by $\hat{q}_{\underline{k}}(t) \rightarrow \hat{q}_{k,in}(t)$. Explicitly,

$$\hat{q}_{k,in}(t) = \frac{1}{2\omega_{\underline{k}}} \left(\hat{a}_{\underline{k},in} e^{-i\omega_{\underline{k}} t} + \hat{a}_{\underline{k},in}^\dagger e^{i\omega_{\underline{k}} t} \right). \quad (31)$$

where $\hat{a}_{\underline{k},in}$ denotes the annihilation operator, whereas $\hat{a}_{\underline{k},in}^\dagger$ is the corresponding creation operator. At late times Eq.(30) also reduces to a homogeneous type. In this case the solution $\hat{q}_{\underline{k}}(t) \rightarrow \hat{q}_{k,out}(t)$ reads

$$\hat{q}_{k,out}(t) = \frac{1}{2\omega_{\underline{k}}} \left(\hat{a}_{\underline{k},out} e^{-i\omega_{\underline{k}} t} + \hat{a}_{\underline{k},out}^\dagger e^{i\omega_{\underline{k}} t} \right). \quad (32)$$

The solution for $\hat{q}_{\underline{k}}(t)$, at times for which $j(\underline{x}, t)$ is active, would then consist of the homogeneous solution plus a term containing the current:

$$\hat{q}_{\underline{k}}(t) = \hat{q}_{k,in}(t) + \frac{1}{\omega_{\underline{k}}} \int_{-\infty}^t dt' \sin[\omega_{\underline{k}}(t - t')] \tilde{j}(\underline{k}, t'), \quad (33)$$

for late times we again go over to the different homogeneous solution $\hat{q}_{k,out}(t)$ and take the integral to $+\infty$.

$$\hat{q}_{k,out}(t) = \hat{q}_{k,in}(t) + \frac{1}{\omega_{\underline{k}}} \int_{-\infty}^{\infty} dt' \sin[\omega_{\underline{k}}(t - t')] \tilde{j}(\underline{k}, t'). \quad (34)$$

Now we will apply a second Fourier transformation in respect to t . The commutation between $\underline{k}, \underline{x}, t, E$ ensures no problem to do these transformations separate and we move the \underline{k} dependence to a index notation.

$$\bar{j}_{\underline{k}}(E) = \int_{-\infty}^{\infty} dt \tilde{j}_{\underline{k}}(t) e^{iEt}. \quad (35)$$

The wave vector \underline{k} and the energy written in terms of the angular frequency ω_k together form the relativistic four-wave-vector $k^\mu = (\omega_{\underline{k}}, k_x, k_y, k_z)$ or just k . This gives us the notation $\bar{j}_{\underline{k}}(\omega_k) = j_k$.

After splitting the sinus function we write:

$$\hat{q}_{\underline{k},out}(t) = \hat{q}_{\underline{k},in}(t) - \frac{i}{2\omega_{\underline{k}}} e^{i\omega_{\underline{k}}t} \bar{j}_{\underline{k}}(-\omega_{\underline{k}}) + \frac{i}{2\omega_{\underline{k}}} e^{-i\omega_{\underline{k}}t} \bar{j}_{\underline{k}}(\omega_{\underline{k}}) \quad (36)$$

From this equation we can obtain the connection between creation and annihilation operators in compact notation.

$$\hat{a}_{\underline{k},out} = \hat{a}_{\underline{k},in} + i j_k \quad (37a)$$

$$\hat{a}_{\underline{k},out}^\dagger = \hat{a}_{\underline{k},in}^\dagger - i j_k \quad (37b)$$

This shows that, in the presence of an external current, the two sets of second quantization operators are not the same. Therefore we need to differ between the corresponding *in* and *out* eigenstates. Especially, it has to be stated that vacua also differ in this scenario. The concept of early and later times to fully solve the full equations will be discussed further in later chapters.

Coming back to the idea of pictures, we incorporate these two sets of solutions as a starting point. Two new pictures should embody the influence of external currents along our boundary conditions we established.

Taking these new results into account, we need to impose for the case $V = j(t)\hat{q}(t)$ from Eq.(11) in beginning a similar approach. On the realm of pictures this means asymptotic conditions on $j(t)$. We again assume it vanishes outside of finite intervals.

$$j(t) \rightarrow 0 \text{ for } t \rightarrow \pm\infty. \quad (38)$$

The effect of these conditions on Eq.(19) leads to

$$\Omega(\pm\infty) \rightarrow 1 \quad (39)$$

The early time condition defines the *In* picture in reminiscence to the first homogeneous equation and it writes :

$$i\hbar \frac{\partial}{\partial t} \hat{\Omega}_{in}(t) = \hat{V}_{in}(t) \hat{\Omega}_{in}(t). \quad (40)$$

Fulfilling:

$$\hat{\Omega}_{in}(-\infty) = 1. \quad (41)$$

The so called *Out* picture will fulfill the condition for late times:

$$i\hbar \frac{\partial}{\partial t} \hat{\Omega}_{out}(t) = \hat{V}_{out}(t) \hat{\Omega}_{out}(t). \quad (42)$$

Stating:

$$\hat{\Omega}_{out}(+\infty) = 1. \quad (43)$$

The Interaction picture as introduced above does not require V to be of any special form but can still be applied for external source terms. To distinguish between the other two pictures above from the condition is chosen :

$$\hat{\Omega}_I(0) = 1. \quad (44)$$

The connecting operator between the pictures is called scattering operator or S-Matrix. It is defined as :¹

$$S = \Omega_I(\infty). \quad (45)$$

3 Scattering operator

3.1 Solutions for the Interaction picture

The verification of statement (45) will make use of explicit expressions of all Ω .

Starting with the Interaction picture depended on t' and integrate the left-hand side of (19) :

$$\int_{t_0}^t dt' i\hbar \frac{\partial}{\partial t'} \Omega_I(t') = i\hbar [\Omega_I(t) - \Omega_I(t_0)], \quad (46a)$$

¹We will now go to natural units $c = \hbar = 1$ and drop the operator hat for convenience sake

here we state $t_0 = 0$ as it is the condition for this picture and for now $t > 0$.

$$\int_0^t dt' i\hbar \frac{\partial}{\partial t'} \Omega_I(t') = i\hbar [\Omega_I(t) - 1]. \quad (46b)$$

Using this formula and the integral over the right-hand side of (19), we find an expression for $\Omega_I(t)$.

$$\Omega_I(t) = 1 - \frac{i}{\hbar} \int_0^t dt' V_I(t') \Omega_I(t') \quad (47)$$

,since the expression has an Ω_I on the other side we will go on by an iterative approach.

$$\Omega_I(t) = 1 - \frac{i}{\hbar} \int_0^t dt' V_I(t') \cdot \left(1 - \frac{i}{\hbar} \int_0^{t'} dt'' V_I(t'') \Omega_I(t'') \right) \quad (48a)$$

$$= 1 - \frac{i}{\hbar} \int_0^t dt' V_I(t') + \left(\frac{i}{\hbar}\right)^2 \int_0^t dt' \int_0^{t'} dt'' V_I(t'') \Omega_I(t''). \quad (48b)$$

The iteration will lead us by single digit incrementation in the power and number of integrals. An problematic aspect of this series are the different integral borders. Each term introduces a new t and keeps the previous as an integral boundary which forces us to solve them in a strict order. To keep the different t as variables over the same integral \int_0^t , we use the chronological time ordering.

$$T(V(t_1), V(t_2)) = V(t_1) V(t_2) \theta(t_1 - t_2) + V(t_2) V(t_1) \theta(t_2 - t_1). \quad (49)$$

In chronological time ordering the Heaviside-Step-function is used to set the earlier times to the right and later to the left by checking the difference. The Step-function is 0 for negative values and 1 for positive.²

For convince sake we drop \hbar again, t will be labelled with numbers instead of primes and time dependency will be indicated by index : $V(t_1) \rightarrow V_1$.

The solution to $\Omega_I(t)$ written as an infinite sum and time ordered :

$$\Omega_I(t) = \sum_{n=0}^{\infty} (-i)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n V_I(t_1) \cdot \dots \cdot V_I(t_n) \quad (50a)$$

$$= \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_n} dt_n T \{V_I(t_1), \dots, V_I(t_n)\} \quad (50b)$$

²Proofs and elaborations for time ordering are to be found in the Appendices

The factorial sum over n converges to exponential function as seen in calculus and this non perturbativ expression reads:

$$\Omega_I(t) = T \left(e^{-i \int_0^t dt' V_I(t')} \right), \text{ for } t > 0. \quad (51)$$

We needed to split the solution of $\Omega_I(t)$, since the condition for the picture is not on the limits of the time span. Assuming $t < 0$ changes the first integral to :

$$\int_t^0 dt' i\hbar \frac{\partial}{\partial t'} \Omega_I(t') = i\hbar [1 - \Omega_I(t)] \quad (52)$$

The sign difference to the expression for $t > 0$ can be fixed by switching the integral borders. This means our infinite sum expression still holds:

$$\Omega_I(t) = \sum_{n=0}^{\infty} (-i)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n V_I(t_1) \cdot \dots \cdot V_I(t_n). \quad (53)$$

The key difference now stands in the negativity of all t and a logical order for them would prefer later times to the right, coming closer to 0. This requires the anti-chronological time ordering :

$$\bar{T}(V(t_1), V(t_2)) = V(t_2) V(t_1) \theta(t_1 - t_2) + V(t_1) V(t_2) \theta(t_2 - t_1). \quad (54)$$

Using it similar as before:

$$\Omega_I(t) = \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n \bar{T} \{V_I(t_1), \dots, V_I(t_n)\}. \quad (55)$$

Converging to the second expression:

$$\Omega_I(t) = \bar{T} \left(e^{-i \int_0^t dt' V_I(t')} \right), \text{ for } t < 0. \quad (56)$$

These two cases make the use of Ω_I safe in a sense of not having to watch out for sign flip in the integral while t runs. Second, the picture condition is clearly stated and not possible to hit while performing integration over t from negative to positive. But to perform the iterative solution ones only needs to state Ω_I being finite in the region of integration. Lowering the integral boundary to $-\infty$ instead of 0. Losing the advantages from above but gaining the single expression:

$$\Omega_I(t) = T \left(e^{-i \int_{-\infty}^t dt' V_I(t')} \right). \quad (57)$$

3.2 Solutions for the In and Out picture

Beginning with the In picture and eq.(40) we reach:

$$\int_{t_0}^t dt' i\hbar \frac{\partial}{\partial t'} \Omega_{in}(t') = i\hbar [\Omega_I(in) - \Omega_{in}(t_0)] \quad (58a)$$

,here we state $t_0 = -\infty$

$$\int_{-\infty}^t dt' i\hbar \frac{\partial}{\partial t'} \Omega_{in}(t') = i\hbar [\Omega_{in}(t) - 1]. \quad (58b)$$

The condition of the In picture being a limit for the time span spares us any splitting. Following the first solution of the Interaction picture:

$$\Omega_{in}(t) = \sum_{n=0}^{\infty} (-i)^n \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 \dots \int_{-\infty}^{t_{n-1}} dt_n V_{in}(t_1) \cdot \dots \cdot V_{in}(t_n) \quad (59a)$$

$$= \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 \dots \int_{-\infty}^{t_n} dt_n T \{V_{in}(t_1), \dots, V_{in}(t_n)\}. \quad (59b)$$

Converging to :

$$\Omega_{in}(t) = T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right). \quad (60)$$

The Out picture on the other hand follows the second solution. Here we argue t being smaller then ∞ needs one change of sign like before and anti-chronological ordering, since t only coming closer to the limit as it runs.

$$\Omega_{out}(t) = \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_{\infty}^t dt_1 \int_{\infty}^{t_1} dt_2 \dots \int_{\infty}^{t_n} dt_n \bar{T} \{V_{out}(t_1), \dots, V_{out}(t_n)\}. \quad (61)$$

Concluding :

$$\Omega_{out}(t) = \bar{T} \left(e^{-i \int_{\infty}^t dt' V_{in}(t')} \right). \quad (62)$$

3.3 Connections

Using this set of solution, the first set of relations to be proven is:

$$S = \Omega_{in}(t) \Omega_{out}(t)^{-1} \quad (63a)$$

$$= \Omega_{in}(\infty) \quad (63b)$$

$$= \Omega_{out}(-\infty)^{-1}. \quad (63c)$$

Writing the first equality out,

$$S = T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) \cdot \bar{T} \left(e^{-i \int_{\infty}^t dt'' V_{out}(t'')} \right) \quad (64a)$$

$$= \sum_n (-i)^n \int_{-\infty}^t dt_1 V_{in}(t_1) \dots \int_{-\infty}^{t_{n-1}} dt_n V_{in}(t_n) \quad (64b)$$

$$\cdot \left(\sum_n (-i)^n \int_{\infty}^t dt'_1 V_{out}(t'_1) \dots \int_{\infty}^{t'_{n-1}} dt'_n V_{out}(t'_n) \right)^{-1} \quad (64c)$$

As an unitar operator $\Omega_{out}(t)^{-1} = \Omega_{out}(t)^\dagger$

$$= T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) \cdot T \left(e^{+i \int_{\infty}^t dt'' V_{out}(t'')} \right) \quad (64d)$$

The lack of overlap in the integral limits allows us to fuse T

$$= T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \cdot e^{+i \int_{\infty}^t dt'' V_{out}(t'')} \right) \quad (64e)$$

Commutation in T allow ease use of the Baker-Campbell-Hausdorff-formula

$$= T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t') - i \int_t^{\infty} dt'' V_{out}(t'')} \right). \quad (64f)$$

In addition the overlap makes S time-independent. Therefore t could be ∞ :

$$S = T \left(e^{-i \int_{-\infty}^{\infty} dt' V_{in}(t') - i \int_{\infty}^{\infty} dt'' V_{out}(t'')} \right) = T \left(e^{-i \int_{-\infty}^{\infty} dt' V_{in}(t')} \right) = \Omega_{in}(\infty). \quad (65)$$

Or we choose $-\infty$:

$$S = T \left(e^{-i \int_{-\infty}^{-\infty} dt' V_{in}(t') - i \int_{-\infty}^{-\infty} dt'' V_{out}(t'')} \right) = T \left(e^{-i \int_{-\infty}^{-\infty} dt' V_{out}(t')} \right) = \Omega_{out}(-\infty)^{-1}. \quad (66)$$

Eq. (65) is very similar to the definition in eq.(45) for Ω_I . This comes from V_I and V_{in} both being picture transformations of V_H in the Heisenberg picture. We discussed the main difference lies in position of the condition on the time span. The time independence and integral over all t makes them equal.

$$S = T \left(e^{-i \int_{-\infty}^{\infty} dt V_I(t)} \right). \quad (67)$$

The next set of relations is :

$$\Omega_{in}(t)\Omega_I(t)^{-1} = S \Omega_I(\infty)^{-1} \quad (68a)$$

$$= \Omega_{in}(0). \quad (68b)$$

The first one is just 1, since the definition was $S = \Omega_I(\infty)$ was Using the exponential formulae:

$$\Omega_{in}(t)\Omega_I(t)^{-1} = T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) \left(\bar{T} \left(e^{-i \int_0^t dt' V_I(t')} \right) \right)^{-1} \quad (69a)$$

$$\stackrel{T \rightarrow \bar{T}}{=} T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) T \left(e^{+i \int_0^t dt' V_I(t')} \right) \quad (69b)$$

$$= T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} e^{+i \int_0^t dt' V_I(t')} \right) \quad (69c)$$

$$\stackrel{CBH}{=} T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t') - i \int_t^0 dt' V_I(t')} \right) \quad (69d)$$

$$= \Omega_{in}(0)\Omega_I(0) \stackrel{\Omega_I(0)=1}{=} \Omega_{in}(0) \quad (69e)$$

This last expression is time independent and is also 1 by looking at the differential equation for Ω_{in} .

The last relation is :

$$\Omega_{in}(t) = \bar{T} \left(e^{i \int_t^\infty dt' V_{in}(t')} \right) S. \quad (70)$$

First it must satisfy the differential equation for Ω_{in} :

$$i\partial_t \Omega_{in}(t) = i\partial_t \left(\sum_n i^n \int_t^\infty dt_1 V_{in}(t_1) \dots \int_{t_{n-1}}^\infty dt_n V_{in}(t_n) \right) S \quad (71a)$$

making an integration by part,

$$= i \left(\sum_n i^n \partial_t [\bar{V}_{in}(\infty) - \bar{V}_{in}(t)] \int_{t_1}^\infty dt_2 V_{in}(t_2) \dots \int_{t_{n-1}}^\infty dt_n V_{in}(t_n) \right) S \quad (71b)$$

partial differentiate in respect to t ,

$$= i \left(\sum_n i^n [0 - V_{in}(t)] \int_{t_1}^\infty dt_2 V_{in}(t_2) \dots \int_{t_{n-1}}^\infty dt_n V_{in}(t_n) \right) S \quad (71c)$$

multiplying and convergence,

$$= V_{in}(t) \sum_n \frac{i^{n-1}}{(n-1)!} \int_{t_1}^{\infty} dt_2 \dots \int_{t_{n-1}}^{\infty} dt_n \bar{T}(V_{in}(t_2) \dots V_{in}(t_n)) S \quad (71d)$$

$$= \bar{T} \left(V_{in}(t) e^{i \int_t^{\infty} dt' V_{in}(t')} \right) S \quad (71e)$$

$$= V_{in}(t) \bar{T} \left(e^{i \int_t^{\infty} dt' V_{in}(t')} \right) S \quad (71f)$$

The last step was based on t being the earliest time in the integral and anti-chronological time ordering.

One also has to provide $\Omega_{in}(-\infty) = 1$:

$$\Omega_{in}(t) = \bar{T} \left(e^{i \int_{-\infty}^{\infty} dt' V_{in}(t')} \right) S \quad (72a)$$

$$= (\Omega_{in}(\infty))^{\dagger} S \quad (72b)$$

$$= S^{\dagger} S = S^{-1} S = 1 \quad (72c)$$

3.4 Unitarity of S

This last step required S to be unitar. Beginning with rewriting it.

$$S = T \left(e^{-i \int_{-\infty}^{\infty} dt V_I(t)} \right) \quad (73a)$$

$$= \sum_n \frac{(-i)^n}{(n)!} \int_{-\infty}^{\infty} dt_1 \dots \int_{-\infty}^{\infty} dt_n T(V_{in}(t_1) \dots V_{in}(t_n)) \quad (73b)$$

$$= \sum_n (-i)^n \int_{-\infty}^{\infty} dt_1 \dots \int_{-\infty}^{\infty} dt_n V_{in}(t_1) \dots V_{in}(t_n). \quad (73c)$$

Without variables in the limits of integration the order is arbitrary

$$= \sum_n \frac{(-i)^n}{(n)!} \int_{-\infty}^{\infty} dt_1 \dots \int_{-\infty}^{\infty} dt_n \bar{T}(V_{in}(t_1) \dots V_{in}(t_n)) \quad (73d)$$

$$= \bar{T} \left(e^{-i \int_{-\infty}^{\infty} dt V_I(t)} \right). \quad (73e)$$

Keeping this in mind we express $1 = 1$ as :

$$1 = T(e^0) \quad (74a)$$

$$= T\left(e^{i \int_{-\infty}^{\infty} dt V_I(t) - i \int_{-\infty}^{\infty} dt V_I(t)}\right) \quad (74b)$$

$$\stackrel{\text{CBH}}{=} T\left(e^{i \int_{-\infty}^{\infty} dt V_I(t)} e^{-i \int_{-\infty}^{\infty} dt V_I(t)} e^{\frac{1}{2} [i \int_{-\infty}^{\infty} dt V_I(t), -i \int_{-\infty}^{\infty} dt V_I(t)]}\right) \quad (74c)$$

$$\stackrel{e^{\frac{1}{2} \cdot 0}}{=} T\left(e^{i \int_{-\infty}^{\infty} dt V_I(t)} e^{-i \int_{-\infty}^{\infty} dt V_I(t)}\right) \quad (74d)$$

$$= T\left(e^{i \int_{-\infty}^{\infty} dt V_I(t)}\right) T\left(e^{-i \int_{-\infty}^{\infty} dt V_I(t)}\right) \quad (74e)$$

$$= T\left(e^{i \int_{-\infty}^{\infty} dt V_I(t)}\right) \bar{T}\left(e^{-i \int_{-\infty}^{\infty} dt V_I(t)}\right) \quad (74f)$$

$$= S \cdot S^\dagger \quad (74g)$$

$$= S \cdot S^{-1} \quad (74h)$$

$$= 1 \quad (74i)$$

4 Gell-Mann Low formula

To motivate Gell-Mann Low formula as the important asset, a common way of application and requirement will be laid out.

First, the formula allows us to transform a polynomial , chronological (or anti-chronological) ordered set of operators in the Heisenberg picture to the three pictures connected to the scattering operator.

This strikes as a fundamental step for dealing with the pictures. Furthermore this transition will be made very early when working on many topics of quantum field theory. As most of the time, one would begin with classical mechanics to get to QFT. Starting with the action of your problem in term of classical fields and then apply second quantisation to promote them to operators in the Heisenberg picture. This would be the point to transition and one needs the Gell-Mann Low formula.

Recalling Eq.(16) we can write:

$$Q_H(t) = \Omega_I(t)^{-1} Q_I(t) \Omega_I(t). \quad (75)$$

Since the proof didn't require any specifications on Ω as it only remains as the difference to the Heisenberg operator U . Allowing us to write:

$$Q_H(t) = \Omega_{in}(t)^{-1} Q_{in}(t) \Omega_{in}(t). \quad (76)$$

Expressing Ω_{in} by S as seen above:

$$Q_H(t) = \left(\bar{T} \left(e^{i \int_t^\infty dt' V_{in}(t')} \right) S \right)^{-1} Q_{in}(t) T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) \quad (77)$$

being a unitary operator, we replace -1 by \dagger and apply hermitian conjugation

$$Q_H(t) = \left(\bar{T} \left(e^{i \int_t^\infty dt' V_{in}(t')} \right) S \right)^\dagger Q_{in}(t) T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) \quad (78a)$$

$$= S^\dagger \left(\bar{T} \left(e^{i \int_t^\infty dt' V_{in}(t')} \right) \right)^\dagger Q_{in}(t) T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right) \quad (78b)$$

$$= S^{-1} T \left(e^{-i \int_t^\infty dt' V_{in}(t')} \right) Q_{in}(t) T \left(e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right). \quad (78c)$$

This step used the unitarity of S itself and the appendix for transition to chronological time ordering. Following that it is allowed to merge T due to zero overlap in the boundaries and correct order

$$Q_H(t) = S^{-1} T \left(e^{-i \int_t^\infty dt' V_{in}(t')} Q_{in}(t) e^{-i \int_{-\infty}^t dt' V_{in}(t')} \right). \quad (79)$$

The position of Q_{in} is still not ideal. Changing it will require commutations. Using the infinite series for e and removing time ordering will reduce the problem to commutations between Q_{in} and V_{in} .

$$Q_H(t) = S^{-1} \left(\sum_n (-i)^n \int_{-\infty}^t dt'_1 \int_t^\infty dt'_1 \dots \int_{-\infty}^{t_{n-1}} dt'_n \int_{t_{n-1}}^\infty dt'_n V_{in}(t'_1) \dots V_{in}(t'_n) Q_{in}(t) \right). \quad (80)$$

We see Q_{in} and V_{in} can commute without any problem since Q_{in} depends on t and $t \neq t'$. Moving it to the left and reapply T as well as e :

$$Q_H(t) = S^{-1} \left(\sum_n (-i)^n Q_{in}(t) \int_{-\infty}^t dt'_1 \int_t^\infty dt'_1 \dots \int_{-\infty}^{t_{n-1}} dt'_n \int_{t_{n-1}}^\infty dt'_n V_{in}(t'_1) \dots V_{in}(t'_n) \right) \quad (81a)$$

$$Q_H(t) = S^{-1} T \left(Q_{in}(t) e^{-i \int_{-\infty}^\infty dt' V_{in}(t')} \right). \quad (81b)$$

Identifying the exponential function as S , the result is:

$$Q_H(t) = S^{-1} T (Q_{in}(t) S). \quad (82)$$

Next will be for more than one operator. Starting with the left side in a non trivial time ordering and applying the transformation for each operator:

$$T(Q_H(t_1)Q_H(t_2)\dots) = T\left(\prod_j \Omega_{in}(t_j)^{-1}Q_{in}(t_j)\Omega_{in}(t_j)\right) \quad (83a)$$

,inserting the expression with S for a Ω depending on t_j ,

$$= T\left(\prod_j S^{-1}T\left(e^{-i\int_{t_j}^{\infty} dt'_j V_{in}(t'_j)}\right)Q_{in}(t_j)T\left(e^{-i\int_{-\infty}^{t_j} dt'_j V_{in}(t'_j)}\right)\right) \quad (83b)$$

,the series and commutation follows the same argumentation as for a single Q ,

$$T(Q_H(t_1)Q_H(t_2)\dots) = T\left(\prod_j S^{-1}Q_{in}(t_j)S\right). \quad (84)$$

Moving S outside of T gives the final form of the Gell-Mann Low formula:

$$T(Q_H(t_1)Q_H(t_2)\dots) = S^{-1}T(Q_{in}(t_1)Q_{in}(t_2)\dots S). \quad (85)$$

To proof S can be outside of chronological time ordering we look at $T(S^{-1})$:

$$T(S^{-1}) = T(S^\dagger) \quad (86a)$$

,using the expression of S in term of the In-picture and apply hermitian conjugation,

$$T(S^{-1}) = T\left(\bar{T}\left(e^{i\int_{-\infty}^{\infty} dt' V_{in}(t')}\right)\right) \quad (86b)$$

, e^x as a series,

$$= T\left(\sum_n \frac{(i)^n}{n!} \int_{-\infty}^{\infty} dt_1 \int_{-\infty}^{\infty} dt_2 \dots \int_{-\infty}^{\infty} dt_n \bar{T}(V_{in}(t_1)\dots V_{in}(t_n))\right) \quad (86c)$$

and removing the anti-chronological time ordering,

$$= T\left(\sum_n \frac{(i)^n}{n!} i^n \int_{-\infty}^{\infty} dt_1 \int_{-\infty}^{\infty} dt_2 \dots \int_{-\infty}^{\infty} dt_n V_{in}(t_1)\dots V_{in}(t_n)\right). \quad (86d)$$

The boundaries make every integral independent from each other so rearranging t_1 to t_n wouldn't change the result

$$\rightarrow T(S^{-1}\dots) = S^{-1}T(\dots)$$

4.1 Vacuum expectation value

The Gell-Mann Low formula eq.(85) plays a central role in many perturbative or non-perturbative calculations. In particular by evaluating it sandwiched between eigenstates to rising and lowering operators in the In- and Out-picture. If they are in the groundstate of the theory aka. the vacuum state, we obtain the vacuum expectation value.

$$\langle 0_{out} | T(Q_H(t_1)Q_H(t_2) \dots) | 0_{in} \rangle = \quad (87a)$$

$$= \langle 0_{out} | S^{-1} T(Q_{in}(t_1)Q_{in}(t_2) \dots S) | 0_{in} \rangle \quad (87b)$$

$$= \langle 0_{out} | S^{-1} T\left(Q_{in}(t_1)Q_{in}(t_2) \dots e^{-i \int_{-\infty}^{\infty} dt' V_{in}(t')}\right) | 0_{in} \rangle \quad (87c)$$

This average (expectation value) of a product of n operators is called the n -point correlation function. For example, using Wick's theorem and expansions in powers of $V_{in}(Q_{in}(t), t)$, we can gain knowledge about processes that involves in total n excitations of the ground state (or vacuum) in both asymptotic states.

At this point one can be in two situations. The ground state can be either stable or unstable. Over time it could spontaneously go into a state of excitation. Mathematically speaking instability leads to :

$$\langle 0_{out} | 0_{in} \rangle \neq 1. \quad (88)$$

Here of course already normalized states. This possibility for excitations stem from external sources and therefore external currents. A stable state would lead to:

$$\langle 0_{out} | S^{-1} = \langle 0_{in} | \quad (89)$$

An asymptotic state would go over time into a asymptotic state of same excitation.

5 Conclusion

In this report we have presented a set of fundamental concepts and methods. These are essential for starting to study Quantum field theory. We began with well-known pictures from quantum mechanics and introduced the In and Out picture as consequences of working with asymptotic states. They lead us to the important connecting element between the pictures, the scattering operator. The connections allowed us to derive the Gell-Mann Low formula, which plays a major part in the second quantisation process one would do in quantum field theories.

5.A Vacuum transition probability under external current

Still, we need a way to transition from one operator to another and from eigenstate to eigenstate. This link is called the scattering operator S .

$$|out\rangle = S^{-1} |in\rangle \quad (90a)$$

$$|in\rangle = S |out\rangle. \quad (90b)$$

The scalar product of the mentioned two vacua can be written as :

$$\langle 0_{out} | 0_{in} \rangle = \langle 0_{out} | S | 0_{out} \rangle \quad (91)$$

These applications for the operator require S to be of the form

$$\begin{aligned} S &= e^{i\Lambda} \\ S^\dagger &= e^{-i\Lambda^\dagger}, \end{aligned} \quad (92)$$

since $S = S^\dagger$ should hold $\rightarrow \Lambda = -\Lambda^\dagger$. We choose for Λ the chronological time-ordered sum $\sum_{\underline{k}} \int dt \bar{j}_{\underline{k}}(t) \hat{q}_{\underline{k},in}(t)$.

Explicitly,

$$S = T \left[\exp \left(i \sum_{\underline{k}} \int dt \bar{j}_{\underline{k}}(t) \hat{q}_{\underline{k},in}(t) \right) \right] \quad (93a)$$

$$\stackrel{\text{Eq.(31)}}{=} T \left[\exp \left(i \sum_{\underline{k}} \int dt \bar{j}_{\underline{k}}(t) \frac{1}{2\omega_{\underline{k}}} \left(\hat{a}_{\underline{k},in} e^{-i\omega_{\underline{k}}t} + \hat{a}_{\underline{k},in}^\dagger e^{i\omega_{\underline{k}}t} \right) \right) \right], \quad (93b)$$

using the the CBH-formula allows us to separate,

$$\begin{aligned} S = & T \left[\exp \left(i \sum_{\underline{k}} \int dt \bar{j}_{\underline{k}}(t) \frac{1}{2\omega_{\underline{k}}} \left(\hat{a}_{\underline{k},in} e^{-i\omega_{\underline{k}}t} \right) \right) \right. \\ & \cdot \exp \left(i \sum_{\underline{k}} \int dt \bar{j}_{\underline{k}}(t) \frac{1}{2\omega_{\underline{k}}} \left(\hat{a}_{\underline{k},in}^\dagger e^{i\omega_{\underline{k}}t} \right) \right) \\ & \cdot \exp \left(-i \sum_{\underline{k}} \int dt \bar{j}_{\underline{k}}(t) \frac{1}{2 \cdot 2\omega_{\underline{k}}} \left[\hat{a}_{\underline{k},in} e^{-i\omega_{\underline{k}}t}, \hat{a}_{\underline{k},in}^\dagger e^{i\omega_{\underline{k}}t} \right] \right) \left. \right]. \end{aligned} \quad (94)$$

The integral over t will Fourier transform $\bar{j}_{\underline{k}}(t)$ into our relativistic notation,

$$S = \left[\exp \left(i \sum_{\underline{k}} \frac{1}{2\omega_{\underline{k}}} j_{-k\mu} \hat{a}_{\underline{k},in} \right) \cdot \exp \left(i \sum_{\underline{k}} \frac{1}{2\omega_{\underline{k}}} j_{k\mu} \hat{a}_{\underline{k},in}^\dagger \right) \cdot \exp \left(-i \sum_{\underline{k}} \frac{1}{4\omega_{\underline{k}}} |j_{k\mu}|^2 \right) \right]. \quad (95)$$

We dropped the time ordering since the transformation made it time independent.

For now we choose S in a more specific form, where we used $\bar{j}_{\underline{k}}(-\omega_{\underline{k}}) = \bar{j}_{\underline{k}}^*(\omega_{\underline{k}})$. The general one will be obtained later:

$$S = e^{\frac{i}{2\omega_{\underline{k}}} \bar{j}_{\underline{k}}(\omega_{\underline{k}}) \hat{a}_{\underline{k},in}^\dagger} e^{\frac{i}{2\omega_{\underline{k}}} \bar{j}_{\underline{k}}(-\omega_{\underline{k}}) \hat{a}_{\underline{k},in}} e^{-\frac{1}{4\omega_{\underline{k}}} |\bar{j}_{\underline{k}}(-\omega_{\underline{k}})|^2} \quad (96)$$

For the operators we can write:

$$S^{-1} \hat{a}_{\underline{k},in} S = \hat{a}_{\underline{k},out} = \hat{a}_{\underline{k},in} + i \bar{j}_{\underline{k}}(\omega_{\underline{k}}) \quad (97)$$

Using this, we obtain first of all the probability for staying in the ground state for one mode:

$$p_{0,k} = |\langle 0_{out} | 0_{in} \rangle|^2 = |\langle 0_{in} | S | 0_{in} \rangle|^2 \quad (98a)$$

$$= \left| \langle 0_{in} | e^{\frac{i}{2\omega_{\underline{k}}} \bar{j}_{\underline{k}}(\omega_{\underline{k}}) \hat{a}_{\underline{k},in}^\dagger} e^{\frac{i}{2\omega_{\underline{k}}} \bar{j}_{\underline{k}}(-\omega_{\underline{k}}) \hat{a}_{\underline{k},in}} e^{-\frac{1}{4\omega_{\underline{k}}} |\bar{j}_{\underline{k}}(-\omega_{\underline{k}})|^2} | 0_{in} \rangle \right|^2. \quad (98b)$$

The annihilation operator will return a 0 in the exponent. Therefore only one factor remain important. The probability while taking to account any mode transition just involves the integral over all k :

$$p_0 = \exp \left\{ - \int \frac{d^3 k}{(2\pi)^3} \left| \frac{\bar{j}_{\underline{k}}(\omega_{\underline{k}})}{\sqrt{2\omega_{\underline{k}}}} \right|^2 \right\}. \quad (99)$$

This confirms our statement about an unstable vacuum. The negative sign in the exponent translates to smaller probability at higher external currents. So the vacuum can change it's state if a current is > 0 .

5.B Chronological time ordering

Let's begin with at a regular function in form of:

$$I(t) = \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V(t_1) V(t_2) \quad (100)$$

We see that although the the right hand side has functions of t_2 and t_1 the left hand side doesn't. The integrals allow us to perform a change of variable and rewrite I as $I(t) = \frac{1}{2}I(t) + \frac{1}{2}I(t)$:

$$I(t) = \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V(t_1) V(t_2) + \frac{1}{2} \int_{-\infty}^t dt_2 \int_{-\infty}^{t_2} dt_1 V(t_2) V(t_1). \quad (101)$$

We define the **chronological time ordering**³ at this point to equalize the boundaries.

Note $V(t_1) \rightarrow V_1$:

$$T(V_1, V_2) = V_1 V_2 \theta(t_1 - t_2) + V_2 V_1 \theta(t_2 - t_1). \quad (102)$$

Being based on the Heaviside step-function it allows us to write the Eq.(100) as:

$$I(t) = \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_1 V_2 \theta(t_1 - t_2) + \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_2 V_1 \theta(t_2 - t_1) \quad (103)$$

$$I(t) = \frac{1}{2!} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 T(V_1, V_2) \quad (104)$$

The Heaviside function allows us to set terms to zero for negative arguments of θ and 1 for positive. By subtracting t_1 and t_2 we will be able to switch between the later and earlier points in time.

³earlier times to the right and later times to the left

for $t_1 > t_2 \rightarrow \theta(t_1 - t_2)$ and for $t_2 > t_1 \rightarrow \theta(t_2 - t_1)$

To expand our concept to cases of more then two t we start by advancing T . In general it consists of a summation of all permutations P of a given set multiplied by Heaviside functions. These Heaviside functions have arguments with the negative summation of the t in the same permutation P .

Definition:

$$T(V_1, V_2, \dots, V_n) = \sum_{j=1}^{n!} P_j [V_1, V_2, \dots, V_n] \cdot \theta \left(P_j \left[t_j - \sum_{i \neq j}^n t_i \right] \right) . \quad (105)$$

But this summation over j would not add up to the same value for $I(t)$ since it started in the order of one element in the sum. In this case we need a normalization in addition to T . From statistics we know a set of n different elements can be linear arranged in $n!$ ways. Coming in as a factor of $\frac{1}{n!}$ in the expressions later.

For a test we use $n = 3$

$$T(V_1, V_2, V_3) = \sum_{j=1}^{3!} P_j [V_1, V_2, V_3] \cdot \theta \left(P_j \left[t_j - \sum_{i \neq j}^3 t_i \right] \right) \quad (106a)$$

$$= V_1 V_2 V_3 \cdot \theta(t_1 - t_2 - t_3) \quad (106b)$$

$$+ V_1 V_3 V_2 \cdot \theta(t_1 - t_3 - t_2) \quad (106c)$$

$$+ V_2 V_1 V_3 \cdot \theta(t_2 - t_1 - t_3) \quad (106d)$$

$$+ V_2 V_3 V_1 \cdot \theta(t_2 - t_3 - t_1) \quad (106e)$$

$$+ V_3 V_1 V_2 \cdot \theta(t_3 - t_1 - t_2) \quad (106f)$$

$$+ V_3 V_2 V_1 \cdot \theta(t_3 - t_2 - t_1) \quad (106g)$$

Now we can apply it to new I and do a proof by Induction based on the number of V . The Induction start is the case of two V . In the Induction step we say in a mathematicians way that it works for at least one unspecified higher order. Let's call it k : (Note: $t_0 = t$)

$$I_k(t) = \prod_{a=1}^k \int_{-\infty}^{t_{a-1}} dt_a V_a = \frac{1}{k!} \left(\prod_{a=1}^k \int_{-\infty}^t dt_a \right) T(V_1, \dots, V_k). \quad (107)$$

Moving one increment higher in our 'chain' $k + 1$,

$$I_{k+1}(t) = \prod_{a=1}^{k+1} \int_{-\infty}^{t_{a-1}} dt_a V_a = \prod_{a=1}^k \int_{-\infty}^{t_{a-1}} dt_a V_a \cdot \int_{-\infty}^{t_k} dt_{k+1} V_{k+1} \quad (108)$$

Using the Induction Step and general definition for T :

$$I_{k+1}(t) = \frac{1}{k!} \left(\prod_{a=1}^k \int_{-\infty}^t dt_a \right) T(V_1, \dots, V_k) \cdot \int_{-\infty}^{t_k} dt_{k+1} V_{k+1} \quad (109)$$

$$I_{k+1}(t) = I_k(t) \cdot \int_{-\infty}^{t_k} dt_{k+1} V_{k+1} \quad (110)$$

This shows that the incrementation of k reduces to a multiplication with one more different element for the set. This increases the possible permutations by a factor of $k + 1$ resulting in $(k + 1)!$ in total. Giving us:

$$I_{k+1}(t) = \frac{1}{(k + 1)!} \left(\prod_{a=1}^{k+1} \int_{-\infty}^t dt_a \right) T(V_1, \dots, V_{k+1}) \quad (111)$$

Time ordering is a very grounded concept in field theory, it appears very naturally for expressing propagators in term of fields. The reason apart from pure mathematics is comprehensible. The different functions, functionals or fields are best organized for summarizing a scattering or interaction event if they are time-like sorted.

Furthermore it comes with a great advantage. All products in $T(\dots)$ do commute. Proof for two elements:

$$T(V_1, V_2) = V_1 V_2 \theta(t_1 - t_2) + V_2 V_1 \theta(t_2 - t_1) \quad (112a)$$

$$T(V_2, V_1) = V_2 V_1 \theta(t_2 - t_1) + V_1 V_2 \theta(t_1 - t_2) \quad (112b)$$

,since terms in sums always commutes,

$$T(V_1, V_2) = T(V_2, V_1) \quad (113)$$

In other words, the commutation relations say whether the subtraction of permutations of elements is zero or not. But in time ordering all permutations appear, we can rearrange the terms so subtraction of equal permutations happens. Therefore commutation in T holds.

5.C Anti-chronological time ordering

On a close inspection of the introduction of the Heaviside-function and its' insertion into the integral, we actually skipped a choice. If we just would have wanted to the overall boundaries the order of t_1 and t_2 in relation to V_1 and V_2 the insertion of θ could have been switched. This secretly led us to the definition of time-ordering or chronological time ordering as it is called more precisely. The other path would have resulted in anti-chronological time ordering:⁴

$$\bar{T}(V_1, V_2) = V_1 V_2 \theta(t_2 - t_1) + V_2 V_1 \theta(t_1 - t_2) \quad (114)$$

This section is not just to satisfy the observed readers but to proof a connection between both involving hermitian conjugation of the integrals, which can easily appear while doing picture transitions.

We choose the same starting point as for T :

$$\begin{aligned} I(t) &= \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_1 V_2 + \frac{1}{2} \int_{-\infty}^t dt_2 \int_{-\infty}^{t_2} dt_1 V_2 V_1 \\ &= \frac{1}{2!} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 T(V_1, V_2) \end{aligned}$$

$$I(t)^\dagger = \left(\frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_1 V_2 + \frac{1}{2} \int_{-\infty}^t dt_2 \int_{-\infty}^{t_2} dt_1 V_2 V_1 \right)^\dagger \quad (115a)$$

$$= \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 (V_1 V_2)^\dagger + \frac{1}{2} \int_{-\infty}^t dt_2 \int_{-\infty}^{t_2} dt_1 (V_2 V_1)^\dagger \quad (115b)$$

$$= \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_2^\dagger V_1^\dagger + \frac{1}{2} \int_{-\infty}^t dt_2 \int_{-\infty}^{t_2} dt_1 V_1^\dagger V_2^\dagger \quad (115c)$$

, we observe that the switch of positions due to hermitian conjugation allows to use Eq.(114)

$$I(t)^\dagger = \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_2^\dagger V_1^\dagger \theta(t_1 - t_2) + \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 V_1^\dagger V_2^\dagger \theta(t_2 - t_1) \quad (115d)$$

$$\Rightarrow \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^{t_1} dt_2 \bar{T}(V_1^\dagger, V_2^\dagger) \quad (115e)$$

⁴earlier times to the left and later times to the right

Now we assume our total Hamiltonian $\hat{H} = \hat{H}_0 + \hat{V}$ is hermitian and \hat{V} won't lead to non fixed ground state energy \rightarrow vacuum instability

$$I(t)^\dagger = \frac{1}{2} \int_{-\infty}^t dt_1 \int_{-\infty}^t dt_2 \bar{T}(V_1, V_2) \quad (115f)$$

References

- [1] S.Randjbar-Daemi. *Course on Quantum Electrodynamics: Introduction to Quantum Field Theory*. The Abdus Salam International Centre for Theoretical Physics , 2007-2008.
- [2] M.Peskin; D.Schroeder. *Quantum field theory*. Perseus Books Publishing, 1995.
- [3] W.Greiner;J.Reinhart. *Quantum electrodynamics*. Verlag Harri Deutsch Thun und Frankfurt am Main, 1984.
- [4] W.Greiner;J.Reinhart. *Feldquantisierung*. Verlag Harri Deutsch Thun und Frankfurt am Main, 1993.