


# Research Review: Do antibullying interventions reduce internalizing symptoms? A systematic review, meta-analysis, and meta-regression exploring intervention components, moderators, and mechanisms

Carolina Guzman-Holst,<sup>1</sup> Mirela Zaneva,<sup>1</sup> Chloe Chessell,<sup>2</sup> Cathy Creswell,<sup>1,3</sup>  and Lucy Bowes<sup>1</sup>

<sup>1</sup>Department of Experimental Psychology, University of Oxford, Oxford, UK; <sup>2</sup>School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK; <sup>3</sup>Department of Psychiatry, University of Oxford, Oxford, UK

**Background:** Effective antibullying interventions may reduce the impact of bullying on young people's mental health. Nevertheless, little is known about their effectiveness in reducing internalizing symptoms such as anxiety or depression, and what factors may influence intervention effects. The aim of this systematic review, meta-analysis, and metaregression is to assess the effects of school-based antibullying interventions on children's and adolescent's internalizing symptoms. The secondary aims are to explore potential moderators, intervention components, and reductions in bullying as mediators of intervention effects on internalizing symptoms. **Methods:** We searched nine databases: PsycINFO, Web of Science, ERIC, SCOPUS, CINAHL, Medline, Embase, ProQuest, and Cochrane Library, and performed an author search of included studies in English from January 1983 to April 2021. We included studies that evaluated school-based antibullying interventions using controlled designs and reporting on both bullying and internalizing outcomes. Random-effects and metaregression models were used to derive Hedges *g* values with pooled 95% CIs as estimates of effect size and to test associations between moderator variables and effect size estimates. Path analysis was used to test potential mediation using effect size measures of victimization, perpetration, and internalizing outcomes. Quality and risk of bias were assessed using Cochrane collaboration tools. **Results:** This review included 22 studies with 58,091 participants in the meta-analysis. Antibullying interventions had a very small effect in reducing overall internalizing symptoms (ES, 0.06; 95% CI, 0.0284 to 0.1005), anxiety (ES, 0.08; 95% CI, 0.011 to 0.158), and depression (ES, 0.06; 95% CI, 0.014 to 0.107) at postintervention. The reduction in internalizing symptoms did not vary significantly across geographic location, grade level, program duration, and intensity. The intervention component 'working with peers' was associated with a significant reduction, and 'using CBT techniques' was associated with a significant increase in internalizing outcomes. Bullying victimization and perpetration did not mediate the relationship between intervention condition and internalizing outcomes. **Conclusions:** Antibullying interventions have a small impact on reducing internalizing symptoms. Ongoing development of antibullying interventions should address how best to maximize their impact on internalizing symptoms to safeguard young people from the damaging mental health outcomes of bullying. **Keywords:** School intervention; antibullying; internalizing; depression; anxiety; children; adolescents.

## Introduction

Depression and anxiety (internalizing) disorders are among the most frequently diagnosed psychiatric conditions in childhood and are the leading causes of disability and burden in young people (Erskine et al., 2015; Ghandour et al., 2019). Longitudinal studies have shown continuity in internalizing problems from childhood to adulthood (Hofstra, Van der Ende, & Verhulst, 2000) and higher lifetime prevalence rates compared to any other psychiatric disorder (Kessler et al., 2005) suggesting the need to identify modifiable risk factors that can be targeted in interventions during early life. Bullying victimization appears to be one of the most tractable risk factors

for internalizing symptoms. Specifically, research has shown that over a quarter of cases of depression at age 18 might be attributable to bullying victimization during early adolescence (Bowes, Joinson, Wolke, & Lewis, 2015). Studies suggest that young people with internalizing difficulties are more likely to be bullied and young people who are bullied are more likely to develop internalizing difficulties, even after adjusting for baseline internalizing difficulties (Arseneault et al., 2008; Singham et al., 2017; Stapinski, Araya, Heron, Montgomery, & Stallard, 2015). If this relationship is causal as it has been suggested, antibullying interventions offer an important way to safeguard young people's mental health.

In relation to its mental health impact, school-based bullying is recognized as a major public health issue. A study by UNESCO (2019) found that 32% of children globally had experienced bullying

Conflict of interest statement: See Acknowledgements for full disclosures.

victimization on one or more days over the past month and 7.3% of young people had experienced bullying on 6 or more days over the same time period (UNESCO, 2019). It is important to highlight that while most children experience low and stable trajectories of peer victimization throughout the school years, other children experience chronic and increasing levels of peer victimization (Barker et al., 2008; Biggs et al., 2010). Indeed, research has shown that this latter group is also the most at risk of prior and subsequent internalizing problems (Sukhawathanakul & Leadbeater, 2020).

A person characterizes bullying repeatedly being exposed to aggressive or negative actions by another person intending to cause physical harm, psychological distress, or humiliation (Olweus, 1993). Different types of bullying include traditional forms such as physical, relational, emotional, and verbal bullying, while other nontraditional forms such as cyber bullying can take place at any time and context. In childhood, traditional bullying victimization and perpetration predominantly occurs within the school context (Department for Education, 2018) and have been associated with higher mean prevalence rates (35%) than nontraditional bullying victimization and perpetration (15%) across 80 studies (Modecki, Minchin, Harbaugh, Guerra, & Runions, 2014). This highlights the important role of schools as grounds for antibullying prevention.

### *Effectiveness of school antibullying interventions*

Many school-based interventions have been devised and implemented to reduce school bullying. Gaffney, Farrington, and Ttofi's (2019) meta-analysis showed that, on average and when compared with school usual practice, these programs decreased bullying by 19%–20% and victimization by 15%–16% (Gaffney et al., 2019). Yet, despite evidence showing an association between bullying victimization and internalizing symptoms (Reijntjes, Kamphuis, Prinzie, & Telch, 2010), not all interventions measure or report their intervention's effects on mental health. For those who do report them, findings have been mixed with some suggesting that interventions can reduce psychological distress while others find no such effects (Bonell et al., 2018; Dempsey, 2009; Hoglund, Hosan, & Leadbeater, 2012; Williford et al., 2012). Recently, Fraguas et al. (2021) conducted a systematic review and meta-analysis assessing the effect of school antibullying interventions on mental health and found an overall effect size using Cohen's *d* of  $-.205$  (95% CI,  $-0.277$  to  $-0.133$ ). However, they only examined randomized controlled trials (RCTs) in published journal articles and used a broad definition of mental health including proxies such as mental health symptoms, quality of life, self-esteem, self-blame, and social skills. It is likely that such broad proxies for mental health resulted in larger effect sizes and more

heterogeneity, but less specificity in identifying intervention effects on internalizing difficulties. Furthermore, they did not look at intervention components or whether reductions in victimization mediated mental health outcomes. This information could aid in determining what specific components improve intervention effects on internalizing and through what mechanisms they potentially operate. A systematic review and meta-analysis to quantify intervention effects on internalizing symptoms is warranted.

### *Moderating and mediating effects of antibullying interventions*

Very little is known about the moderators and mechanisms that play a role in determining intervention effects on internalizing symptoms. Several studies have shown that type of intervention such as whole school versus targeted (Gaffney et al., 2019), age (Fraguas et al., 2021; Yeager, Fong, Lee, & Espelage, 2015), informant type (Shakoor et al., 2012), location (Gaffney et al., 2019), and number of components (Gaffney, Ttofi, & Farrington, 2021a, 2021b) may determine the effectiveness of interventions at reducing bullying. This has not been examined yet in the context of internalizing symptoms and could provide important information about what type of antibullying interventions can improve mental health.

Understanding which specific intervention components are more effective at reducing children's internalizing symptoms can also provide important recommendations on how to maximize the effectiveness of future antibullying programs. In a comprehensive Campbell review, Farrington and Ttofi (2009) found that increased teacher training, increased duration and intensity of the intervention, parental involvement, playground supervision, and using disciplinary methods were associated with a decrease in both bullying victimization and perpetration. Working with peers was associated with an increase in victimization, although this was recently found to not be the case in a follow-up review (Gaffney et al., 2021a, 2021b). This follow-up review found that interventions with a whole-school approach, antibullying policy, classroom rules, information for parents, informal peer involvement, work with victims, cooperative group work, and mental health approaches were associated with larger reductions in bullying perpetration, and informal peer involvement and information for parents were associated with larger reductions in bullying victimization. The *absence* of socioemotional skills components was associated with significantly larger reductions in both perpetration and victimization outcomes. Program richness (i.e. the number of intervention components used) was not significantly associated with changes in perpetration or victimization (Gaffney et al., 2021a, 2021b). It is unclear

whether these components would have similar effects on internalizing outcomes.

In terms of mediating effects, a factor that may explain the difference in intervention's effects on internalizing symptoms is the extent to which the interventions change the prevalence of bullying perpetration and victimization (Palladino, Nocentini, & Menesini, 2019). That is, interventions that achieve greater reductions in bullying victimization should theoretically lead to greater reductions in internalizing symptoms. These mechanisms have not yet been examined across multiple antibullying programs, yet they could inform researchers and public health professionals of how these programs work and target mental health moving forward.

### The current study

This review aims to answer the following questions:

1. Are school-based antibullying interventions associated with improvement or deterioration of internalizing symptoms? Do effect sizes vary according to type of internalizing symptom (anxiety or depression), type of intervention (targeted vs. whole school), age group (children or adolescents), and study quality (low vs. high risk of bias)?
2. Do specific study and participant characteristics such as duration and intensity of intervention, type of informant for internalizing measures, school grade level, location, and number of intervention components moderate the effect of antibullying interventions on internalizing outcomes?
3. Do bullying victimization and perpetration prevalence outcomes mediate the effect of antibullying interventions on internalizing symptoms?
4. Are specific intervention components independently associated with improving or worsening internalizing outcomes?

## Methods

### Protocol and registration

This systematic review and meta-analysis was preregistered with PROSPERO (CRD42020191857) and uses PRISMA reporting guidelines. Deviations from our preregistration protocol are detailed in Appendix S1.

### Search strategy

For this review, we searched nine databases: PsycINFO, Web of Science, ERIC, SCOPUS, CINAHL, Medline, Embase, ProQuest, and Cochrane Library from January 1983, as this is when bullying intervention notably emerged (Olweus, 1993), to May 2021. This search was complemented by an author search and a meta-analysis search to obtain as many evaluations of known intervention programs as possible. To decrease publication bias, we included theses and dissertations in addition to book chapters and journal articles written in English. The specific search strategy is described in Appendix S2.

### Eligibility criteria

We included studies that evaluated an antibullying intervention delivered in schools to children and adolescents aged 4–19 years old. To meet eligibility criteria, the study stated a clear definition of bullying, measured bullying perpetration, or victimization as a primary outcome, and measured internalizing symptoms as either a coprimary or secondary outcome postintervention. Studies reported effect sizes or relevant data that allowed for the computation of effect sizes for bullying and internalizing. Bullying and internalizing outcomes were measured using a validated, standardized, age-appropriate measure (internalizing scales must have been completed by child, parent, or teacher, and show good internal consistency ( $\alpha = .6$  or above), previous evidence of construct validity, and  $\pm 1$  year within the suggested age range). The study design included an experimental group that received the intervention with a control group that did not. The experimental and control group could have been in the same school or in different schools. If there was more than one experimental group (e.g. two groups using different interventions) in the same study, there must have been a control group not receiving any intervention. The types of designs included were as follows: randomized experiments, experimental-control comparisons with pre- and post-measures of bullying and internalizing symptoms between groups, and other experimental-control comparisons. For trials with multiple publications, the most relevant study was included after assessing them against all eligibility criteria. For studies that reported on two or more experimental intervention groups or one or more control groups, we followed Cochrane recommended guidelines (Chapter 16.5.4) and either selected the most relevant group/condition or split the shared group into two or more groups with smaller sample sizes to make independent comparisons.

### Study selection


A flow chart of the study selection process is shown in Figure 1. We identified 10,751 records (9,610 from the first search and 1,141 from the 2020–2011 search) from the electronic databases and an additional 478 records from reviews and author searches from the first search. A total of 5,454 records were retained after duplicates were removed following both searches. Two authors (CGH and MZ) used Raayan Software to independently screen titles and abstracts and identify records for full-text screening. Interrater reliability between the two review authors was calculated and classified as 'almost perfect agreement' ( $k = .85$ ; Landis & Koch, 1977). Disagreements were resolved between two authors (CGH and MZ) and when consensus was not reached, a third review author (LB or CCR) was consulted.

### Data extraction

Two authors (CGH and CCh) independently extracted study characteristics, outcome data, intervention components, and potential moderators from eligible articles. Intervention components were identified based on elements found during article screening in combination with Farrington & Ttofi's (2009) element list. We consulted with the evaluators of the various programs by sending them our coding and description of the elements (see Appendix S3). We received author ratings for 18 of the 22 evaluations included in the meta-analysis (see Table S1). Missing information for key variables from included studies were also requested from authors.

### Assessment of methodological quality (risk of bias)

This review used the Cochrane collaboration's tools for assessing risk of bias for cluster randomized controlled trials (RoB-2



**Figure 1** Prisma flow chart

cluster randomized), individual cluster randomized controlled trial (RoB-2), and cross-over trials (RoB-2 cross-over). Two reviewers (CGH and MZ) independently rated the quality of all included studies. Any discrepancies were initially discussed by the two reviewers and a third review author (LB or CCr) when consensus was not reached.

### Effect measures

Hedge's *g* effect sizes and 95% confidence intervals were considered the primary outcome of the meta-analysis. They were calculated using post-intervention sample size, means, and standard deviation, or regression coefficients from each included study. As many studies involved clustered designs, we corrected the sample size for clustering using Cochrane guidelines. When intraclass correlation coefficients (ICC) were not available, we used an average estimate from the literature (ICC = 0.025). For studies reporting multiple internalizing outcomes, effect sizes, and respective variances were averaged to obtain a single-effect estimate using equations reported elsewhere (López-López, Page, Lipsey, & Higgins, 2018).

### Data synthesis and statistical analysis

For the meta-analysis, main results for overall internalizing symptoms were examined using random-effect models with an inverse-variance approach and DerSimonian-Laird estimator and subgroups were examined using mixed-effect models.

Subgroup analysis included type of internalizing symptom (anxiety or depression), type of intervention (targeted vs. whole school), age group (children or adolescents), and study quality (low vs. high risk of bias). The *Q* and *I*<sup>2</sup> statistics were used to assess heterogeneity between study results. R packages *meta*, *metafor*, and *dmetar* were used.

To assess moderation and intervention component effects, we performed metaregression analysis for each moderator and intervention component determined a priori. For metaregression, it is recommended that each covariate contain at least 10 studies (Borenstein, Hedges, Higgins, & Rothstein, 2011), therefore type of informant was not examined as only 2 studies used a different type of informant that was not self-reported.

Lastly, to examine the extent bullying victimization and perpetration prevalence outcomes mediate intervention effects on internalizing symptoms, we used mean pre-post scores to calculate Hedges' *g* effect sizes in bullying and internalizing outcomes for control and intervention groups, for each study, in a school-level mediation analysis. We calculated the direct effect '*c*' (effect of intervention condition on internalizing), indirect effect '*a\*b*' (effect of intervention on victimization/perpetration and effect of victimization/perpetration on internalizing), and the total effect '*c + (a\*b)*'. To test the significance of the indirect effect, we used bootstrapping procedures. Unstandardized indirect effects were computed for each of 1,000 bootstrapped samples. Two studies (Bonell et al., 2018, and Williford et al., 2012) were not included in the mediation analysis as they did not have baseline measures of internalizing symptoms. These analyses were conducted using the *lavaan* package in R.



## Results

### Search results

A total of 5,454 abstracts were reviewed. Of these, 27 articles were included for full-text review and 22 met criteria for the meta-analysis. The process of study selection is presented in Figure 1.

### Study characteristics

Study characteristics are presented in Table S2. In total, 27 studies conducted between 1999 and 2020 were included in the review. Studies ranged in sample size from 24 students in targeted interventions to 7,741 students in whole-school interventions. Forty-eight percent of studies targeted grades 1–6, and 52% of studies targeted grades 7–12. Participants' mean age was 10.5 years. Fifty-nine percent of studies had interventions delivered by school staff or teachers and all were based in schools. Seventy percent of studies included whole-school interventions, 26% targeted interventions, and 3.7% included both whole-school and targeted components. Most studies came from high-income countries, Australia ( $n = 7$ ), United States ( $n = 5$ ), England ( $n = 4$ ), Canada ( $n = 3$ ), Netherlands ( $n = 2$ ), Finland ( $n = 1$ ), Italy ( $n = 2$ ), and Germany ( $n = 1$ ), with one study from a middle-income country (China), and one from a lower-middle-income country (Pakistan). Only one study (INCLUSIVE trial, Bonell et al., 2018) provided program costs and two others (KiVA, Huitsing et al., 2019; Williford et al., 2012) provided costs from the main trials of the study. Costs for these trials were estimated to be less than £100 per student per year, without including annual costs and once only costs (Huitsing et al., 2019; Persson, Wennberg, Beckman, Salmivalli, & Svensson, 2018).

A summary of study design characteristics is shown in Table S3. Studies varied in terms of study design characteristics, 51.9% ( $n = 14$ ) of studies were cluster randomized, 11.1% ( $n = 3$ ) were individually randomized, 29.6% ( $n = 8$ ) were not randomized and 7.4% ( $n = 2$ ) were cross-over trials. Interventions included in these studies varied considerably in their measure of internalizing symptoms. 44.4% ( $n = 12$ ) of studies had a specific measure of anxiety, 55.5% ( $n = 15$ ) studies had a specific measure of depression and 44.4% ( $n = 12$ ) had a general measure of internalizing symptoms (e.g. SDQ emotional problems subscale). More than half of the studies had more than one measure of internalizing symptoms.

### Quality assessment and risk of bias in included studies

The quality of included studies varied considerably. Ten studies were rated as having a low risk of bias, 6 were rated as having some concerns for risk of


bias, and 11 studies were rated as having a high risk of bias. For studies included in the meta-analysis, eight were rated as having a low risk of bias, four were rated as having some concerns, and seven studies were rated as having a high risk of bias. A detailed quality assessment is shown in Figure S1.

### Meta-analysis and subgroup analysis

Fifteen of the 22 effects (68%) were larger than 0, indicating a reduction in overall internalizing symptoms favoring the intervention. The overall mean effect on internalizing symptoms was 0.06 (95% CI, 0.0284 to 0.1005;  $z = 3.50$ ;  $p < .001$ ), indicating that antibullying interventions had very little effect on overall internalizing symptoms. Our two heterogeneity measures indicated that there was low heterogeneity in our data ( $Q = 21.47$ ,  $df = 21$ ,  $p = .4304$ ;  $I^2 = 2.2\%$ , 95% CI: 0% to 47.4%). Figure 2 shows the forest plot for the overall and individual effect sizes.

### Subgroup analysis

Subgroup analysis for specific internalizing symptoms included 12 studies with measures of depression and 14 studies with measures of anxiety. Within-group effects were small but significant. The mean effect of depression was 0.06 (95% CI, 0.014 to 0.107;  $p = .011$ ,  $Q = 3.06$ ,  $I^2 = 0\%$ ) and for anxiety it was 0.08 (95% CI, 0.11 to 0.158;  $p = .024$ ,  $Q = 23.39$ ,  $I^2 = 44\%$ ). When directly comparing the effect of interventions on anxiety versus depression, between-group differences were not significant ( $Q = 0.31$ ,  $df = 1$ ,  $p = .578$ ). Subgroup analysis for type of intervention included 8 studies implementing targeted and 13 studies implementing whole-school interventions. Within-group effects were small and significant only for studies implementing whole-school interventions. The mean effect of targeted interventions was 0.01 (95% CI,  $-0.094$  to  $0.109$ ;  $p = .88$ ,  $Q = 7.52$ ,  $I^2 = 7\%$ ) and for whole-school interventions it was 0.08 (95% CI, 0.036 to 0.117;  $p < .001$ ,  $Q = 12.16$ ,  $I^2 = 1\%$ ). However, when directly comparing the effect of targeted with whole-school interventions, between-group differences were not significant ( $Q = 1.53$ ,  $df = 1$ ,  $p = .215$ ). Subgroup analysis for age/grade level included 13 studies targeting primary school children and 9 studies targeting secondary school adolescents. Within-group effects were small and significant only for studies targeting primary schools. The mean effect of primary schools was 0.07 (95% CI, 0.023 to 0.109;  $p = .002$ ,  $Q = 12.61$ ,  $I^2 = 5\%$ ) and for secondary schools it was .06 (95% CI,  $-0.014$  to  $0.144$ ;  $p = .107$ ,  $Q = 8.81$ ,  $I^2 = 9\%$ ). When directly comparing the effect of primary with secondary school interventions, between-group differences were not significant ( $Q = 0$ ,  $df = 1$ ,



**Figure 2** Impact of school-based antibullying interventions on overall internalizing symptoms

$p = .979$ ). Lastly, we conducted an additional analysis to examine the effects of methodological quality. Ten studies reporting low risk of bias were compared with 12 studies reporting some to high risk of bias. The mean effect of studies with a low risk of bias was 0.06 (95% CI, 0.014 to 0.113,  $p = .012$ ,  $Q = 12.75$ ,  $I^2 = 29\%$ ) and for studies with some-to-high concerns, 0.07 (95% CI, -0.015 to 0.148,  $p = .110$ ,  $Q = 8.72$ ,  $I^2 = 0\%$ ). There was no statistically significant difference between low-risk studies and higher-risk studies ( $Q = 0.005$ ,  $df = 1$ ,  $p = .946$ ).

### Publication bias

Overall, results were robust to sensitivity analysis examining heterogeneity and publication bias. No outliers were detected using leave-one-out analysis and influence diagnostics. Eggers' test did not indicate presence of funnel plot asymmetry (intercept, 0.073; 95% CI, -0.88 to 0.73;  $p = .861$ ). The SMDs adjusted for publication bias according to Duval and Tweedie trim-and-fill procedure barely changed (SMD, 0.067; 95% CI, 0.0282 to 0.1051;  $p < .001$ ). The funnel plot is shown in Figure S2.

### Metaregression analysis of moderators

We assessed the effect of several moderators separately (determined a priori) on internalizing effect sizes using metaregression analysis. Intervention intensity ( $\beta = .05$ ;  $p = .303$ ,  $R^2 = 14.95\%$ ), duration ( $\beta = .05$ ;  $p = .312$ ,  $R^2 = 5.23\%$ ), location ( $\beta = -.03$ ;  $p = .439$ ,  $R^2 = 0\%$ ), and number of intervention components ( $\beta = 0.002$ ;  $p = .701$ ,  $R^2 = 0\%$ ) were not significantly related to or did they account for significant variation in overall internalizing effect

sizes. Results of the metaregression analysis are presented in Table S4.

### Metaregression analysis of intervention components

Intervention components were tested separately using metaregression analysis. Inclusion of the component, work with peers ( $\beta = .087$ ;  $p = .016$ ,  $R^2 = 100\%$ ), was significantly associated with a greater reduction in internalizing outcomes and the component cognitive-behavioral (CBT) elements ( $\beta = -0.08$ ;  $P = .029$ ,  $R^2 = 100\%$ ) were significantly associated with an increase in internalizing outcomes. No other components were significantly related to, or did they account for significant variation in, overall internalizing effect sizes. Results of the metaregression analysis are presented in Table S5.

### Mediation analysis

Mediation using path analysis was conducted to examine whether reductions in bullying victimization and perpetration mediated intervention effects on overall internalizing symptoms. We found the indirect effect of victimization mediating intervention status on internalizing symptoms ( $b = .058$ ,  $p = .314$ ), and the direct effect of intervention status on internalizing symptoms ( $b = .046$ ,  $p = .189$ ) to be insignificant, implying no mediation effects. We found similar effects with bullying perpetration; the indirect effect of perpetration mediating intervention status on internalizing symptoms ( $b = -.001$ ,  $p = .934$ ) and the direct effect of intervention status on internalizing ( $b = .057$ ,  $p = .052$ ) were also insignificant. Diagrams for both path analyses can be found in Figures S3 and S4.

## Discussion

This systematic review aimed to assess the impact of school-based antibullying interventions on internalizing symptoms in children and adolescents. A meta-analysis with 22 studies found a very small but statistically significant effect on overall internalizing symptoms, favoring interventions compared to controls. Sensitivity analyses and adjustments for quality of studies and bias demonstrated that the overall effect size was robust. Subgroup analyses exploring the effect of interventions on anxiety and depression also found a very small but significant effect. Similar results were found for interventions targeting children in primary versus adolescents in secondary school and for types of interventions using targeted versus whole-school approaches. Despite the effects being statistically significant, it is unclear whether the size of the effects can be considered to be clinically meaningful. Other universal school-based interventions have commonly powered their effect sizes on mental health outcomes being meaningful at around 0.2–0.3 (e.g. Challen, Machin, & Gillham, 2014; Ford et al., 2012; Gillham et al., 2007; Stallard et al., 2014), whereas the mean effect across both universal and targeted bullying interventions on internalizing symptoms found here was only 0.06. To ensure antibullying trials are maximizing their effectiveness, we recommend researchers power their studies to detect meaningful effects on mental health outcomes.

Our results partially concur with previous systematic reviews. Reviews looking at primary outcomes in antibullying interventions have shown small-to-modest reductions in bullying victimization and perpetration (Ferguson, Miguel, Kilburn, & Sanchez, 2007; Merrell, Gueldner, Ross, & Isava, 2008) making it likely that even smaller effects would be observed for secondary outcomes. Likewise, similar to our subgroup analysis, it appears that other meta-analyses have found inconclusive evidence supporting the effectiveness of other educational setting-based interventions at preventing anxiety and depression in young people (Caldwell et al., 2019). It is likely that larger (yet still small) effects observed in a previous meta-analysis are the result of using broader definitions of mental health (Fraguas et al., 2021).

A rather surprising result came from the subgroup analysis examining differences between whole-school and targeted approaches. We found no difference between intervention types; however, whole-school approaches were significantly more effective than usual school practice and had a larger effect size than targeted interventions, which were not significantly more effective than usual practice. We had expected targeted interventions to have a more substantial effect on internalizing symptoms as they are typically more effective at managing anxiety and depression than universal interventions (Calear &

Christensen, 2010; Werner-Seidler, Perry, Calear, Newby, & Christensen, 2017) and most only include pupils with high internalizing symptoms at baseline. It may be the case that whole-school interventions have an advantage because they reach more pupils and often target a wider range of prosocial behaviors and the school climate. Changes in these factors may ultimately help children deal with interpersonal conflicts, foster better relationships, feel safer at school, and thus reduce internalizing symptoms. Indeed, some research suggests that whole-school interventions can have more successful outcomes when they are integrated into daily practice and school culture, engage all staff and students and are reinforced outside the classroom by parents and the community (Goldberg et al., 2019). Evidence also suggests that a whole-school approach compared to individual- or classroom-based target interventions is more likely to result in ‘enduring positive changes’ due to the duration and multicomponent focus of these interventions (Weare & Nind, 2011). However, it has been suggested that embedding targeted interventions within whole-school approaches could be the most effective way to target bullying and potentially internalizing symptoms as this combination would improve overall school climate but also focus on the most vulnerable children and persistent cases of bullying (Garandeau & Salmivalli, 2019). Indeed, more research is needed to untangle the advantages and disadvantages of whole-school versus targeted approaches. It is clear from this systematic review that very few studies, particularly targeted interventions, applied consistent approaches and targeted similar mechanisms of action, thus potentially contributing to the lack of significant effects. Lastly, it is also possible that different types of interventions had symptom-specific effects which were not captured with overall internalizing symptoms. Based on our findings, existing targeted antibullying interventions may not be sufficiently tailored to reduce internalizing symptoms, and thus could benefit from re-examining program aims, content, and design or potentially embedding them in whole-school interventions to improve student’s mental health outcomes.

We also found that interventions were equally effective at reducing internalizing symptoms for children and for adolescents. Findings from previous literature have been inconsistent in relation to the moderating effects of age on the effectiveness of antibullying interventions on reducing bullying outcomes (Smith, Salmivalli, & Cowie, 2012), however, less is known about its moderating effect on internalizing outcomes. A recent review showed that educational intervention effects on internalizing symptoms were not significantly moderated by age (Werner-Seidler et al., 2017). To understand this relationship further, age should be examined in combination with program components, type of



intervention, type of internalizing outcome, and bullying outcomes.

In terms of intervention components, our findings show that interventions that included a work with peers component were associated with larger effects on internalizing symptoms than those that did not use this component. Recent reviews show that work with peers is associated with a decrease in victimization (Gaffney et al., 2021a, 2021b); however, this is the first time it has been looked at in association with internalizing outcomes. Working with peers has been shown to increase student's perception and sense of safety at school (Cowie, Hutson, Oztug, & Myers, 2008; Cowie & Oztug, 2008), which may ultimately lead to overall reductions in internalizing symptoms. Some of the most effective interventions at reducing bullying such as KiVA (Salmivalli, Poskiparta, Ahtola, & Haataja, 2013) and No Trap! (Palladino, Nocentini & Menesini, 2016) involve working with peers in different capacities (e.g. as mediators or by encouraging bystander behavior) and further research should investigate which specific capacities are the most helpful at reducing internalizing symptoms. For example, in a recent review by Gaffney et al. (2021a, 2021b), researchers explored the distinction between informal and formal peer involvement in addition to encouraging bystander behavior for students witnessing bullying. Results showed that informal peer involvement was associated with significant reductions in bullying perpetration and victimization, while formal peer involvement was associated with significant reductions in bullying perpetration only. Surprisingly, the *exclusion* of encouraging bystander behavior was associated with significantly larger reductions in victimization. The difference in results for each, albeit nonmutually exclusive, category shows the need to make these distinctions to try and understand what makes interventions work better. Notably, in past systematic reviews, work with peers had been found to be associated with an *increase* in victimization, and it has been suggested that this might be the case if peer leaders have high levels of externalizing symptoms which might lead to deviancy training and negative outcomes (Dishion, McCord, & Poulin, 1999). Therefore, future work examining baseline levels of internalizing and externalizing in peer leaders, and other potential moderators, should be considered when evaluating program effects on victimization and mental health.

Contrary to expectations, given that some studies suggest that components targeting mental health (including CBT) can be effective at reducing bullying perpetration (Gaffney et al., 2021a, 2021b), the use of CBT components was significantly associated with *smaller* effects on internalizing symptoms compared to when CBT components were not used. This negative effect appeared to be carried by two of the seven interventions which used CBT. These were both targeted interventions: one delivered general

counseling to children in care and the other taught emotional literacy to children identified as bullies by their peers. However, there was no clear pattern in outcomes based on study type or program duration and intensity. Also of note, most of the interventions that implemented CBT components were delivered by school staff; research on programs targeting internalizing symptoms suggests that larger effect sizes are associated with programs delivered by mental health professionals than programs delivered by teachers (Calear & Christensen, 2010; Stallard et al., 2014; Werner-Seidler et al., 2017). Our findings raise reasons to be cautious about the use of CBT for dealing with internalizing symptoms in antibullying contexts and support other reviews showing similar trends from other school-based interventions (Caldwell et al., 2019; Mertens, Deković, Leijten, Van Londen, & Reitz, 2020). Yet, future research should examine whether incorporating specific CBT techniques can potentially promote children's mental health in antibullying interventions, depending on study type, design, their use in combination with other components (e.g. work with peers), their frequency, and their mode of delivery.

We found no evidence that other components examined in this review were effective in reducing internalizing symptoms, but this might reflect the small number of studies that applied consistent approaches and targeted similar mechanisms of action. For example, some universal interventions were based on restorative practice (Bonell et al., 2018) or adventure-based learning (Battey, 2008), while targeted programs varied among CBT-based, personality-targeted approaches (Kelly et al., 2020), social skills training (Fox & Boulton, 2003), or art therapy (Yan, Chen, & Huang, 2019). Going forward it will be important to examine the specific impacts of components grounded in empirical knowledge and theory of change frameworks, both to maximize meaningful clinical impacts and to provide a deeper understanding of any causal pathways between bullying and internalizing symptoms. Lastly, although we examined each intervention component individually, it is important to note that we did not assess for fidelity, and thus future studies should consider measuring adherence to interventions components. It is likely that factors such as the time required for preparation of lessons, duration of activities, and teacher's available time and enthusiasm for individual components throughout the intervention, among other factors, may influence program fidelity.

Our last set of analyses examined the mediating effects of bullying with the aim of exploring a possible causal pathway for intervention effects. Reductions in bullying victimization and perpetration did not account for intervention effects on internalizing outcomes. It is possible that antibullying interventions may be acting in many other ways to reduce internalizing, such as by promoting a



positive school climate, encouraging bystanders to take action, fostering prosocial skills, through peer support or other types of social support, or possibly a combination of elements that work towards improving student's mental health. To our knowledge, this is the first study to investigate potential causal pathways in a meta-analysis of antibullying interventions and sets a framework for prospective research to explore other factors that might explain intervention mechanisms and target them in future interventions designed to decrease bullying and internalizing symptoms. It is important to note that despite using pre-post changes for this analysis to account for baseline levels of bullying and internalizing, small effects and low heterogeneity could have had an impact on this analysis. Furthermore, this analysis only reflects school-level and not student-level mediation effects.

### Limitations and future directions

Our findings should be interpreted in the context of several limitations. In terms of our included studies, quality and risk of bias varied widely. Studies typically scored high on risk of bias due to no randomization taking place, data handling being inadequate, or missing data and attrition not being described. Yet, subgroup analyses looking at overall study quality showed no difference between low and high risk of bias. In terms of statistical analysis, many studies did not report the necessary statistics to compute effect sizes and thus were excluded from the meta-analysis. Additionally, most of our outcomes were based on self-report which could inflate the risk of bias. Unfortunately, we were unable to test type of informant as a moderator due to a lack of studies with informants other than self-reports. In terms of intervention components, we did not pre-register or distinguish between specific types of the 'work with peers' component (e.g. Gaffney et al., 2021a, 2021b) as this could lead to a loss of power and potentially biased results.

There are also other limitations to consider that are specific to this review. Our objective was to examine interventions with a primary aim to reduce bullying in schools. As such, we excluded general conflict resolution and aggression prevention interventions as well as general health promotion interventions which could have an impact on both bullying and internalizing symptoms. This review focused on internalizing symptoms as outcome measures as they are the leading cause of poor mental health in children (Erskine et al., 2015; Ghandour et al., 2019). We recognize that antibullying interventions may also have an impact on other outcomes such as externalizing symptoms, and to a lesser extent general wellbeing. Although we tried to capture a range of internalizing and emotional symptoms measured in a standardized and validated way, we excluded studies if they only presented

measures of self-esteem, loneliness, or guilt as we did not consider them to be fully representative of internalizing symptoms. Standardization of measures should be considered for future intervention studies to promote data synthesis and comparison of findings across studies.

In terms of statistical analysis, there were also some limitations. First and foremost, having average effect sizes to obtain a single measure of internalizing may have resulted in a loss of information for metaregression analyses, particularly in cases where depression and anxiety had opposing effects, or if one outcome was significantly different from the other. Effects were computed at postintervention only, however, it is unlikely that results would drastically change over time as seen in previous reviews looking at intervention effects on anxiety and depression (Werner-Seidler et al., 2017). Other statistical limitations included having to control for clustering with studies that did not report intraclass correlation coefficients (ICC). In these cases, ICCs were drawn from the literature, but might not be fully representative of specific trials. Lastly, it is likely that our lack of heterogeneity contributed to our secondary analysis observing minimal effects.

### Conclusion

Overall, we found insufficient evidence to conclude that antibullying interventions are effective at reducing internalizing problems in children and adolescents. While we are confident that most antibullying programs are effective at reducing the prevalence of bullying victimization and perpetration in schools, this review suggests that more work needs to be done to maximize the mental health benefits of these interventions. Specifically, there is a need to select and measure internalizing outcomes consistently, power studies to be able to detect clinically significant changes in internalizing effect sizes, improve the quality of research designs, and design antibullying interventions using evidence-based components that are likely to enhance positive impacts on mental health. This review suggests that including social or peer support components may be effective at reducing internalizing symptoms and therefore could be a target for intervention. On the other hand, interventions should carefully re-evaluate how CBT components are being employed to optimize their effect. Moving forward, research should prioritize understanding the causal pathways between intervention effects and mental health outcomes to highlight the specific type of support needed in interventions. Targeting internalizing outcomes in schools offers the opportunity to alleviate many typical barriers to accessing treatment such as time, location, and cost. Thus, future researchers, clinicians, and public healthcare professionals should recognize the importance of internalizing outcomes when designing school-based interventions to

maximize their impact and safeguard young people from the damaging outcomes of bullying.

## Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article:

**Appendix S1.** Note on deviations from the Protocol.

**Appendix S2.** Detailed Search Strategy.

**Appendix S3.** Intervention component definitions.

**Table S1.** Rating of intervention components.

**Table S2.** Study and participant characteristics.

**Table S3.** Characteristics associated with study design.

**Table S4.** Moderator Metaregression analysis.

**Table S5.** Intervention components Metaregression analysis.

**Figure S1.** Quality assessment (RoB).

**Figure S2.** Duval and Tweedie Trim-and-Fill Funnel Plot.

**Figure S3.** Mediation model for Bullying Victimization.

**Figure S4.** Mediation model for Bullying Perpetration.

## Acknowledgements

The authors would like to thank all the authors who provided them with data and ratings of intervention components. This research was supported through funding from the Calleva Foundation. L.B. is a primary investigator in the NIH-funded *Stand Together Trial* (ID: PHR 17/92/11), an evaluation of the KiVa antibullying intervention in the United Kingdom. The remaining authors have declared that they have no competing or potential conflicts of interest.

## Correspondence

Carolina Guzman Holst, Magdalen College, Oxford OX1 4AU, UK; Email: carolina.guzmanholst@magd.ox.ac.uk

## Key points

- Little is known about the effectiveness of antibullying interventions in reducing internalizing symptoms such as anxiety or depression, and what factors may influence intervention effects.
- Interventions have a small effect on reducing internalizing symptoms. Social or peer support components may be effective at improving student's mental health and could be a target for intervention, while implementation of CBT components could have negative implications and must be re-evaluated.
- Bullying victimization and perpetration did not mediate the relationship between intervention condition and internalizing outcomes, suggesting that interventions may be acting in many other ways to reduce internalizing symptoms – a priority for future research.
- This study sets a framework for researchers, practitioners, professionals in education and policy-makers wishing to make recommendations about how antibullying programs can improve children's mental health by examining different intervention characteristics, components and mechanism of action that can inform future interventions.

## References

\* These are all the included studies part of the review, meta-analysis. They are cited in the Figure/Supplement.

Arseneault, L., Milne, B.J., Taylor, A., Adams, F., Delgado, K., Caspi, A., & Moffitt, T.E. (2008). Being bullied as an environmentally mediated contributing factor to children's internalizing problems: A study of twins discordant for victimization. *Archives of Pediatrics & Adolescent Medicine*, 162, 145–150.

Barker, E.D., Boivin, M., Brendgen, M., Fontaine, N., Arseneault, L., Vitaro, F., ... & Tremblay, R.E. (2008). Predictive validity and early predictors of peer-victimization trajectories in preschool. *Archives of General Psychiatry*, 65, 1185–1192.

\*Battay (2008). *Can bullies become buddies? Evaluation of and theoretical support for an experiential education bully prevention curriculum with seventh grade student*. PhD thesis. Retrieved from Oregon State University Library Repository.

\*Berry, K., & Hunt, C.J. (2009). Evaluation of an intervention program for anxious adolescent boys who are bullied at school. *The Journal of Adolescent Health*, 45, 376–382.

Biggs, B.K., Vernberg, E., Little, T.D., Dill, E.J., Fonagy, P., & Twemlow, S.W. (2010). Peer victimization trajectories and their association with children's affect in late elementary school. *International Journal of Behavioral Development*, 34, 136–146.

\*Bonell, C., Allen, E., Warren, E., McGowan, J., Bevilacqua, L., Jamal, F., ... & Viner, R.M. (2018). Effects of the Learning Together intervention on bullying and aggression in English secondary schools (INCLUSIVE): A cluster randomised controlled trial. *The Lancet*, 392, 2452–2464.

Borenstein, M., Hedges, L.V., Higgins, J.P., & Rothstein, H.R. (2011). *Introduction to meta-analysis*. Chichester: John Wiley & Sons.

Bowes, L., Jolinson, C., Wolke, D., & Lewis, G. (2015). Peer victimisation during adolescence and its impact on depression in early adulthood: Prospective cohort study in the United Kingdom. *BMJ*, 350, h2469.

Caldwell, D.M., Davies, S.R., Hetrick, S.E., Palmer, J.C., Caro, P., López-López, J.A., Gunnell, D., Kidger, J., Thomas, J., French, C., Stockings, E., Campbell, R., & Welton, N.J. (2019). School-based interventions to prevent anxiety and depression in children and young people: A systematic

- review and network meta-analysis. *The Lancet Psychiatry*, 6, 1011–1020.
- Calear, A.L., & Christensen, H. (2010). Systematic review of school-based prevention and early intervention programs for depression. *Journal of Adolescence*, 33, 429–438.
- Challen, A.R., Machin, S.J., & Gillham, J.E. (2014). The UK Resilience Programme: A school-based universal nonrandomized pragmatic controlled trial. *Journal of Consulting and Clinical Psychology*, 82, 75–89.
- \*Connolly, J., Josephson, W., Schnoll, J., Simkins-Strong, E., Pepler, D., MacPherson, A., ... & Jiang, D. (2015). Evaluation of a youth-led program for preventing bullying, sexual harassment, and dating aggression in middle schools. *The Journal of Early Adolescence*, 35, 403–434.
- Cowie, H., Hutson, N., Oztug, O., & Myers, C. (2008). The impact of peer support schemes on pupils' perceptions of bullying, aggression and safety at school. *Emotional and Behavioural Difficulties*, 13, 63–71.
- Cowie, H., & Oztug, O. (2008). Pupils' perceptions of safety at school. *Pastoral Care in Education*, 26, 59–67.
- \*Cross, D., Shaw, T., Epstein, M., Pearce, N., Barnes, A., Burns, S., ... & Runions, K. (2018). Impact of the *Friendly Schools* whole-school intervention on transition to secondary school and adolescent bullying behavior. *European Journal of Education*, 53, 495–513.
- \*Dempsey, A. (2009). *Aggression and victimization in middle school: A mixed methods analysis of the process and effectiveness of implementing a prevention program*. PhD thesis. Retrieved from Proquest.
- Department for Education. (2018). *Bullying in England April 2013-March 2018: Analysis on 10 to 15 year olds from the Crime Survey for England and Wales*.
- \*DeRosier, M.E. (2004). Building relationships and combating bullying: Effectiveness of a school-based social skills group intervention. *Journal of Clinical Child and Adolescent Psychology*, 33, 196–201.
- Dishion, T.J., McCord, J., & Poulin, F. (1999). When interventions harm. Peer groups and problem behavior. *The American Psychologist*, 54, 755–764.
- Erskine, H.E., Moffitt, T.E., Copeland, W.E., Costello, E.J., Ferrari, A.J., Patton, G., ... & Scott, J.G. (2015). A heavy burden on young minds: The global burden of mental and substance use disorders in children and youth. *Psychological Medicine*, 45, 1551–1563.
- Farrington, D.P., & Ttofi, M.M. (2009). School-based programs to reduce bullying and victimization. *Campbell Systematic Reviews*, 5, i-148.
- \*Fekkes, M., Pijpers, F.I., & Verloove-Vanhorick, S.P. (2006). Effects of antibullying school program on bullying and health complaints. *Archives of Pediatrics & Adolescent Medicine*, 160, 638–644.
- Ferguson, C.J., Miguel, C.S., Kilburn, J.C., & Sanchez, P. (2007). The effectiveness of school-based anti-bullying programs: A meta-analytic review. *Criminal Justice Review*, 32, 401–414.
- \*Finger, L.R. (2009). *Beyond Bullying Primary Schools Program: Implementing an effective whole-school program to manage bullying, enhance prosocial behaviour, and boost student well-being in the upper primary grades*. PhD thesis. Retrieved from U. of Western Sydney Repository.
- \*Fite, P.J., Cooley, J.L., Poquiz, J., & Williford, A. (2019). Pilot evaluation of a targeted intervention for peer-victimised youth. *Journal of Clinical Psychology*, 75, 46–65.
- Ford, T., Edwards, V., Sharkey, S., Ukoumunne, O.C., Byford, S., Norwich, B., & Logan, S. (2012). Supporting teachers and children in schools: the effectiveness and cost-effectiveness of the incredible years teacher classroom management programme in primary school children: A cluster randomised controlled trial, with parallel economic and process evaluations. *BMC Public Health*, 12, 719.
- \*Fox, C.L., & Boulton, M.J. (2003). Evaluating the effectiveness of a social skills training (SST) programme for victims of bullying. *Educational Research*, 45, 231–247.
- Fraguas, D., Díaz-Caneja, C.M., Ayora, M., Durán-Cutilla, M., Abregú-Crespo, R., Ezquiaga-Bravo, I., ... & Arango, C. (2021). Assessment of school anti-bullying interventions: A meta-analysis of randomized clinical trials. *JAMA Pediatrics*, 175, 44–55.
- Gaffney, H., Farrington, D.P., & Ttofi, M.M. (2019). Examining the effectiveness of school-bullying intervention programs globally: A meta-analysis. *International Journal of Bullying Prevention*, 1, 14–31.
- Gaffney, H., Ttofi, M.M., & Farrington, D.P. (2021a). What works in anti-bullying programs? Analysis of effective intervention components. *Journal of School Psychology*, 85, 37–56.
- Gaffney, H., Ttofi, M.M., & Farrington, D.P. (2021b). Effectiveness of school-based programs to reduce bullying perpetration and victimization: An updated systematic review and meta-analysis. *Campbell Systematic Reviews*, 17, e1143.
- Garandeau, C.F., & Salmivalli, C. (2019). Can healthier contexts be harmful? A new perspective on the plight of victims of bullying. *Child Development Perspective*, 13, 147–152.
- Ghandour, R.M., Sherman, L.J., Vladutiu, C.J., Ali, M.M., Lynch, S.E., Bitsko, R.H., & Blumberg, S.J. (2019). Prevalence and treatment of depression, anxiety, and conduct problems in US children. *The Journal of Pediatrics*, 206, 256–267.e3.
- Gillham, J.E., Reivich, K.J., Freres, D.R., Chaplin, T.M., Shatté, A.J., Samuels, B., ... & Seligman, M.E. (2007). School-based prevention of depressive symptoms: A randomized controlled study of the effectiveness and specificity of the Penn Resiliency Program. *Journal of Consulting and Clinical Psychology*, 75, 9–19.
- Goldberg, J.M., Sklad, M., Elfrink, T.R., Schreurs, K.M.G., Bohlmeijer, E.T., & Clarke, A.M. (2019). Effectiveness of interventions adopting a whole school approach to enhancing social and emotional development: A meta-analysis. *European Journal of Psychology Education*, 34, 755–782.
- Hofstra, M.B., Van der Ende, J., & Verhulst, F.C. (2000). Continuity and change of psychopathology from childhood into adulthood: A 14-year follow-up study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39, 850–858.
- \*Hoglund, W., Hosan, N., & Leadbeater, B. (2012). Using your WITS: A 6-year follow-up of a peer victimization prevention program. *School Psychology Review*, 41, 193–214.
- Huitsing, G., Barends, S., & Lokkerbol, J. (2020). Cost benefit analysis of the kiva anti-bullying program in the Netherlands. *International Journal of Bullying Prevention*, 2, 215–224.
- \*Huitsing, G., Lodder, G.M.A., Oldenburg, B., Schacter, H.L., Salmivalli, C., Juvonen, J., & Veenstra, R. (2019). The Healthy Context Paradox: Victims' adjustment during an anti-bullying intervention. *Journal of Child Family Studies*, 28, 2499–2509.
- \*Karmaliani, R., McFarlane, J., Khuwaja, H., Somani, Y., Shehzad, S., Saeed Ali, T., ... & Jewkes, R. (2020). Right To Play's intervention to reduce peer violence among children in public schools in Pakistan: A cluster-randomized controlled trial. *Global Health Action*, 13, 1836604.
- \*Kelly, E.V., Newton, N.C., Stapinski, L.A., Conrod, P.J., Barrett, E.L., Champion, K.E., & Teesson, M. (2020). A Novel Approach to Tackling Bullying in Schools: Personality-targeted intervention for adolescent victims and bullies in Australia. *Journal of the American Academy of Child and Adolescent Psychiatry*, 59, 508–518.e2.
- Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R., & Walters, E.E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National



- Comorbidity Survey Replication. *Archives of General Psychiatry*, 62, 593–602.
- \*Knowler, C., & Frederickson, N. (2013). Effects of an emotional literacy intervention for students identified with bullying behaviour. *Educational Psychology*, 33, 862–883.
- Landis, J., & Koch, G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33, 159–174.
- López-López, J.A., Page, M.J., Lipsey, M.W., & Higgins, J.P.T. (2018). Dealing with effect size multiplicity in systematic reviews and meta-analyses. *Research Synthesis Methods*, 9, 336–351.
- Merrell, K.W., Gueldner, B., Ross, S., & Isava, M.D. (2008). How effective are school bullying intervention programs? A meta-analysis of intervention research. *School Psychology Quarterly*, 23, 26–42.
- Mertens, E., Deković, M., Leijten, P., Van Londen, M., & Reitz, E. (2020). Components of school-based interventions stimulating students' intrapersonal and interpersonal domains: A meta-analysis. *Clinical Child and Family Psychology Review*, 23, 605–631.
- Modecki, K.L., Minchin, J., Harbaugh, A.G., Guerra, N.G., & Runions, K.C. (2014). Bullying prevalence across contexts: A meta-analysis measuring cyber and traditional bullying. *The Journal of Adolescent Health*, 55, 602–611.
- Neil, A.L., & Christensen, H. (2009). Efficacy and effectiveness of school-based prevention and early intervention programs for anxiety. *Clinical Psychology Review*, 29, 208–215.
- \*Nocentini, A., Menesini, E., & Pluess, M. (2018). The personality trait of environmental sensitivity predicts children's positive response to school-based antibullying intervention. *Clinical Psychological Science*, 6, 848–859.
- Olweus, D. (1993). *Bullying at school: What we know and what we can do (understanding children's worlds)*. Oxford: Blackwell Publishing.
- Palladino, B.E., Nocentini, A., & Menesini, E. (2012). Online and offline peer led models against bullying and cyberbullying. *Psicothema*, 24, 634–639.
- Palladino, B.E., Nocentini, A., & Menesini, E. (2016). Evidence-based intervention against bullying and cyberbullying: Evaluation of the NoTrap! program in two independent trials. *Aggressive Behavior*, 42, 194–206.
- \*Palladino, B.E., Nocentini, A., & Menesini, E. (2019). How to stop victims' suffering? Indirect effects of an anti-bullying program on internalizing symptoms. *International Journal of Environmental Research and Public Health*, 16, 2631.
- \*Parada, R.H., Craven, R.G., & Marsh, H.W. (2008). The beyond bullying secondary program: An innovative program empowering teachers to counteract bullying in schools. In H.W. Marsh, R.G. Craven, & D.M. McInerney (Eds.), *Self-processes, learning, and enabling human potential: Dynamic new approaches* (pp. 373–400). Charlotte, NC: Information Age Publishing.
- Persson, M., Wennberg, L., Beckman, L., Salmivalli, C., & Svensson, M. (2018). The cost-effectiveness of the kiva antibullying program: Results from a decision-analytic model. *Prevention Science*, 19, 728–737.
- \*Pintabona, Y. *Frequently bullied students: Outcomes of a universal school-based bullying preventive intervention on peer victimization and psychological health*. PhD thesis. Retrieved from Curtin's institutional repository.
- \*Pryce, S., & Frederickson, N. (2013). Bullying behaviour, intentions and classroom ecology. *Learning Environments Research*, 16, 183–199.
- \*Rahey, L., & Craig, W. (2002). Evaluation of an ecological program to reduce bullying in schools. *Canadian Journal of Counselling*, 36, 281–296.
- \*Rapee, R.M., Shaw, T., Hunt, C., Bussey, K., Hudson, J.L., Mihalopoulos, C., ... & Cross, D. (2020). Combining whole-school and targeted programs for the reduction of bullying victimization: A randomized, effectiveness trial. *Aggressive Behavior*, 46, 193–209.
- Reijntjes, A., Kamphuis, J.H., Prinzie, P., & Telch, M.J. (2010). Peer victimization and internalizing problems in children: A meta-analysis of longitudinal studies. *Child Abuse & Neglect*, 34, 244–252.
- Salmivalli, C., Poskiparta, E., Ahtola, A., & Haataja, A. (2013). The implementation and effectiveness of the KiVa antibullying program in Finland. *European Psychologist*, 18, 79–88.
- Shakoor, S., Jaffee, S.R., Bowes, L., Ouellet-Morin, I., Andreou, P., Happé, F., ... & Arseneault, L. (2012). A prospective longitudinal study of children's theory of mind and adolescent involvement in bullying. *Journal of Child Psychology and Psychiatry*, 53, 254–261.
- Singham, T., Viding, E., Schoeler, T., Arseneault, L., Ronald, A., Cecil, C.M., ... & Pingault, J.B. (2017). Concurrent and longitudinal contribution of exposure to bullying in childhood to mental health: The role of vulnerability and resilience. *JAMA Psychiatry*, 74, 1112–1119.
- Smith, P.K., Salmivalli, C., & Cowie, H. (2012). Effectiveness of school-based programs to reduce bullying: A commentary. *Journal of Experimental Criminology*, 8, 433–441.
- Stallard, P., Skryabina, E., Taylor, G., Phillips, R., Daniels, H., Anderson, R., & Simpson, N. (2014). Classroom-based cognitive behaviour therapy (FRIENDS): A cluster randomised controlled trial to Prevent Anxiety in Children through Education in Schools (PACES). *The Lancet Psychiatry*, 1, 185–192.
- Stapinski, L.A., Araya, R., Heron, J., Montgomery, A.A., & Stallard, P. (2015). Peer victimization during adolescence: concurrent and prospective impact on symptoms of depression and anxiety. *Anxiety, Stress, and Coping*, 28, 105–120.
- Sukhawathanakul, P., & Leadbeater, B. (2020). Trajectories of peer victimization in elementary school children: Associations with changes in internalizing, externalizing, social competence, and school climate. *Journal of Community Psychology*, 48, 1751–1769.
- UNESCO. (2019). *Behind the numbers: ending school violence and bullying*, 2019. Available from: <https://unesdoc.unesco.org/ark:/48223/pf0000366483> [last accessed 21 February, 2022].
- \*Van Ryzin, M.J., & Roseth, C.J. (2018). Cooperative learning in middle school: A means to improve peer relations and reduce victimization, bullying, and related outcomes. *Journal of Educational Psychology*, 110, 1192–1201.
- Weare, K., & Nind, M. (2011). Mental health promotion and problem prevention in schools: What does the evidence say? *Health Promotion International*, 26, 29–69.
- Werner-Seidler, A., Perry, Y., Calear, A.L., Newby, J.M., & Christensen, H. (2017). School-based depression and anxiety prevention programs for young people: A systematic review and meta-analysis. *Clinical Psychology Review*, 51, 30–47.
- \*Williford, A., Boulton, A., Noland, B., Little, T.D., Kärnä, A., & Salmivalli, C. (2012). Effects of the KiVa anti-bullying program on adolescents' depression, anxiety, and perception of peers. *Journal of Abnormal Child Psychology*, 40, 289–300.
- \*Yan, H., Chen, J., & Huang, J. (2019). School bullying among left-behind children: The efficacy of art therapy on reducing bullying victimization. *Frontiers in Psychiatry*, 10, 40.
- Yeager, D.S., Fong, C.J., Lee, H.Y., & Espelage, D.L. (2015). Declines in efficacy of anti-bullying programs among older adolescents: Theory and a three-level meta-analysis. *Journal of Applied Developmental Psychology*, 37, 36–51.
- \*Zagorscak, P., Schultze-Krumbholz, A., Heinrich, M., Wölfer, R., & Scheithauer, H. (2019). Efficacy of cyberbullying prevention on somatic symptoms-randomized controlled trial applying a reasoned action approach. *Journal of Research on Adolescence*, 29, 908–923.

Accepted for publication: 28 March 2022