

Hands-on workshop on brain criticality

Jacek Grela, Jakub Janarek
CNA2023 W3 workshop session

bioNN



Fundacja na rzecz
Nauki Polskiej



Funded by
the European Union




UNIVERSITAS
JAGELLONICA
CRACOVENSIS

Basic org info

- (15:30 - 18:30) 3 hours with two-three 15 min breaks
- Venue: room G-01-09 or on-line
- Workshop language: **python**
a crash-course **python_crash_course.ipynb** is provided
- The working notebook
 - open <http://bit.ly/critical-brain> and use **google colab**
 - open <https://bit.ly/critical-brain-github>, clone git repo and open **exercises.ipynb**
- Outline
 - The concept of criticality
 - Towards a brain model
 - Criticality in the Haimovici model
 - Playing around



The concept of criticality

- a fascinating statistical physics concept connected with phases of matter





The concept of criticality

- a fascinating statistical physics concept connected with phases of matter
- also present in magnetism, liquid-gas mixtures; **the critical state** (long-range corr, rich dynamics)





The concept of criticality

- a fascinating statistical physics concept connected with phases of matter
- also present in magnetism, liquid-gas mixtures; **the critical state** (long-range corr, rich dynamics)
- beyond typical physics - earthquakes, traffic jams (power laws)



The concept of criticality

- a fascinating statistical physics concept connected with phases of matter
- also present in magnetism, liquid-gas mixtures; **the critical state** (long-range corr, rich dynamics)
- beyond typical physics - earthquakes, traffic jams (power laws)
- complexity, self-organization (Saturday lecture L17 by prof. Chialvo for more!)




The concept of criticality

- Ising model: a toy model of magnetism

The concept of criticality

- Ising model: a toy model of magnetism
 - spins on a 2D grid, each with 4 neighbours

$$\sigma_k \in \{+1, -1\}$$




The concept of criticality

- Ising model: a toy model of magnetism
 - spins on a 2D grid, each with 4 neighbours
- energy of the system

$$E(\sigma; J) = -J \sum_{\langle ij \rangle} \sigma_i \sigma_j$$

$$\sigma_k \in \{+1, -1\}$$



The concept of criticality

- Ising model: a toy model of magnetism
 - spins on a 2D grid, each with 4 neighbours
- energy of the system


$$E(\sigma; J) = -J \sum_{\langle ij \rangle} \sigma_i \sigma_j$$

← sum over neighbours

- statistical model -> temperature T as a control parameter


$$P(\sigma; T) \sim \exp \left(-\frac{E(\sigma; J)}{T} \right)$$

$$\sigma_k \in \{+1, -1\}$$



The concept of criticality


- control parameter T is changed; three regimes!



The concept of criticality

- control parameter T is changed; three regimes!
- a single critical T value

$$T_c = \frac{2J}{\ln(\sqrt{2} + 1)}$$




The concept of criticality

- control parameter T is changed; three regimes!
- a single critical T value

$$T_c = \frac{2J}{\ln(\sqrt{2} + 1)}$$

- but what really changes near T_c ?



The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**

The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**
 - pick a node σ_i (at random or in some order)

The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**
 - pick a node σ_i (at random or in some order)
 - **do a spin-flip** and find its energy change: $\Delta E = E(-\sigma_i) - E(\sigma_i)$

The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**
 - pick a node σ_i (at random or in some order)
 - **do a spin-flip** and find its energy change: $\Delta E = E(-\sigma_i) - E(\sigma_i)$
 - **accept** new configuration when $\Delta E < 0$

The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**
 - pick a node σ_i (at random or in some order)
 - **do a spin-flip** and find its energy change: $\Delta E = E(-\sigma_i) - E(\sigma_i)$
 - **accept** new configuration when $\Delta E < 0$
 - if $\Delta E \geq 0$, accept with probability $\exp(-\Delta E/T)$


The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**
 - pick a node σ_i (at random or in some order)
 - **do a spin-flip** and find its energy change: $\Delta E = E(-\sigma_i) - E(\sigma_i)$
 - **accept** new configuration when $\Delta E < 0$
 - if $\Delta E \geq 0$, accept with probability $\exp(-\Delta E/T)$
 - repeat

The concept of criticality

- how to run a simulation ? **Metropolis-Hastings algorithm**
 - pick a node σ_i (at random or in some order)
 - **do a spin-flip** and find its energy change: $\Delta E = E(-\sigma_i) - E(\sigma_i)$
 - **accept** new configuration when $\Delta E < 0$
 - if $\Delta E \geq 0$, accept with probability $\exp(-\Delta E/T)$
 - repeat
- dynamics are given by a transition graph

$$p_{I \rightarrow A} = 1 + \left(e^{-\Delta E/T} - 1 \right) \theta(\Delta E)$$



The concept of criticality

Your tasks are:

- **1.1 Magnetization**
Run the Ising model simulation for a set of temps T ; find magnetization as a function of T
- **1.2 Snapshots**
Plot snapshots in three regimes: subcritical, critical and supercritical.
- **1.3 Binomial model (*)**
Run a “binomial model” simulation. Plot again the magnetization and snapshots of the dynamics. What is the main difference between the Ising and this model?

Towards a brain model

- ...but where's the brain? Consider the Ising energy...

$$E(\sigma; J) = -J \sum_{\langle ij \rangle} \sigma_i \sigma_j$$

Towards a brain model

- ...but where's the brain? Consider the Ising energy...
- with a weighted grid

$$E(\sigma; J) = - \sum_{\langle ij \rangle} J_{ij} \sigma_i \sigma_j$$

Towards a brain model

- ...but where's the brain? Consider the Ising energy...
- with a weighted grid
- grid -> a general graph G

$$E(\sigma; J) = -J \sum_{(i,j) \in E(G)} w_{ij} \sigma_i \sigma_j$$


Towards a brain model

- ...but where's the brain? Consider the Ising energy...
- with a weighted grid
- grid -> a general graph G
- ... but need some neuron-like behavior

$$E(\sigma; J) = -J \sum_{(i,j) \in E(G)} w_{ij} \sigma_i \sigma_j$$


Towards a brain model

- transition graph is modified accordingly:




Towards a brain model

- transition graph is modified accordingly:




Towards a brain model

- transition graph is modified accordingly:




Towards a brain model

- transition graph is modified accordingly:



Towards a brain model

- transition graph is modified accordingly:




Towards a brain model

- transition graph is modified accordingly:

$$p_{I \rightarrow A} = 1 + (r_1 - 1) \theta(\Delta E)$$

$$\Delta E = \mathcal{T} - \sum_{j \text{ active}} w_{ij}$$




Towards a brain model

- **Haimovici model** of the brain
 - defined on a graph (connectome)
 - 3-state system (active - inactive - refractory)
 - out-of-equilibrium (no energy func)

$$p_{I \rightarrow A} = 1 + (r_1 - 1) \theta(\Delta E)$$

$$\Delta E = \mathcal{T} - \sum_{j \text{ active}} w_{ij}$$



Towards a brain model

Your tasks are:

- **2.1 Haimovici model**

Run the Haimovici model simulation with Hagmann connectome for a set of thresholds T . Find “magnetizations” for each neuron sub-population (“active” = “excited”, “refractory”, “inactive” = “susceptible”)

- **2.2 Temperature or threshold?**

Investigate how the threshold parameter differ from the temperature of the Ising model?
Tip: How do the sub-critical-super regimes behave? Inspect temporal dynamics or “magnetizations”.

- **2.3 Artificial connectomes**



Investigate artificial connectomes, try Watts-Strogatz or others. Look at magnetizations and different parameter regimes.

Criticality in the Haimovici model

- But what about criticality?

Criticality in the Haimovici model

- But what about criticality? Magnetization is not enough...



Criticality in the Haimovici model

- But what about criticality? Magnetization is not enough...
- Other indicators:
 - Variance of magnetization
 - Autocorrelation
 - Sizes of the largest clusters

Criticality in the Haimovici model

- But what about criticality? Magnetization is not enough...
- Other indicators:
 - Variance of magnetization
 - Autocorrelation
 - **Sizes of the largest clusters** (explicitly depend on the connectome!)

Criticality in the Haimovici model

- But what about criticality? Magnetization is not enough...
- Other indicators:
 - Variance of magnetization
 - Autocorrelation
 - **Sizes of the largest clusters** (explicitly depend on the connectome!)

Definition:

Maximal sets of **connected nodes** sharing the same type of activity.

We typically focus on the **largest** cluster and the **second-largest** cluster.

Criticality in the Haimovici model

Definition:

Maximal sets of **connected nodes** sharing the same type of activity.

We typically focus on the **largest** cluster and the **second-largest** cluster.



Towards a brain model

Your tasks are:

- **3.1 Own clustering algorithm (*)**
Make your own clustering algorithm using the networkx library.
- **3.2 Clusters in the Ising model**
Use an Ising model snapshot near T_c and plot the largest and the second-largest cluster. What is different near $T = 0$?
- **3.3 Criticality indicators in the Haimovici model**
Find cluster sizes in the Haimovici model and investigate other indicators as well (st. dev. of activity and autocorrelation).
- **3.4 Detective work**
We give you simulated data and a set of derived criticality indicators. Is the data taken from a system poised at criticality?

Playing around

- So far: the healthy brain is posed at criticality (Hagmann)

Playing around

- So far: the healthy brain is posed at criticality (Hagmann)
- What about changes in the brain?

Playing around

- So far: the healthy brain is posed at criticality (Hagmann)
- What about changes in the brain?
 - strokes
 - epileptic seizures
 - drugs
 - etc

Playing around

Your tasks are:

- **4.1 Lobotomy**

Take a healthy brain (Hagmann connectome) and create an artificial lobotomy-like procedure. Show cluster sizes.

- **4.2 Stroke**

What happens if you disconnect a single RSN from the brain? Show cluster sizes.

- **4.3 Epilepsy**

Model epilepsy by rescaling/translating connectome weights. What happens then?