

Problem Set 4

Instructions: For full credit, turn in 5 problems in a pdf file and a .m2 file. You are welcome to work together with your classmates on all the problems, and I will be happy to give you hints or discuss the problems with you, but you should write up your solutions by yourself. You cannot use any resources besides me, your classmates, our course notes, and the Macaulay2 documentation.

Problem 1. Let R be a noetherian ring and let I be a proper ideal in R .

a) Show that there exists an integer n such that $\sqrt{I}^n \subseteq I$.

b) Show that if R is local, then

$$\bigcap_{n \geq 0} I^n = (0).$$

Hint: reduce to the case when $I = \mathfrak{m}$, the unique maximal ideal. In that case, let $J := \bigcap_{n \geq 0} \mathfrak{m}^n$, and show that if $\mathfrak{m}J = Q_1 \cap \cdots \cap Q_s$ is a primary decomposition, then $J \subseteq Q_i$ for all i .

c) Show that if R is a domain, then

$$\bigcap_{n \geq 0} I^n = (0).$$

d) What if R is not local nor a domain?

Problem 2. For each of the following rings R determine whether the inclusion $\mathbb{C}[x] \subseteq R$ is integral, with justification. You do not need to justify that $\mathbb{C}[x] \subseteq R$ is an inclusion.

a) $R = \mathbb{C}[x, y, z]/(z^2 - xy)$

b) $R = \mathbb{C}[x, y, z]/(y^3 - x^2, z^2 - xy)$

c) $R = \mathbb{C}[x, y, z]/(x^2 - yz, y^2 - xz, z^2 - xy)$

If the answer is no, you might find it helpful to consider properties of the induced map on spectra.

Problem 3. For each of the following rings, compute their dimension in Macaulay2 for some particular field k of your choice, and then prove that the answer is correct for all fields k .

a) $R = \frac{k[x, y, z]}{(x^3, x^2y, xyz)}$.

b) $R = \frac{k[x, y, z, u, v]}{(x^3u^2 + y^3uv + z^3v^2)}$.

c) $R = k[x^2u, xyu, y^2u, x^2v, xyv, y^2v] \subseteq k[x, y, u, v]$.

d) $R = \mathbb{C}[x, y, z]/(x^2 - yz, y^2 - xz, z^2 - xy)$