

Problem Set 2

Problem 1. Let I be an ideal with no embedded primes in a noetherian ring R . Show that there exists an ideal J , which we can take to be principal, such that $I^{(n)} = (I^n : J^\infty)$ for all $n \geq 1$.

Problem 2. Let $R = \mathbb{Q}[x_1, x_2, x_3, x_4]$, and consider the ideal I you can define in Macaulay2 as follows:

```
M = matrix{{0,-x_1,-x_3,x_2,-x_1,x_4,-x_3},{x_1,0,-x_3,x_2,x_1,-x_4,-x_1},
{x_3,x_3,0,0,-x_3,x_1,x_4},{-x_2,-x_2,0,0,-x_4,x_2,0},
{x_1,-x_1,x_3,x_4,0,-x_3,x_1},{-x_4,x_4,-x_1,-x_2,x_3,0,x_2},{x_3,x_1,x_4,0,-x_1,x_2,0}}
I = pfaffians(6,M)
```

Using only the packages that come preloaded with Macaulay2, compute $I^{(2)}$ and $I^{(3)}$. Is $I^{(2)} = I^2$? Is $I^{(3)} = I^3$?

Problem 3. Find all the symbolic powers of $I = (6)$ in $\mathbb{Z}[\sqrt{-5}]$.

Problem 4. Let x be a regular element in a ring R , meaning that $xa = 0 \implies a = 0$.

- a) Show that $(x^n : x^{n-1}) = (x)$ for all $n \geq 1$.
- b) Show that $\text{Ass}(x^n) = \text{Ass}(x)$ for all $n \geq 1$.

Problem 5. Let $R = k[x, y, z]/(xy - z^c)$ where k is a field and $c \geq 2$, and let $P = (x, z)$.

- a) Show that (x^n) is a primary ideal for all $n \geq 1$. What is its radical?
- b) Prove that $P^{(cn)} = (x^n)$ for all $n \geq 1$.
- c) Prove that $P^{(n)} \neq P^n$ for all $n \geq 2$.

Problem 6. Height and dimension.

- a) Find the height of $J = (ab, bc, cd, ad)$ in $k[a, b, c, d]$ over any field k , and the dimension of $k[a, b, c, d]/J$.
- b) Find the dimension of the ring $S = \mathbb{Q}[x^3y^3, x^3y^2z, x^2z^3] \subseteq \mathbb{Q}[x, y, z]$.
- c) Let I be the defining ideal of the curve parametrized by (t^{13}, t^{42}, t^{73}) over \mathbb{Q} . Find the height of I , and notice that $\text{height}(I) < \mu(I)$.
- d) Let $R = \mathbb{Q}[x, y, z]$, and $I = (x^3, x^2y, x^2z, xyz)$. Find the dimension of R/I and the height of I .
- e) Find the dimension of the module I/I^2 , where $I = (xz)$ in $R = \mathbb{C}[x, y, z]/(xy, yz)$.