Topology
Home Work 7

Student: 26-712-224

October 29, 2026

"I wrote these solutions fully myself und would like to receive feedback on them"

Exercise 1.

i) Prove that a topological space is compact if and only if the following holds. Whenever $\{C_i : i \in I\}$ is a collection of closed sets such that for all $n \in \mathbb{N}$ and $\{i_1, i_2, \cdots, i_n\} \subset I$ we have $\bigcap_{j=1}^{n} C_{i_j} \neq \varnothing$, then $\bigcap_{i \in I} C_i \neq \varnothing$.

[Hint. Use contrapositive and De Morgan's law.]

ii) Show that if $(X, \mathcal{U})$ is compact, then for any infinite collection of points $\{x_n : n \in \mathbb{N}\}$, we have

$$\bigcap_{k \geq 1} \overline{\bigcup_{n \geq k} \{x_n\}} \neq \varnothing.$$


Conclude that any set of infinite points admits a limit point (or accumulation point). (Recall that x is a limit point of the set S if every neighborhood of x contains at least one point of S that is different from x itself.)

Exercise 2.

1. **(Gluing Lemma.)** Let $(X, \mathcal{U})$, $(Y, \mathcal{V})$ be topological spaces and C_1, C_2 closed subsets of X such that $C_1 \cup C_2 = X$. Let $f_i : C_i \to Y$ for $i = 1, 2$ be continuous functions such that $f_1|_{C_1 \cap C_2} = f_2|_{C_1 \cap C_2}$. (in other words, the restriction of f_1 and f_2 to $C_1 \cap C_2$ coincide.) Prove that one may define a function $f : X \to Y$ such that $f|_{C_i} = f_i$ for $i = 1, 2$ and prove that f is continuous.


   [Hint. A function is continuous iff the preimage of every closed set is closed.]

2. Recall the representation of a torus by a square S with opposite sides identified as in the figure below. We denote the equivalence relation given by this identification by $\sim$, and we let $p : S \to S/\sim$ denote the quotient map. Let the points $P \sim P'$ be as in the figure, and use the gluing lemma to find a neighborhood of $p(P) = p(P')$ in $S/\sim$ with the quotient topology that is homeomorphic to a disk.




Exercise 3. Which of the following topological spaces are manifolds? Which are manifolds with boundary? Which are not manifolds at all? You do NOT have to provide a rigorous proof, but do give a brief explanation hinting at the reason for your answer.

i) The closed ball $\bar{B}_{\mathbb{R}^3}(0,1) = \{p \in \mathbb{R}^3;\ \|p\| \leq 1\}$, with the topology induced by the Euclidean topology of $\mathbb{R}^3$.




ii) $\{(x,y) \in \mathbb{R}^2;\ x \cdot y = 0\}$, with the topology induced by the Euclidean topology of $\mathbb{R}^2$.

iii) The closed half cone $\{(x,y,z) \in \mathbb{R}^3;\ x^2 + y^2 = z^2,\ z \leq 0\}$, with the topology induced by the Euclidean topology of $\mathbb{R}^3$.



iv) The full cone $\{(x,y,z) \in \mathbb{R}^3;\ x^2 + y^2 = z^2\}$, with the topology induced by the Euclidean topology of $\mathbb{R}^3$.




v) $\mathbb{R}^2 \setminus \{(1/n, 0) \in \mathbb{R}^2;\ n \in \mathbb{N} \setminus \{0\}\}$, with the topology induced by the Euclidean topology of $\mathbb{R}^2$.


Exercise 4. Let M be any topological 2-dimensional manifold. Show that the connected sum of M and S^2 is homeomorphic to M itself.

[Hint. Use the gluing lemma and the fact that if $D \subset S^2$ is homeomorphic to a disk, then $S^2 \setminus D$ is also homeomorphic to a disk.]

Exercise 5. Find a subset of the projective plane $\mathbb{RP}^2$ that is (homeomorphic to) a Möbius band, then show that by removing this Möbius band, the remaining subspace is homeomorphic to a square. (Hint, use the following square representation of the projective plane.)




Exercise 6. Recall that the Möbius band M is the space obtained by gluing a square with the following identification:



i) Take a strip of paper and glue it into a Möbius band. (Much easier with a long thin strip – don't try it with a square!) First check that it has only 1 face by drawing a line with a pen along the middle line. Now take a pair of scissors and cut it along the middle lengthwise (see left picture below). Observe what you get.

ii) Make another Möbius band, but now instead of cutting in the middle, cut close to the edge, say at height $\frac{1}{3}$ as in the right picture below. (Keep cutting until you get back to where you started!) What do you get this time?



iii) Think of M as the square with identifications again. Let $n \geq 2$. If you cut it at heights $\frac{k}{n}$ for $1 \leq k \leq n-1$, how many connected components do you get? Do they have 1 or 2 faces? What are they homeomorphic to?