



Property-matched decoys that BEAT the 6TD3 gate, beside the real glue each most resembles

Left = decoy, right = its nearest known CDK12-DDB1 glue. Every decoy is < 0.35 Tanimoto to EVERY active by construction, so these are the closest anything gets. A more negative ΔV_{ina} is a better "glue" score; the gate is -2.0. These are the strongest false positives— the gate admits them.


DECOY ΔV_{ina} -4.52 [carries a purine] · Tanimoto 0.19 · nearest REAL GLUE ΔV_{ina} -2.41




DECOY ΔV_{ina} -4.23 · Tanimoto 0.20 · nearest REAL GLUE ΔV_{ina} +0.68



DECOY ΔV_{ina} -4.20 · Tanimoto 0.16 · nearest REAL GLUE ΔV_{ina} -1.83



DECOY ΔV_{ina} -3.72 · Tanimoto 0.16 · nearest REAL GLUE ΔV_{ina} -1.16



Shared substructure haloed in red. 97% of these decoys carry NO purine warhead — 5 of 160 do, because the matcher enforces global dissimilarity, not absence of a fragment (and 1 of the 4 pairs above is one of them). Purine content is not the cause of. Structurally unrelated molecules out-score real glues on the differential — which is why the gate admits 31% of them.