



Property-matched decoys the oracle REJECTS most confidently, beside their nearest real glue

Left = decoy, right = its nearest known CDK12-DDB1 glue. Every decoy is < 0.35 Tanimoto to EVERY active by construction, so these are the closest anything gets. A more negative ΔV_{ina} is a better "glue" score; the gate is -2.0. These sit at the OPPOSITE end: the 4 weakest of 160 decoys, far from passing.


DECOY ΔV_{ina} +0.39 · Tanimoto 0.13 · nearest REAL GLUE ΔV_{ina} -3.37




DECOY ΔV_{ina} +0.00 · Tanimoto 0.16 · nearest REAL GLUE ΔV_{ina} -1.75



DECOY ΔV_{ina} -0.07 · Tanimoto 0.13 · nearest REAL GLUE ΔV_{ina} -0.31



DECOY ΔV_{ina} -0.11 · Tanimoto 0.19 · nearest REAL GLUE ΔV_{ina} -2.04



Shared substructure haloed in red. 97% of these decoys carry NO purine warhead — 5 of 160 do, because the matcher enforces global dissimilarity, not absence of a fragment. These are the clean negatives: their differentials sit at roughly ZERO, so the oracle does separate a population it is confident about. The failure is not that it rejects nothing — it is that the 31% at the other end are indistinguishable from real glues on this signal.