

Machine Learning


Prof. Barbara Caputo

Dip. Ingegneria Informatica, Automatica e Gestionale, Roma




SAPIENZA
UNIVERSITÀ DI ROMA

Support Vector Machines




Support Vector Machines

$$\underset{w, b}{\text{minimize}} \frac{1}{2} \|w\|^2 \quad \text{subject to } y_i [\langle x_i, w \rangle + b] \geq 1$$



Support Vector Machines

$$\underset{w, b}{\text{minimize}} \frac{1}{2} \|w\|^2 \quad \text{subject to } y_i [\langle x_i, w \rangle + b] \geq 1$$




$$\underset{\alpha}{\text{maximize}} -\frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle + \sum_i \alpha_i$$

$$\text{subject to } \sum \alpha_i y_i = 0 \text{ and } \alpha_i \geq 0$$

Support Vector Machines


$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 \quad \text{subject to } y_i [\langle x_i, w \rangle + b] \geq 1$$



$$\text{subject to } \sum \alpha_i y_i = 0 \text{ and } \alpha_i \geq 0$$

Support Vectors

$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 \quad \text{subject to } y_i [\langle x_i, w \rangle + b] \geq 1$$



Karush Kuhn Tucker

Optimality condition

$$\alpha_i [y_i [\langle w, x_i \rangle + b] - 1] = 0$$




$$\alpha_i = 0$$


$$\alpha_i > 0 \implies y_i [\langle w, x_i \rangle + b] = 1$$

Properties

$$w = \sum_i y_i \alpha_i x_i$$




Properties



- Weight vector w as weighted linear combination of instances


Properties



- Weight vector w as weighted linear combination of instances
- Only points on margin matter (ignore the rest and get same solution)

Properties


$$w = \sum_i y_i \alpha_i x_i$$



- Weight vector w as weighted linear combination of instances
- Only points on margin matter (ignore the rest and get same solution)
- Only inner products matter

Properties


$$w = \sum_i y_i \alpha_i x_i$$



- Weight vector w as weighted linear combination of instances
- Only points on margin matter (ignore the rest and get same solution)
- Only inner products matter
 - Quadratic program

Properties


$$w = \sum_i y_i \alpha_i x_i$$



- Weight vector w as weighted linear combination of instances
- Only points on margin matter (ignore the rest and get same solution)
- Only inner products matter
 - Quadratic program
 - We can replace the inner product by a kernel


Properties

$$w = \sum_i y_i \alpha_i x_i$$




- Weight vector w as weighted linear combination of instances
- Only points on margin matter (ignore the rest and get same solution)
- Only inner products matter
 - Quadratic program
 - We can replace the inner product by a kernel
- Keeps instances away from the margin

Example




Example


Number of Support Vectors: 3 (-ve: 2, +ve: 1) Total number of points: 15



Why large margins?




Why large margins?




- Maximum robustness relative to uncertainty

Why large margins?




- Maximum robustness relative to uncertainty
- Symmetry breaking

Why large margins?



- Maximum robustness relative to uncertainty
- Symmetry breaking
- Independent of correctly classified instances


Why large margins?



- Maximum robustness relative to uncertainty
- Symmetry breaking
- Independent of correctly classified instances
- Easy to find for easy problems

Soft Margin Classifiers


Large Margin Classifier



linear function

$$f(x) = \langle w, x \rangle + b$$


Large Margin Classifier



linear function

$$f(x) = \langle w, x \rangle + b$$

Large Margin Classifier




linear function


$$f(x) = \langle w, x \rangle + b$$

linear separator
is impossible


Adding slack variables




Adding slack variables



Adding slack variables




Adding slack variables




Convex optimization problem

Adding slack variables



Convex optimization problem

Adding slack variables



Adding slack variables

- Hard margin problem

$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 \text{ subject to } y_i [\langle w, x_i \rangle + b] \geq 1$$

Adding slack variables

- Hard margin problem

$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 \text{ subject to } y_i [\langle w, x_i \rangle + b] \geq 1$$

- With slack variables

$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$
$$\text{subject to } y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i \text{ and } \xi_i \geq 0$$

Adding slack variables

- Hard margin problem

$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 \text{ subject to } y_i [\langle w, x_i \rangle + b] \geq 1$$

- With slack variables

$$\underset{w,b}{\text{minimize}} \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

$$\text{subject to } y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i \text{ and } \xi_i \geq 0$$

Problem is always feasible.

Dual Problem

- Primal optimization problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$

Dual Problem

- Primal optimization problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$

- Lagrange function

$$L(w, b, \alpha) = \frac{1}{2} \|w\|^2 + C \sum_i \xi_i - \sum_i \alpha_i [y_i [\langle x_i, w \rangle + b] + \xi_i - 1] - \sum_i \eta_i \xi_i$$

Dual Problem

- Primal optimization problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$

- Lagrange function

$$L(w, b, \alpha) = \frac{1}{2} \|w\|^2 + C \sum_i \xi_i - \sum_i \alpha_i [y_i [\langle x_i, w \rangle + b] + \xi_i - 1] - \sum_i \eta_i \xi_i$$

Optimality in w, b, ξ is at saddle point with α, η

- Derivatives in w, b, ξ need to vanish

Dual Problem

- **Lagrange function**

$$L(w, b, \alpha) = \frac{1}{2} \|w\|^2 + C \sum_i \xi_i - \sum_i \alpha_i [y_i [\langle x_i, w \rangle + b] + \xi_i - 1] - \sum_i \eta_i \xi_i$$

Dual Problem

- **Lagrange function**

$$L(w, b, \alpha) = \frac{1}{2} \|w\|^2 + C \sum_i \xi_i - \sum_i \alpha_i [y_i [\langle x_i, w \rangle + b] + \xi_i - 1] - \sum_i \eta_i \xi_i$$

- **Derivatives in w, b need to vanish**

$$\partial_w L(w, b, \xi, \alpha, \eta) = w - \sum_i \alpha_i y_i x_i = 0$$

$$\partial_b L(w, b, \xi, \alpha, \eta) = \sum_i \alpha_i y_i = 0$$

$$\partial_{\xi_i} L(w, b, \xi, \alpha, \eta) = C - \alpha_i - \eta_i = 0$$

Dual Problem

- Lagrange function

$$L(w, b, \alpha) = \frac{1}{2} \|w\|^2 + C \sum_i \xi_i - \sum_i \alpha_i [y_i [\langle x_i, w \rangle + b] + \xi_i - 1] - \sum_i \eta_i \xi_i$$

- Derivatives in w, b need to vanish

$$\partial_w L(w, b, \xi, \alpha, \eta) = w - \sum_i \alpha_i y_i x_i = 0$$

$$\partial_b L(w, b, \xi, \alpha, \eta) = \sum_i \alpha_i y_i = 0$$

$$\partial_{\xi_i} L(w, b, \xi, \alpha, \eta) = C - \alpha_i - \eta_i = 0$$

- Plugging terms back into L yields

$$\underset{\alpha}{\text{maximize}} - \frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle + \sum_i \alpha_i$$

$$\text{subject to } \sum_i \alpha_i y_i = 0 \text{ and } \alpha_i \in [0, C]$$


bound
influence

Karush Kuhn Tucker Conditions

$$\underset{\alpha}{\text{maximize}} -\frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle + \sum_i \alpha_i$$

$$\text{subject to } \sum_i \alpha_i y_i = 0 \text{ and } \alpha_i \in [0, C]$$

$$w = \sum_i y_i \alpha_i x_i$$




Karush Kuhn Tucker Conditions

$$\underset{\alpha}{\text{maximize}} -\frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle + \sum_i \alpha_i$$

$$\text{subject to } \sum_i \alpha_i y_i = 0 \text{ and } \alpha_i \in [0, C]$$

$$w = \sum_i y_i \alpha_i x_i$$



$$\alpha_i [y_i [\langle w, x_i \rangle + b] + \xi_i - 1] = 0$$

$$\eta_i \xi_i = 0$$



$$\alpha_i = 0 \implies y_i [\langle w, x_i \rangle + b] \geq 1$$

$$0 < \alpha_i < C \implies y_i [\langle w, x_i \rangle + b] = 1$$

$$\alpha_i = C \implies y_i [\langle w, x_i \rangle + b] \leq 1$$

5 minutes break

Nonlinear Separation

The Kernel Trick

- Linear soft margin problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$

The Kernel Trick

- Linear soft margin problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$

- Dual problem

$$\underset{\alpha}{\text{maximize}} \quad -\frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle + \sum_i \alpha_i$$

subject to $\sum \alpha_i y_i = 0$ and $\alpha_i \in [0, C]$

The Kernel Trick

- Linear soft margin problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, x_i \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$

- Dual problem

$$\underset{\alpha}{\text{maximize}} \quad -\frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle + \sum_i \alpha_i$$

subject to $\sum_i \alpha_i y_i = 0$ and $\alpha_i \in [0, C]$

- Support vector expansion

$$f(x) = \sum_i \alpha_i y_i \langle x_i, x \rangle + b$$

The Kernel Trick

- Linear soft margin problem

$$\underset{w,b}{\text{minimize}} \quad \frac{1}{2} \|w\|^2 + C \sum_i \xi_i$$

subject to $y_i [\langle w, \phi(x_i) \rangle + b] \geq 1 - \xi_i$ and $\xi_i \geq 0$


- Dual problem


$$\underset{\alpha}{\text{maximize}} \quad -\frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j k(x_i, x_j) + \sum_i \alpha_i$$


subject to $\sum_i \alpha_i y_i = 0$ and $\alpha_i \in [0, C]$


- Support vector expansion


$$f(x) = \sum_i \alpha_i y_i k(x_i, x) + b$$
























































That's all!