

without skips
with identity skips
no normalization
pre-RMSNorm, $\varepsilon = 10^{-5}$
hypercube
clusters
satimage
segmentation
s_i/s_1
skips add $L\,d_{\text{model}}^2 = 48$
rank 168 and 216, $P = 232$, 5 seeds
skips add $L\,d_{\text{model}}(d_{\text{model}} - 1)/2 = 18$
rank 208 and 226, $P = 232$, 5 seeds
skips add $L\,d_{\text{model}}^2 = 48$
rank 180 and 228, $P = 244$, 5 seeds
skips add $L\,d_{\text{model}}(d_{\text{model}} - 1)/2 = 18$
rank 220 and 238, $P = 244$, 5 seeds
skips add $L\,d_{\text{model}}^2 = 48$
rank 296 and 344, $P = 360$, 5 seeds
skips add $L\,d_{\text{model}}(d_{\text{model}} - 1)/2 = 18$
rank 336 and 354, $P = 360$, 5 seeds
skips add $L\,d_{\text{model}}^2 = 48$
rank 228 and 276, $P = 292$, 5 seeds
skips add $L\,d_{\text{model}}(d_{\text{model}} - 1)/2 = 18$
rank 268 and 286, $P = 292$, 5 seeds
singular value index (largest first)
singular value index (largest first)