


$(1 + 1) - D$ QCA Algorithm To Simulate Quantum TASEP Dynamics Using Matrix Product States




UDDHAV SEN

Contact: senu@coventry.ac.uk

The System

- ▶ The **Totally Asymmetric Simple Exclusion Process (TASEP)** is a fundamental stochastic model used to describe **non-equilibrium transport phenomena**[1].
- ▶ It consists of **particles hopping in one direction** (totally asymmetric) on a **discrete lattice**, subject to the **exclusion principle**—each site can be occupied by at most one particle.
- ▶ We start with nearest neighbour interactions given by the Hamiltonian $\mathcal{H} = -\sum_{\langle \mu, \nu \rangle} J_{\mu\nu} \sigma_{\mu}^{\dagger} \sigma_{\nu}^{-} + h.c.$
- ▶ The jump operators for the Quantum TASEP are given by $L_{pump} = \sigma^{\dagger}, L_{drain} = \sigma^{-}, L_{bulk} = \sigma^{-} \sigma^{\dagger}$.

The Lattice




- ▶ Lattice Diagram for a $N = 4$ system.
- ▶ We create a reverse L-shaped perceptron structure and sweep through the lattice.
- ▶ The bulk perceptron $G_k = D_k U_k$ has U_k acting on $c_k - r_k$ and D_k on $c_k - t_k$ qubits respectively.
- ▶ The left and right gates $G_{pump/drain}$ are added to the left and right boundaries.

Quantum Cellular Automata

- ▶ QCA can be seen as an extension of **Classical Cellular Automata**; where each cells hold a **binary** state.
- ▶ QCA follows some fundamental quantum principles:
 1. **Superposition**
 2. **Entanglement**
 3. **Unitary Evolution**
 4. **Locality**
- ▶ They are Quantum **Turing Complete**[2]!
- ▶ Tensor Network methods like **TEBD** and **DMRG** are widely used to investigate many-body quantum systems[3].
- ▶ QCAs modeled with Tensor Networks have recently found increasing popularity to simulate similar systems out of equilibrium.
- ▶ They can also be easily extended to higher dimensions; on top of the simplicity of their structure.

Tensor Network Construction



- ▶ MPO for the L-shaped lattice.
- ▶ The alternating full lattice MPO from the ladder construction for a $N = 8$ system.

The Perceptron Dynamics

- ▶ The system evolves as

$$\rho(t+1) = \text{tr}_{\bar{p}}(\mathcal{G}(\rho(t) \otimes |0\rangle \langle 0|_{anc}) \mathcal{G}^{\dagger}).$$
- ▶ More explicitly we have,

$$\rho(t+1) = \text{tr}_{\bar{p}}(D_{pump} \prod_k \mathcal{G}_k D_{drain}(\rho(t) \otimes |0\rangle \langle 0|_{anc}) D_{drain}^{\dagger} \prod_k \mathcal{G}_k^{\dagger} D_{pump}^{\dagger}).$$
- ▶ Here we have the following operators;
 1. $U = e^{-i\theta \sum_k h_{k,k+1}} = e^{-i\theta \sum_k h_{k,k+1}}$
 2. $D_{pump} = e^{i\theta \sum_k (\sigma_k^{\dagger} \otimes \sigma_k^{\dagger} + \sigma_k^{-} \otimes \sigma_k^{-})}$
 3. $D_{drain} = e^{i\theta \sum_k (\sigma_k^{\dagger} \otimes \sigma_k^{-} + \sigma_k^{-} \otimes \sigma_k^{\dagger})}$
 4. $D_{bulk} = e^{i\theta \sum_k (\sigma_k^{-} \otimes \sigma_k^{\dagger} \otimes \sigma_{k+1}^{\dagger} + \sigma_k^{\dagger} \otimes \sigma_k^{-} \otimes \sigma_{k+1}^{-})}$
- ▶ $\mathcal{G}_k := D_k U_k$ is defined as the **bulk perceptron** of the QCA.
- ▶ Our choice of operators recovers the **Lindblad Master equation** with the choice of $\theta = \sqrt{\gamma dt}$ in $O(dt)$ limit[4],

$$\partial_t \rho = -i[\mathcal{H}, \rho] + \sum_{\mu} A_{\mu} \rho A_{\mu}^{\dagger} - \frac{1}{2} \{A_{\mu}^{\dagger} A_{\mu}, \rho\}$$
- ▶ We create a QCA perceptron with these equations and evolve the MPO-MPS system till we hopefully reach steady state!!

Conclusions And Future Work

- ▶ The Poster provides a framework to simulate a quantised version of TASEP with **open boundary conditions** with MPS.
- ▶ We are currently simulating large-scale systems on the HPC and looking for possible **phase transitions** for various order parameters like current density etc.
- ▶ This is an essential building block for **Quantum Active Matter** and looking for interesting kinds of quantum/classical correlations.

Plots




Figure: Verification of Q-TASEP with QuTIP.




Figure: Individual Lattice Simulation of $N=6$ with $|100000\rangle$.




Figure: Full Lattice Simulation of $N=6$ with $|100000\rangle$.




Figure: Full Lattice Occupation Number Density Plot For various In/Out Rates.

References

- [1] Frank Spitzer. "Interaction of Markov processes". In: *Advances in Mathematics* 5.2 (1970), pp. 246–290. ISSN: 0001-8708. DOI: [https://doi.org/10.1016/0001-8708\(70\)90034-4](https://doi.org/10.1016/0001-8708(70)90034-4). URL: <https://www.sciencedirect.com/science/article/pii/0001870870900344>.
- [2] Carlos A. Pérez-Delgado and Donny Cheung. "Local unitary quantum cellular automata". In: *Physical Review A* 76.3 (Sept. 2007). ISSN: 1094-1622. DOI: 10.1103/PhysRevA.76.032320. URL: <http://dx.doi.org/10.1103/PhysRevA.76.032320>.
- [3] Guifré Vidal. "Efficient Simulation of One-Dimensional Quantum Many-Body Systems". In: *Phys. Rev. Lett.* 93 (4 July 2004), p. 040502. DOI: 10.1103/PhysRevLett.93.040502. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.93.040502>.
- [4] Edward Gillman, Federico Carollo, and Igor Lesanovsky. "Using $(1+1)D$ quantum cellular automata for exploring collective effects in large-scale quantum neural networks". In: *Phys. Rev. E* 107 (2 Feb. 2023), p. L022102. DOI: 10.1103/PhysRevE.107.L022102. URL: <https://link.aps.org/doi/10.1103/PhysRevE.107.L022102>.