

# BAYESIAN HIERARCHICAL MODELS

Linda Maokomatanda  
Tim Mensinger  
Markus Schick

University of Bonn

1. *Bayesian Thinking*

2. *Hierarchical Models*

3. *Monte Carlo Study*

4. *Application*

---

# *Bayesian Thinking*

---

## *Bayes' Theorem*

$$p(\theta \mid \text{data}) = \frac{p(\text{data} \mid \theta)p(\theta)}{p(\text{data})} \propto p(\text{data} \mid \theta)p(\theta)$$

## *Bayes' Theorem*

$$p(\theta \mid \text{data}) = \frac{p(\text{data} \mid \theta)p(\theta)}{p(\text{data})} \propto p(\text{data} \mid \theta)p(\theta)$$

$$\text{posterior} = \frac{\text{likelihood} \times \text{prior}}{\text{evidence}} \propto \text{likelihood} \times \text{prior}$$

## *Solving for the Posterior Analytically*

*Setting:*  $\{y_i : i = 1, \dots, n\}$  with  $y_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$   
and  $\sigma^2$  known

## *Solving for the Posterior Analytically*

*Setting:*  $\{y_i : i = 1, \dots, n\}$  with  $y_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$   
and  $\sigma^2$  known

*Likelihood:*  $p(y \mid \mu) = \prod_i p(y_i \mid \mu)$

## *Solving for the Posterior Analytically*

*Setting:*  $\{y_i : i = 1, \dots, n\}$  with  $y_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$   
and  $\sigma^2$  known

*Likelihood:*  $p(y \mid \mu) = \prod_i p(y_i \mid \mu)$

*Prior:*  $p(\mu)$



## *Solving for the Posterior Analytically*

*Setting:*  $\{y_i : i = 1, \dots, n\}$  with  $y_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$   
and  $\sigma^2$  known

*Likelihood:*  $p(y \mid \mu) = \prod_i p(y_i \mid \mu)$

*Prior:*  $p(\mu)$

*Posterior:*  $p(\mu \mid y) \propto p(y \mid \mu)p(\mu)$

## *Solving for the Posterior Analytically*

*Setting:*  $\{y_i : i = 1, \dots, n\}$  with  $y_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$   
and  $\sigma^2$  known

*Likelihood:*  $p(y \mid \mu) = \prod_i p(y_i \mid \mu)$

*Prior:*  $p(\mu)$

*Posterior:*  $p(\mu \mid y) \propto p(y \mid \mu)p(\mu)$

*Goal:* Infer distribution of  $\mu \mid y$

## *Solving for the Posterior Analytically*

*Setting:*  $\{y_i : i = 1, \dots, n\}$  with  $y_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$   
and  $\sigma^2$  known

*Likelihood:*  $p(y \mid \mu) = \prod_i p(y_i \mid \mu)$

*Prior:*  $p(\mu)$

*Posterior:*  $p(\mu \mid y) \propto p(y \mid \mu)p(\mu)$

*Goal:* Infer distribution of  $\mu \mid y$   
Why?

## *Conjugate Prior*

Let  $\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$

## *Conjugate Prior*

Let  $\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$

Then

$$p(\mu \mid y) \propto p(y \mid \mu)p(\mu)$$

## Conjugate Prior

Let  $\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$

Then

$$\begin{aligned} p(\mu \mid y) &\propto p(y \mid \mu)p(\mu) \\ &\propto \exp\left(\frac{-n}{2\sigma^2}(\mu - \bar{y})^2\right) \exp\left(\frac{-1}{2\sigma_0^2}(\mu - \mu_0)^2\right) \end{aligned}$$

## Conjugate Prior

Let  $\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$

Then

$$\begin{aligned} p(\mu \mid y) &\propto p(y \mid \mu)p(\mu) \\ &\propto \exp\left(\frac{-n}{2\sigma^2}(\mu - \bar{y})^2\right) \exp\left(\frac{-1}{2\sigma_0^2}(\mu - \mu_0)^2\right) \\ &\propto \exp\left(-\frac{1}{2\sigma_\mu}(\mu - \bar{\mu})^2\right) \end{aligned}$$

## Conjugate Prior

Let  $\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$

Then

$$\begin{aligned} p(\mu \mid y) &\propto p(y \mid \mu)p(\mu) \\ &\propto \exp\left(\frac{-n}{2\sigma^2}(\mu - \bar{y})^2\right) \exp\left(\frac{-1}{2\sigma_0^2}(\mu - \mu_0)^2\right) \\ &\propto \exp\left(-\frac{1}{2\sigma_\mu^2}(\mu - \bar{\mu})^2\right) \\ \implies \mu \mid y &\sim \mathcal{N}(\bar{\mu}, \sigma_\mu^2) \end{aligned}$$



## *Conjugate Prior*

$$\mu \mid y \sim \mathcal{N} \left( \bar{\mu}, \sigma_{\mu}^2 \right), \text{ with}$$

## Conjugate Prior

$$\mu \mid y \sim \mathcal{N}(\bar{\mu}, \sigma_\mu^2), \text{ with}$$

$$\sigma_\mu = \left( \frac{1}{\sigma^2/n} + \frac{1}{\sigma_0^2} \right)^{-1}$$

## Conjugate Prior

$$\mu \mid y \sim \mathcal{N}(\bar{\mu}, \sigma_\mu^2), \text{ with}$$

$$\sigma_\mu = \left( \frac{1}{\sigma^2/n} + \frac{1}{\sigma_0^2} \right)^{-1}$$

$$\bar{\mu} = \sigma_\mu \left( \frac{1}{\sigma^2/n} \bar{y} + \frac{1}{\sigma_0^2} \mu_0 \right)$$

## Conjugate Prior

$$\mu \mid y \sim \mathcal{N}(\bar{\mu}, \sigma_\mu^2), \text{ with}$$

$$\sigma_\mu = \left( \frac{1}{\sigma^2/n} + \frac{1}{\sigma_0^2} \right)^{-1}$$

$$\begin{aligned}\bar{\mu} &= \sigma_\mu \left( \frac{1}{\sigma^2/n} \bar{y} + \frac{1}{\sigma_0^2} \mu_0 \right) \\ &= \alpha \bar{y} + (1 - \alpha) \mu_0\end{aligned}$$

A normal model with known variance and no features, really?



# *Sampling from the Posterior*

*Object of interest:*  $\theta$  | data

# *Sampling from the Posterior*

*Object of interest:*  $\theta \mid \text{data}$

*Quantity of interest:*  $\mathbb{E} [ h(\theta) \mid \text{data} ] =: \mathbb{E}_{\theta}[h]$

## *Sampling from the Posterior*

*Object of interest:*  $\theta \mid \text{data}$

*Quantity of interest:*  $\mathbb{E} [h(\theta) \mid \text{data}] =: \mathbb{E}_\theta[h]$

*Estimation:* Let  $\theta^{(1)}, \dots, \theta^{(n)} \stackrel{\text{iid}}{\sim} p(\theta \mid \text{data})$ ,  
then

$$\frac{1}{\sqrt{n}} \left( \sum_i h(\theta^{(i)}) - \mathbb{E}_\theta[h] \right) \xrightarrow{d} \mathcal{N}(0, \omega)$$



## *Sampling from the Posterior*

But how do we sample from  $p(\theta \mid \text{data})$ ?

## *Sampling from the Posterior*

But how do we sample from  $p(\theta \mid \text{data})$ ?

*Problems:*  $p(\theta \mid \text{data})$  might be

## *Sampling from the Posterior*

But how do we sample from  $p(\theta \mid \text{data})$ ?

*Problems:*  $p(\theta \mid \text{data})$  might be

(i.) of unknown form

## *Sampling from the Posterior*

But how do we sample from  $p(\theta \mid \text{data})$ ?

*Problems:*  $p(\theta \mid \text{data})$  might be

(i.) of unknown form

(ii.) very complex

## *Sampling from the Posterior*

But how do we sample from  $p(\theta \mid \text{data})$ ?

*Problems:*  $p(\theta \mid \text{data})$  might be

(i.) of unknown form

(ii.) very complex

(iii.) only known up to an integration const.

# Markov Chain Monte Carlo

---

## Algorithm Metropolis-Hastings (1953, 1970)

---

**Input:**  $(\pi, q, T) = (\text{target}, \text{proposal}, \text{no. of samples})$

1: initialize  $x_0$  in supp  $q$

2: **for**  $t = 0, \dots, T$  **do**

3:     candidate:  $y \sim q(\cdot \mid x_t)$


4:     acceptance prob.:  $\mathcal{A} \leftarrow \min \left\{ \frac{\pi(y)}{\pi(x_t)} \frac{q(x_t \mid y)}{q(y \mid x_t)}, 1 \right\}$

5:     update:  $x_{t+1} \leftarrow \begin{cases} y & , \text{with prob. } \mathcal{A} \\ x_t & , \text{with remaining prob.} \end{cases}$


6: **return**  $\{x_t : t = 1, \dots, T\}$

---

# Markov Chain Monte Carlo




# Markov Chain Monte Carlo






# Markov Chain Monte Carlo



# Markov Chain Monte Carlo



---

# *Hierarchical Models*

---

# *Structure of HM - Setup*

*Hierarchical Data:*

# *Structure of HM - Setup*

*Hierarchical Data:*

*Individual Level:*  $(y_i, x_i)$  for  $i = 1, \dots, n$

# Structure of HM - Setup

*Hierarchical Data:*

*Individual Level:*  $(y_i, x_i)$  for  $i = 1, \dots, n$

*Group Level:*  $u_j$  for  $j = 1, \dots, J$

# Structure of HM - Setup

## *Hierarchical Data:*

*Individual Level:*  $(y_i, x_i)$  for  $i = 1, \dots, n$

*Group Level:*  $u_j$  for  $j = 1, \dots, J$


## *Example:*

*Test Outcome:*  $y_i$

*Parental Income:*  $x_i$

*Num. of Teachers:*  $u_j$

## Structure of HM





# *The Prior Revisited*

*Before:*

*Model:*  $p(\text{data} \mid \theta)$

*Prior:*  $p(\theta)$

# The Prior Revisited

*Before:*

*Model:*  $p(\text{data} \mid \theta)$

*Prior:*  $p(\theta)$

*Now:*

*Model:*  $p(\text{data} \mid \theta, \phi) = p(\text{data} \mid \theta)$

*Prior:*  $p(\theta \mid \phi)$

*Hyperprior:*  $p(\phi)$

# *The Posterior Revisited*

*Posterior:*

$$p(\theta, \phi \mid \text{data}) \propto p(\text{data} \mid \theta)p(\theta, \phi)$$

# *The Posterior Revisited*

*Posterior:*

$$\begin{aligned} p(\theta, \phi \mid \text{data}) &\propto p(\text{data} \mid \theta)p(\theta, \phi) \\ &\propto p(\text{data} \mid \theta)p(\theta \mid \phi)p(\phi) \end{aligned}$$

# *The Posterior Revisited*

*Posterior:*

$$\begin{aligned} p(\theta, \phi \mid \text{data}) &\propto p(\text{data} \mid \theta)p(\theta, \phi) \\ &\propto p(\text{data} \mid \theta)p(\theta \mid \phi)p(\phi) \end{aligned}$$

$$p(\phi \mid \text{data}) \propto \int p(\theta, \phi \mid \text{data})d\theta$$

## *Varying Slopes, Varying Intercepts*

*Setup:* Individual  $i$  in group  $j$

## *Varying Slopes, Varying Intercepts*

*Setup:* Individual  $i$  in group  $j$

*Individual Level:*  $y_i = \alpha_{j[i]} + \beta_{j[i]}x_i + \epsilon_i$

## *Varying Slopes, Varying Intercepts*

*Setup:* Individual  $i$  in group  $j$

*Individual Level:*  $y_i = \alpha_{j[i]} + \beta_{j[i]}x_i + \epsilon_i$

*Group Level:*  $\begin{bmatrix} \alpha_j \\ \beta_j \end{bmatrix} = \gamma_0 + \gamma u_j + \eta_j$



## Varying Slopes, Varying Intercepts

*Setup:* Individual  $i$  in group  $j$

*Individual Level:*  $y_i = \alpha_{j[i]} + \beta_{j[i]}x_i + \epsilon_i$

*Group Level:*  $\begin{bmatrix} \alpha_j \\ \beta_j \end{bmatrix} = \gamma_0 + \gamma u_j + \eta_j$

*Priors on:*  $\gamma_0, \gamma, \epsilon_i, \eta_j$

## Varying Slopes, Varying Intercepts

*Setup:* Individual  $i$  in group  $j$

*Individual Level:*  $y_i = \alpha_{j[i]} + \beta_{j[i]}x_i + \epsilon_i$

*Group Level:*  $\begin{bmatrix} \alpha_j \\ \beta_j \end{bmatrix} = \gamma_0 + \gamma u_j + \eta_j$

*Priors on:*  $\gamma_0, \gamma, \epsilon_i, \eta_j$

---

# *Monte Carlo Study*

---

# *Stan*

*What is Stan?* C++ package (fast run times)

# *Stan*

*What is Stan?* C++ package (fast run times)

*Use cases:* Bayesian/Maximum Likelihood estimation of statistical models

# Stan

*What is Stan?* C++ package (fast run times)

*Use cases:* Bayesian/Maximum Likelihood estimation of statistical models

*Interfaces:* PyStan, Rstan, Stan.jl,...

## *Bayesian Models with Stan*

$$y_i = \alpha + \beta_{j[i]} x_i + \epsilon_i,$$

$$\epsilon \sim \mathcal{N}(0, 1)$$

## *Bayesian Models with Stan*

$$y_i = \alpha + \beta_{j[i]}x_i + \epsilon_i,$$

$$\epsilon \sim \mathcal{N}(0, 1)$$

$$\beta_j = \gamma_0 + \gamma_1 u_j + \eta_j,$$

$$\eta \sim \mathcal{N}(0, 1)$$



## Bayesian Models with Stan

$$y_i = \alpha + \beta_{j[i]}x_i + \epsilon_i,$$

$$\epsilon \sim \mathcal{N}(0, 1)$$

$$\beta_j = \gamma_0 + \gamma_1 u_j + \eta_j,$$

$$\eta \sim \mathcal{N}(0, 1)$$

```
data {  
  vector[N] y;  
  vector[N] x;  
  vector[N] u;  
  int<lower=0> J;  
  int<lower=0> N;  
  int<lower=1,upper=J>  
    group[N];  
}
```

## Bayesian Models with Stan

$$y_i = \alpha + \beta_{j[i]}x_i + \epsilon_i,$$
$$\epsilon \sim \mathcal{N}(0, 1)$$

$$\beta_j = \gamma_0 + \gamma_1 u_j + \eta_j,$$
$$\eta \sim \mathcal{N}(0, 1)$$

```
parameter {  
  real alpha;  
  real gamma_0;  
  real gamma_1;  
  vector[J] eta_b;  
  real<lower=0> sigma_b;  
  real<lower=0> sigma_y;  
}
```

## Bayesian Models with Stan

$$y_i = \alpha + \beta_{j[i]}x_i + \epsilon_i,$$
$$\epsilon \sim \mathcal{N}(0, 1)$$

$$\beta_j = \gamma_0 + \gamma_1 u_j + \eta_j,$$
$$\eta \sim \mathcal{N}(0, 1)$$

```
# model
for (i in 1:N) {
  beta[i] = gamma_0 +
    u[i] * gamma_1 +
    eta[group[i]]

  y_hat[i] = alpha +
    x[i] * beta[i];
}
y ~ normal(
  y_hat, sigma_y);
```

## Bayesian Models with Stan

$$y_i = \alpha + \beta_{j[i]}x_i + \epsilon_i,$$

$$\epsilon \sim \mathcal{N}(0, 1)$$

$$\beta_j = \gamma_0 + \gamma_1 u_j + \eta_j,$$

$$\eta \sim \mathcal{N}(0, 1)$$

*# priors*

`gamma_0 ~ normal(1, 1);`


`gamma_1 ~ normal(1, 1);`

`eta ~ normal(0, sigma_b);`


`sigma_y ~ cauchy(0, 5);`

`sigma_b ~ cauchy(0, 5);`


*How can we be sure that we sample from the right distribution?*




*How can we be sure that we sample from the right distribution?*



*How can we be sure that we sample from the right distribution?*




# Monitoring Convergence






# Monitoring Convergence




## Monitoring Convergence



*Variance of a single chain:*

$$s_m^2 = \frac{1}{N-1} \sum_{n=1}^N (\theta_m^{(n)} - \bar{\theta}_m)^2$$

## Monitoring Convergence



*Average within chain variance:*

$$W = \frac{1}{M} \sum_{m=1}^M s_m^2$$

## *Brooks and Gelman convergence criterium*

*Average Variance between chains:*

$$B/N = \frac{1}{M-1} \sum_{m=1}^M (\bar{\theta}_m - \bar{\theta})^2$$

## *Brooks and Gelman convergence criterium*

*Average Variance between chains:*

$$B/N = \frac{1}{M-1} \sum_{m=1}^M (\bar{\theta}_m - \bar{\theta})^2$$

*Total Variance:*

$$\widehat{\text{Var}}^+(\theta \mid y) = \frac{N-1}{N}W + \frac{1}{N}B$$

# Monitoring Convergence


*Scale Reducing Factor:*

$$\hat{R} = \sqrt{\frac{\widehat{\text{Var}}^+(\theta | y)}{W}}$$

# Monitoring Convergence

Scale Reducing Factor:


$$\hat{R} = \sqrt{\frac{\widehat{\text{Var}}^+(\theta | y)}{W}}$$



# Monitoring Convergence

Scale Reducing Factor:

$$\hat{R} = \sqrt{\frac{\widehat{\text{Var}}^+(\theta | y)}{W}}$$





## *Prior Design*

*Good Prior:*  $\gamma_0 \sim \mathcal{N}(1, 1), \gamma_1 \sim \mathcal{N}(1, 1)$

## *Prior Design*

*Good Prior:*  $\gamma_0 \sim \mathcal{N}(1, 1), \gamma_1 \sim \mathcal{N}(1, 1)$

*Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 1), \gamma_1 \sim \mathcal{N}(2, 1)$

## Prior Design

*Good Prior:*  $\gamma_0 \sim \mathcal{N}(1, 1), \gamma_1 \sim \mathcal{N}(1, 1)$

*Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 1), \gamma_1 \sim \mathcal{N}(2, 1)$

*Weak, Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 3), \gamma_1 \sim \mathcal{N}(2, 3)$

## Prior Design

*Good Prior:*  $\gamma_0 \sim \mathcal{N}(1, 1), \gamma_1 \sim \mathcal{N}(1, 1)$

*Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 1), \gamma_1 \sim \mathcal{N}(2, 1)$

*Weak, Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 3), \gamma_1 \sim \mathcal{N}(2, 3)$

*Flat Prior:*  $\gamma_0 \sim \mathcal{U}(-\infty, \infty), \gamma_1 \sim \mathcal{U}(-\infty, \infty)$

## Prior Design

*Good Prior:*  $\gamma_0 \sim \mathcal{N}(1, 1), \gamma_1 \sim \mathcal{N}(1, 1)$


*Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 1), \gamma_1 \sim \mathcal{N}(2, 1)$

*Weak, Bad Prior:*  $\gamma_0 \sim \mathcal{N}(2, 3), \gamma_1 \sim \mathcal{N}(2, 3)$

*Flat Prior:*  $\gamma_0 \sim \mathcal{U}(-\infty, \infty), \gamma_1 \sim \mathcal{U}(-\infty, \infty)$


*In all models:*  $\sigma_y, \sigma_b \sim \text{half-Cauchy}(0, 5)$

## *Posterior Distribution - good prior*




**Figure:** Posterior Draws of  $\gamma_1$  with  $N=200$ ,  $J=10$  and 300 simulations

*What happens if we decrease the number of levels  $J$ ?*



**Figure:** Posterior Draws of  $\gamma_1$  with  $N=50$ ,  $J=5$  and 300 simulations


## *Posterior Distribution - bad prior*



**Figure:** Posterior Draws of  $\gamma_1$  with  $N=200$ ,  $J=10$  and 300 simulations




## *Posterior Distribution - weak, bad prior*




**Figure:** Posterior Draws of  $\gamma_1$  with  $N=200$ ,  $J=10$  and 150 simulations

## *Not a good idea: Uniform Prior*



**Figure:** Posterior Draws of  $\gamma_1$  with  $N=200$ ,  $J=10$  and 300 simulations

*Increase sample size dramatically*



**Figure:** Single posterior draw for model with wrong prior and  $N=500$ ,  $J=50$

---

# *Application*

---

## *The Data*

*Description:* General Certificate of Secondary Education (GCSE) exam scores of 1,905 students from 73 schools in England on a science subject

*Variables of interest:* school identifier, student identifier, gender, total score on written paper and total score of course work.

---

*ML and Bayesian Approach  
Application*

---

## Comparison

*The model:* Varying intercept and slope model with a single predictor

$$y_{ij} = \alpha_j + \beta_j x_{ij} + \epsilon_{ij}, \quad (1)$$

$$\alpha_j = \mu_\alpha + u_j, \quad (2)$$

$$\beta_j = \mu_\beta + v_j, \quad (3)$$

$$y_{ij} = \mu_\alpha + \mu_\beta x_{ij} + u_j + v_j x_{ij} + \epsilon_{ij} \quad (4)$$

# Comparison

## Maximum Likelihood (ML) Estimation

*Package: lmer*

*The lmer Package:* combines of ML estimation of model parameters and empirical Bayes (EB) predictions of the varying intercepts and/or slopes resulting in the Best Linear Unbiased Predictions (BLUPs) of the model parameters.

*Why lmer??* allows for comparison between parameter estimates



# Comparison

*Bayesian Estimation:*

*Package:* `stan`

*Priors:* weekly informative normally distributed priors for hyperparameters

## Results

Dependent variable: Course test score		
	ML	Bayes
<i>A. Random effects</i>		
Intercept	– (10.146)	– (10.249)
Female	– (6.924)	– (7.099)
<i>B. Fixed effects</i>		
Intercept	69.425 (1.352)	69.413 (1.287)
Female	7.128 (1.131)	7.132 (1.165)
<hr/>		
N		
Students	1725	1725
Schools	73	73

(i.) point estimates  
almost the same


(ii.) Bayes standard  
deviations for  
random effects may  
be higher because  
ML does not take  
into account group  
level variance

---


*Bayesian Approach: Further Analysis  
Application*

---


## *Posterior distribution ranking:*




## *School specific regression lines and pooling:*




## *Making comparisons between individual schools:*



# Convergence



# Convergence





[https://github.com/timmens/  
bayesian-hierarchical-models](https://github.com/timmens/bayesian-hierarchical-models)

<http://mfviz.com/hierarchical-models/>