


M. Hohle:

Physics 77: Introduction to Computational Techniques in Physics





<u>Week</u>	<u>Date</u>	<u>Topic</u>
1	June 8th	Programming Environment & UIs for Python, Programming Fundamentals
2	June 15th	Basic Types in Python
3	June 22nd	Parsing, Data Processing and File I/O, Visualization
4	June 29th	Functions, Map & Lambda
5	July 6th	Random Numbers & Probability Distributions, Interpreting Measurements
6	July 13th	Numerical Integration and Differentiation
7	July 20th	Root finding, Interpolation
8	July 27th	Systems of Linear Equations, Ordinary Differential Equations (ODEs)
9	Aug 3rd	Stability of ODEs, Examples
10	Aug 10th	Final Project Presentations



finding the intersection of **two lines**:

$$y_1 = a_1x_1 + c_1$$

$$y_2 = a_2x_2 + c_2$$


$$x_1 = x_2$$

$$y_1 = y_2$$

$$a_2x + c_2 = a_1x + c_1$$

$$x = \frac{c_2 - c_1}{a_1 - a_2}$$

$$y = a_1 \frac{c_2 - c_1}{a_1 - a_2} + c_1$$



finding the intersection of **three planes**:

$$y_1 = a_{11}x_{11} + a_{12}x_{12} + c_1$$


$$y_2 = a_{21}x_{21} + a_{22}x_{22} + c_2$$

$$y_3 = a_{31}x_{31} + a_{32}x_{32} + c_3$$

$$x_{11} = x_{21} = x_{31} = x_1$$

$$x_{12} = x_{22} = x_{32} = x_2$$

$$y_1 = y_2 = y_3 = y$$





more general:

$$x_{11} = x_{21} = x_{31} = x_1$$

$$x_{12} = x_{22} = x_{32} = x_2$$

$$y_1 = y_2 = y_3 = y$$

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = c_1$$


$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = c_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 + \dots + a_{3n}x_n = c_3$$

...

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n = c_m$$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}$$






more general:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}$$

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}}_A \vec{x} = \vec{c}$$

$$A\vec{x} = \vec{c}$$





more general:


$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}$$

$\vec{x} \quad \vec{c}$

$$A\vec{x} = \vec{c}$$

general set of solutions

for $n = m \rightarrow$ solution is unique: a point






more general:

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}$$

$\vec{x} \quad \vec{c}$

$$A\vec{x} = \vec{c}$$



general set of solutions

for $n = m \rightarrow$ solution is unique: a point

for $n > m$ (more variables than equations)
 $\rightarrow$ solution is not unique: line, hyperplane


for $n < m$ (more equations than variables)
 $\rightarrow$ no solution



more general:

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix}}_{\vec{x}} = \underbrace{\begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}}_{\vec{c}}$$

$$A\vec{x} = \vec{c}$$



general set of solutions

for $n = m \rightarrow$ solution is unique: a point

for $n > m$ (more variables than equations)
 $\rightarrow$ solution is not unique: line, hyperplane

for $n < m$ (more equations than variables)
 $\rightarrow$ no solution

exceptions!






more general:

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}$$

$\vec{x} \quad \vec{c}$

$$A\vec{x} = \vec{c}$$




general set of solutions

for $n = m \rightarrow$ solution is unique: a point

for $n > m$ (more variables than equations)
 $\rightarrow$ solution is not unique: line, hyperplane

for $n < m$ (more equations than variables)
 $\rightarrow$ no solution

exceptions!





more general:

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{bmatrix}$$

$\vec{x} \quad \vec{c}$

$$A\vec{x} = \vec{c}$$

$$\vec{x} = ?$$

for $n = m$

$$A = [a_{ij}]$$

$$A^{-1}A\vec{x} = A^{-1}\vec{c}$$

$$\vec{x} = A^{-1}\vec{c}$$

inverse:

$$A^{-1}A = I$$

identity:

$$I M = M$$

transpose:

$$[a_{ij}]^T = [a_{ji}]$$

symmetry:

$$[a_{ij}] = [a_{ji}]$$

conjugate transpose:

$$A^+$$

unitary:

$$A^{-1} = A^+$$

idempotency:

$$AA = A \rightarrow A^n = A$$

normal:

$$A^+A = AA^+$$



solving for x:

$$A\vec{x} = \vec{c}$$

→ need to calculate A^{-1}



→ need to calculate a quantity called

determinant of A, $\det(A)$

$$A^{-1} \sim \frac{1}{\det(A)}$$

- if $\det(A) = 0 \rightarrow$ no solution

- $|\det(A)|$: volume spanned by the vectors in A



inverse:

$$A^{-1}A = I$$

identity:

$$I M = M$$

transpose:

$$[a_{ij}]^T = [a_{ji}]$$

symmetry:

$$[a_{ij}] = [a_{ji}]$$

conjugate transpose:

$$A^+$$

unitary:

$$A^{-1} = A^+$$

idempotency:

$$AA = A \rightarrow A^n = A$$

normal:

$$A^+A = AA^+$$



solving for x:


$$A\vec{x} = \vec{c}$$

→ need to calculate A^{-1}

→ need to calculate a quantity called

determinant of A, $\det(A)$

$$A^{-1} \sim \frac{1}{\det(A)}$$



Kmhkmh

$$\det(A) = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{31}a_{22}a_{13} - a_{32}a_{23}a_{11} - a_{33}a_{21}a_{12}$$

inverse:

$$A^{-1}A = I$$

identity:

$$I M = M$$

transpose:

$$[a_{ij}]^T = [a_{ji}]$$

symmetry:

$$[a_{ij}] = [a_{ji}]$$

conjugate transpose:

$$A^+$$

unitary:

$$A^{-1} = A^+$$

idempotency:

$$AA = A \rightarrow A^n = A$$

normal:

$$A^+A = AA^+$$



solving for x:

$$A\vec{x} = \vec{c}$$

→ need to calculate A^{-1}

→ need to calculate a quantity called

determinant of A, $\det(A)$

$$A^{-1} \sim \frac{1}{\det(A)}$$

N x N matrix:

$$\det(A) = \sum_{i_1, i_2, \dots, i_n} \varepsilon_{i_1 \dots i_n} a_{1, i_1} \dots a_{n, i_n}$$

where

$$\varepsilon_{i_1 \dots i_n} = \prod_{1 \leq \mu < \vartheta \leq n} \text{sgn}(i_\vartheta - i_\mu)$$

(Levi-Civita symbol)

changing indices
does not change $|\det(A)|$

inverse:

$$A^{-1}A = I$$

identity:

$$I M = M$$

transpose:

$$[a_{ij}]^T = [a_{ji}]$$

symmetry:

$$[a_{ij}] = [a_{ji}]$$

conjugate transpose:

$$A^+$$

unitary:

$$A^{-1} = A^+$$

idempotency:

$$AA = A \rightarrow A^n = A$$


normal:

$$A^+A = AA^+$$



determinant of A, $\det(A)$

$$A\vec{x} = \vec{c}$$



inverse:

$$A^{-1}A = I$$

identity:

$$I M = M$$

transpose:

$$[a_{ij}]^T = [a_{ji}]$$

symmetry:

$$[a_{ij}] = [a_{ji}]$$

conjugate transpose:

$$A^+$$

unitary:

$$A^{-1} = A^+$$

idempotency:

$$AA = A \rightarrow A^n = A$$

normal:

$$A^+A = AA^+$$

$$\varepsilon_{i_1 \dots i_n} = \prod_{1 \leq \mu < \vartheta \leq n} \text{sgn}(i_\vartheta - i_\mu)$$

$$V = \left| \det \begin{pmatrix} 230 & 0 & 115 \\ 0 & 230 & 115 \\ 0 & 0 & 140 \end{pmatrix} \right| \frac{1}{3} = \frac{230 * 230 * 140 + 0 + 0 - 0 - 0 - 0}{3} = 2,468,666 \text{ m}^3$$



determinant of A, $\det(A)$

$$\varepsilon_{i_1 \dots i_n} = \prod_{1 \leq \mu < \nu \leq n} \text{sgn}(i_\nu - i_\mu)$$




$$V = \left| \det \begin{pmatrix} 230 & 0 & 115 \\ 0 & 230 & 115 \\ 0 & 0 & 140 \end{pmatrix} \right| \frac{1}{3} = \frac{230 * 230 * 140 + 0 + 0 - 0 - 0 - 0}{3} = 2,468,666 \text{ m}^3$$

volume does not
depend on **where**
I put my **coord**
origin...

$$V = \left| \det \begin{pmatrix} 0 & 230 & 115 \\ 230 & 0 & 115 \\ 0 & 0 & 140 \end{pmatrix} \right| \frac{1}{3} = \left| \frac{0 + 0 + 0 - 140 * 230 * 230 - 0 - 0}{3} \right| = 2,468,666 \text{ m}^3$$

...or how I **turn**
the object!

$$V = \left| \det \begin{pmatrix} 115 & 230 & 0 \\ 115 & 0 & 230 \\ 140 & 0 & 0 \end{pmatrix} \right| \frac{1}{3} = \left| \frac{0 + 230 * 230 * 140 + 0 - 0 - 0 - 0}{3} \right| = 2,468,666 \text{ m}^3$$



$$V = \left| \det \begin{pmatrix} 140 & 0 & 0 \\ 115 & 230 & 0 \\ 115 & 0 & 230 \end{pmatrix} \right| \frac{1}{3} = \left| \frac{140 * 230 * 230 + 0 + 0 - 0 - 0 - 0}{3} \right| = 2,468,666 \text{ m}^3$$



linear regression

Goal 1: finding a model that tells us how we can **predict y_k from all the x_i**

	x_1	x_2	x_3	x_4	x_5	y_k
Index	molecular_weight	electronegativity	bond_lengths	num_hydrogen_bonds	logP	toxicity_score
0	341.704	2.65585	3.09407	2	9.11147	80.9281
1	335.951	3.22262	2.89039	7	8.92848	83.4911
2	235.203	2.44115	2.48203	1	6.49731	61.8406
3	246.505	2.76656	2.71547	7	7.45089	57.0538
4	437.939	3.4801	3.59569	3	10.9156	131.326
	336.453	2.81474	3.11	9	8.55696	?

kth row

Goal 2: once we have a model: **predict y_k from new data set**



linear regression

idea: data point y_k in N dimensional space

$$\rightarrow y_k = f(x_1, \dots, x_n, \dots, x_N) + \epsilon \quad \text{for each data point } k$$



ansatz:

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$

linear combination

- y : response
- x : regressors (assumed to be independent)
- β : factors (how a regressor contributes to the response)
- β_0 : intercept
- ϵ : error (stochasticity of the data, assumed to be normally dist.)

Finding β_n is the model!







general: linear refers to the **factors**


$$y_k = \beta_0 + \beta_1 x_1 + \beta_2 x_2 \quad 2D \text{ plane in } 3D \text{ space}$$

$$y_k = \beta_0 + \beta_1 x_1^2 + \beta_2 x_2^2 \quad 2D \text{ parabolic}$$

$$y_k = \beta_0 + \beta_1 x_1^2 - \beta_2 x_2^2 \quad 2D \text{ hyperbolic}$$



...and many more...



y :	response
x :	regressors
β :	factors
β_0 :	intercept
ϵ :	error


all linear

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$



for K data points in N dimensional space

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$



y :	response
x :	regressors
β :	factors
β_0 :	intercept
ϵ :	error

$$\underbrace{\begin{pmatrix} y_1 \\ \vdots \\ y_k \\ \vdots \\ y_K \end{pmatrix}}_Y = \underbrace{\begin{pmatrix} 1 & x_{11} & x_{12} & \dots & x_{1n} & \dots & x_{1N} \\ \vdots & \vdots & \vdots & & \vdots & & \vdots \\ 1 & x_{k1} & & & x_{kn} & & \\ \vdots & \vdots & & & \vdots & & \vdots \\ 1 & \vdots & & & \vdots & & \vdots \\ 1 & x_{K1} & x_{K2} & \dots & x_{Kn} & \dots & x_{KN} \end{pmatrix}}_X \underbrace{\begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_n \\ \vdots \\ \beta_N \end{pmatrix}}_{\beta} + \underbrace{\begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_k \\ \vdots \\ \epsilon_K \end{pmatrix}}_{\epsilon}$$

$$Y = X\beta + \epsilon$$

fitting: finding the best β in terms of minimizing the errors

$$(Y - X\beta)^T (Y - X\beta) = \sum_k \epsilon_k^2$$

the model

$$\frac{\partial}{\partial \beta} \sum_k \epsilon_k^2 = 0 \longrightarrow \beta_{best} = \hat{\beta} = (X^T X)^{-1} X^T Y \longrightarrow \hat{Y} = X\hat{\beta} = X(X^T X)^{-1} X^T Y$$


X and Y are all
observables



for K data points in N dimensional space

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$

$$Y = X\beta + \epsilon$$



y : response
 x : regressors
 β : factors
 β_0 : intercept
 ϵ : error

check out `Walk_Through_LinRegression.ipynb`

predicting **toxicity of molecules** based on their features

	Index	molecular_weight	electronegativity	bond_lengths	num_hydrogen_bonds	logP	toxicity_score
	0	341.704	2.65585	3.09407	2	9.11147	80.9281
	1	335.951	3.22262	2.89039	7	8.92848	83.4911
	2	235.203	2.44115	2.48203	1	6.49731	61.8406
	3	246.505	2.76656	2.71547	7	7.45089	57.0538
	4	437.939	3.4801	3.59569	3	10.9156	131.326



see `Walk_Through_LinRegression.ipynb`

```
import numpy as np
```

```
import pandas as pd
```

```
import matplotlib.pyplot as plt
```

```
import seaborn as sns
```

```
import pylab
```

```
import scipy.stats as stats
```

```
import statsmodels.api as sm
```

```
from statsmodels.formula.api import ols
```

```
from sklearn.preprocessing import MinMaxScaler
```

reading .xlsx
.csv
.txt
...

standard plots

fancy plots:
here a pair-
plot

Q-Q plot

the actual
super tool for
superb data
analysis

scaling and normalizing



```
Train = pd.read_csv("molecular_train_gbc.csv")
```

```
Test = pd.read_csv("molecular_test_gbc.csv")
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

	x_1	x_2	x_3	x_4	x_5	y_k
Index	molecular_weight	electronegativity	bond_lengths	num_hydrogen_bonds	logP	toxicity_score
0	341.704	2.65585	3.09407	2	9.11147	80.9281
1	335.951	3.22262	2.89039	7	8.92848	83.4911
2	235.203	2.44115	2.48203	1	6.49731	61.8406
3	246.505	2.76656	2.71547	7	7.45089	57.0538
4	437.939	3.4801	3.59569	3	10.9156	131.326

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$


y: toxicity_score
 x_n : molecular_weight, electronegativity,
bond_lengths, num_hydrogen_bonds, logP



```
Train = pd.read_csv("molecular_train_gbc.csv")  
Test  = pd.read_csv("molecular_test_gbc.csv")
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

```
out = sns.pairplot(Train, kind = "kde", \  
                  plot_kws = {'color': [176/255, 224/255, 230/255]}, \  
                  diag_kws = {'color': 'black'})  
out.map_offdiag(plt.scatter, color = 'black')
```





```
Train = pd.read_csv("molecular_train_gbc.csv")  
Test  = pd.read_csv("molecular_test_gbc.csv")
```

```
out = sns.pairplot(Train, kind = "kde", \  
                  plot_kws = {'color':[176/255, 224/255, 230/255]}, \  
                  diag_kws = {'color': 'black'})  
out.map_offdiag(plt.scatter, color = 'black')
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

Scaling the data because unit system is arbitrary!

Large numerical values dominate the optimization! → rescaling!

molecular_weight	electronegativity	bond_lengths
341.704	2.65585	3.09407

```
scaler = MinMaxScaler(feature_range = (0, 1))  
TrainS = scaler.fit_transform(Train)  
TestS  = scaler.transform(Test)
```

the scaler returns an np.array
→ convert back to data frame

```
TrainS = pd.DataFrame(TrainS, columns = Train.columns)  
TestS  = pd.DataFrame(TestS, columns = Train.columns)
```



```
TrainS = pd.DataFrame(TrainS, columns = Train.columns)
TestS  = pd.DataFrame(TestS,  columns = Train.columns)
```

```
equation = 'toxicity_score ~ ' + '+'.join(Train.columns[:-1])
print(equation)
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$

```
toxicity_score ~      molecular_weight + electronegativity +
                      bond_lengths + num_hydrogen_bonds + logP
```

```
my_model = ols(equation, data = TrainS).fit()
my_model.summary()
```

OLS (ordinary least squares)



more accurate: determining **the p-values for the factors using ANOVA** for the corresponding residuals

```
table = sm.stats.anova_lm(my_model, typ = 1)
print(table)
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$

	df	sum_sq	mean_sq	F	PR(>F)	vs from t-test
molecular_weight	1.0	13.346285	13.346285	2847.525516	8.024085e-265	0.0000
electronegativity	1.0	0.640388	0.640388	136.631363	3.085962e-29	0.0000
bond_lengths	1.0	0.000684	0.000684	0.145954	7.025342e-01	0.6766
num_hydrogen_bonds	1.0	0.000703	0.000703	0.150055	6.985866e-01	0.6473
logP	1.0	0.013917	0.013917	2.969353	8.524510e-02	0.0852
Residual	794.0	3.721459	0.004687	NaN	NaN	




```
residuals = my_model.resid
```

```
plt.hist(residuals, color = 'w', edgecolor = 'black')  
plt.title('fit residuals')  
plt.ylabel('#')  
plt.xlabel('value')  
plt.show()
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$



residuals approx.
normally distributed
around $\mu = 0$



```
residuals = my_model.resid
```


```
plt.hist(residuals, color = 'w', edgecolor = 'black')  
plt.title('fit residuals')  
plt.ylabel('#')  
plt.xlabel('value')  
plt.show()
```

```
stats.probplot(residuals, dist = "norm", plot = pylab)  
pylab.show()
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

residuals approx.
normally distributed
around $\mu = 0$

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$





```
Ypred = my_model.predict(TestS)
```

```
higher = np.max([Ypred, TestS.toxicity_score])
```

```
lower = np.min([Ypred, TestS.toxicity_score])
```

```
plt.plot([lower, higher], [lower, higher], c = [0, 0, 0, 0.2],\  
         linewidth = 4)
```

```
plt.scatter(TestS.toxicity_score, Ypred, marker = '.', c = 'k')
```


```
plt.ylabel('prediction')
```

```
plt.xlabel('toxicity score')
```

```
plt.show()
```

- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$





```
Ypred = my_model.predict(TestS)
```

```
higher = np.max([Ypred, TestS.toxicity_score])
```

```
lower = np.min([Ypred, TestS.toxicity_score])
```


```
plt.plot([lower, higher], [lower, higher], c = [0, 0, 0, 0.2], linewidth = 4)
```

```
plt.scatter(TestS.toxicity_score, Ypred, marker = '.', c = 'k')
```

```
plt.ylabel('prediction')
```

```
plt.xlabel('toxicity score')
```

```
plt.show()
```



- 1) loading data
- 2) plotting data
- 3) scaling data
- 4) fitting model
- 5) evaluating model

$$y_k = \beta_0 + \sum_{n=1}^N \beta_n x_n + \epsilon$$

```
mean_dev = np.sum( abs(TestS.toxicity_score - Ypred) )/len(Ypred)  
print(mean_dev)
```

5%

M. Hohle:

Thank you for your attention!

