

Spring 2020 MAT303 Recitations

Week of 4/20/20: Sections 4.1 and 4.2

Section 4.1: First-order systems and applications

A system of differential equation consists of a finite collection of differential equations in indeterminates $x_1(t), x_2(t), \dots, x_n(t)$, depending on a parameter t . In this chapter, we will deal with linear systems.

Section 4.1: First-order systems and applications

A system of differential equation consists of a finite collection of differential equations in indeterminates $x_1(t), x_2(t), \dots, x_n(t)$, depending on a parameter t . In this chapter, we will deal with linear systems.

One's first encounter with systems of differential equations arises from higher-order scalar equations. One way of solving them is by reducing such equations to systems of first-order problems, as we shall see next. .

Section 4.1: First-order systems and applications

This example is extracted from problem 4.1.3 in our textbook.
Consider the third-order equation

$$tx^{(3)} - 2t^2x'' + 3tx' + 5x = \ln(t)$$

By using the substitutions $x_1 = x$, $x_2 = x'$, $x_3 = x''$, we can rewrite this equation as a system in three variables

$$tx_3' - 2t^2x_3 + 3tx_2 + 5x_1 = \ln(t)$$

$$x_2' = x_3$$

$$x_1' = x_2$$

Section 4.1: First-order systems and applications

The following example is extracted from problem 4.1.17.

Section 4.1: First-order systems and applications

The following example is extracted from problem 4.1.17. Consider the system of differential equations

$$\begin{aligned}x' &= y \\ y' &= -x\end{aligned}$$

Section 4.1: First-order systems and applications

The following example is extracted from problem 4.1.17. Consider the system of differential equations

$$\begin{aligned}x' &= y \\ y' &= -x\end{aligned}$$

By elimination, we find

$$x'' = y' = -x.$$

Section 4.1: First-order systems and applications

The following example is extracted from problem 4.1.17. Consider the system of differential equations

$$\begin{aligned}x' &= y \\ y' &= -x\end{aligned}$$

By elimination, we find

$$x'' = y' = -x.$$

This second-order equation on x can be solved by the method of characteristic equations,

$$x(t) = A \cos(t) + B \sin(t).$$

Section 4.1: First-order systems and applications

The following example is extracted from problem 4.1.17. Consider the system of differential equations

$$\begin{aligned}x' &= y \\ y' &= -x\end{aligned}$$

By elimination, we find

$$x'' = y' = -x.$$

This second-order equation on x can be solved by the method of characteristic equations,

$$x(t) = A \cos(t) + B \sin(t).$$

Section 4.1: First-order systems and applications

To solve for y , we use the first equation in our system,


$$\begin{aligned}y(t) &= x'(t) \\ &= -A \sin(t) + B \cos(t).\end{aligned}$$

Section 4.1: First-order systems and applications

To solve for y , we use the first equation in our system,

$$\begin{aligned}y(t) &= x'(t) \\ &= -A \sin(t) + B \cos(t).\end{aligned}$$

Below is a plot of the slope field for this equation, with several solution curves outlined.



Section 4.1: First-order systems and applications

In this example, extracted from problem 4.1.23, we have the system

$$x' = y$$

$$y' = 6x - y,$$

Section 4.1: First-order systems and applications

In this example, extracted from problem 4.1.23, we have the system

$$x' = y$$

$$y' = 6x - y,$$

with initial conditions $x(0) = 1, y(0) = 2$.

Section 4.1: First-order systems and applications

In this example, extracted from problem 4.1.23, we have the system

$$x' = y$$

$$y' = 6x - y,$$

with initial conditions $x(0) = 1$, $y(0) = 2$. We can turn this system into a second-order equation for x as follows,

$$x'' = y' = 6x - y = 6x - x',$$

Section 4.1: First-order systems and applications

In this example, extracted from problem 4.1.23, we have the system

$$\begin{aligned}x' &= y \\ y' &= 6x - y,\end{aligned}$$

with initial conditions $x(0) = 1$, $y(0) = 2$. We can turn this system into a second-order equation for x as follows,

$$x'' = y' = 6x - y = 6x - x',$$

that is

$$x'' + x' - 6x = 0.$$

Section 4.1: First-order systems and applications

In this example, extracted from problem 4.1.23, we have the system

$$\begin{aligned}x' &= y \\ y' &= 6x - y,\end{aligned}$$

with initial conditions $x(0) = 1$, $y(0) = 2$. We can turn this system into a second-order equation for x as follows,

$$x'' = y' = 6x - y = 6x - x',$$

that is

$$x'' + x' - 6x = 0.$$

The characteristic polynomial of this equation is

$r^2 + r - 6 = (r - 2)(r + 3)$, with roots $r = 2$, $r = -3$. The general solution takes the form

$$x(t) = Ae^{2t} + Be^{-3t}$$

Section 4.1: First-order systems and applications

To find the appropriate values for the constants, we need to set up the initial conditions in terms of x .

Section 4.1: First-order systems and applications

To find the appropriate values for the constants, we need to set up the initial conditions in terms of x . The initial condition on y yields

$$x'(0) = y(0) = 2,$$

Section 4.1: First-order systems and applications

To find the appropriate values for the constants, we need to set up the initial conditions in terms of x . The initial condition on y yields

$$x'(0) = y(0) = 2,$$

thus we find

$$A + B = 1$$

$$2A - 3B = 2,$$

Section 4.1: First-order systems and applications

To find the appropriate values for the constants, we need to set up the initial conditions in terms of x . The initial condition on y yields

$$x'(0) = y(0) = 2,$$

thus we find

$$A + B = 1$$

$$2A - 3B = 2,$$

whose solutions are $A = 1, B = 0$.

Section 4.1: First-order systems and applications

To find the appropriate values for the constants, we need to set up the initial conditions in terms of x . The initial condition on y yields

$$x'(0) = y(0) = 2,$$

thus we find

$$A + B = 1$$


$$2A - 3B = 2,$$

whose solutions are $A = 1, B = 0$. As a result, $x(t) = e^{2t}$, and

$$y(t) = x'(t) = 2e^{2t}$$

Section 4.1: First-order systems and applications

Below is a plot of the direction field, as well as the solution curve corresponding to the given initial condition (in red).



Section 4.2: The method of elimination

Consider the following system, extracted from problem 4.2.5,

$$x' = -3x - 4y$$

$$y' = 2x + y.$$

Section 4.2: The method of elimination

Consider the following system, extracted from problem 4.2.5,

$$x' = -3x - 4y$$

$$y' = 2x + y.$$

We will rewrite the system by using the short-hand notation D to mean derivatives relative to time, $D = \frac{d}{dt}$,

$$(D + 3)x + 4y = 0$$

$$-2x + (D - 1)y = 0$$

Section 4.2: The method of elimination

Consider the following system, extracted from problem 4.2.5,

$$x' = -3x - 4y$$

$$y' = 2x + y.$$

We will rewrite the system by using the short-hand notation D to mean derivatives relative to time, $D = \frac{d}{dt}$,

$$(D + 3)x + 4y = 0$$

$$-2x + (D - 1)y = 0$$

In order to eliminate one of the variables from the system, we will perform the following operations:

1. Apply the operator $(D - 1)$ to the first equation, that is, differentiate it and subtract the original equation from the result.

Section 4.2: The method of elimination

Consider the following system, extracted from problem 4.2.5,

$$x' = -3x - 4y$$

$$y' = 2x + y.$$

We will rewrite the system by using the short-hand notation D to mean derivatives relative to time, $D = \frac{d}{dt}$,

$$(D + 3)x + 4y = 0$$

$$-2x + (D - 1)y = 0$$

In order to eliminate one of the variables from the system, we will perform the following operations:

1. Apply the operator $(D - 1)$ to the first equation, that is, differentiate it and subtract the original equation from the result.
2. Multiply the second equation by (-4) .

Section 4.2: The method of elimination

Consider the following system, extracted from problem 4.2.5,

$$x' = -3x - 4y$$

$$y' = 2x + y.$$

We will rewrite the system by using the short-hand notation D to mean derivatives relative to time, $D = \frac{d}{dt}$,

$$(D + 3)x + 4y = 0$$

$$-2x + (D - 1)y = 0$$

In order to eliminate one of the variables from the system, we will perform the following operations:

1. Apply the operator $(D - 1)$ to the first equation, that is, differentiate it and subtract the original equation from the result.
2. Multiply the second equation by (-4) .

Section 4.2: The method of elimination

The resulting (equivalent) system is

$$(D - 1)(D + 3)x + 4(D - 1)y = 0$$

$$8x - 4(D - 1)y = 0$$

Section 4.2: The method of elimination

The resulting (equivalent) system is

$$\begin{aligned}(D - 1)(D + 3)x + 4(D - 1)y &= 0 \\ 8x - 4(D - 1)y &= 0\end{aligned}$$

By adding the two equations we find

$$(D - 1)(D + 3)x + 8x = 0,$$

which in usual notation for derivatives is

$$\begin{aligned}(D - 1)(D + 3)x + 8x &= 0 \\ (D - 1)(x' + 3x) + 8x &= 0 \\ (x' + 3x)' - (x' + 3x) + 8x &= 0 \\ x'' + 3x' - x' - 3x + 8x &= 0 \\ x'' + 2x' + 5x &= 0\end{aligned}$$

Section 4.2: The method of elimination

This equation can be solved bby the method of characteristic equations, and I will leave the details to you. Its solutions take the form

$$x(t) = e^{-t}(A \cos(2t) + B \sin(2t)).$$

Section 4.2: The method of elimination

This equation can be solved by the method of characteristic equations, and I will leave the details to you. Its solutions take the form

$$x(t) = e^{-t}(A \cos(2t) + B \sin(2t)).$$

To find the solution $y(t)$ we may use the first equation of the system,

$$\begin{aligned} y(t) &= -\frac{x' + 3x}{4} \\ &= -\frac{e^{-t}[(2B - A) \cos(2t) - (2A + B) \sin(2t)] + 3e^{-t}(A \cos(2t) + B \sin(2t))}{4} \\ &= -\frac{e^{-t}[(A + B) \cos(2t) + (B - A) \sin(2t)]}{2} \end{aligned}$$